

Local Genetic Structure on Breeding Grounds of a Long-Distance Migrant Passerine: The Bluethroat (*Luscinia svecica*) in Spain

FERNANDO ALDA, JAVIER GARCÍA, JESÚS T. GARCÍA, AND SUSANA SUÁREZ-SEOANE

From the Instituto de Investigación en Recursos Cinegéticos (IREC, CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13005 Ciudad Real, Spain; and Área de Ecología, Facultad de Ciencias Biológicas y Ambientales, Universidad de León, Campus de Vegazana s/n, 24071 León, Spain.

Address correspondence to Fernando Alda, Smithsonian Tropical Research Institute, Apartado 0843-03092 Balboa, Ancón, Republic of Panama, or e-mail: alda.fernando@gmail.com

Abstract

Breeding site fidelity can be determined by environmental features, which depending on their heterogeneous distribution may shape the genetic landscape of a population. We used 10 microsatellite loci to study the genetic variation of 83 bluethroats (*Luscinia svecica azuricollis*) across 14 localities within the Spanish breeding population and assess the relative influence of different habitat characteristics (physiography and vegetation) on genetic differentiation. Based on the genetic variation of this population, we identified 3 geographically consistent genetic clusters that on average showed a higher genetic differentiation than among other north European populations, even those belonging to different subspecies. The inferred genetic clusters occurred in geographic areas that significantly differed in elevation. The highest genetic differentiation was observed between sites at different mountain ranges, as well as between the highest altitude sites in the northeastern locale, whereas vegetation type did not explain a significant percentage of genetic variation. The lack of correlation between geographic and genetic distances suggests that this pattern of genetic structure cannot be explained as a consequence of isolation by distance. Finally, we discuss the importance of preserving areas encompassing high environmental and genetic variation as a means of preserving evolutionary processes and adaptive potential.

Key Words: *breeding site selection, environmental factors, genetic structure, Luscinia svecica, microsatellites, Spain*

The interplay between gene flow and local habitat selection and its influence on species diversification constitutes a long-lasting research topic in evolutionary biology (Wright 1940; Felsenstein 1976; Hedrick 1986; Hedrick 2006). The occurrence of a species at a particular site largely depends on environmental variability, which is ultimately determined by the range of suitable habitats according to their spatial configuration and seasonal variation (Bell et al. 1993; Dufour et al. 2006). The spatial variation of ecological factors, linked both to habitat heterogeneity and quality, may also shape levels of genetic variability in wild populations (Frankham 1995; Foll and Gaggiotti 2006; Pitra et al. 2011). As a consequence, genetic differentiation among populations depends not only on the strength of habitat selection on each local population but also on the relative importance of dispersal. Therefore, it is expected that if habitat preferences are stronger than dispersal among local populations, local adaptation may arise in such populations

even if this geographic scale is much smaller than the scale of dispersal (Wright 1940; Blondel et al. 2006). Strong habitat selection in heterogeneous landscapes may cause local populations to evolve traits that provide advantages under their local habitat characteristics (Kawecki and Ebert 2004). However, several factors may hamper local adaptation. In this context, gene flow is the most important factor, since the exchange of genes between populations homogenizes allele frequencies and thus prevents genetic differentiation (Balloux and Lugon-Moulin 2002). Therefore, it is generally assumed that at small spatial scales, intraspecific variation does not occur in highly vagile organisms such as birds. This assumption would be valid if gene flow was spatially random, but evidence suggests that birds may show dispersal biases with respect to habitat (Davis and Stamps 2004; Blondel et al. 2006; Hull et al. 2008; Alda et al. 2011).

Birds breeding in heterogeneous landscapes may choose territories with different environmental qualities, which can

affect demographic parameters and genetic diversity of populations (Penteriani et al. 2004; Porlier et al. 2009). For example, birds with migratory behavior might differ in their degrees of fidelity to their breeding and wintering sites (i.e., migratory connectivity; Esler 2000). This philopatric behavior has been associated with key features of the environment that are patchily distributed or difficult to locate, such as specialized breeding locations or food resources (Van Bekkum et al. 2006; Clark et al. 2008; Hull et al. 2008). Hence, migratory connectivity is directly related to gene flow, which in turn determines the geographical pattern of genetic variation within a species. Consequently, it would be expected that high levels of genetic and morphological variation among populations with strong migratory connectivity are due to low gene flow and local adaptations (Webster et al. 2002).

The bluethroat *Luscinia svecica* (Linnaeus 1758) is a long-distance migratory passerine that breeds throughout Europe, Asia, and Alaska. There are 10 subspecies that constitute a subspecies complex described on the basis of body size and plumage coloration of males and on differences of their breeding habitats, migration routes, and wintering areas (Cramp 1988). However, these subspecies are not recognized according to mitochondrial DNA differentiation and only a shallow divergence exists between the northern and southern subspecies, suggesting a recent divergence of these populations (Questiau et al. 1998; Zink et al. 2003). In addition, faster evolving microsatellite markers indicate restricted gene flow among some subspecies in *L. svecica*, particularly among southern populations, which generally are more differentiated than northern populations. Furthermore, the southern group of subspecies, which includes the Spanish and French subspecies, is morphologically distinct in showing white or no throat spots, in contrast with the northern group of chestnut-spotted populations. Thus, because the Spanish subspecies *L. s. azuricollis* is clearly genetically differentiated, it and the French *L. s. namnetum* populations are proposed to be ancestral to the other European subspecies (Johnsen et al. 2006). In general, bluethroats show high fidelity to their migratory routes between wintering and breeding areas (Markovets and Yosef 2005; Hellgren et al. 2008), so the observed genetic heterogeneity among regions in Europe could be either due to isolation processes or a consequence of local adaptations of southern populations (Johnsen et al. 2006).

Spanish bluethroats are believed to winter south of the Sahara (Arizaga et al. 2006) and breed in the northwestern mountains of Iberian Peninsula (Tellería 1999; Gómez-Manzanares 2003). In the Iberian mountains, *L. s. azuricollis* occurs in a variety of habitat types greatly differing in vegetation structure and composition, altitude, and orientation. These differences can be observed at a very small spatial scale (only a few kilometers apart), providing a framework for habitat choice and some degree of local genetic divergence (Guschanski et al. 2008). However, there is limited knowledge of the genetic variation among bluethroat populations at such small geographic scales, with the exception of *L. s. svecica* in Scandinavia (Hellgren et al. 2008). Thus, the bluethroat breeding population in Spain constitutes a good model to evaluate the relationships between this site fidelity

and the environmental features shaping the genetic structure at a local scale in a wide-ranging species.

The main aim of this study is to examine the genetic variation of bluethroats within the Spanish breeding population, in order to determine: 1) the extent of genetic differentiation at the local scale and 2) whether landscape features have a direct influence on the genetic structure of local populations. Different habitat characteristics (physiography and vegetation) might imply different adaptations or selection patterns for breeding individuals. Thus, we would expect to observe significant genetic differentiation among breeding sites if bluethroats are preferentially selecting certain habitat conditions. If this selection is strong, it might imply a low capability of adaptation to different environments. On the other hand, a lack of genetic differentiation could be a consequence of extensive gene flow and therefore suggest a lack of habitat selection.

Materials and Methods

Study Sites and Sampling

Breeding bluethroats were sampled across the species distribution range in northwestern Spain, from the southern slope of the Cantabrian Mountains to the Mountains of León (León province), ranging from 800 to 1900 m above sea level (Figure 1A). This area spans the putative limit of 2 major European biogeographic regions, the Atlantic and the Mediterranean, and features a wide diversity of habitats. Fourteen localities were sampled during the breeding season between April 2009 and August 2010 and classified on the basis of the main environmental characteristics that could directly or indirectly influence the selection of breeding sites by bluethroats (Table 1).

Localities were assigned to the mountain range where they were sampled (Cantabrian Mountains and Mountains of León). The Cantabrian Mountains run on an east–west axis and are on average higher in altitude than the Mountains of León. They are also more influenced by the Atlantic climate and have higher precipitation than the Mountains of León. Most sampling localities were found along valley bottoms and foothills (800–1200 m) and mountain ridges (1500–1900 m) (Figure 1B) and were further differentiated into low- and high-altitude sites, respectively. Three main habitats were defined according to their vegetation type: brooms, mainly composed by *Cytisus* spp. and *Genista* spp.; heathlands, constituted by *Erica* spp. and *Calluna vulgaris*; and holm oak shrublands, consisting of *Quercus rotundifolia* and *Cistus* spp. (Table 1, Figure 1A, B).

Bluethroats were captured with tape-lured mistnets and clap-traps baited with mealworms. Blood samples from all individuals were obtained by venipuncture of the brachial vein and stored in absolute ethanol until they were analyzed. All animals were released unharmed.

DNA Extraction and Microsatellite Genotyping

Total genomic DNA was extracted from blood using a standard ammonium acetate precipitation protocol (Perbal




Figure 1. (A) Map illustrating the 14 bluethroat localities sampled in northwestern Spain. Gray layers, from light to dark, correspond to elevations 400–800 m, 800–1200 m, 1200–1600 m, and 1600–2600 m. Black lines represent province limits and blue lines are main rivers in the area. Numbers refer to localities in Table 1. (B) Schematic representation of the relief profile of the study region. Mountain range, altitude classes, and vegetation type for each locality is indicated. Colors represent genetic clusters to which localities were assigned; black (red): cluster K-NE, northeastern localities; medium gray (blue): cluster K-NW, northwestern and central areas; and light gray (green): K-S, southern sites. Colors between parentheses refer to the color version of the figure.

1988) following Proteinase K digestion. All samples were genotyped for 12 microsatellite loci: Aar8, Ase19, C μ 4, C μ 10, Fhu2, Hru7, Mcy4, PAT MP 2-43, Pdo5, Phtr2, PmaC25, and Ppi2 (Ellegren 1992; Primmer et al. 1996; Double et al. 1997; Fridolfsson et al. 1997; Otter et al. 1998; Gibbs et al. 1999; Martínez et al. 1999; MacColl et al. 2000; Richardson et al. 2000; Saladin et al. 2003). The microsatellites were co-amplified in 4 multiplex polymerase chain reactions (PCRs; Mix1: Fhu2, PmaC25, Ptc2; Mix2:

Ase19, C μ 4, PAT MP 2-43; Mix3: C μ 10, Hru7, Mcy4; Mix4: Aar8, Pdo5, Phtr2), following the QIAGEN Multiplex PCR kit protocol for 30 cycles and 3 different annealing temperatures (60 °C for Mix1, 57 °C for Mix2 and 48 °C for Mix3 and 4). Reactions were prepared in a final volume of 7 μ L including: 3.5 μ L of Qiagen 2X PCR Master Mix, 0.7 μ L of 10X primer mix (2 μ M each), 1 μ L DNA (ca. 25 ng/ μ L) and 1.8 μ L of RNase-free H₂O. Fluorescently labeled PCR products were analyzed on an ABI3130x/ DNA Analyzer

Table 1 Sampling localities of bluethroat (*Luscinia s. azuricollis*)

	Locality	<i>n</i>	Mountain range	Altitude class	Vegetation	Altitude (m)	Latitude	Longitude
1	Genicera	14	Cantabrian Mountains	High	Brooms	1777.9	42.95°	-5.49°
2	Rodillazo	2	Cantabrian Mountains	High	Brooms	1640.5	42.92°	-5.51°
3	Meroy	2	Cantabrian Mountains	High	Brooms	1592.0	42.97°	-6.22°
4	La Cueta	5	Cantabrian Mountains	High	Brooms	1566.0	43.01°	-6.18°
5	La Majúa	2	Cantabrian Mountains	High	Brooms	1895.0	42.98°	-6.02°
6	Ferreras de Cepeda	17	Mountains of León	Low	Heathlands	973.1	42.65°	-6.03°
7	La Seca	1	Cantabrian Mountains	Low	Heathlands	1122.0	42.74°	-5.60°
8	Corcos	9	Cantabrian Mountains	Low	Heathlands	1012.7	42.67°	-5.08°
9	Pobladura de la Sierra	2	Mountains of León	High	Brooms	1676.5	42.42°	-6.44°
10	Molinaferrera	1	Mountains of León	Low	Heathlands	1138.0	42.39°	-6.36°
11	Palacios de la Valduerna	13	Mountains of León	Low	Holm oak shrublands	809.4	42.33°	-5.94°
12	Villar de Golfer	3	Mountains of León	Low	Heathlands	974.3	42.35°	-6.19°
13	Bustos	8	Mountains of León	Low	Holm oak shrublands	834.0	42.38°	-6.02°
14	Toralino de la Vega	4	Mountains of León	Low	Holm oak shrublands	834.0	42.37°	-5.97°

Number of individuals sampled in each locality, classes based on physiographic and ecological characteristics, mean altitude and coordinates are indicated.

(Applied Biosystems) and allele sizes were determined using GeneMapper 3.7 software (Applied Biosystems).

Data Analysis

Data were checked for null alleles and genotyping errors using MICRO-CHECKER 2.2.3 (van Oosterhout et al. 2004). We estimated the following genetic diversity parameters: number of alleles (N_A), allelic richness permuted by the lowest number of individuals genotyped in a locality (A_R), observed and expected heterozygosity (H_o and H_e) and inbreeding coefficient (F_{IS}) using FSTAT 2.9.3 (Goudet 1995). Departures from Hardy–Weinberg equilibrium were assessed in GenoDive 2.0b20 (Meirmans and Van Tienderen 2004).

To investigate the genetic structure and spatial location of genetic discontinuities within the breeding population, we first employed a Bayesian clustering method without prior assignment to their locations of origin. For that purpose, we used GENELAND 3.2.2 (Guillot et al. 2005; Guillot et al. 2008), which utilizes both genetic information and geographic coordinates from each individual to infer population structure. We initially ran 10 independent Markov Chain Monte Carlo (MCMC) simulations using the following parameters: 5×10^5 iterations, maximum rate of Poisson process fixed at 50, maximum number of nuclei in the Poisson–Voronoi tessellation fixed at 150, and the Dirichlet model for allele frequencies. Since the number of genetic populations was unknown, we allowed the number of clusters (K) to vary on a wide range from $K = 1$ to $K = 10$. Next, we determined the best number of clusters from the highest likelihood number of K obtained from these runs and ran the MCMC 20 times with K fixed to the value identified in the first step. We then computed the posterior probability of population membership for each pixel of the spatial domain (150×150 pixels) and for each individual for each of the 20 runs (with a burn-in of 5×10^4 iterations).

Spatial patterns of genetic differentiation across the full landscape were visualized using the “Genetic Landscape

Shape interpolation” analysis implemented in Alleles in Space 1.0 (Miller 2005). This analysis infers a genetic surface based on interindividual distances of sampled individuals and on interpolated distances in areas where individuals were not sampled. Across the genetic landscape, the peaks and troughs indicate high and low genetic distances between individuals, respectively.

To test genetic differentiation among all sampling localities and to assess whether the inferred genetic clusters, the physiographic or habitat characteristics (i.e., mountain range, altitude, and vegetation) explained a higher percentage of the genetic variance, we performed an analysis of molecular variance (AMOVA) in GenoDive 2.0b20. Moreover, we calculated the genetic diversity parameters previously explained for each group of localities obtained from the best partition in AMOVA.

In addition, we tested the effect of geographic distance on the observed genetic differentiation of the bluethroat. We calculated Euclidean and altitudinal distances between localities and individuals, and tested their correlation with their genetic distance (pairwise $F_{ST}/1-F_{ST}$ between localities and Smouse & Peakall distances between individuals; Smouse & Peakall 1999, using Mantel tests; Mantel 1967). We used partial Mantel tests (Smouse et al. 1986) to assess the association between altitudinal and genetic distances while controlling for the influence of Euclidean geographic distances and vice versa (i.e., the association between geographic and genetic distances controlled by altitudinal distances). These analyses were performed in GenoDive 2.0b20 and their statistical significance was assessed by 10 000 randomizations.

Further relationships of altitude of sampling localities with genetic diversity parameters (N_A , A_R , H_o , H_e) were tested by Pearson correlations. Statistical support for the hypothesis that localities with different habitat features differ in genetic diversity was tested using a type-III analysis of variance (ANOVA), with altitudinal block (high or low) and mountain range (Cantabrian Mountains or Mountains of León) as factors and each of the genetic diversity parameters as response variables. Finally, to address if the assignment of

Table 2 Genetic diversity of bluethroat based on microsatellite loci for the whole population and for each of the three genetic clusters (K-NE, K-NW and K-S) inferred in GENELAND

Locus	Ase19	Cuμ4	Cuμ10	Hru7	Mcy4	PAT MP 2-43	PmaC25	Ppi2	Ptc2	Phtr2	Pdo5*	Aar8*	Mean (SD)
K-NE (<i>n</i> = 16)													
N_A	4	5	3	7	5	4	3	5	2	8	5	1	4.636 (1.747)
A_R	3.597	4.818	2.988	6.613	4.812	3.682	2.786	5.000	2.000	7.316	4.734	1.000	4.395 (1.604)
H_o	0.500	0.938	0.250	0.750	0.688	0.750	0.286	0.545	0.286	0.857	0.214	0.000	0.551 (0.079)
H_e	0.606	0.729	0.425	0.760	0.644	0.631	0.508	0.773	0.516	0.835	0.541	0.000	0.634 (0.039)
F_{IS}	0.212	-0.183	0.600	-0.027	-0.123	-0.122	0.19	0.231	0.323	-0.007	0.508	na	0.090 (0.081)
K-NW (<i>n</i> = 27)													
N_A	5	4	3	7	6	4	3	5	3	11	7	1	5.273 (2.412)
A_R	4.390	3.963	2.394	6.496	5.227	3.344	2.984	4.963	2.653	8.729	5.729	1.000	4.625 (1.890)
H_o	0.556	0.593	0.115	0.923	0.852	0.593	0.500	0.731	0.519	0.889	0.200	0.000	0.588 (0.078)
H_e	0.652	0.706	0.245	0.814	0.748	0.607	0.520	0.795	0.520	0.875	0.562	0.000	0.640 (0.054)
F_{IS}	0.050	0.097	0.053	-0.060	-0.092	-0.091	0.021	0.192	0.046	0.144	0.695	na	0.033 (0.042)
K-S (<i>n</i> = 40)													
N_A	5	6	3	10	8	6	3	7	4	11	6	1	6.273 (2.611)
A_R	4.074	5.274	2.579	7.307	6.582	3.952	2.983	5.228	2.769	8.504	4.622	1.000	4.897 (1.927)
H_o	0.650	0.625	0.250	0.895	0.850	0.450	0.579	0.605	0.462	0.775	0.176	0.000	0.574 (0.069)
H_e	0.637	0.751	0.267	0.850	0.793	0.421	0.596	0.763	0.535	0.864	0.491	0.000	0.634 (0.058)
F_{IS}	-0.025	0.356	-0.04	-0.111	-0.081	0.180	0.149	0.111	0.117	-0.037	0.683	na	0.052 (0.036)
ALL (<i>n</i> = 83)													
N_A	6	6	3	10	8	6	3	7	4	13	8	1	6.727 (3.003)
A_R	4.185	5.025	2.638	6.911	5.996	3.692	2.977	5.173	2.587	8.577	5.179	1.000	4.813 (1.870)
H_o	0.590	0.675	0.207	0.875	0.819	0.554	0.500	0.640	0.450	0.827	0.192	0.000	0.571 (0.070)
H_e	0.635	0.740	0.289	0.833	0.755	0.563	0.561	0.783	0.527	0.876	0.517	0.000	0.636 (0.049)
F_{IS}	0.070	0.089	0.282	-0.051	-0.085	0.016	0.109	0.182	0.146	0.056	0.629	na	0.058 (0.039)

n: number of samples, N_A : number of alleles, A_R : allelic richness standardized to the minimum sample size, H_o : observed heterozygosity, H_e : expected heterozygosity, F_{IS} : inbreeding index. Bold values indicate significant departures from Hardy–Weinberg equilibrium ($P < 0.05$). *indicates loci that were not included in the analyses.

birds to each of the inferred genetic clusters was independent of altitude, vegetation, and mountain range of their sampling localities, a log-linear analysis of frequencies was performed. The log-linear analysis is considered an ANOVA-like design of frequency data. Specifically, it is used to test the different factors that are used in a cross-tabulation with categorical factors and their interactions for statistical significance (StatSoft-Inc. 2007). All these analyses were performed in STATISTICA 8.0 (StatSoft-Inc. 2007).

Results

Eighty-three bluethroats were captured and genotyped for 12 microsatellite loci. Evidence of null alleles was found for locus Pdo5 and consequently it was not included in further analyses. Also, Aar8 turned out to be monomorphic and was removed. Overall, the number of alleles ranged from 3 for loci PmaC25 and Cuμ10 to 13 for locus Phtr2 (average $N_A = 6.727 \pm 3.003$ standard deviation [SD]). Observed heterozygosity per locus ranged from 0.207 to 0.875 with an average value of $H_o = 0.571 \pm 0.070$ SD (Table 2).

The Bayesian clustering analysis performed with GENELAND suggested an optimum structure of three genetic clusters in over 85% of the MCMC iterations. One cluster (K-NE) consisted of the individuals from northeastern localities of Genicera and Rodillazo. The second cluster

(K-NW) was formed by the northwestern and central localities: Meroy, La Cueta, La Majúa, Ferreras de Cepeda, and La Seca. The third cluster (K-S) included the southernmost localities (Pobladura de la Sierra, Molinaferrera, Villar de Golfer, Bustos, Toralino, and Palacios de la Valduerna) but also the most eastern one (Corcos) (Figure 1B and Figure 2). The three clusters showed similar and significant pairwise F_{ST} values, such as $F_{ST} = 0.025$ ($P = 0.007$) between K-NE and K-NW, $F_{ST} = 0.024$ ($P = 0.004$) between K-NE and K-S, and $F_{ST} = 0.020$ ($P = 0.000$) between K-NW and K-S. All individuals were assigned with high probabilities (>80%) and none of the sampled localities contained individuals assigned to more than one genetic cluster.

The genetic surface obtained in the Genetic Landscape Shape interpolation analysis showed sharper “ridges” in the southwestern part of the range, indicating the greatest genetic distances between localities from Mountains of León and western Cantabrian Mountains (Figure 3). Furthermore, this analysis indicated that genetic distances decreased in areas to the east of the main genetic discontinuity, with the exception of the localities in the northeastern Cantabrian Mountains, which also indicated high genetic differentiation. Qualitatively similar results were obtained regardless of the grid size or distance weighting parameters chosen. Likewise, use of raw genetic distances or residual genetic distances had no effect on the relative shape of the landscape surface.




Figure 2. Maps of the posterior probabilities to belong to each genetic cluster inferred in GENELAND. Color gradient represents high (white) to low (gray) posterior probabilities.

The AMOVA analyses indicated that most of the molecular variation resided among individuals within the breeding population ($F_{IT} = 0.919$). The remaining genetic variation was best explained by differences among the three genetic clusters inferred in GENELAND ($F_{CT} = 0.026$, $P < 0.001$), and no significant differences were found among localities within clusters (Table 3). Partitions according to altitude classes and mountain ranges explained significant although lower percentages of genetic variation, but vegetation was nonsignificant (Table 3).

Genetic diversity parameters were very similar among the 3 inferred clusters (ANOVA, all $P > 0.104$) and compared

with the whole population, although lower genetic variability was found in cluster K-NE (Table 2). Furthermore, none of the genetic diversity parameters were significantly correlated with the altitude of the sampling localities (all P values > 0.148) or were significantly different between mountain ranges (all P values > 0.157).

On the other hand, H_o values were almost significantly different between altitude classes (ANOVA $F_{1,11} = 3.488$, $P = 0.088$), suggesting a tendency for lower genetic diversity in localities at a higher altitude. Furthermore, the altitude at which individuals were sampled was significantly different among the 3 genetic clusters, after controlling for




Figure 3. Genetic Landscape Shape interpolation based on a 50×50 grid and a distance weighting value (α) of 0.2. Surface plot heights are proportionate to genetic distances.

Table 3 Analysis of molecular variance performed between the bluethroat localities analyzed

Partition tested	% variation among groups	F_{CT}	F_{SC}	F_{ST}	F_{IS}
Among localities (All) (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14)	2.2			0.022**	0.054*
Between mountain ranges (Cantabrian Mt.) vs (Mt. León) (1, 2, 3, 4, 5, 7, 8) vs (6, 9, 10, 11, 12, 13, 14)	0.1	0.001*	0.025**		0.089**
Between altitude classes (High) vs (Low) (1, 2, 3, 4, 5, 9) vs (6, 7, 8, 10, 11, 12, 13, 14)	1	0.010*	0.020*		0.089**
Among vegetation types (Brooms) vs (Heathlands) vs (Shrublands) (1, 2, 3, 4, 5, 9) vs (6, 7, 8, 10, 12) vs (11, 13, 14)	0	0	0.025*		0.089**
Among genetic clusters (K-NE) vs (K-NW) vs (K-S) (1, 2) vs (3, 4, 5, 6, 7) vs (8, 9, 10, 11, 12, 13, 14)	2.6	0.026**	0.004		0.054*

F_{IS} : variation among individuals within localities, F_{ST} : variation among localities within the population, F_{SC} : variation of localities within groups, F_{CT} : variation among groups within the population.

*values indicate significant probabilities at $P < 0.05$ and **values indicate significant probabilities at $P < 0.01$. Numbers correspond to locality codes in Table 1.

their geographic position (i.e., latitude and longitude; ANOVA $F_{2,78} = 116.252$, $P < 0.001$), with K-NE at the highest altitude (post-hoc Tukey Test: $P = 0.0002$ for K-NE vs. K-NW and $P = 0.0002$ for K-NE vs. K-S) and K-S at the lowest (post-hoc Tukey Test: $P = 0.0002$ for K-S vs. K-NW). The log-linear analysis indicated that the best model for sample distribution did not include any interaction involving the variable genetic cluster (all P values > 0.501). Only the interaction between the variables genetic cluster and mountain range was close to significance ($\chi^2_2 = 5.457$, $P = 0.065$), indicating a trend for samples from cluster K-S to be more frequent in the Mountains of León than in the Cantabrian Mountains. As

expected for these highly correlated variables, the interaction between vegetation and altitude was significant in the model ($\chi^2_2 = 6.306$, $P = 0.043$), indicating that samples belonging to broom-type vegetation were more frequent at high altitudes and samples in shrub lands were more frequent at low altitudes.

The Mantel test found a nonsignificant correlation between geographic or altitudinal distances and genetic distances between bluethroat localities (Mantel's $r = 0.061$, $P = 0.319$ and $r = 0.007$, $P = 0.456$, respectively), indicating that geographic distance between localities has no effect on their genetic differentiation. On the other hand, correlations were significant when individuals instead of localities

were considered (Mantel's $r = 0.051$, $P = 0.017$ for the geographic distances and $r = 0.060$, $P = 0.025$ for the altitudinal distances). However, when the effect of altitude was controlled by Euclidean geographic distances and vice versa, correlations were not significant (Partial Mantel's $r = 0.014$, $P = 0.317$ and $r = 0.023$, $P = 0.239$).

Discussion

Higher Genetic Structure but Lower Diversity in Spanish Than in European Bluethroat Populations

Three genetic clusters were identified within the Spanish breeding range of *L. s. azuricollis* (Figure 1B and Figure 2), which were almost equally divergent from each other, indicating the existence of well-delimited genetic groups at a local spatial scale and restricted effective dispersal (gene flow) (Clark et al. 2008). Our work provides additional evidence for a significant and much stronger genetic structure in Spain than in northern Europe, considering that the observed values were one order of magnitude greater than those found among all bluethroat populations in Scandinavia ($F_{ST} = 0.002$; Heggren et al. 2008). Furthermore, the levels of genetic differentiation within the Spanish subspecies were in the range of those obtained among distinct bluethroat subspecies across Europe (significant pairwise $F_{ST} = -0.004$ to 0.174, average pairwise $F_{ST} = 0.044 \pm 0.043$ SD). Indeed, at the continental scale, the highest values of genetic differentiation between bluethroat subspecies were those involving comparisons with *L. s. azuricollis*, whereas the lowest were those comparing the subspecies with a northern distribution (Johnsen et al. 2006; Heggren et al. 2008).

Our data were congruent with previous studies, with 9 out of 10 microsatellite loci in common but lower sampling size, indicating that *L. s. azuricollis* is the subspecies with the lowest genetic variability. On average, the Spanish population holds $38.6\% \pm 21.6$ SD of all the species alleles, although ranging from 76.9% to 16.6% depending on the locus considered (Johnsen et al. 2006). One possibility is that the low genetic diversity of bluethroats breeding in Spain is a consequence of their geographic and genetic isolations, because the associated effects of genetic drift may both decrease genetic diversity and increase differentiation (Frankham et al. 2002).

In addition, the apparently high philopatry and low gene flow at local scales compared with northern European populations (Heggren et al. 2008), and the fact that *L. s. azuricollis* is basal to the remaining European subspecies (Johnsen et al. 2006), might also support an isolation of Spanish breeding bluethroats and suggest a relatively independent evolution for this subspecies. This might explain their pattern of greater genetic differentiation, because besides the effect of geographic distance, the isolation of local populations would promote more rapid evolutionary change within the breeding population, and thus more rapid differentiation from the European populations from which it is isolated (Wright 1940). Furthermore, this pattern of genetic variation agrees with a nonmutually exclusive

hypothesis proposing an inverse relationship between population differentiation and latitude (Martin and McKay 2004). Our results support the arguments of several authors that increased seasonal variation in climatic conditions at higher latitudes may result in broader tolerance of northern organisms to environmentally changing conditions. Thus, a greater adaptation capability could reduce costs of dispersing between populations, resulting in relaxed philopatric behavior and also in higher levels of gene flow and reduced genetic differentiation among high latitude populations (Martin and McKay 2004; Croteau et al. 2007; Berg et al. 2010). In contrast, strong fidelity to breeding sites at lower latitudes would prevent gene flow among different populations and might reduce genetic variation for dispersal behavior (Both and Visser 2001).

Environmental Factors Shaping Genetic Structure and Diversity

Our study helps identify some of the key factors conditioning species dispersal and distribution, and contributes to a growing body of work that suggests that landscape features influence dispersal and gene flow among bird populations (Bruggeman et al. 2010; Coulon et al. 2010; Milá et al. 2010; Thomassen et al. 2010; Alda et al. 2011). As has been described in previous studies, we found that geographic distance by itself is not a factor determining genetic differentiation in the bluethroat, neither at a local nor at a continental scale (Johnsen et al. 2007). In this case, altitude and mountain range of the localities explained significant percentages of genetic variance (Table 3) and were likely responsible for the observed genetic differentiation, as revealed by the significant differences in altitude among clusters, as well as the almost significant association observed between mountain ranges and the inferred genetic clusters. Indeed, these factors were clearly reflected in the landscape analyses of genetic structure, which showed genetic differentiation of the localities in Mountains of León, as well as those in the highest northeastern localities (Figures 2 and 3). Moreover, these areas that encompass high environmental and genetic variations are particularly important for maximizing adaptive diversity and consequently should be prioritized for conservation (Thomassen et al. 2010). In the end, we must be aware that the variables defined for this study are correlated with ultimate factors, such as climate, which will condition phenology and habitat availability. Therefore, we must keep in mind the combined effect of multiple factors on avian habitat selection that consequently gives rise to the observed genetic structure (Milá et al. 2010).

Limited or differential availability of those features selected by a species across its range distribution may not only explain genetic structure but also differences in population sizes and consequently in genetic diversity (Salvi et al. 2009). We observed a general, although nonsignificant, tendency for lower genetic diversity at high altitude localities. Such patterns of differentiation in altitude are expected in organisms with low dispersal abilities but are remarkable in species with high potential for dispersal, especially given the small geographic

scale of our study (Martínez-Solano and González 2008; Milá et al. 2010). Although our limited sampling size precludes drawing definite conclusions regarding this issue, we might deduce, based on this trend and the genetic differentiation of some high-altitude sites (e.g., cluster K-NE), that a limited number of individuals reach these regions. We further hypothesize that climate variables, such as time differences in the melting of snow at increasing altitudes, might limit habitat availability and thus hinder colonization of breeders and eventually gene flow (Santos González et al. 2010). Our results suggest that the environmental differences across the range explain the putatively neutral genetic variation, rather than by isolation by distance, which further indicates that this pattern of genetic structure might likely be shaped by adaptive differentiation (Salvi et al. 2009; Thomassen et al. 2010). However, the mechanisms underlying the observed genetic structure remain unknown. In our case, genetic differentiation between low- and high-altitude sites could be associated with differences in life-history traits. These differences could be the result of divergent selection pressures, which could have a role in restricting gene flow and leading to local adaptations and differentiation (Milá et al. 2010). On the other hand, under a high migration connectivity scenario, birds arriving from different wintering areas or at different times could select different breeding sites depending on their ecological characteristics. In other species, this pattern has been detected on the basis of genetic differences in birds arriving or breeding at different times in the same place (Moore et al. 2005; Casagrande et al. 2006; Porlier et al. 2009). Nevertheless, for the bluethroat, it is still unknown whether Spanish breeding birds show a pattern of temporal genetic differentiation or originate from different wintering areas (Arizaga et al. 2006). Further research with broader geographical sampling and additional genetic and morphological markers would be necessary to test these hypotheses, as adaptive changes in morphology often evolve at a faster rate than neutral genetic markers and may reflect noncongruent patterns of differentiation (Marthinsen et al. 2007; Milá et al. 2009).

Implications for Conservation

The strength of local selection informs how a species might react in diverse and dynamic environments and influences its potential for adaptation in the face of future climate change (Walther et al. 2002; Thomassen et al. 2010). In this respect, it is necessary to bear in mind that in the Iberian Peninsula, there is no suitable habitat for the bluethroat further north of the Cantabrian Mountains. Consequently, under a global warming scenario, the northward expansion of the Spanish subspecies would be limited (Walther et al. 2002; Förtschler et al. 2011). It remains unclear if the proposed site selection and philopatry is strong enough to hamper the adaptation of individuals from clusters K-NE and K-NW to a southern and more Mediterranean habitat under a global warming scenario. On the contrary, if lowland Mediterranean habitats were to expand under such climatic scenario, bluethroats might expand their populations from those already extant in

those regions (K-S). Ultimately, all of the above strengthen the importance of preserving the evolutionary potential held in these areas encompassing both high environmental and genetic variations.

Funding

University of León (2009/00131/00); CSIC/MICINN Proyecto Intramural Especial-PIE (201030I019).

Acknowledgement

We thank María Calero-Riestra for her help with the statistical analyses and two anonymous referees for helpful suggestions. We also thank Sara Lipshutz for kindly reviewing the English text. Nacho Rodríguez, Eva Álvarez, and other ringers of the Grupo Ibérico de Anillamiento helped with fieldwork.

References

- Alda F, Sastre P, de la Cruz-Cardiel PJ, Doadrio I. 2011. Population genetics of the endangered Cantabrian capercaillie in northern Spain. *Animal Conserv.* 14:249–260.
- Arizaga J, Campos F, Alonso D. 2006. Variations in wing morphology among subspecies might reflect different migration distances in bluethroat. *Ornis Fennica.* 83:162–169.
- Balloux F, Lugon-Moulin N. 2002. The estimation of population differentiation with microsatellite markers. *Mol Ecol.* 11:155–165.
- Bell G, Lechowicz MJ, Appenzeller A, Chandler M, Deblois E, Jackson L, Mackenzie B, Preziosi R, Schallenberg M, Tinker N. 1993. The spatial structure of the physical environment. *Oecologia.* 96:114–121.
- Berg MP, Kiers ET, Driessen G, van der Heijden M, Kooi BW, Kuenen F, Liefing M, Verhoef HA, Eilers J. 2010. Adapt or disperse: understanding species persistence in a changing world. *Glob Change Biol.* 16:587–598.
- Blondel J, Thomas DW, Charmantier A, Perret P, Bourgault P, Lambrechts MM. 2006. A thirty-year study of phenotypic and genetic variation of blue tits in Mediterranean habitat mosaics. *BioScience.* 56:661–673.
- Both C, Visser ME. 2001. Adjustment to climate change is constrained by arrival date in a long-distance migrant bird. *Nature.* 411:296–298.
- Bruggeman DJ, Wiegand D, Fernández N. 2010. The relative effects of habitat loss and fragmentation on population genetic variation in the red-cockaded woodpecker (*Picoides borealis*). *Mol Ecol.* 19:3679–3691.
- Casagrande S, Dell’Omo G, Costantini D, Tagliavini J. 2006. Genetic differences between early- and late-breeding Eurasian kestrels. *Evol Ecol Res.* 8:1029–1038.
- Clark RW, Brown WS, Stechert R, Zamudio KR. 2008. Integrating individual behaviour and landscape genetics: the population structure of timber rattlesnake *hibernacula*. *Mol Ecol.* 17:719–730.
- Coulon A, Fitzpatrick JW, Bowman R, Lovette IJ. 2010. Effects of habitat fragmentation on effective dispersal of Florida scrub-jays. *Conserv Biol.* 24:1080–1088.
- Cramp S. 1988. The Bluethroat. In: Cramp S, editor. *The Birds of the Western Palearctic Handbook of Birds in Europe, the Middle East and Africa.* Oxford: Oxford University Press. p. 645–661.
- Croteau EK, Loughheed SC, Krannitz PG, Mahony NA, Walker BL, Boag PT. 2007. Genetic population structure of the sagebrush Brewer’s sparrow, *Spizella breweri breweri*, in a fragmented landscape at the northern range periphery. *Conserv Genet.* 8:1453–1463.
- Davis JM, Stamps JA. 2004. The effect of natal experience on habitat preferences. *Trends Ecol Evol.* 19:411–416.

- Double MC, Dawson D, Burke T, Cockburn A. 1997. Finding the fathers in the least faithful bird: A microsatellite-based genotyping system for the superb fairy-wren *Malurus cyaneus*. *Mol Ecol*. 6:691–693.
- Dufour A, Gadallah F, Wagner HH, Guisan A, Buttler A. 2006. Plant species richness and environmental heterogeneity in a mountain landscape: effects of variability and spatial configuration. *Ecography*. 29:573–584.
- Ellegren H. 1992. Polymerase chain reaction (PCR) analysis of microsatellites—a new approach to studies of genetic relationships in birds. *Auk*. 109:886–895.
- Esler D. 2000. Applying metapopulation theory to conservation of migratory birds. *Conserv Biol*. 14:366–372.
- Felsenstein J. 1976. The theoretical population genetics of variable selection and migration. *Annu Rev Genet*. 10:253–280.
- Foll M, Gaggiotti O. 2006. Identifying the environmental factors that determine the genetic structure of populations. *Genetics*. 174:875–891.
- Förschler MI, Senar JC, Borrás A, Cabrera J, Björklund M. 2011. Gene flow and range expansion in a mountain-dwelling passerine with a fragmented distribution. *Biol J Linn Soc*. 103:707–721.
- Frankham R. 1995. Conservation genetics. *Annu Rev Genet*. 29:305–327.
- Frankham R, Ballou JD, Briscoe DA, McInness KH. 2002. Introduction to conservation genetics. Cambridge (UK): Cambridge University Press.
- Fridolfsson AK, Gyllensten UB, Jakobsson S. 1997. Microsatellite markers for paternity testing in the willow warbler *Phylloscopus trochilus*: high frequency of extra-pair young in an island population. *Hereditas*. 126:127–132.
- Gibbs HL, Tabak LM, Hobson K. 1999. Characterization of microsatellite DNA loci for a neotropical migrant songbird, the Swainson's thrush (*Catharus ustulatus*). *Mol Ecol*. 8:1551–1552.
- Gómez-Manzanique A. 2003. Pechiazul *Luscinia svecica*. In: Martí R, del Moral JC, editors. Atlas de las aves reproductoras de España. Madrid: Dirección General de Conservación de la Naturaleza-SEO/BirdLife. p. 420–421.
- Goudet J. 1995. FSTAT (version 1.2): a computer program to calculate F-statistics. *J Hered*. 86:485–486.
- Guillot G, Mortier F, Estoup A. 2005. GENELAND: a computer package for landscape genetics. *Mol Ecol Notes*. 5:712–715.
- Guillot G, Santos F, Estoup A. 2008. Analysing georeferenced population genetics data with Geneland: a new algorithm to deal with null alleles and a friendly graphical interface. *Bioinformatics*. 24:1406–1407.
- Guschanski K, Caillaud D, Robbins MM, Vigilant L. 2008. Females shape the genetic structure of a gorilla population. *Curr Biol*. 18:1809–1814.
- Hedrick PW. 1986. Genetic polymorphism in heterogeneous environments: a decade later. *Annu Rev Ecol Syst*. 17:535–566.
- Hedrick PW. 2006. Genetic polymorphism in heterogeneous environments: the age of genomics. *Annu Rev Ecol Syst*. 37:67–93.
- Hellgren O, Bensch S, Hobson KA, Lindström A. 2008. Population structure and migratory directions of Scandinavian bluethroats *Luscinia svecica*—a molecular, morphological and stable isotope analysis. *Ecography*. 31:95–103.
- Hull JM, Hull AC, Sacks BN, Smith JP, Ernest HB. 2008. Landscape characteristics influence morphological and genetic differentiation in a widespread raptor (*Buteo jamaicensis*). *Mol Ecol*. 17:810–824.
- Johnsen A, Andersson S, García Fernández J, Kempnaers B, Pavel V, Questiau S, Raess M, Rindall E, Liffield JT. 2006. Molecular and phenotypic divergence in the bluethroat (*Luscinia svecica*) subspecies complex. *Mol Ecol*. 15:4033–4047.
- Johnsen A, Fidler AE, Kuhn S, Carter KL, Hoffman A, Barr IR, Biard C, Charmanier A, Eens M, Korsten P, et al. 2007. Avian *Clock* gene polymorphism: evidence for a latitudinal cline in allele frequencies. *Mol Ecol*. 16:4867–4880.
- Kawecki TJ, Ebert D. 2004. Conceptual issues in local adaptation. *Ecol Lett*. 7:1225–1241.
- MacColl ADC, Piertney S, Moss R, Lambin X. 2000. Spatial arrangement of kin affects recruitment success in young male red grouse. *Oikos*. 90:261–270.
- Mantel NA. 1967. The detection of disease clustering and a generalized regression approach. *Cancer Res*. 27:209–220.
- Markovets M, Yosef R. 2005. Phenology, duration and site fidelity of wintering bluethroat (*Luscinia svecica*) at Eilat, Israel. *J Arid Environ*. 61:93–100.
- Marthinsen G, Wennerberg L, Liffield JT. 2007. Phylogeography and subspecies taxonomy of dunlins (*Calidris alpina*) in western Palearctic analysed by DNA microsatellites and amplified fragment length polymorphism markers. *Biol J Linn Soc*. 92:713–726.
- Martin PR, McKay JK. 2004. Latitudinal variation in genetic divergence of populations and the potential for future speciation. *Evolution*. 2004:5.
- Martínez J, Soler J, Soler M, Møller A, Burke T. 1999. Comparative population structure and gene flow of a brood parasite, the great spotted cuckoo (*Clamator glandarius*), and its primary host, the magpie (*Pica pica*). *Evolution*. 53:269–278.
- Martínez-Solano I, González EG. 2008. Patterns of gene flow and source-sink dynamics in high altitude populations of the common toad *Bufo bufo* (Anura: Bufonidae). *Biol J Linn Soc*. 95:824–839.
- Meirmans PG, Van Tienderen PH. 2004. GENOTYPE and GENODIVE: two programs for the analysis of genetic diversity of asexual organisms. *Mol Ecol Notes*. 4:792–794.
- Milá B, Warren BH, Heeb P, Thébaud C. 2010. The geographic scale of diversification on islands: genetic and morphological divergence at a very small spatial scale in the Mascarene grey white-eye (Aves: *Zosterops borbonicus*). *BMC Evol Biol*. 10:158.
- Milá B, Wayne RK, Fitze P, Smith TB. 2009. Divergence with gene flow and fine-scale phylogeographical structure in the wedge-billed wood-creeper, *Chyborhynchus spirurus*, a Neotropical rainforest bird. *Mol Ecol*. 18:2979–2995.
- Miller MP. 2005. Alleles In Space (AIS): Computer software for the joint analysis of interindividual spatial and genetic information. *J Hered*. 96:722–724.
- Moore IT, Bonier F, Wingfield JC. 2005. Reproductive asynchrony and population divergence between two tropical bird populations. *Behav Ecol*. 16:755–762.
- Otter K, Ratcliffe L, Michaud D, Boag PT. 1998. Do female black-capped chickadees prefer high-ranking males as extra-pair partners? *Behav Ecol Sociobiol*. 43:25–36.
- Penteriani V, Delgado MM, Gallardo M, Ferrer M. 2004. Spatial heterogeneity and structure of bird populations: a case example with the eagle owl. *Popul Ecol*. 46:185–192.
- Perbal BA. 1988. A practical guide to molecular cloning. New York: Wiley.
- Pitra C, Suárez-Seoane S, Martín CA, Streich W-J, Alonso JC. 2011. Linking habitat quality with genetic diversity: a lesson from great bustards in Spain. *Eur J Wildl Res*. 57:411–419.
- Porlier M, Bélisle M, Garant D. 2009. Non-random distribution of individual genetic diversity along an environmental gradient. *Proc R Soc Lond B*. 364:1543–1554.
- Primmer C, Møller A, Ellegren H. 1996. New microsatellites from the pied flycatcher *Ficedula hypoleuca* and the swallow *Hirundo rustica* genomes. *Hereditas*. 124:281–284.
- Questiau S, Eybert M-C, Gaginckaya AR, Gielly L, Taberlet P. 1998. Recent divergence between two morphologically differentiated subspecies of bluethroat (Aves: Muscicapidae: *Luscinia svecica*) inferred from mitochondrial DNA sequence variation. *Mol Ecol*. 7:239–245.
- Richardson DS, Jury FL, Dawson DA, Salgueiro P, Komdeur J, Burke T. 2000. Fifty Seychelles warbler (*Acrocephalus sechellensis*) microsatellite loci polymorphic in Sylviidae species and their cross-species amplification in other passerine birds. *Mol Ecol*. 9:2226–2231.
- Saladin V, Bonfils D, Binz T, Richner H. 2003. Isolation and characterization of 16 microsatellite loci in the Great Tit *Parus major*. *Mol Ecol Notes*. 3:520–522.

- Salvi D, Capula M, Bombi P, Bologna MA. 2009. Genetic variation and its evolutionary implications in a Mediterranean island endemic lizard. *Biol J Linn Soc.* 98:661–676.
- Santos González J, Redondo Vega JM, Gómez Villar A, González Gutiérrez RB. 2010. Dinámica actual de los nichos de nivación del Alto Sil (Cordillera Cantábrica). *Cuadernos de Investigación Geográfica.* 36:87–106.
- Smouse PE, Long JC, Sokal RR. 1986. Multiple regression and correlation extensions of the Mantel test of matrix correspondence. *Syst Zool.* 35:627–632.
- Smouse PE, Peakall R. 1999. Spatial autocorrelation analysis of individual multiallele and multilocus genetic structure. *Heredity.* 82:561–573.
- StatSoft-Inc. 2007. STATISTICA (data analysis software system) version 8. Available from: www.statsoft.com.
- Tellería JL. 1999. Aves Ibéricas. Vol. II. Paseriformes. Madrid: J. M. Reyero.
- Thomassen HA, Buermann W, Milá B, Graham CH, Cameron SE, Schneider CJ, Pollinger JP, Saatchi S, Wayne RK, Smith TB. 2010. Modeling environmentally associated morphological and genetic variation in a rainforest bird, and its application to conservation prioritization. *Evol Appl.* 3:1–16.
- Van Bekkum M, Sagar PM, Stahl J-C, Chambers GK. 2006. Natal philopatry does not lead to population genetic differentiation in Buller's albatross (*Thalassarche bulleri bulleri*). *Mol Ecol.* 15:73–79.
- van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. 2004. MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Mol Ecol Notes.* 4:535–538.
- Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin J-M, Hoegh-Guldberg O, Bairlein F. 2002. Ecological response to recent climate change. *Nature.* 416:389–395.
- Webster MS, Marra PP, Haig SM, Bensch S, Holmes RT. 2002. Links between worlds: unraveling migratory connectivity. *Trends Ecol Evol.* 17:76–83.
- Wright S. 1940. Breeding structure of populations in relation to speciation. *Am Nat.* 74:232–248.
- Zink RM, Drovetski SV, Questiau S, Fadeev IV, Nesterov EV, Westberg MC, Rohwer S. 2003. Recent evolutionary history of the bluethroat (*Luscinia svecica*) across Eurasia. *Mol Ecol.* 12:3069–3075.

Received December 13, 2011; Revised July 11, 2012;
Accepted July 17, 2012

Corresponding Editor: Jennifer Jackson