

1 **Is the interaction between rabbit hemorrhagic disease and**  
2 **hyperpredation by raptors a major cause of the red-legged partridge**  
3 **decline in Spain?**

4

5

6 José A. BLANCO-AGUIAR\*<sup>1,2</sup>, Miguel DELIBES-MATEOS<sup>1</sup>, Beatriz ARROYO<sup>1</sup>, Pablo FERRERAS<sup>1</sup>,  
7 -Fabián CASAS<sup>1,4</sup>, Raimundo REAL<sup>3</sup>, Mario J. VARGAS<sup>3</sup>, Rafael VILLAFUERTE<sup>1</sup> & Javier  
8 VIÑUELA<sup>1</sup>.

9

10 <sup>1</sup> *Instituto de Investigación en Recursos Cinegéticos (IREC, CSIC-UCLM-JCCM), Ronda de Toledo s/n,*  
11 *E-13071 Ciudad Real, Spain*

12 <sup>2</sup> *CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto,*  
13 *Campus Agrário de Vairão, 4485-661 Vairao, Portugal*

14 <sup>3</sup> *Departamento de Biología Animal, Facultad de Ciencias, Universidad de Málaga, 29071 Málaga,*  
15 *Spain*

16 <sup>4</sup> *School of Biological Sciences, University of Aberdeen Zoology Building, Tillydrone Avenue, Aberdeen*  
17 *AB24 2TZ, Scotland, United Kingdom*

18

19 \*Corresponding author.

20 Email: jota.blanco@gmail.com

21

22

1 **Abstract**

2 ~~Hyperpredation can be defined~~

3 ~~Hyperpredation can be described as a restrictive case of apparent competition, where an increased number~~  
4 ~~of primary prey species indirectly induces the decrease of the secondary prey species through numerical~~  
5 ~~response of predators to the primary prey dynamics, as an enhanced predation pressure on a secondary~~  
6 ~~prey due to either an increase of predator population supported by an increase of its main prey, or by a~~  
7 ~~sudden drop in the abundance of the main prey.~~ It has been proposed that Rabbit Hemorrhagic Disease  
8 (RHD), which decimated populations of European wild rabbit (*Oryctolagus cuniculus*) in Spain, led to  
9 prey switching by raptors towards red-legged partridges (*Alectoris rufa*), causing declines in their  
10 populations, ~~as consequence of hyperpredation process,~~

11 We have reviewed field studies that disagree with estimates (based on bag records) of recent increases of  
12 rabbit and partridge numbers in Spain. Because of an increase in releases of farm-reared animals, there is  
13 doubt about the use of bag records to accurately estimate population trends in recent decades. We also  
14 provide new data and discuss some analytical considerations related to the temporal and spatial scales that  
15 might affect the interpretation of data. Finally, we discuss why studies associated with raptor food habits,  
16 predator population dynamics and predator distribution suggest that diet data are not sufficient to link  
17 patterns and processes. Although we agree that the RHD outbreak has markedly affected rabbit  
18 populations, which has indirectly affected many other species in the Iberian Peninsula, we consider that  
19 hyperpredation mediated by raptors has not been clearly demonstrated. In contrast, endorsing the  
20 hypothesis of raptor-mediated hyperpredation without sufficient proof may have conflictive consequences  
21 if we consider the ~~increased intensification of the~~ persecution of raptors ~~occurred during the last~~ in recent  
22 decades in Spain.

23

24 **Keywords:** *Alectoris*; emerging disease; hunting; Mediterranean ecosystem; *Oryctolagus*;  
25 predator persecution; shooting

26

Con formato: Inglés (Estados Unidos)

Con formato: Fuente: (Predeterminado) Times New Roman, 10 pto

Con formato: Espacio Antes: Automático, Después: 0 pto, Dividir palabras

Con formato: Fuente: 10 pto

Con formato: Fuente: 12 pto, Inglés (Estados Unidos)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

## Introduction

Over the past decades, the red-legged partridge (*Alectoris rufa*) has notably declined in the Iberian Peninsula, where the species probably originated and where its wild populations reach the highest densities within its current distribution range (Blanco-Aguilar et al. 2004). Excessive hunting pressure, changes in pesticide management and livestock densities, hybridization with *Alectoris chukar*, diseases associated to farm-reared game releases, and the intensification of agriculture have been cited as possible causes of partridges decline (e.g. Blanco-Aguilar 2007; Blanco-Aguilar et al. 2008; Villanúa et al. 2008; Casas and Viñuela 2010; Díaz-Sánchez et al. 2011). ~~In addition, it~~ has been recently proposed that hyperpredation could have been one of the main causes explaining partridge decline in the Iberian Peninsula. Hyperpredation process has been described as a restrictive case of apparent competition, where an increased number of primary prey species indirectly induces the decrease of the secondary prey species through numerical response of predators to the primary prey dynamics (e.g. Smith and Quin 1996; Courchamp et al. 2000). However, ~~a~~ according to Moleón et al. (2008), hyperpredation refers to an enhanced predation pressure on a secondary prey due to either an increase in the abundance of a predator population or a sudden drop in the abundance of the main prey ~~the hyperpredation effect (Moleón et al. 2008).~~ Following only the latter approach, the Rabbit Hemorrhagic Disease (RHD) outbreak, which decimated populations of European wild rabbit (*Oryctolagus cuniculus*), forced raptors to switch from eating rabbits to eating red-legged partridges. In turn, this is hypothesized to have further declined or limited partridge populations. More recently, a post immunization phase with a reduction in rabbit mortality due to the immunization, led again to medium-high rabbit population densities, which would have led to partridge population increases, as raptors would have stopped concentrating on secondary prey once their main prey was recovered.

We assessed the potential role of hyperpredation (or enhanced predation) on partridge decline. In particular: (1) whether or not rabbit and wild partridge populations have indeed increased during the last decade, as suggested by Moleón et al. (2008), (2) whether the analytical approach used in Moleón et al. (2008) is adequate to support the hyperpredation hypothesis, and (3) whether the functional response of the three raptor species mentioned by Moleón et al. (2008), which generally are found at low population densities, can cause the suggested large-scale effects on the dynamics of partridge populations. Finally,

- Con formato: Fuente: 10 pto
- Con formato: Fuente: (Predeterminado) Times New Roman, 10 pto
- Con formato: Fuente: (Predeterminado) Times New Roman, 10 pto
- Con formato: Fuente: (Predeterminado) Times New Roman, 10 pto
- Con formato: Fuente: (Predeterminado) Times New Roman, 10 pto
- Con formato: Fuente: 10 pto

1 we discuss the potential consequences of hyperpredation for the conflict between hunting and  
2 biodiversity.

3

4

## 5 **Material and methods**

6 To evaluate whether or not partridge and rabbit populations have recently increased, we reviewed the  
7 results of field studies on rabbit and partridge population trends in the Iberian Peninsula. Using this  
8 information, we discuss the reliability of current hunting bags as an indicator of game species abundance.  
9 We also reviewed some limitations of data and analyses used to test the hyperpredation hypothesis, and  
10 incorporated new data that questioned this hypothesis. In this respect, we assessed whether or not the  
11 predator community fed differently before and after the outbreak of RHD. In addition, we discuss the  
12 potential consequences of prey populations by predators according to the functional response of raptors  
13 and the density dependence of productivity theories (Sinclair and Pech 1996; Valkama et al. 2005).  
14 Finally, Moleón et al. (2008) used information from national bag records to describe partridge population  
15 trends, whereas they used only four local raptor studies (figure 1 in reference Moleón et al. 2008) to infer  
16 processes causing these patterns (i.e. predation). However, the patterns that we observe in nature, and the  
17 processes that cause them, can change with the scale of our observation (Wiens 1989; Levin 1992). For  
18 this reason, we used the same hunting bags data, from the Spanish Ministry of Agriculture, Fisheries and  
19 Food, to evaluate whether rabbit and partridge densities follow the same trends at national scale as in the  
20 provinces where Moleón et al. (2008) analyzed raptor diet. Provincial (and national) harvest numbers are  
21 based on questionnaires that all managers of hunting lands are required to complete each year and send to  
22 the Ministry of Agriculture [et al.](#) This agency calculates (and publishes) a summary value of the harvest in  
23 each province by summing the data from each hunting land in each province for each year. Because  
24 hunting effort can influence the number of rabbits and partridges hunted, we divided total harvest by the  
25 number of hunting licenses in each province and year to obtain a reliable estimate of population size  
26 (Cattadori et al. 2003, Virgós et al. 2007).

## 27 **Results and Discussion**

28 Did rabbit and partridge populations increase during the last decade in Spain?

1  
2 Monitoring programmes are among the most important data sources for assessing temporal changes in  
3 biodiversity (e.g. Fewster et al. 2000; van Strien et al. 2001; Gregory et al. 2005). Most monitoring data  
4 available in the literature show that both partridges and rabbits have followed stable or even negative  
5 trends since the mid 1990s, which clearly contradicts the pattern suggested by national bag records. For  
6 instance, wild partridge population trends between 1998 and 2006 estimated from the Spanish monitoring  
7 programme of common breeding birds (SACRE), showed a fairly stable trend, -0.2% annual population  
8 growth rate (SEO/BIRDLIFE 2006). Also, a recent review of wild rabbit trends during the last decade in  
9 different Iberian regions showed a negative trend in five of six independent regional field studies  
10 (covering more than 300 study plots; Delibes-Mateos et al. 2009b).

11 The contradiction between the findings of Moleón et al. (2008) and other published studies  
12 regarding rabbit and partridge demographic trends comes apparently from the data source used by these  
13 authors to estimate population trends. They used hunting bag data from the Spanish Ministry of  
14 Agriculture, Fisheries and Food. However, bag records can be useful to estimate population trends *only* if  
15 the relationship between harvesting and population density remains unchanged throughout the time period  
16 studied. During the last decade, the Spanish Red-legged partridge bag records are likely inflated by the  
17 releases of farm-raised partridges, and the authors did not critically address this issue. There are not  
18 accurate records about releases in Spain, but Blanco-Aguilar et al. (2008) documented, using partridge-  
19 farm advertisement records in hunting journals, an exponential growth of partridge-farm facilities and  
20 releases, which started in the mid 1980s but that was particularly marked from the late 1990s. It is  
21 estimated that since 2000 between 4 and 5 million farm-bred partridges have been released every year in  
22 Spain for hunting (Gortázar et al. 2000; REGHAB 2002). Unfortunately, Spanish bag records cannot  
23 discriminate between wild-born and released partridges, but it is highly probable that released partridges  
24 have been inflating bag records since the 1990s, particularly in those provinces where releases are  
25 frequent. For instance, in Navarra province, the only Spanish province where partridge releases are  
26 forbidden since 1991, field surveys of red-legged partridges have shown stable or declining trends  
27 (Gobierno de Navarra 2006). Therefore, changes in game management practices could have distorted the  
28 value of partridge hunting bags as indicators of population size since late 1990s.

29 Similarly, Ferreira et al. (2010) have recently observed a discrepancy between rabbit trends  
30 obtained from field data and hunting statistics in Portugal. These authors argued that hunting bags may be

Comentario [B1]: Ver comment en referencias

1 influenced by sporadic management operations undertaken by hunters and the lack of systematic  
2 procedures in data collection, and therefore hunting statistics are probably not representative of real  
3 changes in rabbit populations.

4 On the other hand, when national bag records per license are analyzed from 1973 it is possible to  
5 observe that rabbit populations had some fluctuations before RHD arrival (1973-1988). In contrast,  
6 partridge populations showed a marked decline in that same period (Fig. 1; see also Blanco-Aguiar 2007),  
7 and therefore it is not possible to discard that the same factors that were affecting partridge populations  
8 before RHD arrival were also working after RHD arrival. In fact, other possible causes of [red-legged](#)  
9 [partridge RLP](#) decline in Spain have been suggested. These include disease, habitat loss, agricultural  
10 intensification, earlier cereal harvesting, and modification of game management (including changes in  
11 shooting intensity) (Blanco-Aguiar et al. 2004; Buenestado et al. 2008; Villanúa et al. 2008; Buenestado  
12 et al. 2009; Delibes-Mateos et al. 2009a; Casas and Viñuela 2010).

13

14 Are feeding data enough to link patterns and processes?

15

16 *Assumptions associated with raptor food habits.* Moleón et al. (2008) tested six alternative models to  
17 explain the temporal series of hunting bags, taking into account a climate effect, but not temporal series of  
18 predation data. This is probably due to the fact that predation data are not available at this scale. But this  
19 is precisely the point: there are no available data to check whether predation has a role in the temporal  
20 changes of bags. Insightful papers on the theoretical dynamics of hyperpredation already exist (e.g.  
21 Courchamp et al. 2000). However, the cross correlation (or second order density dependent model)  
22 described between both species is not enough to link pattern and hyperpredation caused by raptors. The  
23 RHD- mediated hyperpredation hypothesis is based on the assumption that the predator community fed  
24 differently before and after the outbreak of RHD (Moleón et al. 2008). Thus, Moleón et al. (2008)  
25 maintain that they showed the existence of hyperpredation by pointing to the decline of rabbits in the  
26 local diet of three predators, which was accompanied by a relative increase in the percentage of partridge  
27 in the diet. However, this increase was statistically significant in the case of the Golden eagle (*Aquila*  
28 *chrysaetos*), not so in the diet of Bonelli's eagle (*Hieraaetus fasciatus*), and in only one out of two

1 analysed cases of the Northern goshawk (*Accipiter gentilis*). In this way, two of the four cases analysed  
2 showed no significant increase in partridge predation and the statistical significance of the other two cases  
3 was not corrected to control for the increase in type I error due to multiple test. Thus, the evidence in  
4 favour of the suggested mechanism is weak.

5 In addition, the decline of rabbit consumption due to the decrease of rabbit populations  
6 automatically entails the increase in the percentage of occurrence of the other species in the diet, even if  
7 no absolute increase in the consumption of these species occurred. Thus, to invoke the existence of  
8 hyperpredation, the authors should demonstrate that the decrease of rabbit in the diet resulted in a  
9 differential increase in partridge consumption (in relation to population densities), rather than in a higher  
10 consumption of other prey species. This differential increase in partridge presence in the diet is certainly  
11 not occurring in the case of Bonelli's eagle (the opposite is true), is non-significant for the northern  
12 goshawk in heavily forested areas, and of moderate importance for the Golden eagle.

13 Furthermore, the revision of raptor diet studies by Moleón et al. (2008) seems to be skewed,  
14 since there are other raptor species, such as the Eagle Owl (*Bubo bubo*), that have maintained a similar  
15 diet based on rabbits before and after RHD arrival (Tella and Mañosa 1993; Serrano 2000; Martínez and  
16 Zuberogoitia 2001; Martínez and Calvo 2001). The main change detected in the diet of this species after  
17 RHD arrival is an increased predation on other diurnal and nocturnal raptors, and its effect on the avian  
18 predator community may have been relevant (Tella and Mañosa 1993; Serrano 2000). Thus, increased  
19 intraguild predation by eagle owl could have even reduced overall predation pressure in the ecosystems  
20 studied.

21

22 *Predator population dynamics data.* [Early definitions and models of hyperpredation effect \(e.g. Smith](#)  
23 [and Quin 1996, and Courchamp et al. 2000\) did not consider ~~that~~ hyperpredation processes to include](#)  
24 [indirect effects mediated by predator's functional response. These studies defined the hyperpredation](#)  
25 [process as the process of indirect effects mediated by the predator's numerical response. ~~Therefore, in~~](#)  
26 [the strictest sence. ~~being strictest this present case not should not be categorized as a hyperpredation~~](#)  
27 [process.](#) In any case, even if the number of partridges consumed by the raptors had increased, it would not  
28 be possible yet with those data to prove that partridge populations are limited by raptors, without knowing  
29 the functional response of raptors to partridge numbers, total predator abundance (or predator abundance

Con formato: Fuente:  
(Predeterminado) Times New Roman,  
10 pto

1 change), and the density dependence of productivity (Sinclair and Pech 1996; Valkama et al. 2005). Even  
2 a high percentage of gamebirds in the diet of a predator does not necessarily mean that this predator will  
3 have a negative impact on gamebird population densities (Valkama et al. 2005). The existence of  
4 limitation due to predation would require that the per capita predation rate would overcome the per capita  
5 production rate (Newton 1998), and this point is unknown in the case of red-legged partridge and raptors.  
6 Although we do not discard this possibility, the data shown by Moleón et al. (2008) do not demonstrate  
7 limitation of partridge populations by raptors.

8

9 On the contrary, it has been clearly demonstrated in the last few decades that productivity of many raptor  
10 species in Spain may have been strongly reduced by low abundance of rabbits (e.g. Martínez and  
11 Zuberogitia 2001, Margalida et al. 2007), even for raptor species considered generalist predators, such as  
12 golden eagles or Black kites (*Milvus migrans*) (Viñuela and Veiga 1992; Fernández 1993). Thus, what  
13 may have occurred during the last decades is that raptor population parameters have been negatively  
14 affected by rabbit scarcity, and raptors have also suffered increased intraguild predation (Serrano 2000,  
15 Lourenço et al. 2011).

16 *P*

17 *Predator distribution.* It appears that Moleón et al. (2008) assume that the local rabbit and partridge  
18 densities follow the same trends as those observed at national scale. These authors reviewed *Hieraaetus*  
19 *fasciatus* and *Accipiter gentilis* studies that had been conducted in Catalonia (Lleida, Tarragona,  
20 Barcelona, and Girona provinces) while the *Aquila chrysaetos* study had been conducted in Navarra  
21 province. All these provinces are located in North-eastern Spain, whereas hunting bags reflect mainly the  
22 values recorded in South-western Spain, where partridge and rabbits are hunted in higher numbers  
23 (Blanco-Aguiar et al. 2004; Virgós et al. 2007). Population patterns of both rabbits and partridges do not  
24 necessarily follow the same trends all through the Spanish territory (Blanco-Aguiar et al. 2004; Delibes-  
25 Mateos et al. 2009b), and their density show patchy patterns associated with habitat quality. In addition,  
26 molecular data show the existence in Spain of different evolutive management unit for rabbits (Carneiro  
27 et al. 2010) and red-legged partridges (Ferrero et al. 2011), thus [the assumption that both species are](#)  
28 [homogeneous throughout Spain territory might unfounded.](#) ~~consider that both species are homogeneous~~  
29 ~~along Spanish territory might be an unfounded assumption.~~

Con formato: Punto de tabulación:  
11,25 cm, Izquierda



1           It is possible to compare if partridge trends in the same provinces where raptor diet was analyzed  
2 followed the three dynamic phases from Moleón et al. (2008) predictions. Only in Navarra province both  
3 partridge populations showed a decline after RHD outbreak (Fig. 2). However, in Catalonia the red-  
4 legged partridge bag records showed no decline after the arrival of the RHD outbreak (Fig. 2). If we  
5 consider that bag records constitute an accurate source of population trends, then predictions of the model  
6 at a regional scale in three of the four diet studies (using the same data source) were not adjusted to the  
7 assumptions of the general model of hyperpredation hypothesis on partridges. Otherwise, if we consider  
8 that game releases have altered the records of hunting bags in these provinces, then it is possible that diet  
9 analyses might be also corrupted by these releases.

10

## 11 **Conclusions**

12

13 We agree that the RHD outbreak has markedly affected rabbit populations, which has indirectly affected  
14 many other species in the Iberian Peninsula (Delibes-Mateos et al. 2008a); this might include partridge  
15 populations (Blanco-Aguilar et al. 2004). However, caution must be taken before assuming that red-legged  
16 partridge decline was caused mainly by this factor, without taking into account other longer-term factors  
17 of decline. In our view, data presented in Moleón et al. (2008) do not entirely support that hyperpredation  
18 by raptors is a main process influencing partridge population trends at a national scale. First, caution must  
19 be taken when assuming bag records in recent years are a reliable estimate of wild population size  
20 because of the effect of farm-bred bird releases. On the other hand, the relations between predation rate  
21 and partridge density are unclear. A density-dependent relationship could be due to the opportunistic  
22 nature of certain predators such as red foxes (*Vulpes vulpes*; Delibes-Mateos et al. 2008b) or to the  
23 decrease in the density and/or detectability of wild rabbits (Ontiveros et al. 2005). In any case, there are  
24 not clear data supporting a major large-scale effect of predation on red-legged partridge population  
25 dynamics.

26           It is always important to evaluate alternative hypotheses, but particularly so in cases when the  
27 conclusions of a study may be used to determine management criteria for the recovery of gamebirds. If  
28 hyperpredation is offered as a significant reason for partridge limitation in Spain (which we doubt), some

1 | might advocate of control raptors populations to increase partridge populations. Hunters and game  
2 | managers in Spain tend to consider ~~ing~~ predator control as a valuable tool to increase game populations  
3 | (Angulo 2003; Delibes-Mateos et al. 2008c); in fact, the persecution of raptors has been intensified during  
4 | the last decades in Spain (Villafuerte et al. 1998). Thus, researchers must be careful before arguing that  
5 | this has been a general pattern at a national scale without serious criticism of data analyzed.

6  
7 | We suggest that anthropogenic influence (through changes in habitat quality or game  
8 | management and hunting practices, potentially due also to the decline in rabbit numbers) might explain  
9 | the patterns observed in quarry populations. However, more studies and efforts are necessary to conclude  
10 | which factors are working at this wide scale.

11  
12 | **Acknowledgements.** J.A. Blanco-Aguilar was supported by Fundação para a Ciência e a Tecnologia  
13 | (SFRH/BPD/65464/2009) grant. M. Delibes-Mateos was supported by a Juan de la Cierva research  
14 | contract awarded by the Ministerio de Ciencia e Innovación – Fondo European Social Fund. F. Casas was  
15 | supported by a post-doctoral grant of the Junta de Comunidades de Castilla la Mancha (JCCM). We thank  
16 | the comments provided by Drs. E. Fernández-Juricic M. Moleón, G. Nugent, and two anonymous  
17 | reviewers on previous versions of this manuscript.

## 18 19 | **References**

20 | Angulo E (2003) Factores que Afectan a la Distribución y Abundancia del Conejo en Andalucía. PhD  
21 | Thesis, Universidad Complutense de Madrid.

22 | Blanco-Aguilar JA, Virgós E, Villafuerte R (2004) Perdiz roja, *Alectoris rufa*. In: Madroño A, González  
23 | C, Atienza JC (eds), Libro Rojo de las Aves de España. Dirección General para la  
24 | Biodiversidad-SEO/Birdlife, pp. 182-185.

25 | Blanco-Aguilar JA (2007) Variación Espacial en la Biología de la Perdiz Roja (*Alectoris rufa*): Una  
26 | Aproximación Multidisciplinar. PhD Thesis, Universidad Complutense de Madrid.

- 1 Blanco-Aguiar JA, González-Jara P, Ferrero ME, Sánchez-Barbudo I, Virgós E, Villafuerte R, Dávila JA  
2 (2008) Assessment of game restocking contributions to anthropogenic hybridization: the case of  
3 the Iberian red-legged partridge. *Anim Conserv* 11: 535-545.
- 4 Buenestado FJ, Ferreras P, Delibes-Mateos M, Tortosa FS, Blanco-Aguiar JA, Villafuerte R (2008)  
5 Habitat selection and home range size of red-legged partridges in Spain. *Agric Ecosyst Environ*  
6 126: 158-162.
- 7 Buenestado FJ, Ferreras P, Blanco-Aguiar JA, Tortosa FS, Villafuerte R (2009) Survival and causes of  
8 mortality among wild Red-legged Partridges *Alectoris rufa* in southern Spain: Implications for  
9 conservation. *Ibis* 151: 720-730.
- 10 Carneiro M, Blanco-Aguiar JA, Villafuerte R, Ferrand N, Nachman MW (2010) Speciation in the  
11 european rabbit (*Oryctolagus cuniculus*): Islands of differentiation on the X chromosome and  
12 autosomes. *Evolution*: 64, 3443-3460
- 13 Casas F, Viñuela J (2010) Agricultural practices or game management: which is the key to improve red-  
14 legged partridge nesting success in agricultural landscapes?. *Environ Conserv* 37 : 177-186.
- 15 Cattadori I M, Haydon D T, Thirgood S J, Hudson P J (2003) Are indirect measures of abundance a  
16 useful index of population density? The case of red grouse harvesting. *Oikos* 100: 439-446.
- 17 Courchamp, F, Langlais M., Sugihara G. (2000) Rabbits killing birds: Modelling the hyperpredation  
18 process. *J Anim Ecol* 69: 154-164.
- 19 Delibes-Mateos M., Delibes M., Ferreras P., Villafuerte R. (2008a) Key role of European rabbits in the  
20 conservation of the western Mediterranean Basin hotspot. *Conserv Biol* 22: 1106-1117.
- 21 Delibes-Mateos M, Fernandez de Simón J, Villafuerte R, Ferreras P (2008b) Feeding responses of the red  
22 fox (*Vulpes vulpes*) to different wild rabbit (*Oryctolagus cuniculus*) densities: a regional  
23 approach. *Eur J Wildl Res* 54: 71-78.
- 24 Delibes-Mateos M, Ferreras P, Villafuerte R (2008c) Rabbit populations and game management: The  
25 situation after 15 years of rabbit haemorrhagic disease in central-southern Spain. *Biodivers*  
26 *Conserv* 17: 559-574.

- 1 Delibes-Mateos M, Farfán MA, Olivero J, Márquez AL, Vargas JM (2009a) Long-Term Changes in  
2 Game Species Over a Long Period of Transformation in the Iberian Mediterranean Landscape.  
3 Environ Manag 43: 1256-1268.
- 4 Delibes-Mateos M, Ferreras P, Villafuerte R (2009b) European rabbit population trends and associated  
5 factors: A review of the situation in the Iberian Peninsula. Mamm Rev 39: 124 -140.
- 6 Díaz-Sánchez S., Mateo Moriones A, Casas F. and Höfle U. (2011) Prevalence of Escherichia coli,  
7 Salmonella sp. and Campylobacter sp. in the intestinal flora of farm-reared, restocked and wild  
8 red-legged partridges (*Alectoris rufa*): is restocking using farm-reared birds a risk? Eur J Wildl  
9 Res DOI: 10.1007/s10344-011-0547-5
- 10 Fernández C (1993) Effect of the viral haemorrhagic pneumonia of the wild rabbit on the diet and  
11 breeding success of the golden eagle *Aquila chrysaetos* (L.). Rev Ecol Terre Vie 48: 323-329.
- 12 Ferreira C, Paupério J, Alves PC (2010) The usefulness of field data and hunting statistics in the  
13 assessment of wild rabbit (*Oryctolagus cuniculus*) status in Portugal. Wildl Res 37: 223–229.
- 14 Ferrero ME, Blanco-Aguilar JA , Loughheed SC , Sánchez-Barbudo I, De Nova PJG, Villafuerte R, Dávila  
15 JA (2011) Phylogeography and genetic structure of the red-legged partridge (*Alectoris rufa*):  
16 more evidence for refugia within the Iberian glacial refugium. Mol Ecol 20: 2628-2642
- 17 Fewster RM, Buckland ST, SiriwardenaGM, Baillie SR, Wilson JD (2000) Analysis of population trends  
18 for farmland birds using generalized additive models. Ecology 81: 1970-1984.
- 19 Gobierno de Navarra (2006) Memoria de Actividades 2006: Seguimiento Caza y Pesca. Departamento de  
20 Desarrollo Rural y Medio Ambiente. Gobierno de Navarra.
- 21 Gortázar C, Villafuerte R, Martín M (2000) Success of traditional restocking of red-legged partridge for  
22 hunting purposes in areas of low density of northeast Spain Aragon. Z Jagdwiss 46: 23-30.
- 23 Gregory RD, Strien AV, Vorisek P, Meyling AWG, Noble DG, Foppen RPB, Gibbons DW (2005)  
24 Developing indicators for European birds. Phil Trans R Soc B Biol Sci 360: 269-288.

- 1 Levin SA, (1992) The problem of pattern and scale in ecology. *Ecology* 73: 1943-1967.
- 2 Lourenço R, Santos SM, Rabaça JE, Penteriani V (2011) Superpredation patterns in four large European  
3 raptors. *Popul Ecol* 53:175–185
- 4 Margalida A, Gonzalez LM, Sanchez R, Oria J, Prada L, Caldera J, Aranda A, Molina JI (2007) A long-  
5 term large-scale study of the breeding biology of the spanish imperial eagle (*Aquila adalverti*). *J*  
6 *Ornithol* 148: 309-322
- 7 Martínez JA, Zuberogitia I (2001) The response of the Eagle Owl (*Bubo bubo*) to an outbreak of the  
8 rabbit haemorrhagic disease. *J Ornithol* 142: 204-211.
- 9 Martínez JE, Calvo JF (2001) Diet and breeding success of eagle owl in southeastern Spain: effect of  
10 rabbit haemorrhagic disease. *J Raptor Res* 35: 259-262.
- 11 Moleón M, Almaraz P, Sánchez-Zapata JA (2008) An emerging infectious disease triggering large-scale  
12 hyperpredation. *PLoS ONE* 3: e2307.doi:2310.1371/journal.pone.0002307.
- 13 Newton I (1998). *Population limitation in Birds*. Academic Press
- 14 Ontiveros D, Pleguezuelos JM, Caro J (2005) Prey density, prey detectability and food habits: the case of  
15 Bonelli's eagle and the conservation measures. *Biol Conserv* 123: 19-25.
- 16 REGHAB (2002) Reconciling Gamebird Hunting and Biodiversity. V Framework Program of the  
17 European Union. Proposal number: EKV-2000-00637.
- 18 SEO/BIRDLIFE (2006) Monitoring Common Breeding Birds in Spain. The SACRE Programme Report  
19 1996-2005. SEO/BIRDLIFE.
- 20 Serrano D (2000) Relationship between raptors and rabbits in the diet of Eagle Owls in southwestern  
21 Europe: Competition removal or food stress? *J Raptor Res* 34: 305-310.
- 22 Sinclair ARE, Pech RP (1996) Density dependence, stochasticity, compensation and predator regulation.  
23 *Oikos* 75: 164-173.

**Comentario [B2]:** Esto es un proyecto, no una referencia. Yo creo que deberías cambiarlo por

Arroyo, B. & Beja, P. (2002) *Impact of hunting management practices on biodiversity*. Report on Workpackage 2 of the European Project REGHAB N° EKV-2000-00637.

- 1 Smith AP, Quin DG (1996) Patterns and causes of extinction and decline in Australian conilunire rodents  
2 Biol Conser 77: 243–267.
- 3 Tella JL, Mañosa S (1993) Eagle owl predation on egyptian vulture and northern goshawk-possible effect  
4 of a decrease in european rabbit availability. J Raptor Res 27: 111-112.
- 5 Valkama J, Korpimäki E, Arroyo B, Beja P, Bretagnolle V, Bro E, Kenward R, Mañosa S, Redpath SM,  
6 Thirgood S, Viñuela J (2005) Birds of prey as limiting factors of gamebird populations in  
7 Europe: A review. Biol Rev 80: 171-203.
- 8 van Strien AJ, Pannekoek J, Gibbons DW (2001) Indexing European bird population trends using results  
9 of national monitoring schemes: a trial of a new method. Bird Study 48: 200-213.
- 10 Villafuerte R, Viñuela J, Blanco JC (1998) Extensive predator persecution caused by population crash in  
11 a game species: The case of red kites and rabbits in Spain. Biol Conserv 84: 181-188.
- 12 Villanúa D, Pérez-Rodríguez L, Casas F, Alzaga V, Acevedo P, Viñuela J, Gortázar C (2008) Sanitary  
13 risks of red-legged partridge releases: introduction of parasites. Eur J Wildl Res 54: 199–204.
- 14 Viñuela J, Veiga JP (1992) Importance of rabbits in the diet and reproductive success of black kites in  
15 southwestern Spain. Ornis Scandinavica 23: 132-138.
- 16 Virgós E, Cabezas-Díaz S, Lozano J (2007) Is the wild rabbit (*Oryctolagus cuniculus*) a threatened  
17 species in Spain? Sociological constraints in the conservation of species. Biodivers Conserv 16:  
18 3489-3504.
- 19 Wiens JA (1989) Spatial scaling in ecology. Funct Ecol 3: 385-397.

1 | **Figure 1.** Trends in numbers of red-legged partridges (*Alectoris rufa*) and rabbits (*Oryctolagus cuniculus*)  
2 | hunted per licence in Spain. The dashed line represents the year ~~in~~ which rabbit hemorrhagic disease  
3 | (RHD) arrived in Spain.


Con formato: Fuente: Cursiva

Con formato: Fuente: Cursiva

4

5 | **Figure 2.** Number of red-legged partridge and rabbit hunted per provincial licence and fitted (only for  
6 | graphical purposes) with a robust locally weighted regression. a) Barcelona. b) Girona. c) Lleida. d)  
7 | Tarragona e) Navarra. The dashed line represents the year in which rabbit hemorrhagic disease (RHD)  
8 | arrived ~~in~~ Spain.

1 Figure 1



2

3


4

5

6



1 Figure 2



2