

Irrigation of *Myrtus communis* plants with reclaimed water: morphological and physiological responses to different levels of salinity

---

By J.R. ACOSTA-MOTOS<sup>1</sup>, S. ÁLVAREZ<sup>1</sup>, J.A. HERNÁNDEZ<sup>2</sup> and M.J. SÁNCHEZ-BLANCO<sup>1,3\*</sup>

<sup>1</sup>Departamento de Riego, Centro de Edafología y Biología Aplicada del Segura (CSIC), PO. Box 164, Espinardo E-30100 Murcia, Spain

<sup>2</sup>Grupo de Biotecnología de Frutales, Departamento de Mejora Vegetal, Centro de Edafología y Biología Aplicada del Segura (CSIC), P.O. Box 164, Espinardo E-30100 Murcia, Spain

<sup>3</sup>Horticultura Sostenible en Zonas Áridas, Unidad Asociada al CSIC-CEBAS, Paseo Alfonso XIII, 52, Cartagena 30203, Spain

(e-mail: [quechu@cebas.csic.es](mailto:quechu@cebas.csic.es))

(Accepted xxxx 2014)

Running head: *Response of M. communis to reclaimed water*

\* Author for correspondence.

## SUMMARY

The influence of irrigation with different sources of reclaimed water on physiological and morphological changes in *Myrtus communis* plants was investigated to evaluate their adaptability to such conditions. *M. communis* plants, growing in a growth chamber, were subjected to four irrigation treatments over 4 months (120 d): a control [tap water, 0.8 dS m<sup>-1</sup>, leaching 10% (v/v) of the applied water] and three reclaimed water irrigation treatments: 1.5 dS m<sup>-1</sup> leaching 25% (v/v) of the applied water (RW1), 4.0 dS m<sup>-1</sup> leaching 40% (v/v) of the applied water (RW2), and 8.0 dS m<sup>-1</sup> leaching 55% (v/v) of the applied water (RW3). After this, all plants were irrigated as for the control plants for a further 2 months (60 d). At the end of the first period (4 months), none of the myrtles plants showed any adverse changes in biomass and the average total dry weight (DW) increased by 53% in treatment RW2. However, at the end of recovery period (six months), accumulations of Cl<sup>-</sup> ions and especially Na<sup>+</sup> ions negatively affected the growth of all RW3 plants. Plants irrigated with all three reclaimed water samples had greater difficulty in taking-up water from the substrate (i.e., lower leaf water potential and relative water content values). RW2 plants showed a better response in their gas exchange parameters. The use of reclaimed water decreased leaf K<sup>+</sup>/Na<sup>+</sup> and Ca<sup>2+</sup>/Na<sup>+</sup> ratios, but no chlorosis or necrosis were observed. The three reclaimed water samples had different effects on the plants depending on the specific chemical properties of the water. Leaching was found to be important to minimise the negative effects of salinity in the irrigation water.

Reclaimed water is water that has previously been used, suffered a loss in quality, but has been treated to a point where it is suitable for additional use. Use of this water in agriculture is a common practice in many areas of the World, especially in arid and semi-arid environments where access to water is a limiting factor (Yermiyahu *et al.*, 2008). Several studies have reported environmental and agronomic interest in using waste water for irrigation in different crops (Parson *et al.*, 2001; Pedrero and Alarcón, 2009; Pedrero *et al.*, 2010). Treated municipal waste water can be regarded as an alternative source of water and as fertilisation for the production of landscape plants, since it contains nutrients which can reduce the application of fertiliser, thus reducing costs and risks of environmental pollution (Gori *et al.*, 2000; Gomez-Bellot *et al.*, 2013). In spite of these potential benefits, reclaimed waste water is usually of poor quality compared to fresh water. Depending on its source and treatment, reclaimed waste water may contain high concentrations of salts, heavy metals, and/or pathogenic organisms. Nevertheless, the potential physical, chemical, or biological problems associated with the application of waste water to irrigate crops are of less concern for landscape plant production (Gori *et al.*, 2000).

A high concentration of salts in the irrigation water causes water stress due to the decrease in the water potential of the rooting medium (an osmotic effect). In addition, specific ions such as  $\text{Na}^+$  and  $\text{Cl}^-$  can accumulate in plants, where they can reach toxic levels (ion toxicity) and induce nutritional imbalances with those mineral elements that are essential for the correct functioning of the plant. In some cases, reclaimed water also contains high concentrations of boron (B; Feigin *et al.*, 1991) and significant quantities of toxic heavy metals (Barar *et al.*, 2000; Yadav *et al.*, 2002).

Salinity affects the establishment, growth, and development of plants, leading to significant losses in productivity (Giri *et al.*, 2003; Katerji *et al.*, 2003; Mathur *et al.*, 2007; Álvarez *et al.*, 2012), and may also affect the ornamental quality of both cultivated and wild

species (Morales *et al.*, 2001). In the case of landscape plants, maximum growth is not always essential and visual quality may or may not be related to biomass production and/or photosynthetic responses (Zollinger *et al.*, 2007; Álvarez *et al.*, 2011). Another way to determine the effect of salinity would be to study plant responses during a recovery period after salinity stress. Recovery from water stress is generally characterised by an increase in leaf water potential, followed by a recovery of stomatal conductance (Chaves *et al.*, 2011). However, the physiological mechanisms involved in the recovery of plants subjected to high salinity are still poorly understood.

To minimise crop losses, it is necessary to identify new irrigation management strategies such as increased leaching to maintain high and constant substrate humidity (Bañón *et al.*, 2011), or to use salt-tolerant plants, or to develop salt-tolerant crops through breeding programmes (Wu and Dodge, 2005).

*Myrtus communis* L. is a sclerophyllus evergreen shrub (Mendes *et al.*, 2001) of interest for ornamental use in re-vegetation projects in semi-arid and degraded land, and in landscaping (Romani *et al.*, 2004). Although *M. communis* is a typical Mediterranean species, with good adaptability to environmental stresses, it may, under natural conditions, suffer from abiotic stresses (Navarro *et al.*, 2009). Nevertheless, little is known about the growth and physiological responses of *M. communis* to irrigation with recycled water of different quality. Many research studies have been conducted on the effects of waste water on the physiology of ornamental species, with contradictory results, probably due to the different cultivation techniques, environments, and species used (Parnell *et al.*, 1998; Gori *et al.*, 2000; Schuch, 2005; Bañón *et al.*, 2011).

The objective of this paper was to study the negative and positive impacts that reclaimed water of different origin and composition could have on the development and quality of myrtle plants. The aim was to evaluate whether reclaimed water with a high level of

salinity could be used as an alternative source of water and nutrients for the production of *M. communis* plants. The responses of physiological parameters related to water status, photosynthetic efficiency, and nutrient content were also considered. The present study was conducted under controlled environment conditions to avoid other possible effects due to climatic variables. The information generated by this study would be valuable for both landscape and nursery irrigation management.

## MATERIALS AND METHODS

### *Plant material and experimental conditions*

Single rooted cuttings (120) of native myrtle (*Myrtus communis* L) were transplanted into 14 cm x 12 cm pots (1.2 l) filled with an 8:7:1 (v/v/v) mixture of coconut fibre, sphagnum peat, and perlite and amended at 2 g l<sup>-1</sup> substrate with Osmocote Plus (Scotts Australia Pty Ltd, The Hills Shire, New South Wales, Australia; 14:13:13 N, P, K plus microelements). The experiment was conducted in a controlled environment growth chamber, set to simulate natural conditions. The temperature in the canopy was 23°C during the 16 h photoperiod, and 18°C during darkness. Relative humidity (RH) ranged from 55 - 70%. A mean photosynthetically active radiation (PAR) level of 350  $\mu\text{mol m}^{-2} \text{s}^{-1}$  at canopy height was supplied from 08.00-00.00. Although the level of radiation in the growth chamber was lower in the field, we assumed that the PAR level used was of secondary importance compared with the different irrigation treatments.

### *Treatments*

*M. communis* plants (n = 30 per treatment) were exposed to four irrigation treatments for 4 months (120 d; Period I) using water from different sources. The irrigation treatments consisted of a control, where the electrical conductivity (EC) of the tap water was 0.8 dS m<sup>-1</sup>

(indicating no use-restrictions or only slight restrictions according to FAO classifications; FAO, 2003) and three reclaimed water treatments. The latter used water from three sewage treatment plants located in the Province of Murcia (Spain), namely: RW1 (EC 1.7 dS m<sup>-1</sup>) from Jumilla, RW2 (4.0 dS m<sup>-1</sup>) from Campotejar and RW3 (8.0 dS m<sup>-1</sup>) from Mazarrón. FAO classifications indicated severe restrictions on the use of the latter two types of water. All three waste water treatment plants applied a conventional activated-sludge process, followed by ultraviolet radiation as the tertiary treatment. At the start of the experimental period the concentrations of Na<sup>+</sup>, Cl<sup>-</sup> and B<sup>3+</sup> ions in each irrigation water were analysed. The results are shown in Table I.

After 4 months (120 d; Period I), all plants were exposed to a 2-month (60 d) recovery period (Period II) in which the plants were irrigated with the same tap water used for the control plants. Throughout the 6 months (180 d) of the experiment, all plants were irrigated twice a week to above-container capacity. To determine the maximum water-holding capacity of the substrate, medium was uniformly mixed and packed to a bulk density of 0.165 g cm<sup>-3</sup> in all pots. Each substrate surface was covered with aluminium foil to prevent water evaporation and the lower part of each pot was submerged to half its height in a water bath, then left to equilibrate overnight. The next day, the pots were removed and left to drain freely until the drainage became negligible. The fresh weight was then recorded and calculated for each individual pot and considered as the weight at field capacity (WFC). The volume of irrigation water applied to be applied was determined for each treatment as the point at which the leaching fraction reached 10% (v/v) of the water applied in the control treatment, 25% in RW1, 40% in RW2, or 55% of the applied water in RW3. Each plant was weighed before each irrigation event and the volume of irrigation water required to refill the pot to its threshold level (i.e., its WFC plus its pre-determined level of leaching, depending on treatment) was calculated and added to each plant.

### *Growth and colour measurements*

At the ends of Period I and Period II, the substrate was gently washed from the roots of eight plants per treatment and each plant was divided into leaves, stems and roots. These were oven-dried at 80°C until they reached a constant weight to measure their respective dry weights (DW). Leaf numbers and leaf areas (cm<sup>2</sup>) were determined for the same plants before drying, using a leaf area meter (AM 200; ADC BioScientific Ltd., Hoddesdon, UK). The root:shoot DW ratio was determined for each plant by dividing root DW by leaf DW.

At the ends of Period I and Period II, plant heights were measured for 20 plants per treatment and leaf colour and relative chlorophyll concentration (RCC) were measured at the mid-point of a mature leaf using three leaves from each plant and six plants per treatment. Plant height was taken as the vertical distance from the surface of the substrate to the node of the highest leaf. Leaf colour was measured using a CR-10 colorimeter (Konica Minolta Sensing Inc., Osaka, Japan), which provided values for the colour co-ordinates lightness (L\*), chroma (C\*), and hue angle (h°; McGuire, 1992). RCC was estimated using a SPAD-502 chlorophyll meter (Konica Minolta Sensing Inc., Osaka, Japan).

### *Plant water relations and gas exchange*

At the ends of Period I and Period II, changes in leaf water potential ( $\Psi_1$ ), relative water content (RWC), stomatal conductance ( $g_s$ ) and the net rate of photosynthesis ( $P_n$ ) were determined in six plants per treatment midway through the photoperiod.  $\Psi_1$  was estimated according to Scholander et al. (1965), using a pressure chamber (Model 3000; Soil Moisture Equipment Co., Santa Barbara, CA, USA). Each leaf was placed in the chamber within 20 s of collection and pressurised at a rate of 0.02 MPa s<sup>-1</sup> (Turner, 1988). The RWC of leaves was

measured according to Barrs (1968).  $g_s$  and  $P_n$  were determined in attached leaves using a gas exchange system (LI-6400; LI-COR Inc., Lincoln, NE, USA).

#### *Mineral concentrations and water and substrate analyses*

At the end of the salinity treatment and recovery periods (Period I and Period II), eight plants per treatment were separated into leaves, stems, and roots, washed with distilled water, dried at 70°C, and stored at room temperature for inorganic solute analyses. The concentrations of  $Cl^-$  ions were measured using a chloride analyser (Model 926; Sherwood Scientific Ltd., Cambridge, UK) in aqueous extracts obtained by mixing 100 mg DW of each tissue sample with 40 ml of water, shaking for 30 min, and filtering. The concentrations of  $Na^+$ ,  $B^{3+}$ ,  $K^+$ , and  $Ca^{2+}$  ions were determined by digesting 100 mg of tissue powder with 50 ml of a 2:1 (v/v) mix of 14M  $HNO_3$ :12M  $HClO_4$  and using an inductively-coupled plasma-optical emission spectrometer (ICP-OES, IRIS Intrepid II XDL, Thermo Fisher Scientific Inc., Loughborough, UK).

The inorganic solute concentrations and EC values of each irrigation water sample were measured at the start of the experiment by collecting 100 ml in glass bottles and storing them at 5°C until being processed for chemical analyses. EC values were measured using a multirange, Cryson-HI8734 conductivity meter (Crisom Instruments S.A., Barcelona, Spain).  $Na^+$  and  $B^{3+}$  ion concentrations were determined using an ICP-OES (IRIS Intrepid II XDL, Thermo Fisher Scientific Inc.) and  $Cl^-$  ion concentrations were measured using a Metrohm Chromatograph (Metrohm Ltd., Herisau, Switzerland).

Eight samples of the substrate were collected per treatment and sent for analysis to an external laboratory (Antonio Abellán Caravaca S.L., Murcia, Spain) at the ends of Period I and Period II. The substrate was dried at room temperature for 1 week.  $Na^+$ ,  $Ca^{2+}$ , and  $Mg^{2+}$  ion concentrations were then determined using an ICP-OES (IRIS Intrepid II XDL, Thermo



Fisher Scientific Inc.) in a saturated soil extract.  $\text{Cl}^-$  ion concentrations were measured by chromatography. EC was measured in a saturated soil paste using a Cryson-HI8734 conductivity meter (Crisom Instruments S.A.).

#### *Statistical analysis of the data*

Thirty plants were attributed at random to each of the four treatments. The data were analysed by one-way ANOVA using Statgraphics Plus for Windows 5.1 software (Manugistics Ltd., Rockville, MD, USA). Root:shoot ratio data were subjected to an arcsine square-root transformation before statistical analysis to ensure homogeneity of variance. Treatment means were separated using Duncan's Multiple Range Test at  $P \leq 0.05$ .

## RESULTS

The EC of the substrate at the end of Period I increased in line with the increase EC of the irrigation water applied due to the accumulation of  $\text{Cl}^-$  and  $\text{Na}^+$  ions, although no significant differences were observed between RW2 and RW3 (Table II). The latter treatments also gave the highest concentrations of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  ions, especially RW2. After irrigating with reclaimed water, all plants were irrigated with low conductivity ( $0.8 \text{ dS m}^{-1}$ ) tap water in Period II. At the end of Period II, although the EC of the substrate was similar in all treatments, substrate  $\text{Na}^+$  ion concentrations were higher in RW2 and RW3. In general, at the end of Period II, a greater accumulation of salts was observed in the substrate of the control treatment than at the end of Period I. However,  $\text{Na}^+$  and  $\text{Cl}^-$  ions decreased in the substrate of RW2 and RW3 treatments compared to the values recorded after Period I.

At the end of Period I, the growth of all 120 myrtle plants showed no adverse changes after the four irrigation treatments (Table III). Surprisingly, total DWs were higher in plants subjected to RW2 than in control plants. This was due to an increase in the biomass in all

parts of the plant, up to 38% in leaves, 56% in stems and 69% in roots. Leaf areas and numbers of leaves per plant were also significantly higher in RW2 plants compared to the other treatments. Growth parameters of the aerial parts of plants irrigated with RW3 showed no significant changes compared with the control plants, although root DWs increased. Root : shoot DW ratios were higher in RW3 plants, which were shorter, than in plants from the control treatment (Table III). At the end of Period II, when all plants had been watered with the same water as was used for the control plants (Table III), RW2 plants had the highest values for all growth parameters studied, although the differences in leaf and stem DWs ,leaf numbers and leaf areas compared with the controls were not significant. After the recovery period (Period II), plants that had been irrigated at highest salinity level (RW3) had lower shoot DWs and lower leaf area than the control plants, and again had the highest root:shoot ratios. As regards plant height, the differences between control and RW3 plants observed after Period I were maintained at the end of the Period II (180 d; Table III).

At the end of Period I, leaf water potential values ( $\Psi_l$ ) became more negative as the level of salinity increased. Thus, RW3 plants had the lowest values (- 1.0 MPa) and control plants had the highest values (-0.6 MPa), with intermediate values for plants irrigated with RW1 or RW2 (Figure 1A). Relative water content values (RWC) showed a similar behaviour to that observed for  $\Psi_l$ , with RW3 plants having the lowest values (82%; Figure1B). However, the corresponding values for the plants irrigated with RW2 were slightly higher than those shown by plants treated with RW1. At the end of Period II, plants from the most saline treatment (RW3) did not reach the RWC values recorded for the other treatments (Figure 1B). At the end of Period I, lower stomata conductance ( $g_s$ ) and net photosynthesis ( $P_n$ ) values were observed in all 90 plants irrigated with waste water compared to the 30 control plants (Figure 1C, D). When RW1 and RW2 plants were compared, gas exchange

values ( $g_s$  and  $P_n$ ) were higher after the RW2 treatment. At the end of Period II, myrtle plants from the RW2 treatment had similar  $P_n$  values to controls plants (Figure 1D).

Relative chlorophyll concentration values (RCC) did not change at any point during the experiment in any of the treatments studied (Figure 2A). In contrast, leaf colour parameters ( $L^*$ ,  $C^*$ , and  $h^\circ$ ) were affected by the different irrigation treatments (Figure 2B-D). At the end of the recovery period (Period II), RW2 and RW3 plants had similar  $L^*$  and  $C^*$  values, which were lower than those in control plants and RW1, while their  $h^\circ$  values were higher (Figure 2B, C). The higher  $h^\circ$  values and lower  $L^*$  and  $C^*$  values recorded for the leaves of RW2 and RW3 plants confirmed the visibly darker and less-vivid green colour of the foliage compared to the control plants.

At the end of Period I, control and RW1 plants had similar  $Cl^-$ ,  $Na^+$ ,  $B^{3+}$ ,  $K^+$  and  $Ca^{2+}$  ion concentrations in their leaves, stems and roots (Table IV).  $Cl^-$  ion concentrations at the end of Period I in all parts of the plant were similar for RW2 and RW3 and were significantly higher in the RW3 treatment than in the controls.  $Na^+$  ion concentrations in leaves were higher in RW2 than in control plants, while the highest  $Na^+$  concentrations in all parts of the plant were found using RW3. Boron accumulation in leaves was higher in plants irrigated with all three reclaimed water samples. Similarly,  $B^{3+}$  ions accumulated in the stems of RW2 plants and in the roots of RW3 plants. RW2 plants had the highest  $K^+$  ion concentrations and RW3 plants had the lowest values. However, the highest  $Ca^{2+}$  ion concentrations were found in the leaves and stems of plants irrigated with RW2 or RW3.

At the end of Period II, plants from all treatments had similar  $Cl^-$  ion concentrations in all parts (leaves, stems and root). RW1 and RW3 plants had higher  $Na^+$  ion concentrations in their leaves and roots than in the controls and RW2 plants.  $B^{3+}$  ion concentrations were higher than in the controls in the three reclaimed water irrigation treatments, especially RW1 and

RW3. In general, the  $K^+$  ion concentrations were lower in plants watered with reclaimed water than in control plants. RW3 plants had the lowest  $K^+$  ion concentrations.

Leaf  $K^+/Na^+$  and  $Ca^{2+}/Na^+$  ion ratios at the end of Period I were lower in plants irrigated with all three reclaimed water samples, especially RW3, which produced the lowest values (Table V). At the end of Period II, the highest  $K^+/Na^+$  and  $Ca^{2+}/Na^+$  ion ratios in leaves were found in RW1 and RW3, while the values measured in RW2 plants did not differ from those in the controls.

## DISCUSSION

Treated waste waters have a variable salt content that depends on their origin, which makes their use problematic when irrigation strategies are unsuitable. One important aspect when using low quality reclaimed water is the technique used for plant culture, which affects the development and agronomic performance of the crop (Bañón *et al.*, 2012). For example, regulating the drainage is considered to be a valid tool to reduce the problems associated with salinity. The lower the quality of the water, the higher the drainage necessary to prevent the accumulation of salts in the substrate (Evans, 2004). In this experiment, adjusting the amount of drainage according to the EC of the irrigation water applied reduced the toxic negative effects of the salts. Even after 4 months (120 d) of applying reclaimed waste water with EC values of 1.5-8.0 dS  $m^{-1}$ , no reduction in growth parameters was observed in the myrtle plants. Moreover, using RW2 resulted in higher shoot and root DWs than the other three treatments.

One possible advantage of using reclaimed waste water can be the composition of the water, which often has higher organic matter and nutrient contents than fresh water (Janssen *et al.*, 2005). However, it is important to know the concentrations of solutes in the irrigation water, since high concentrations of  $Na^+$  and  $Cl^-$  ions may be offset by the beneficial effects of

other solutes such as  $Mg^{2+}$ ,  $K^+$ ,  $PO_4^{3-}$ ,  $Ca^{2+}$  ions. Analysis of the treated waste waters used here identified high levels of these elements, meaning that their concentrations in the myrtle plants were not diminished by the effect of NaCl, and were even increased, as in the case of P (data not shown; Gómez-Bellot *et al.*, 2013). The highest levels of  $Ca^{2+}$  and  $Mg^{2+}$  ions were found in the substrate of the RW2 treatment (Table II).

Some differences in plant growth parameters were observed between the different treatments after the recovery period (day-180). For example, plants irrigated with RW3 treatment had lower biomass and leaf areas, suggesting that, although they were irrigated with good quality water during the recovery Period II, the accumulation of toxic ions such as  $Na^+$  and  $B^{3+}$  had a negative effect on plant growth. This did not occur in RW2-irrigated plants, which generally had a higher biomass than RW1 plants.

The low  $\Psi_1$  and RWC values of plants irrigated with RW3 (the highest salinity) reflect the increased difficulty for plants to take-up water from the substrate due to the high accumulation of salts (Álvarez *et al.*, 2012). Despite the availability of water in the substrate, the osmotic effect of the salts in the root zone limit the absorption of water (Hardikar and Pandey, 2008), as reflected in the water status of the plants (Figure 1). This behaviour has been observed in other ornamental species grown under similar conditions (Navarro *et al.*, 2007; Miralles *et al.*, 2011). However, the most significant response was the decrease in  $g_s$  values in all plants treated with reclaimed water, which acted as a mechanism to prevent excessive loss of water by transpiration (Muns and Tester, 2008; Figure 1C).  $P_n$  values were also affected. The highest  $P_n$  values among the reclaimed water treatments were observed in RW2 plants, which correlated with their higher DW, increased leaf area, and greater numbers of leaves (Table III). Although recovery after a period of salinity was characterized by an increase in leaf water parameter values (Chaves *et al.*, 2009), this was not observed in plants irrigated with RW3.

In many studies, the effects of salinity on  $P_n$  and  $g_s$  have been shown to depend on species, salinity level and the duration of the saline stress imposed (Tattini *et al.*, 2002; Álvarez and Sánchez-Blanco, 2014). Another parameter used to detect differences in the salt-tolerance of different species used for landscaping is RCC. In some species, it has been observed that reductions in leaf RCC values due to high salt levels reflect a low degree of stress tolerance (Cabrera, 2003). However, under our conditions, RCC values did not change significantly in the four different treatments applied.

Aesthetic value is an important trait in ornamental plants and an absence of visible leaf damage such as chlorosis, necrosis, or premature leaf drop is critical to the evaluation of plant quality. None of these symptoms were observed in our experiments. Controlled environmental conditions (i.e., light, temperature and humidity) and irrigation practices can affect plant responses (Fox *et al.*, 2005). The inhibition of photosynthesis observed at the end of Period II (day-180) led to a reduction of photo-assimilates and less dry matter production in RW3 plants (i.e., the lowest total DW, stem DW and height). This could be related to higher concentrations of toxic ions, especially  $Na^+$  and  $B^{3+}$ , in the leaves of RW3 plants compared with the other treatments (Álvarez *et al.*, 2012). Plants in the RW2 treatment had similar  $Cl^-$  ion concentrations to those in the controls and were the least affected. Thus,  $Cl^-$  ion concentrations were similar in all treatments, whereas  $Na^+$  and  $B^{3+}$  ions accumulated more in RW3 plants compared to the controls.

High concentrations of  $B^{3+}$  ions are another problem associated with the use of reclaimed water, and high levels of  $B^{3+}$  ions were observed in leaves and roots, especially in RW3 plants. However, no  $B^{3+}$ -related toxicity symptoms were observed, perhaps because the higher concentrations of  $Na^+$  ions interfered with the absorption of  $B^{3+}$  ions (El-Motaium *et al.*, 1994; Edelstein *et al.*, 2005). Moreover, symptoms associated with the accumulation of  $Na^+$  and  $Cl^-$  ions may mitigate the damage typically caused by excess  $B^{3+}$  ions (Bañón *et al.*,

2012). While salinity has been seen to aggravate B toxicity symptoms in wheat plants (Wimmer *et al.*, 2003), it has also been reported that the addition of B to the nutrient solution can prevent the reduction of NaCl-induced plant growth in pea plants (El-Handaui *et al.*, 2003).

High levels of salinity reduce the absorption of  $K^+$  and  $Ca^{2+}$  ions in many species (Niu *et al.*, 1995; Chaparzadeh *et al.*, 2003). In our study, the  $K^+/Na^+$  and  $Ca^{2+}/Na^+$  ion ratios decreased in the leaves of all plants irrigated with all reclaimed waters, but to the greatest extent with RW3. RW2 plants showed similar ion ratios to the controls after 60 d of irrigation with low EC, tap water (the recovery phase). Many species exhibit some degree of tolerance to salinity (Heuer and Ravina, 2004), which appears to be related to a higher selective uptake of  $K^+$  ions than  $Na^+$  ions (Heuer and Ravina, 2004; Colmer *et al.*, 2006). The severe reduction in growth, even at relatively low salt levels (2.0 to 3.0  $dSm^{-1}$ ), has been attributed to increases in  $Na^+$  and  $Cl^-$  ions, accompanied by a major reduction in  $Ca^{2+}$  and  $K^+$  ion concentrations in plant tissues (Valdez-Aguilar *et al.*, 2009). However, these effects did not occur in our study. In addition, plants irrigated with reclaimed water showed relatively high  $K^+/Na^+$  and  $Ca^{2+}/Na^+$  ion ratios, especially in leaves, which correlated with their response to salinity. In this sense,  $K^+$  and  $Ca^{2+}$  ions not only play important roles in plant growth and development, but are also vital for osmotic adjustment and the maintenance of cell turgor (Osakabe *et al.*, 2014).

In conclusion, the three reclaimed water samples had different effects on *M. communis* plants, depending on their chemical properties. This was more evident in the ability of RW-treated plants to recover from salinity. Reclaimed water of moderate conductivity (EC = 4.0  $dS m^{-1}$ ; RW2) was able to maintain the quality of the ornamental plants and could be regarded as safe for a nutrient management strategy. None of the problems associated with reclaimed water, such as salinity, were seen in RW2 treatment. However, *M. communis* plants irrigated with reclaimed water of high EC (RW3; 8.0  $dS m^{-1}$ ) were stunted and showed reductions in

their gas exchange parameters, which did not recover after a 2 month period of irrigating with low EC water. EC values, the different salts present in the irrigation water, and the extent of leaching fraction, must all be considered when using reclaimed water for irrigating purposes.

This work was supported by the Spanish Ministry of Science and Innovation (Project No. AGL 2011-30022-C02-01-02) and Fundación Séneca (Project No.15356/PI/10).

## REFERENCES

- ÁLVAREZ, S. and SÁNCHEZ-BLANCO, M.J. (2014). Long-term effect of salinity on plant quality, water relations, photosynthetic parameters and ion distribution in *Callistemon citrinus*. *Plant Biology*, **16**, doi:10.1111/plb.12106.
- ÁLVAREZ, S., NAVARRO, A., NICOLÁS, E. and SÁNCHEZ-BLANCO, M.J. (2011). Transpiration, photosynthetic responses, tissue water relations and dry mass partitioning in *Callistemon* plants during drought conditions. *Scientia Horticulturae*, **129**, 306-312.
- ÁLVAREZ, S., GÓMEZ-BELLOT, M.J., CASTILLO, M., BAÑÓN, S. and SÁNCHEZ-BLANCO, M.J. (2012). Osmotic and saline effect on growth, water relations, and ion uptake and translocation in *Phlomis purpurea* plants. *Environmental and Experimental Botany*, **78**, 138- 145.
- BAÑÓN, S., MIRALLES, J., OCHOA, J., FRANCO, J.A. and SÁNCHEZ-BLANCO, M.J. (2011). Effects of diluted and undiluted treated wastewater on the growth, physiological aspects and visual quality of potted lantana and polygala plants. *Scientia Horticulturae*, **129**, 869- 876.
- BAÑÓN, S., MIRALLES, J., OCHOA, J. and SÁNCHEZ-BLANCO, M.J. (2012). The effect of salinity and high boron on growth, photosynthetic activity and mineral contents of two ornamental shrubs. *Horticultural Science*, **39**, 188-194.



- BARAR, M.S., MAHLI, S.S., SINGH, A.P., AROROA, C.L. and GILL, K.S. (2000). Sewer water irrigation effects on some potentially toxic trace elements in soil and potato plants in north-western India. *Canadian Journal of Soil Science*, **80**, 465-471.
- BARRS, H.D. (1968). *Determination of Water Deficit in Plant Tissues. Water Deficits and Plant Growth*, Volume 1. Academic Press, New Delhi, India. 268 pp.
- CABRERA, RI. (2003). Growth, quality and nutrient responses of azalea hybrids to salinity. *Acta Horticulturae*, **609**, 241-245.
- CHAPARZADEH, N., KHAVARI- NEJAD, R.A., NAVARI –IZZO, F. and IZZO, R. (2003). Water relations and ionic balance in *Calendula officinalis* L. under saline conditions. *Agrochimica*, **XLVII (1-2)**, 69-79.
- CHAVES, M.M., FLEXAS, J. and PINHEIRO, C. (2009). Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. *Annals of Botany*, **103**, 551-560.
- CHAVES, M.M., COSTA, M.J. and MADEIRA SAIBO, N.J. (2011). Recent advances in photosynthesis under drought and salinity. *Advances in Botanical Research*, **57**, 49–104.
- COLMER, T.D., FLOWERS, T.J. and MUNNS, R. (2006). Use of wild relatives to improve salt tolerance in wheat. *Journal of Experimental Botany*, **57**, 1059–1078.
- EDELSTEIN, M., BEN-HUR, M., COHEN, R., BURGER, Y. and RAVINA, I. (2005). Boron and salinity effects on grafted and non-grafted melon plants. *Plant and Soil*, **269**, 273-284.
- EL-HANDAUI, A., REDONDO-NIETO, M., TORRALBA, B., RIVILLA, R., BONILA, I. and BOLAÑOS, L. (2003). Influence of boron and calcium on the tolerance to salinity of nitrogen-fixing pea plants. *Plant and Soil*, **251**, 93–103.

- EL-MOTAUM, R., HU, H. and BROWN, P.H. (1994). The relative tolerance of six *Prunus* rootstocks to boron and salinity. *Journal of the American Society for Horticultural Science*, **119**, 1169-1175.
- EVANS, R. (2004). Hands-on irrigation training for nursery growers. *Growing Points*, **8**, 4-5.
- FAO (Food and Agriculture Organization of the United Nations). (2003). *Review of World Water Resources by Country*. Water Reports 23, FAO, Rome, Italy, 110 pp.
- FEIGIN, A., RAVINA, I. and SHALHEVET, J. (1991). *Irrigation with Treated Sewage Effluent*. Springer-Verlag. Berlin, Germany. 224 pp.
- FOX, L.J., GROSE, N., APPLETON, B.L. and DONOHUE, S.J. (2005). Evaluation of treated effluent as an irrigation source for landscape plants. *Journal of Environmental Horticulture*, **23**, 174-178.
- GIRI, B., KAPOOR, R. and MUKERJI, K.G. (2003). Influence of arbuscular mycorrhizal fungi and salinity on growth, biomass and mineral nutrition of *Acacia auriculiformis*. *Biology and Fertility of Soils*, **38**, 170-175.
- GÓMEZ-BELLOT, M.J., ÁLVAREZ, S., CASTILLO, M., BAÑÓN, S., ORTUÑO, M.F. and SÁNCHEZ-BLANCO, M.J. (2013). Water relations, nutrient content and developmental responses of *Euonymus* plants irrigated with water of different degrees of salinity and quality. *Journal of Plant Research*, **126**, 567-576. DOI 10.1007/s10265-012-0545-z.
- GORI, R., FERRINI, F., NICESE, F.P. and LUBELLO, C. (2000). Effect of reclaimed wastewater on the growth and nutrient content of three landscape shrubs. *Journal of Environmental Horticulture*, **18**, 108-114.
- HARDIKAR, S.A. and PANDEY, A.N. (2008). Growth, water status and nutrient accumulation of seedling of *Acacia senegal* (L.) Willd. in response to soil salinity. *Anales de Biología*, **30**, 17-28.

- HEUER, B. and RAVINA, I. (2004). Growth and development of stock [*Matthiola incana* (L.) R. Brown] under salinity. *Australian Journal of Agricultural Research*, **55**, 907-910.
- JANSSEN, B.H., BOESVELD, H. and JUSTO-RODRIGUEZ, M. (2005). Some theoretical considerations on evaluating wastewater as a source of N, P and K for crops. *Irrigation and Drainage*, **54**, S35-S47.
- KATERJI, N., VAN HOORN, J.W., HAMDY, A. and MASTRORILLI, M. (2003). Salinity effect on crop development and yield, analysis of salt tolerance according to several classification methods. *Agricultural Water Management*, **62**, 37-66.
- MATHUR, N., SINGH, J., BOHRA, S. and VYAS, A. (2007). Arbuscular mycorrhizal status of medicinal halophytes in saline areas of Indian Thar Desert. *International Journal of Soil Science*, **2**, 119-127.
- McGUIRE, R.G. (1992). Reporting of objective colour measurements. *HortScience*, **27**, 1254-1255.
- MENDES, M.M., GAZARINI, L.C. and RODRIGUES, M.L. (2001). Acclimation of *Myrtus communis* to contrasting Mediterranean light environments - effects on structure and chemical composition of foliage and plant water relations. *Environmental and Experimental Botany*, **45**, 165-178.
- MIRALLES, J., VÁLDÉS, R., FRANCO, J.A., SÁNCHEZ-BLANCO, M.J. and BAÑÓN, S. (2011). Irrigation of hydrangea with saline reclaimed wastewater: effects of fresh water flushing. *Acta Horticulturae*, **1000**, 229-236.
- MORALES, M.A., OLMOS, E., TORRECILLAS, A., SÁNCHEZ-BLANCO, M.J. and ALARCÓN, J.J. (2001). Differences in water relations, leaf ion accumulation and excretion rates between cultivated and wild species of *Limonium* sp. grown in conditions of saline stress. *Flora*, **196**, 345-352.

- MUNNS, R. and TESTER, M. (2008). Mechanisms of salinity tolerance. *Annual Review of Plant Biology*, **59**, 651-681.
- NAVARRO, A., BAÑÓN, S., OLMOS, E. and SÁNCHEZ-BLANCO, M.J. (2007). Effects of sodium chloride on water potential components, hydraulic conductivity, gas exchange and leaf ultrastructure of *Arbutus unedo* plants. *Plant Science*, **172**, 473- 480.
- NAVARRO, A., ÁLVAREZ, S., CASTILLO, M., BAÑÓN, S. and SÁNCHEZ-BLANCO, M.J. (2009). Changes in tissue-water relations, photosynthetic activity, and growth of *Myrtus communis* plants in response to different conditions of water availability. *Journal of Horticultural Science & Biotechnology*, **84**, 541-547.
- NIU, X., BRESSAN, R.A. HASEGAWA, P.M. and PARDO, J.M. (1995). Ion homeostasis in NaCl stress environments. *Plant Physiology*, **109**, 735-742.
- OSAKABE, Y., YAMAGUCHI-SHINOZAKI, K., SHINOZAKI, K. and PHAN TRAN, L.S. (2014). ABA control of plant macroelement membrane transport systems in response to water deficit and high salinity. *The New Phytologist*, **202**, 35-49.
- PARNELL, J.R. (1998). *Project Greenleaf Final Report*. Public Utilities Department. City of St. Petersburg, FL, USA. 500 pp
- PARSON, L.R., WHEATON, T.A. and CASTLE, W.S. (2001). High application rates of reclaimed water benefit citrus tree growth and fruit production. *HortScience*, **36**, 1273-1277.
- PEDRERO, F. and ALARCÓN, J.J. (2009). Effects of treated wastewater irrigation on lemon trees. *Desalination*, **246**, 631-639.
- PEDRERO, F., KALAVROUZOTIS, I., ALARCÓN, J.J., KOUKOULAKIS, P. and ASANO T. (2010). Use of treated municipal wastewater in irrigated agriculture - Review of some practices in Spain and Greece. *Agricultural Water Management*, **97**, 1233-1241.

- ROMANI, A., COINU, R., CARTA, S., PINELLI, P., GALARDI, C., VINCIERI, F.F. and FRANCONI, F. (2004). Evaluation of antioxidant effect of different extracts of *Myrtus communis* L. *Free Radical Research*, **38**, 97-103.
- SCHOLANDER, P.F., HAMMEL, H.T., BRADSTREET, E.D. and HEMMINGSEN, E.A. (1965). Sap pressure in vascular plants. *Science*, **148**, 339-346.
- SCHUCH, U. (2005). Effect of reclaimed water and drought on salt-sensitive perennials. *HortScience*, **40**, 1095.
- TATTINI, M., MONTAGNI, G. and TRAVERSI, M.L. (2002). Gas exchange, water relations and osmotic adjustment in *Phillyrea latifolia* grown at various salinity concentrations. *Tree Physiology*, **22**, 403-412.
- TURNER, N.C. (1988). Measurement of plant water status by the pressure chamber technique. *Irrigation Science*, **9**, 289-308.
- VALDEZ-AGUILAR, L.A., GRIEVE, C.M., POSS, J. and MELLANO, M.A. (2009). Hypersensitivity of *Ranunculus asiaticus* to salinity and alkalinity in irrigation water in sand cultures. *HortScience*, **44**, 138–144.
- WIMMER, M. A., MÜHLING, K. H., LÄUCHLI, A., BROWN, P. H. and GOLDBACH, H. E. (2003). The interaction between salinity and boron toxicity affects the subcellular distribution of ions and proteins in wheat leaves. *Plant, Cell and Environment*, **26**, 1267–1274.
- WU, L. and DODGE, L. (2005). *Landscape Plant Salt Tolerance Selection Guide for Recycled Water Irrigation. A Special Report for the Elvenia J. Slosson Endowment Fund*. University of California, Davis. CA, USA. 40 pp.
- YADAV, R.K., GOYAL, B., SHARMA, R.K., DUBEY, S.K. and MINHAS, P.S. (2002). Post-irrigation impact of domestic sewage effluent on composition of soils, crops and ground water - a case study. *Environment International*, **28**, 481-486.

YERMIYAHU, U., BEN-GAL, A., KEREN, R. and REID, R.J. (2008). Combined effect of salinity and excess boron on plant growth and yield. *Plant and Soil*, **304**, 73-87.

ZOLLINGER, N., KJELGREN, R., CERNY-KOENIG, T., KOPP, K. and KOENIG, R. (2006). Drought responses of six ornamental herbaceous perennials. *Scientia Horticulturae*, **109**, 267-274.

TABLE I

*Chemical analyses of the waters samples used for the different irrigation treatments.*

Ion	Control <sup>‡</sup>	RW1 <sup>§</sup>	RW2 <sup>§</sup>	RW3 <sup>§</sup>
Na <sup>+</sup> (mg l <sup>-1</sup> )	140	260	362	1,492
Cl <sup>-</sup> (mg l <sup>-1</sup> )	184	720	862	1,557
B <sup>3+</sup> (mg l <sup>-1</sup> )	0.13	0.18	0.55	1.26

<sup>‡</sup>Control, tap water 0.8 dS m<sup>-1</sup>.

<sup>§</sup>RW1, reclaimed water irrigation treatment: at 1.5 dS m<sup>-1</sup> leaching 25% (v/v) of the applied water; RW2, reclaimed water irrigation treatment: at 4.0 dS m<sup>-1</sup> leaching 40% (v/v) of the applied water; RW3, reclaimed water irrigation treatment: at 8.0 dS m<sup>-1</sup> leaching 55% (v/v) of the applied water.

TABLE II

*Influence of four irrigation treatments on the physico-chemical properties of the substrate collected from M. communis plants at the end of*

*Period I and Period II<sup>‡</sup>*

Period	Parameter	Control	RW1 <sup>§</sup>	RW2 <sup>§</sup>	RW3 <sup>§</sup>
I (4 months; 120 d)	EC (dS m <sup>-1</sup> )	1.83 ± 0.22a <sup>†</sup>	3.30 ± 0.33b	8.57 ± 0.38c	8.33 ± 0.23c
	Cl <sup>-</sup> (mmol kg <sup>-1</sup> DW)	7.53 ± 1.11a	18.11 ± 1.84b	55.65 ± 2.92c	63.44 ± 2.17d
	Ca <sup>2+</sup> (mmol kg <sup>-1</sup> DW)	1.03 ± 0.09a	1.50 ± 0.13a	6.26 ± 0.40c	3.90 ± 0.28b
	Mg <sup>2+</sup> (mmol kg <sup>-1</sup> DW)	0.83 ± 0.11a	1.34 ± 0.16a	8.99 ± 0.43c	6.59 ± 0.23b
	Na <sup>+</sup> (mmol kg <sup>-1</sup> DW)	9.83 ± 1.21a	21.61 ± 2.12b	59.13 ± 2.76c	63.57 ± 1.75c
II (4+2 months; 180d)	EC (dS m <sup>-1</sup> )	3.21 ± 0.28a	4.08 ± 0.34a	4.32 ± 0.37a	4.16 ± 0.24a
	Cl <sup>-</sup> (mmol kg <sup>-1</sup> DW)	11.95 ± 1.41a	16.31 ± 1.62ab	19.50 ± 2.80b	20.38 ± 1.37b
	Ca <sup>2+</sup> (mmol kg <sup>-1</sup> DW)	1.61 ± 0.19a	1.96 ± 0.16a	2.51 ± 0.24b	1.84 ± 0.12a
	Mg <sup>2+</sup> (mmol kg <sup>-1</sup> DW)	1.41 ± 0.18a	1.60 ± 0.15a	3.03 ± 0.32b	2.51 ± 0.24b
	Na <sup>+</sup> (mmol kg <sup>-1</sup> DW)	20.87 ± 1.73a	27.85 ± 2.32b	31.46 ± 2.36bc	34.84 ± 2.05c

<sup>‡</sup>Period I, a 4-month (120 d) period with control or waste water irrigation; Period II, a 2-month (60 d) period of recovery with control (low EC)

tap water irrigation after Period I.

<sup>§</sup>RW1, reclaimed water irrigation treatment: at 1.5 dS m<sup>-1</sup> leaching 25% (v/v) of the applied water; RW2, reclaimed water irrigation treatment: at 4.0 dS m<sup>-1</sup> leaching 40% (v/v) of the applied water; RW3, reclaimed water irrigation treatment: at 8.0 dS m<sup>-1</sup> leaching 55% (v/v) of the applied water.

<sup>†</sup>Mean values (n = 8) ± SD in each row followed by a different lower-case letter are significantly different according to Duncan's Multiple Range Test at P ≤ 0.05.



TABLE III

*Influence of four irrigation treatments on the growth of M. communis plants at the end of Period I and Period II<sup>‡</sup>*

Period	Parameter (units)	Control	RW1 <sup>§</sup>	RW2 <sup>§</sup>	RW3 <sup>§</sup>
I (4 months; 120 d)	Leaf DW (g plant <sup>-1</sup> )	6.00 ± 0.63a <sup>†</sup>	5.53 ± 0.31a	8.27 ± 0.34b	5.84 ± 0.56a
	Stem DW (g plant <sup>-1</sup> )	5.19 ± 0.55a	5.47 ± 0.46a	8.08 ± 0.57b	5.00 ± 0.42a
	Aerial DW (g plant <sup>-1</sup> )	11.20 ± 1.15a	11.00 ± 0.73a	16.35 ± 0.74b	10.83 ± 0.90a
	Root DW (g plant <sup>-1</sup> )	5.45 ± 0.69a	5.14 ± 0.63a	9.19 ± 0.21c	7.22 ± 0.19b
	Total DW (g plant <sup>-1</sup> )	16.64 ± 1.74a	16.14 ± 1.31a	25.54 ± 0.56b	18.06 ± 0.89a
	Root:shoot ratio	0.91 ± 0.07a	0.92 ± 0.08a	1.13 ± 0.07ab	1.32 ± 0.15b
	Leaf number	685 ± 36a	691 ± 31a	918 ± 34b	557 ± 50a
	Leaf area (cm <sup>2</sup> )	1109 ± 98.54a	1119 ± 87.67a	1640 ± 55.76b	975 ± 52.54a
	Plant height (cm)	36.8 ± 1.0b	34.4 ± 0.7ab	35.4 ± 1.1ab	32.9 ± 1.0a
	II (4 + 2 months; 180 d)	Leaf DW (g plant <sup>-1</sup> )	5.89 ± 0.36ab	5.43 ± 0.51a	7.21 ± 0.72b
Stem DW (g plant <sup>-1</sup> )		7.47 ± 0.64b	6.70 ± 0.60b	7.26 ± 0.64b	4.41 ± 0.56a
Aerial DW (g plant <sup>-1</sup> )		13.36 ± 0.82b	12.14 ± 1.05ab	14.48 ± 1.24b	9.60 ± 0.89a
Root DW (g plant <sup>-1</sup> )		8.65 ± 0.71a	8.26 ± 0.43a	11.65 ± 1.04b	8.91 ± 1.04a
Total DW (g plant <sup>-1</sup> )		22.02 ± 1.28ab	20.40 ± 1.19a	26.13 ± 2.25b	18.51 ± 1.85a
Root:shoot ratio		1.50 ± 0.14a	1.63 ± 0.15a	1.66 ± 0.11a	1.76 ± 0.16b
Leaf number		629 ± 35ab	610 ± 48a	759 ± 60b	616 ± 40a
Leaf area (cm <sup>2</sup> )		2127 ± 135b	2042 ± 159b	2431 ± 191b	1561 ± 114a
Plant height (cm)		38.2 ± 1.3b	36.2 ± 0.9ab	36.8 ± 1.4ab	33.8 ± 1.2a

<sup>‡</sup>Period I, a 4-month (120 d) period with control or waste water irrigation; Period II, a 2-month (60 d) period of recovery with control (low EC)

tap water irrigation after Period I.

§RW1, reclaimed water irrigation treatment: at 1.5 dS m<sup>-1</sup> leaching 25% (v/v) of the applied water; RW2, reclaimed water irrigation treatment: at 4.0 dS m<sup>-1</sup> leaching 40% (v/v) of the applied water; RW3, reclaimed water irrigation treatment: at 8.0 dS m<sup>-1</sup> leaching 55% (v/v) of the applied water.

†Mean values (n = 8; except in plant height, when n = 30) ± SD in each row followed by a different lower-case letter are significantly different according to Duncan's Multiple Range Test at  $P \leq 0.05$ .

TABLE IV

*Influence of four irrigation treatments on Na<sup>+</sup>, Cl<sup>-</sup>, B<sup>3+</sup>, K<sup>+</sup> and Ca<sup>2+</sup> ion concentration (in mmol kg<sup>-1</sup> DW tissue) in M. communis plants at the end of Period I and Period II<sup>‡</sup>*

Period	Ion	Tissue	Control	RW1 <sup>§</sup>	RW2 <sup>§</sup>	RW3 <sup>§</sup>
I (4 months; 120 d)	Cl <sup>-</sup>	Leaf	74.37 ± 10.96a <sup>†</sup>	94.84 ± 12.42ab	104.41 ± 7.84ab	120.19 ± 12.41b
		Stem	66.85 ± 4.99ab	57.84 ± 7.40a	91.27 ± 9.97bc	99.34 ± 13.71c
		Root	64.60 ± 1.90a	88.64 ± 12.24ab	121.50 ± 13.52b	123.19 ± 17.48b
	Na <sup>+</sup>	Leaf	25.38 ± 1.69a	36.80 ± 3.38ab	59.35 ± 6.41b	128.10 ± 17.40c
		Stem	60.38 ± 6.25a	94.26 ± 19.85a	86.72 ± 7.63a	167.85 ± 18.69b
		Root	162.13 ± 16.53a	188.51 ± 12.74a	184.88 ± 18.67a	355.83 ± 19.43b
	B <sup>3+</sup>	Leaf	8.02 ± 0.16a	9.28 ± 0.27ab	10.46 ± 1.36b	10.58 ± 0.34b
		Stem	5.63 ± 0.13a	6.21 ± 0.28a	8.79 ± 0.84b	6.13 ± 0.13a
		Root	6.95 ± 0.40a	7.29 ± 0.47ab	6.80 ± 0.55a	8.53 ± 0.29b
	K <sup>+</sup>	Leaf	449.15 ± 15.75b	456.52 ± 26.24b	558.21 ± 19.69c	333.28 ± 8.20a
		Stem	305.28 ± 30.59a	311.52 ± 51.30a	331.45 ± 23.99b	232.69 ± 16.82a
		Root	176.21 ± 17.13ab	135.72 ± 19.12a	202.93 ± 11.15b	134.19 ± 5.97a
	Ca <sup>2+</sup>	Leaf	108.73 ± 6.90a	112.22 ± 5.14a	161.32 ± 15.59b	159.64 ± 2.27b
		Stem	113.04 ± 12.21a	120.43 ± 17.88a	120.06 ± 16.54b	230.85 ± 5.28b
		Root	119.60 ± 11.56	101.30 ± 17.86	111.66 ± 7.70	99.09 ± 0.11
II (4 + 2 months; 180 d)	Cl <sup>-</sup>	Leaf	69.86 ± 14.89	75.72 ± 12.11	84.73±15.55	96.00 ± 14.52
		Stem	71.89 ± 16.28	82.70 ± 12.05	83.38±13.58	77.97 ± 9.24
		Root	75.72 ± 14.06	58.82 ± 5.67	59.61±7.03	68.96 ± 7.81
	Na <sup>+</sup>	Leaf	31.62 ± 4.57a	94.85 ± 15.66b	39.92±8.52a	75.47 ± 13.10b
		Stem	98.80 ± 17.59a	124.75 ± 9.78ab	109.43±21.10a	167.60 ± 16.13b
		Root	190.09 ± 15.37a	282.83 ± 13.48b	209.44±14.52a	274.95 ± 22.27b
	B <sup>3+</sup>	Leaf	10.26 ± 0.37a	12.96 ± 0.55c	11.54±0.39b	14.11 ± 0.41c

	Stem	6.18 ± 0.14a	7.67 ± 0.19c	7.08±0.18b	7.79 ± 0.19c
	Root	6.55 ± 0.19a	9.09 ± 0.22d	7.75±0.19b	8.45 ± 0.15c
K <sup>+</sup>	Leaf	401.14 ± 6.85b	314.63 ± 13.52a	419.56±17.44b	310.82 ± 10.49a
	Stem	298.96 ± 16.77c	226.44 ± 11.67b	203.87±16.06ab	181.85 ± 5.86a
	Root	122.65 ± 8.89b	99.13 ± 7.13a	98.91±8.01a	87.58 ± 7.40a
Ca <sup>2+</sup>	Leaf	116.24 ± 4.89ab	131.76 ± 6.89bc	105.24±5.62a	135.96 ± 4.76c
	Stem	139.74 ± 13.46b	136.11 ± 10.84b	95.38±8.25a	139.14 ± 11.49b
	Root	152.87 ± 9.57a	202.81 ± 17.07b	154.73±9.82a	164.18 ± 13.10a

<sup>‡</sup>Period I, a 4-month period (120 d) with control or waste water irrigation; Period II, a 2-month period (60 d) of recovery with control (low EC)

tap water irrigation after Period I.

<sup>§</sup>RW1, reclaimed water irrigation treatment: at 1.5 dS m<sup>-1</sup> leaching 25% (v/v) of the applied water; RW2, reclaimed water irrigation treatment: at 4.0 dS m<sup>-1</sup> leaching 40% (v/v) of the applied water; RW3, reclaimed water irrigation treatment: at 8.0 dS m<sup>-1</sup> leaching 55% (v/v) of the applied water.

<sup>†</sup>Mean values (n = 8) ± SD in each row followed by a different lower-case letter are significantly different according to Duncan's Multiple Range Test at P ≤ 0.05.

TABLE V

*Influence of four irrigation treatments on  $K^+/Na^+$  and  $Ca^{2+}/Na^+$  ion ratios in leaves of *M. communis* plants at the end of Period I and Period II<sup>‡</sup>*

Period	Ratio	Control	RW1 <sup>§</sup>	RW2 <sup>§</sup>	RW3 <sup>§</sup>
I (4 months; 120 d)	$K^+/Na^+$	18.19 ± 1.63c <sup>†</sup>	12.99 ± 1.42b	10.14 ± 1.45b	2.84 ± 0.36a
	$Ca^{2+}/Na^+$	4.18 ± 0.36c	3.23 ± 0.26b	3.01 ± 0.33b	1.23 ± 0.23a
II (4 + 2 months; 180 d)	$K^+/Na^+$	14.20 ± 1.28b	4.14 ± 0.70a	14.25 ± 2.44b	5.05 ± 0.70a
	$Ca^{2+}/Na^+$	4.21 ± 0.48b	1.72 ± 0.26a	3.83 ± 0.76b	2.19 ± 0.28a

<sup>‡</sup>Period I, a 4-month period (120 d) with control or waste water irrigation; Period II, a 2-month period (60 d) of recovery with control (low EC) tap water irrigation after Period I.

<sup>§</sup>RW1, reclaimed water irrigation treatment: at 1.5 dS m<sup>-1</sup> leaching 25% (v/v) of the applied water; RW2, reclaimed water irrigation treatment: at 4.0 dS m<sup>-1</sup> leaching 40% (v/v) of the applied water; RW3, reclaimed water irrigation treatment: at 8.0 dS m<sup>-1</sup> leaching 55% (v/v) of the applied water.

<sup>†</sup>Mean values (n = 8) ± SD in each row followed by a different lower-case letter are significantly different according to Duncan's Multiple Range Test at  $P \leq 0.05$ .

FIG.1

Leaf water potential ( $\Psi_l$ ; Panel A), relative water content (RWC; Panel B), stomatal conductance ( $g_s$ ; Panel C) and rate of net photosynthesis ( $P_n$ ; Panel D), at the ends of Period I [a 4-month (120 d) period with control or waste water irrigation] and Period II [a 2-month (60 d) period of recovery after Period I (120 d) with control tap water irrigation] in *M. communis* plants under four different irrigation treatments: control, RW1, RW2 and RW3 (see Material and Methods). Values are means ( $n = 6$ ) and vertical bars indicate  $\pm$  SE.

FIG.2

Relative chlorophyll concentration (RCC in SPAD meter values; Panel A), lightness ( $L^*$ ; Panel B), chroma ( $C^*$ ; Panel C) and hue angle ( $h^\circ$ ; Panel D) at the ends of Period I [a 4-month (120 d) period with control or waste water irrigation] and Period II [a 2-month (60 d) period of recovery after Period I (120 d) with control tap water irrigation] in *M. communis* plants under four different irrigation treatments: control, RW1, RW2 and RW3 (see Material and Methods). Values are means ( $n = 6$ ) and vertical bars indicate  $\pm$  SE.




FIG.1




FIG.2