

1 **Physiological and biochemical mechanisms of the ornamental *Eugenia myrtifolia* L. plants for**
2 **coping with NaCl stress and recovery**

3
4
5 **Running title:** Salt stress and recovery in *Eugenia myrtifolia* L.

6
7 José-Ramón Acosta-Motos¹, Pedro Diaz-Vivancos², Sara Álvarez¹, Nieves Fernández-García³, María
8
9 Jesús Sanchez-Blanco¹, José Antonio Hernández^{2*}

10
11
12
13 ¹Irrigation Department, CEBAS-CSIC, Campus Universitario de Espinardo, Murcia, P.O. Box 164, E-
14
15 30100, Spain.

16
17 ²Fruit Tree Biotechnology Group, Dept. of Plant Breeding, CEBAS-CSIC, Campus Universitario de
18
19 Espinardo, Murcia, P.O. Box 164, E-30100, Spain.

20
21 ³Department of Abiotic Stress and Plant Pathology, CEBAS-CSIC, Murcia, P.O. Box 164, E-30100,
22
23 Spain.

24
25
26
27
28
29
30 *Corresponding author

31 Dr. José Antonio Hernández

32 Centro de Edafología y Biología Aplicada del Segura (CEBAS-CSIC)

33
34 Campus Universitario de Espinardo, Murcia, P.O. Box 164, E-30100 SPAIN

35
36 Phone number: 34 968 396319

37
38 FAX: 34 968 396213

39
40 E-mail: jahernan@cebas.csic.es
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Main Conclusion

1
2
3 We studied the response of *Eugenia myrtifolia* L. plants, an ornamental shrub native to tropical and
4 subtropical areas, to salt stress in order to facilitate the use of these plants in Mediterranean areas for
5 landscaping. *E. myrtifolia* plants implement a series of adaptations to acclimate to salinity, including
6 morphological, physiological and biochemical changes. Furthermore, the post-recovery period seems to
7 be detected by *Eugenia* plants as a new stress situation.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

1
2
3 Different physiological and biochemical changes in *Eugenia myrtifolia* L. plants after being
4 subjected to NaCl stress for up to 30 days (Phase I) and after recovery from salinity (Phase II) were
5 studied. Eugenia plants proved to be tolerant to NaCl concentrations between 44 and 88 mM, displaying a
6 series of adaptative mechanisms to cope with salt-stress, including the accumulation of toxic ions in roots.
7
8 Plants increased their root/shoot ratio and decreased their leaf area, leaf water potential and stomatal
9 conductance in order to limit water loss. In addition, they displayed different strategies to protect the
10 photosynthetic machinery, including the limited accumulation of toxic ions in leaves, increase in
11 chlorophyll content, changes in chlorophyll fluorescence parameters, leaf anatomy and antioxidant
12 defence mechanisms. Anatomical modifications in leaves, including an increase in palisade parenchyma
13 and intercellular spaces and decrease in spongy parenchyma, served to facilitate CO₂ diffusion in a
14 situation of reduced stomatal aperture. Salinity produced oxidative stress in Eugenia plants as evidenced
15 by oxidative stress parameters values and a reduction in APX and ASC levels. Nevertheless, SOD and
16 GSH contents increased. The post-recovery period is detected as a new stress situation, as observed
17 through effects on plant growth and alterations in chlorophyll fluorescence and oxidative stress
18 parameters.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 **Keywords:** ASC-GSH cycle, Gas exchange, Leaf anatomy, Oxidative stress, Recovery capacity, Water
35 relations.
36
37
38

39 **Abbreviations:** APX: ascorbate peroxidase; ASC: Ascorbate reduced form; DHAR: dehydroascorbate
40 reductase; GR: Glutathione reductase; GSH: glutathione reduced form; GSSG: glutathione oxidised form;
41 MDHAR: monodehydroascorbate reductase; POX: peroxidase; SOD: superoxide dismutase.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

Mediterranean areas are characterised by limited water availability. Therefore, the use of non-conventional water resources is a common strategy for efficient water management. Saline waters can be an option in irrigation strategies particularly for ornamental shrubs in landscaping (Cassaniti et al. 2009)

Under saline conditions, plants have to activate different physiological and biochemical mechanisms in order to cope with the resulting stress. Such mechanisms include changes in water relations, photosynthesis, respiratory metabolism, the hormonal profile, toxic ion distribution and the antioxidative metabolism response (Hernández et al. 2001; Parida and Das 2005; Álvarez et al. 2012, 2014; Ashraf and Harris 2013). Physiological constraints imposed by salt stress include osmotic stress and ion toxicity, leading to a nutrient imbalance as well as a disruption of the plant's metabolism (Marschner 1995). Furthermore, and as previously reported, salt stress is also manifested as an oxidative stress at the subcellular level (Corpas et al. 1993; Hernández et al. 1995). These three factors mentioned above can all contribute to the negative effects produced by salinity in plants.

Salt-induced reductions in plant growth are associated with decreases in the net photosynthesis rate. It is known that salinity affects the photosynthetic process due to stomatal and non-stomatal limitations, including stomatal closure, a reduction in chlorophyll content, the inhibition of Calvin cycle enzymes and the degradation of membrane-associated proteins in the photosynthetic apparatus (Parida and Das 2005; Mittal et al. 2012; Shu et al. 2013). Many authors have reported the decrease in net photosynthesis and stomatal conductance resulting from short-term and long-term exposure to salinity. However, the reductions in these parameters have been found to be less marked in salt-tolerant than in salt-sensitive plants (Moradi and Ismail 2007; Duarte et al. 2013). Moreover, salt stress has been shown to produce a decrease in the photochemical quenching parameters in different plant species, suggesting inhibition of PSII electron transport (Moradi and Ismail 2007; Mehta et al. 2010). In addition, salt stress has been observed to produce either increases or reductions in the non-photochemical parameters, depending on the plant species studied (Moradi and Ismail 2007; Iqbal et al. 2014).

A correlation between salt stress tolerance and an improved oxidative stress response has been observed by different authors (Hernández et al. 2001; Moradi and Ismail 2007; Duarte et al. 2013; Gil et al. 2014), although increases in antioxidative enzymatic activities have also been described in some salt-sensitive species (Arbona et al. 2003; Lee et al. 2013). Different authors have reported that NaCl-tolerant plants either induce or show higher constitutive levels of antioxidant defences (Gueta-Dahan et al. 1997; Hernández et al. 2000, 2003; Mittova et al. 2003). In fact, it has been observed that halophytes present a

1 higher antioxidant capacity than glycophytes, suggesting that this may be one of the reasons why
2 halophytes tolerate high salinity levels (Ozgun et al. 2013; Bose et al. 2014; Gil et al. 2014).
3

4 The effect of salt stress on crop plants has been extensively studied. However, few authors have
5 focused their attention on the effect of salinity on ornamental shrubs. Saline waters can be an option in
6 irrigation strategies for ornamental shrubs in landscaping and is of particular interest in Mediterranean
7 areas. Yet salinity may affect the aesthetic value of plants, which is a very important aspect when working
8 with ornamental plants (Acosta-Motos et al. 2014a, 2014b).
9

10 In this work, we used *E. myrtifolia* plants, an ornamental shrub native to tropical areas in Asia
11 and Oceania and subtropical areas in South America. One of our goals was to study the response of this
12 plant species to NaCl stress with the hypothesis that it would be a good candidate for use in
13 Mediterranean environments for landscaping. The effect of moderate NaCl levels on plant growth and
14 toxic ion distribution in different ornamental plants, including *Eugenia myrtifolia* L., has been studied in a
15 previous work (Cassaniti et al. 2009) but no further analyses have been performed.
16

17 Based on the working hypothesis, the effect of different NaCl treatments at 15 and 30 days on
18 plant growth, gas exchange, water relation, mineral nutrition, chlorophyll fluorescence, leaf anatomy and
19 antioxidative metabolism in *Eugenia myrtifolia* L. plants was studied. Furthermore, the relevance of
20 studying the plants' capacity for recovery following salinity relief was also taken into account. Current
21 information regarding the response of plants to recovery from salt stress is scarce, and the physiological
22 mechanisms involved in this recovery process remain poorly understood (Chaves et al. 2011). We have
23 also investigated a possible relationship between Na⁺ and Cl⁻ uptake and partitioning among organs in
24 order to evaluate if the plant response might be related to the retention of these ions in the roots.
25

26 **Material and methods**

27 **Plant and experimental conditions**

28 Single rooted cuttings (120) of native *Eugenia myrtifolia* L. plants were transplanted into 14 x 12
29 cm pots (1.2 L) filled with a mixture of coconut fibre, sphagnum peat and perlite (8:7:1) and amended
30 with Osmocote plus (2 g L⁻¹ substrate) (14:13:13 N, P, K + microelements) [Agrosolmen S.L., Lorca
31 (Murcia), Spain]. The experiment was conducted in a controlled environment growth chamber set to
32 simulate natural conditions as described in Acosta-Motos et al. (2014b). The temperature in the chamber
33 was 23°C during the light phase (16 h photoperiod) and 18°C during darkness. Relative humidity (RH)
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 values ranged between 55 and 70%. A mean photosynthetic active radiation (PAR) of $350 \mu\text{mol m}^{-2} \text{s}^{-1}$ at
2 canopy height was supplied during the light phase (08:00h-00:00h) by cold white fluorescent lamps.
3

4 Experimental design and treatments

5
6
7 Once *E.mytilifolia* plants were adapted to chamber conditions, they were exposed for up to 30
8 days (Phase I) to the following four irrigation treatments. Control plants were watered with a mixture of
9 distilled water and tap water with an electrical conductivity (EC) = 0.3 dS/m. Saline treatments were
10 designed as control treatment plus NaCl added specifically for each treatment: S4 (4 dS/m), S8 (8 dS/m)
11 and S12 (12 dS/m), corresponding to 44, 88 and 132 mM NaCl, respectively. The EC of the different
12 treatments was evaluated with a multirange Cryson-HI8734 electrical conductivity meter (Cryson
13 Instruments, S.A., Barcelona, Spain) at the beginning of and throughout the experimental period. Before
14 starting the experimental period, the maximum water holding capacity of the soil was determined for each
15 individual pot and was considered as the weight at field capacity (WFC). Throughout the experiment, all
16 pots were irrigated three times a week below the WFC in order to avoid drainage, favouring an increase in
17 soil salinity due to time and the severity of the saline treatments. After the stress phase (Phase I), all
18 plants were exposed to a 16-day recovery period (Phase II) in which they were irrigated with the same
19 solution used for the control plants. During the first three days of the recovery period, all plants were
20 exposed to a further irrigation event with leaching with the same solution used for the control plants (a
21 mixture of distilled water and tap water). The leaching fraction reached 10% (v/v) of the water applied in
22 the control treatments, 27% of the water applied in S4 treatments, 50% of the water applied in S8
23 treatments, and 72% of the water applied in S12 treatments, respectively.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 Growth, inorganic solutes and ionic absorption rate determinations

44
45 At the beginning and end of the salinity period (Phase I) and during the recovery period (Phase II),
46 the soil was gently washed from the roots of six plants per treatment and each plant harvested was divided
47 into shoots (leaves and stem) and roots, and the different organs were washed with distilled water. The
48 leaf fresh weights (FW) and leaf relative water content were measured. Then, leaves, stems and roots
49 were oven-dried at 80°C until they reached a constant weight in order to measure their respective dry
50 weights (DW). Leaf areas (cm^2) were determined for the same plants before drying using a leaf area
51 meter (AM 200; ADC BioScientific Ltd., Hoddesdon, UK).
52
53
54
55
56
57
58

59 At the beginning and end of Phase I and during Phase II the same plants used for growth
60 measurements were also used to determine the inorganic solutes and ionic absorption rate. Plant material
61
62
63
64
65

that had been previously oven-dried at 80°C until it reached a constant weight, was ground to obtain dry vegetable powder. The concentrations of Cl⁻ were analysed by a chloride analyser (Chloride Analyser Model 926, Sherwood Scientific Ltd.) in the aqueous extracts obtained by mixing 100 mg of dry vegetable powder with 40 mL of water before shaking for 30 min and filtering. The concentrations of Na⁺, K⁺, and Ca²⁺ ions were determined in a digestion extract with HNO₃:HClO₄ (2:1, v/v) by Inductively Coupled Plasma optical emission spectrometer (ICPOES IRIS INTREPID II XDL). The absorption rate of Cl⁻, Na⁺, K⁺, and Ca²⁺ (J) by the root system was calculated considering the total salt content, expressed as mmol of Cl⁻, Na⁺, K⁺, and Ca²⁺ and the mean root weight, using the formula described by Pitman (1975):

$$J = (M_2 - M_1) / (WR * t)$$

where M₁ and M₂ correspond to a concentration of Cl⁻, Na⁺, K⁺, and Ca²⁺ in mmol in the total plant at the beginning, at the end of the salinity (Phase I) and at the end of the recovery periods (Phase II), respectively. In this formula, t corresponds to the time in days and WR is the logarithmic mean root biomass, calculated as (WR₂-WR₁)/Ln (WR₂/WR₁), with WR₁ and WR₂ representing the dry weight of roots at the beginning and at the end of Phase I or at the end of Phase II, respectively.

Plant water measurements and gas exchange

The soil water potential at the root surface (Ψ_r), leaf water potential (Ψ_l), leaf osmotic potential (Ψ_s), leaf turgor potential (Ψ_t), and leaf osmotic potential at full turgor (Ψ_{100s}) were estimated in six plants per treatment during the central hours of illumination at middle (15 d) and end of Phase I and once Phase II was finished.

The soil water potential was estimated using the method described by Jones (1983), which assumes that $\Psi_r = 0$ for control plants. To calculate Ψ_r for NaCl treatments we used the following equation:

$$\Psi_r = \Psi_{NaCl} - (\Psi_c \times g_{SNaCl}) / g_{Sc}$$

where Ψ_c and Ψ_{NaCl} correspond to the mean value of leaf water potential in the control and NaCl treatments, respectively, while g_{Sc} and g_{SNaCl} correspond to the mean value of stomatal conductance in the respective treatments. Leaf water potential was estimated using a pressure chamber (Model 3000; Soil Moisture Equipment Co., Santa Barbara, CA, USA) in which leaves were placed in the chamber within 20 s of collection and pressurised at a rate of 0.02 MPa s⁻¹. Leaves from the Ψ_t measurements were

frozen in liquid nitrogen (-196°C) and stored at -30°C . After thawing, the osmotic potential (Ψ_s) was measured in the extracted sap using a WESCOR 5520 vapour pressure osmometer (Wescor Inc., Logan, UT, USA). Ψ_t was estimated as the difference between leaf water potential (Ψ_l) and leaf osmotic potential (Ψ_s). Leaf osmotic potential at full turgor (Ψ_{100s}) was estimated as indicated above for Ψ_s , using excised leaves with their petioles placed in distilled water overnight to reach full saturation.

The contribution of ions to total Ψ_{100s} was calculated according to Munns and Weir (1981). From the relative dry weight (RDW, kg m^{-3}) (dry weight/leaf water content), the solute concentration on a dry-weight basis (C, g kg^{-1}), the molecular weight of each solute (M, g mol^{-1}) and the van't Hoff relation (using a RT value for 25°C of $0.002479 \text{ m}^3 \text{ MPa mol}^{-1}$, Nobel 1983) for six plant per treatment. It is assumed that ions behaved as ideal osmotic

$$\Psi_{100s \text{ calculated}} = -0.002479 \times \text{RDW} \times \text{C} \times 1/\text{M}$$

The proline in leaf samples was analysed at middle (15 d) and end of Phase I and once Phase II was finished as described in Pérez-Clemente et al. (2012). Briefly, 0.1 g of frozen plant tissue (leaves) was homogenized with 5 mL of 3% sulfosalicylic acid using a tissue homogenizer (Ultra-Turrax). After extraction, homogenates were centrifuged to pellet cell debris at 4°C at 12.000 g for 10 min and 1 mL aliquot of the supernatant was combined with an equal volume of glacial acetic acid and ninhydrin reagent. This mixture was boiled in a water bath for 1 h and then cooled in an ice bath (at least 5 min). The solution was partitioned against 2 mL of toluene and absorbance at 520 nm measured in this organic layer. A calibration curve was performed using commercial proline as a standard.

Evapotranspiration (ET) was measured gravimetrically during Phase I in 30 plants per treatment, based on the difference in weights (weight after irrigation and weight before irrigation again), using a balance (Analytical Sartorius, Model 5201; capacity 5.2 kg and accuracy of 0.01 g).

Leaf stomatal conductance (gs) and leaf photosynthetic rate (P_N) in attached leaves in six plants per treatment during the central hours of illumination were determined at middle (15 d) and end of Phase I and once Phase II was finished using a gas exchange system (LI-6400; LI-COR Inc., Lincoln, NE, USA). Intrinsic water-use efficiency (WUE) was calculated based on the P_N/gs balance registered. For leaf chlorophyll determination, 30 mg of fresh leaves from the central region, avoiding the main vein, were used. Leaf samples were incubated in 3 mL of N, N-dimethylformamide in darkness at least for 72 h. The absorbance was read at 645 nm and 664.5 nm with a Thermo Spectronic (model Helios alpha, UVA No. 092009) and used to calculate chlorophyll content ($\text{mg g}^{-1} \text{ FW}$) according to Romero-Trigueros et al. (2014).

Measurement of chlorophyll fluorescence

Chlorophyll fluorescence was measured in detached leaves from control and salt-treated *Eugenia* plants with a chlorophyll fluorimeter (IMAGIM-PAM M-series, Heinz Walz, Effeltrich, Germany). After a dark incubation period (20 min), the minimum and the maximal fluorescence yields of the plants were monitored. Kinetic analyses were carried out with actinic light ($81 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ PAR) and repeated pulses of saturating light at $2700 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ PAR for 0.8 s at intervals of 20 s. The following parameters were also analysed: effective PSII quantum yield [Y(II)]; the quantum yield of regulated energy dissipation [Y(NPQ)]; the non-photochemical quenching (NPQ); the maximal PSII quantum yield (Fv/Fm); the coefficients of non-photochemical quenching (qN); and the photochemical quenching (qP) (Maxwell and Johnson 2000).

Enzyme extraction and analysis

All operations were performed at 4°C. For the enzymatic determinations, plants were sampled at 15 and 30 days of stress and after 16 days of recovery.

Leaf samples (1 g) were homogenized with an extraction medium (1/3, w/v) containing 50 mM Tris-acetate buffer (pH 6.0); 0.1 mM EDTA; 2 mM cysteine; 1 % (w/v) PVP; 1% (w/v) PVPP; and 0.2% (v/v) Triton X-100. For the APX activity, 20 mM of sodium ascorbate was added to the extraction buffer. The extracts were filtered through two layers of nylon cloth and centrifuged at 10000 g for 15 min. The supernatant fraction was filtered on Sephadex NAP-10 columns (GE Healthcare) equilibrated with the same buffer used for homogenisation and used for the enzymatic determinations. For the APX activity, 2 mM of sodium ascorbate was added to the equilibration buffer. APX (EC 1.11.1.11) was determined at 290 nm following the ASC oxidation by H₂O₂ (Hossain and Asada 1984). MDHAR (EC 1.6.5.4) was assayed by the decrease at 340 nm due to the NADH oxidation (Arrigoni et al. 1981). Monodehydroascorbate was generated by the ascorbate/ascorbate oxidase system (Arrigoni et al. 1981). To determine the MDHAR activity, the rate of monodehydroascorbate-independent NADH oxidation (without ascorbate and ascorbate oxidase) was subtracted from the initial monodehydroascorbate-dependent NADH oxidation rate (with ascorbate and ascorbate oxidase). DHAR (EC 1.8.5.1) was determined by following the increase at 265 nm due to ascorbate formation (Dalton et al. 1993). The reaction rate was corrected for the nonenzymatic reduction of DHA by GSH. GR (EC 1.6.4.2) was assayed by the decrease at 340 nm to the NADPH oxidation, as described by Edwards et al. (1990). The

1 reaction rate was corrected for the small, nonenzymatic oxidation of NADPH by GSSH. SOD (EC
2 1.15.1.1) was assayed by the ferricytochrome c method using xanthine/xanthine oxidase as the source of
3 superoxide radicals (McCord and Fridovich 1969). CAT (EC 1.11.1.6) was measured following the
4 decrease of absorbance of H₂O₂ at 240 nm (Aebi 1984). POX activity (EC. 1.11.1.7) was analysed
5 following the oxidation of 4-methoxy- α -naphthol at 593 nm according to Ros-Barceló (1998).
6
7
8
9

10 11 12 Oxidative stress parameters

13
14 The rate of passive electrolyte leakage from stress-sensitive plant tissue can be used as a measure
15 of alterations in membrane permeability. Ion leakage was estimated at 15 and 30 days in Phase I and at
16 the end of Phase II. Leaf discs (2 mm diameter) were incubated in 10 mL of 0.3 M mannitol in 50-mL
17 plastic centrifuge tubes and the conductivity of the solutions was measured after 24 h with a conductivity-
18 meter (Crison Mod. 524). Tubes containing the mannitol solution and the tissue were weighed and heated
19 to boiling for 5 min. After cooling to room temperature with shaking, deionized water was added to make
20 their initial weight, and the total conductivity was measured after an additional 0.5 h of shaking. Ratios of
21 ion leakage are expressed as percentage of the total conductivity per hour (Acosta-Motos et. al. 2014b).
22
23
24
25
26
27
28
29

30 The extent of lipid peroxidation was estimated by determining the concentration of thiobarbituric
31 acid-reactive substances (TBARS). Briefly, leaf material (400 mg) was homogenized in 1 M perchloric
32 acid solution. The homogenate was centrifuged at 15000 g for 10 min and 0.5 mL of the supernatant
33 obtained was added to 1.5 mL 0.5% TBA in 1M perchloric acid. The mixture was incubated at 90°C in a
34 shaking water bath for 20 min, and the reaction was stopped by placing the reaction tubes in an ice water
35 bath. Then, the samples were centrifuged at 10000 g for 5 min, and the absorbance of the supernatant was
36 read at 532 nm. The value for non-specific absorption at 600 nm was subtracted. The amount of TBARS
37 (red pigment) was calculated from the extinction coefficient 155 mM⁻¹ cm⁻¹ (Hernández and Almansa
38 2002).
39
40
41
42
43
44
45
46
47
48
49

50 51 Ascorbate and glutathione analyses

52
53 Leaf samples (four replicates per treatment) were snap-frozen in liquid nitrogen and then ground to a
54 fine powder and extracted in 1 mL of 1 M HClO₄. Homogenates were centrifuged at 12000 g for 10 min. The
55 supernatant was neutralized with 5 M K₂CO₃ to pH 5.5–6. The homogenate was centrifuged at 12000 g for 1
56 min to remove KClO₄. The supernatant obtained was used to determine ascorbate and glutathione content
57 (Diaz-Vivancos et al. 2010). Reduced ascorbate was measured by the change in absorption at 265 nm,
58
59
60
61
62
63
64
65

1 where ascorbate was determined via oxidation to DHA in the presence of ascorbate oxidase (Pellny et al.
2 2009). Glutathione (GSH, GSSG) were analysed using dithio-bis-2- nitrobenzoic acid and glutathione
3
4 reductase in the presence of NADPH (Pellny et al. 2009).
5
6
7

8 Light microscopy and morphometrical analysis 9

10 Leaves sections (1×1 mm from the most recent fully expanded leaves) from the central region
11 of *Eugenia* leaves, avoiding the main vein, were used for light microscopy. These samples were fixed and
12 postfixed according to Fernández-García et al. (2013). Semi-thin sections (0.5-0.7 μ m thick were cut with
13 a Leica EM UC6 ultramicrotome. The sections were stained with 0.5% toluidine blue, mounted in DPX
14 and observed with a Leica DMR light microscope. For morphometric analysis, 10 different sections from
15 each treatment (3 plants of each treatment), were studied. The percentages of area occupied by palisade
16 parenchyma (PP), spongy parenchyma (SP) and intercellular spaces (IS) in leaves from *E. myrtifolia*
17 plants were measured and expressed as the % of total area using Adobe Photoshop CS4 Extended
18 software.
19
20
21
22
23
24
25
26
27
28
29

30 Statistical analyses of data 31

32 In the experiment, 30 plants were randomly attributed to each treatment. The data were analysed
33 by one-way ANOVA using Statgraphics Plus for Windows 5.1 software. Treatment means were separated
34 with Duncan's Multiple Range Test ($P \leq 0.05$).
35
36
37
38
39

40 Results 41

42 Effect of NaCl on plant growth 43 44

45 At the end of Phase I, 4 dS/m NaCl (S4) stimulated the foliar area in *Eugenia* plants, whereas 8
46 dS/m NaCl (S8) did not affect the studied growth parameters. In addition, control, S8 and S12 plants lost
47 leaf area between Phases I and II. In general, the highest NaCl levels (S12) induced a significant decrease
48 in biomass production as could be observed by the 40% reduction in leaf and stem DW (Table 1).
49 Although salt stress produced no statistically significant changes in the root DW, a concentration-
50 dependent decrease in this parameter was observed, leading to an increase in the DW root/DW shoot ratio
51 in plants treated with the highest NaCl level (Table 1).
52
53
54
55
56
57
58

59 After the recovery phase (Phase II), plants previously subjected to the S4 treatment displayed the
60 best performance, showing higher values in foliar area as well as in the leaf and stem DW than control
61
62
63
64
65

1 plants (Table 1). However, in plants previously irrigated with 8 dS/m NaCl, a reduction in foliar area was
 2 observed after the recovery period. The plants subjected to the S12 treatment did not show any signs of
 3 recovery, and a decrease of about 40% was recorded in the growth parameters of these plants in relation
 4 to the control (Table 1).
 5
 6

7 8 9 Nutritional changes

10 Salt stress increased the uptake rate for Cl^- in a concentration-dependent manner. At the end of
 11 Phase I, these values increased 2-, 5.3- and 7-fold in S4, S8 and S12 plants, respectively, in relation to
 12 control plants (Fig. 1a). The absorption rate for Na^+ did not show statistically significant changes in S4
 13 plants, whereas similar increases were produced in S8 and S12 plants. In contrast, the uptake rate of K^+
 14 by roots significantly decreased in all NaCl-treated plants, whereas an increase in the Ca^{2+} uptake rate
 15 was observed in plants irrigated with 8 and 12 dS/m NaCl (Fig. 1a, b, c).
 16
 17
 18
 19
 20
 21
 22

23 After Phase II, the uptake rate of Cl^- decreased in Eugenia plants, mainly in plants previously
 24 subjected to S8 and S12 treatments, although the values were still much higher than those observed for
 25 control plants. No statistically significant changes were observed for the Na^+ absorption rate, whereas the
 26 behaviour of K^+ uptake was similar to that observed in Phase I. Finally, similar to Cl^- absorption, Ca^{2+}
 27 uptake values decreased in all cases, but the data were higher in plants subjected to salt stress than in
 28 control plants (Fig. 1).
 29
 30
 31
 32
 33
 34

35 Concerning the distribution of the different ions, at the end of Phase I, Cl^- accumulated mainly in
 36 roots from S8 and S12 plants, and the Cl^- concentration was more limited in the aerial part of the plants
 37 (Fig. 2a). Similarly, Na^+ also accumulated in roots from plants subjected to the S8 and S12 treatments. No
 38 important changes were observed in leaves, whereas Na^+ only accumulated in the stems of S12 plants
 39 (Fig. 2b). After Phase II, Cl^- and Na^+ levels were much lower than those observed in Phase I. During the
 40 recovery period, even though the drainage conditions applied reduced Na^+ and Cl^- uptake, both ions still
 41 accumulated in roots. Na^+ concentration also increased in leaves and stems (Fig. 2a, b). At the end of
 42 Phase I, K^+ concentration dropped in all parts of NaCl-treated plants (Fig. 2c). After the recovery period,
 43 K^+ levels decreased in leaves from S4 plants as well as in all organs from plants previously irrigated with
 44 8 and 12 dS/m (Fig. 2c). A significant increase in Ca^{2+} concentrations was produced in all parts of the
 45 plants in both phases of the experiment (Fig. 2d). NaCl had a similar effect on the absolute Na^+ and Cl^-
 46 contents as it had on Na^+ and Cl^- concentrations. S8 and S12 plants thus presented both a higher root
 47 content and concentration of Na^+ and Cl^- in Phase I (Fig. 2, Suppl. Fig. S1). However, in Phase II, due to
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65

drainage, a decrease in Na^+ and Cl^- contents also occurred in roots. In addition, a mobilisation of both toxic ions occurred in the canopy (Suppl. Fig. S1).

Plant water relations

Table 2 shows the effect of NaCl on plant-water relations. During Phase I the soil water potential at the root surface (Ψ_r) decreased in parallel with the severity of the saline treatments (Table 2). However, at the end of the recovery period, these values increased in relation to the data observed in Phase I (Table 2). Leaf water potential (Ψ_l) experienced a progressive decline with the severity of the NaCl treatments. At 15 and 30 days of salt-treatment *Eugenia* plants presented significantly more negative Ψ_l values than control plants, especially those treated with 8 and 12 dS/m NaCl (Table 2). At the end of Phase II, the Ψ_l values of stressed plants increased but did not reach the control values (Table 2).

Regarding leaf turgor potential (Ψ_t), only at the end of Phase I were significant differences observed among control and S8 and S12 plants (Table 2). Nevertheless, no differences in Ψ_t values were observed at the end of Phase II (Table 2).

During Phase I, the osmotic potential at maximum saturation (Ψ_{100s}) values were more negative in S8 plants, indicating an osmotic adjustment process (Table 2). However, the values were more negative at the end of the recovery period than after Phase I, and all previously stressed plants displayed lower Ψ_{100s} values than control plants (Table 2). At the end of Phase I the contribution of the ions to the level of osmotic adjustment differed with the saline treatment. The importance of Na^+ , Cl^- and Ca^{2+} increased with increases in the NaCl level, whereas the importance of K^+ decreased (Suppl. Table S1). At the end of Phase II, the contribution of Na^+ and Cl^- to osmotic adjustment was more important than at the end of the saline period (see data in bracket in Suppl. Table S1, see also Ψ_{100s} in Table 2).

In parallel to the water relation parameters, we studied the effect on NaCl on the proline levels during the experiment (Table 2). In general, during Phase I, proline contents were always higher in plants treated with the most severe NaCl treatments. At the end of this period, only plants previously subjected to the 8 and 12 dS/m treatments showed higher levels of proline than control plants, as occurred also after Phase II (Table 2).

Gas exchange and chlorophyll fluorescence parameters

Eugenia plants showed unchanged or increased levels of total chlorophyll under saline conditions (Table 3), the effect being more evident in S12 plants at 15 days of stress and in S8 and S12 plants at the end of the recovery phase (Table 3).

Evapotranspiration (ET) was higher in control plants throughout the experimental period, and values fell proportionally with respect to increasing NaCl treatments (Suppl. Fig. S2). At 15 days of salt stress, a NaCl-dependent fall in g_s occurred. In this case, the g_s values decreased by about 32%, 46% and 59% in S4, S8 and S12 plants, respectively (Table 2). Regarding P_N values, a 35% decrease occurred in S12 plants (Table 3). The g_s decrease produced a rise in WUE values (Table 3). After 30 days of stress, S4 plants appear to have developed an ability to acclimate to the stress conditions, showing similar g_s values to control plants (Table 3), whereas S8 and S12 plants showed decreased g_s values (Table 3). At the end of Phase I, P_N values only decreased in S8 and S12 plants (Table 3).

At the end of Phase II, g_s values slightly increased in all treatments with respect to the values observed after Phase I. For example, g_s values increased by up to 70% in S4 plants in relation to control plants, and, as a consequence, there was a significant increase in P_N as well (Table 3).

After 15 days of NaCl-stress, plants irrigated with 8 and 12 dS/m showed decreased photochemical quenching parameters [qP and $Y(II)$] and increased non-photochemical quenching parameters [qN , NPQ and $Y(NPQ)$]. However, at 30 days of salt-stress, an inverse response took place: the photochemical quenching parameters increased in salt-treated plants, whereas the non-photochemical quenching parameters decreased (Table 4, Suppl. Fig. S1). After Phase II, an alteration in the chlorophyll fluorescence parameters occurred, particularly in plants previously irrigated with 8 and 12 dS/m. In these plants, a decrease in qP as well as in qN and NPQ was recorded (Table 4 and Suppl. Fig. S3).

Anatomical changes

Salt stress induced changes in the leaf anatomy of Eugenia plants, and such changes were most evident in plants irrigated with 8 dS/m NaCl. Accordingly, at the end of Phase I, plants treated with 8 dS/m NaCl showed an increase in the percentage of palisade parenchyma and intercellular spaces but a decrease in spongy parenchyma (Table 5 and Suppl. Fig. S2). Changes produced in S4 and S12 plants were related to an increase in intercellular spaces (Table 5 and Suppl. Fig. S4).

After Phase II, anatomical modifications were observed for all treatments, especially in S4 plants. In these plants, an increase in palisade parenchyma and intercellular spaces as well as a decrease in spongy parenchyma could be observed. Plants previously treated with 8 dS/m maintained increases in the

percentage of palisade parenchyma and decreases in spongy parenchyma, and similar changes occurred in S12 plants after the recovery period (Table 5 and Suppl. Fig. S4).

Antioxidative metabolism

The NaCl treatment induced oxidative stress in *Eugenia* plants as evidenced by electrolyte leakage (EL) and lipid peroxidation (LP), indicative of membrane damage. Such effects were most noticeable in S12 plants. After Phase II, these plants still presented membrane damage as evidenced by an increase in the oxidative stress parameters (Table 6). In the case of S8 plants, although EL data returned to control values, there was nevertheless an increase in LP in relation to control plants (Table 6).

The effect of NaCl on the activity of some antioxidant enzymes was studied in plants subjected to 4 dS/m NaCl and 8 dS/m NaCl. At 15 days of salt stress, CAT activity increased in NaCl-treated plants, especially in S4 plants. This CAT increase was accompanied by a decrease in APX activity. In addition, a 2-fold increase in SOD as well as a strong decrease in POX activity was produced in S8 plants (Table 7). At 30 days of NaCl-stress, S4 plants showed an increase in GR and SOD and a drop in APX (Table 7). In S8 plants, we observed increases in MDHAR and SOD but significant decreases in APX and POX activities (Table 7).

At the end of Phase II, CAT activity increased and APX activity reached control values in stressed plants (Table 7). In contrast, MDHAR and GR decreased in S4 plants and SOD increased in both treatments. A general decrease in POX activity was produced in Phase II, but S8 plants displayed a significant increase in this enzymatic activity in relation to control plants (Table 7).

After 15 days of NaCl treatment, a strong increase in GSH was observed in *Eugenia* plants. Furthermore, this increase was much higher in S4 plants (5-fold) than in S8 (2.5-fold) plants with respect to the control, but no accumulation of GSSG occurred (Table 8). This response produced an increase in the redox state of GSH. At 30 days of NaCl stress, irrigated S8 plants maintained a significant increase in GSH. At the end of Phase I, an accumulation of GSSG was observed, producing a decrease in the redox state of glutathione in all cases (Table 8). After Phase II, control plants maintained GSH levels, whereas S4 plants displayed duplicate GSH values, and the data were three times higher in S8 plants (Table 8). In this period, GSSG values were much higher in control than in salt-stressed plants, which displayed a higher redox state of glutathione.

No oxidized ascorbate was detected in *Eugenia* plants under our experimental conditions. At 15 days of NaCl stress, reduced ascorbate (ASC) levels showed no significant differences among the

1 treatments, although values were higher in NaCl-treated plants. However, at 30 days of NaCl irrigation,
2 decreased ASC levels were observed in plants subjected to both saline treatments. After Phase II, ASC
3 content increased dramatically in all treatments. Nevertheless, plants previously subjected to NaCl
4 displayed lower ASC levels (3-fold in S4 and 4-fold in S8) than control plants (Table 8).
5
6
7

8 9 10 **Discussion**

11 Our data suggested that *E. myrtifolia* plants could be used for landscaping projects in
12 Mediterranean areas. This plant species implements a series of adaptations to acclimate to salinity at the
13 physiological level (plant growth, ion accumulation, water relations, gas exchange, chlorophyll
14 fluorescence and anatomical changes), and at the biochemical level (antioxidative metabolism).
15 Furthermore, the post-recovery period seems to be detected by *Eugenia* plants as a new stress situation, as
16 observed through effects on plant growth and alterations in chlorophyll fluorescence and oxidative stress
17 parameters.
18
19
20
21
22
23
24
25

26 Growth and ion accumulation

27
28 Tolerance to salt stress is a complex phenomenon that enables plants to adapt via different
29 physiological and biochemical processes (Stepien and Johnson 2009). One of the most prominent effects
30 of salt stress is the reduction in plant growth (Parida and Das 2005). However, the reduction in leaf area
31 as well as the increase in the root DW / shoot DW ratio can be viewed as adaptive mechanisms to salt
32 stress. The reduction in leaf area produces an indirect benefit, because plants can thus limit water loss by
33 transpiration, which in turn can favour the retention of toxic ions in roots, limiting the accumulation of
34 these ions in the aerial part of the plant (Munns and Tester 2008), as occurred in the most severe NaCl
35 treatments. The ability of plants to control salt concentration in their aerial parts, either by salt
36 accumulation in roots, by reduced salt uptake rates and/or by controlled translocation to leaves, can
37 constitute an important mechanism of plant survival under saline conditions (Colmer et al. 2005;
38 Cassaniti et al. 2009). This was the case of *Eugenia* plants, which accumulated high concentrations of Na⁺
39 and Cl⁻ in roots. According to this response, *Eugenia* plants behaved as tolerant to NaCl concentrations
40 up to 44 and 88 mM, especially if we consider that the saline irrigation treatments applied were carried
41 out without any drainage. Our findings agree with a previous study performed by (Cassaniti et al. 2009),
42 who classified *Eugenia* plants as tolerant up to 70 mM NaCl after two months of treatment according to
43 the relative growth rate parameter. After the recovery period, and although the analysed roots were not
44 subjected to “free-space washing”, the concentration of root Na⁺ and especially Cl⁻ strongly decreased.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Other authors also used the same methodology to study ion content and/or concentration in roots. In these
2 cases, roots were washed to remove surface ions (Cassaniti et al., 2009; Álvarez et al. 2012; Acosta-
3 Motos et al. 2014b).

4
5 One of the risks of growing plants in small containers under salt stress conditions is the
6 accumulation of Na^+ and Cl^- ions in the substrate, which can bring about an excessive accumulation of
7 toxic ions in all parts of the plant (Álvarez et al. 2012). In addition, salt stress produced an increase in
8 Ca^{2+} in the different parts of the *Eugenia* plants. The increase in Ca^{2+} concentrations in response to
9 salinity has been reported in other plant species such as *Vicia faba* L. and *Myrtus communis* L. (Gadallah
10 1999; Acosta-Motos et al. 2014b). Although Ca^{2+} concentrations increased in *Eugenia* roots by effect of
11 saline stress, an increase in $\text{Na}^+/\text{Ca}^{2+}$ ratio occurred that could induce an increase in membrane
12 permeability, favouring passive Cl^- and Na^+ transport inside the roots (Greenway and Munns et al. 1980).
13 In contrast, and despite the fact that salt stress reduces K^+ concentrations in all parts of the plants, this
14 decrease was about 30%. The observed increase in Ca^{2+} along with the limited decline in K^+ can be
15 considered important in the response of *Eugenia* plants to salinity conditions in view of the importance of
16 both nutrients in plant growth and development. As well as, in the stomatal response, cellular turgor, cell
17 wall and membrane stability, enzyme activation and cell signalling (Marschner 1995; Osakabe et al.
18 2014).

33 Plant water relations

34
35 The decrease in water potential in NaCl-treated plants can reflect an adaptation in water uptake
36 during the beginning of the stress period as a result of the greater accumulation of salts in the substrate
37 (Álvarez et al. 2012). Such accumulation was more evident in the S8 and S12 treatments. Despite the
38 availability of water in the substrate, salts can promote an osmotic effect in the soil, limiting water uptake
39 (Hardikar and Pandey 2008). This behaviour has been observed in other ornamental species grown under
40 the same conditions (Koyro 2006; Acosta-Motos et al. 2014b). As a response to this osmotic effect, a
41 reduction in evapotranspiration and stomatal conductance occurred during the stress period, acting as a
42 mechanism to prevent excessive loss of water (Munns and Tester 2008), particularly in the plants
43 subjected to the highest saline concentrations. Ψ_r data reflected the accumulation of toxic ions on the root
44 surface and may have direct effects on the reduction of Ψ_1 in order to guarantee water transport to the
45 leaves.

46
47 The contribution of the ions to osmotic adjustment was different, but the contribution of Na^+ and
48 Cl^- was the most important in NaCl-treated plants. This adjustment by toxic ion accumulation can be
49

positive only if plants have the ability to compartmentalise the ions (Alarcón et al. 1999; Koyro 2006).

This response has also been described in other ornamental plants subjected to salt stress (Sánchez-Blanco et al. 1998; Navarro et al. 2008).

However, a role for proline in osmotic adjustment, although limited, cannot be ruled out. It has been described that proline can act as an osmoprotectant as well as an antioxidant molecule, protecting different macromolecules during dehydration and reducing power storage (Ashraf and Foolad 2007; Planchet et al. 2014).

Gas exchange and chlorophyll fluorescence

As mentioned above, the aerial parts of the *Eugenia* plants studied were reduced, but chlorophyll levels on the other hand increased as a strategy to protect the photosynthetic machinery. It is known that salt-tolerant species show increased or unchanged chlorophyll content under saline conditions but that chlorophyll levels decrease in salt-sensitive species, suggesting this parameter as a biochemical marker of salt tolerance in plants (Stepien and Johnson 2009; Ashraf and Harris 2013).

At 15 days of stress, an increase in WUE was observed, mainly due to decreased g_s values. However, at longer-term (30 days of stress) S4 plants appeared to adapt to the salinity conditions. Decreases in g_s during the stress period can be also considered as an adaptative mechanism of salt tolerance (Flowers and Yeo 1981). After Phase II, the gas exchange parameters of plants seemed to stabilise, and P_N and g_s even increased in plants previously treated with 4 dS/m NaCl.

Studies investigating the capacity for photosynthetic recovery after a salinity period are very scarce, yet this capacity can determine a plant's resilience to salt stress. Recovery depends on the intensity of photosynthesis decline during the stress period (Chaves et al. 2009). In our data, S12 plants did not show a significant decline in photosynthesis after the recovery period. This response likely allowed these plants to recover photosynthetic rates. However, S12 plants displayed a reduction in plant growth after Phase II, and a role for the accumulation of Na^+ and Cl^- in disturbing cell metabolism cannot be ruled out.

The response of *Eugenia* plants to NaCl was also reflected in the chlorophyll fluorescence parameter, data that were parallel with P_N and g_s changes. In general, salt-sensitive plants show a drop in photochemical quenching parameters but an increase in non-photochemical quenching parameters (Moradi and Ismail 2007; Lee et al. 2013; Ikbal et al. 2014). However, and depending on the plant species and the severity of the stress, a decrease in photochemical and non-photochemical quenching parameters

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

can take place. In *Eugenia* plants, after 15 days of salt treatments, plants subjected to 8 and 12 dS/m NaCl responded to the imposed stress with decreases in qP and Y(II) and a concomitant increase in the non-photochemical quenching parameters, a mechanism for safely dissipating excess light energy and minimising ROS generation (Maxwell and Johnson 2000). At 30 days of stress, the increase in qP and Y(II) and the decrease in the non-photochemical quenching parameters observed in salt-treated plants paralleled the response observed in gas exchange parameters, indicating an adaptative response to the imposed stress conditions. The recovery period was detected by plants as a new challenge, as evidenced by an alteration in the fluorescence parameters, especially in S8 and S12 plants. The observed decrease in qP as well as the drop in non-photochemical quenching parameters in this period suggested the generation of ROS in the chloroplasts as well as photooxidative damage (Foyer and Harbison 1994), a response similar to that in NaCl-sensitive plants.

Anatomical changes

It is known that prolonged water and salt stress may cause changes in leaf anatomy (Olmos et al. 2007; Fernández-García et al. 2014). In this study, the observed morphological changes at 30 days of stress (increased root/canopy ratio) were accompanied by leaf anatomical changes. For example, there was an increase in the percentage of intercellular spaces observed in all stressed plants, which allows for better CO₂ diffusion. In addition, S8 plants experienced an increase in palisade parenchyma, involving an increase in the number of chloroplasts and a reduction in spongy parenchyma, making it easier for CO₂ to reach the chloroplasts present in the palisade parenchyma. These changes were reflected in the P_N and g_s values. After 30 days of stress, although g_s decreased in plants treated with 8 and 12 dS/m NaCl, the anatomical changes made it possible for CO₂ to reach the chloroplast in a more efficient manner in a situation of reduced stomatal aperture. These alterations seem to be another strategy to protect the photosynthetic process. The same anatomical changes also took place in Phase II, especially in S4 plants. These changes correlated with the best P_N performance in the recovery period.

Information regarding the effect of salinity on the leaf anatomy of ornamental plants is very scarce. One study found that the leaf structure of *Rosmarinus officinalis* L. plants was modified in response to water stress, including a reduction in the intercellular spaces in the spongy mesophyll (Olmos et al. 2007). Salt stress also produced anatomical alterations in other shrub species. In *Lawsonia inermis* L. plants, a 150 mM NaCl treatment produced a significant increase in leaf thickness due to a higher mesophyll cell area as an strategy to maximise photosynthesis potential (Fernández-García et al. 2014).

Antioxidative metabolism

In this study, salt stress was found to produce oxidative stress, as evidenced by damage in membranes, ROS accumulation and changes in antioxidative metabolism. Nevertheless, the response of S4 and S8 plants to salt stress was somewhat different. At 30 days of stress, the induction of an H₂O₂-generating enzyme (SOD) was observed in S8 plants in addition to a decrease in H₂O₂-scavenging enzymes (APX, POX and CAT), which would entail the accumulation of H₂O₂. However, DAB-staining did not show significant H₂O₂ accumulation in leaves. In fact, only S12 plants showed some H₂O₂ staining in leaves (data not shown). The S4 plants showed a more balanced ASC-GSH cycle than S8 plants with higher APX activity, unchanged MDHAR levels and an increase in GR activity. In addition, S4 plants presented similar SOD values to S8 plants, but higher CAT and POX activities, suggesting tightly controlled ROS generation. In general, salt-tolerant plants show increased levels of antioxidant mechanisms, including enzymatic and non-enzymatic defences, whereas salt-sensitive species display a decreased response in antioxidative defences (Hernández et al. 1995; Moradi and Ismail 2007; Diaz-Vivancos et al. 2013; Lee et al. 2013; Shu et al. 2013; Ikbal et al. 2014).

Salt stress affects the ASC content, but an increase in reduced glutathione (GSH) occurred. Different authors (Hernández et al. 1999, 2000; Mittova et al. 2003; Diaz-Vivancos et al. 2013) have suggested a role for ASC in salt tolerance. In addition to playing a significant role in the protection and regulation of photosynthesis, ASC also plays an important role as a co-factor of many enzymes (Gest et al. 2013). At 30 days of stress, ASC decreased by up to 30% in S4 plants. In S8 plants there was a nearly 21% decrease in ASC, which correlated with an increase in the ASC-recycling enzyme MDHAR. In salt-tolerant plants, ASC levels can also suffer a decrease ranging from 30-35% due to salinity, as observed in salt-tolerant pea plants or in salt-tolerant transgenic plum lines (Hernández et al. 2000; Diaz-Vivancos et al. 2013). Eugenia plants seemed to use GSH instead of ASC to tackle salt stress. Reduced glutathione can be used not only in H₂O₂ elimination but also to eliminate other peroxides (lipid peroxides or hydroperoxides) by GST and/or GPX enzymes (Noctor et al. 2012). It has been reported that glutathione-dependent enzymes, such as GST and GPX, play a crucial role in the limitation of oxidative processes under salt stress conditions (Roxas et al. 2000; Naliwajski and Skłodowska 2014). It is important to remark that in Eugenia plants the increase in GSH was not accompanied by changes in GR activity, suggesting that GSH biosynthesis could be enhanced. In contrast, in Phase II, Eugenia plants seemed to use both ASC and GSH to respond to the new imposed growth conditions. It is important to highlight the

1 strong increase in ASC levels as well as the restoration of APX activity in recovered plants in relation to
2 Phase I. Recovered plants could use both ASC-dependent and GSH-dependent mechanisms to control
3 ROS metabolism.
4

5
6 Surprisingly, after stress release (Phase II), plants previously treated with 8 or 12 dS/m NaCl
7 behaved as salt-sensitive according to the foliar area, root DW and the lipid peroxidation data. It is likely
8 that the new irrigation conditions produce hypoosmotic stress, leading to an oxidative burst inducing cell
9 damage (Cazalé et al. 1998). This response may be due the fact that plants, once adapted to NaCl stress,
10 can detect new growth conditions as a new challenge. However, literature regarding the removal of salt
11 stress is scarce. This response has also been described in pea leaves in response to short-term salt stress
12 and after 8 h of the post-stress period, suggesting that plants can perceive the removal of NaCl as another
13 stress situation (Hernández and Almansa 2002).
14
15
16
17
18
19
20
21

22 In response to the new conditions, previously stressed plants exhibited the highest values for
23 CAT and SOD activity and recovered APX activity values. In pea plants recovered from drought or salt
24 stress, an increase in APX, SOD and GR has also been described (Mittler and Zilinskas 1994; Hernández
25 and Almansa 2002). Increased CAT and SOD values were a common response in salt-stressed *Eugenia*
26 plants, especially in recovered plants. The response of CAT activity suggested that the photo-respiratory
27 pathway can be induced under salinity conditions, whereas SOD is considered to act as the 'first line of
28 defence' against oxidative stress in plants (Alscher et al. 2002). Photorespiration can supply electron
29 acceptors to PSI and CO₂ for the chloroplast from the decarboxylation of glycine in the mitochondria
30 (Halliwell and Gutteridge 2000). In addition, a close correlation between CAT activity and the
31 photosynthetic rate has been described. Increased CAT activity has been found to reduce the
32 photorespiratory loss of CO₂ by limiting the H₂O₂-dependent decarboxylation of the keto-acids glyoxylate
33 and hydroxypyruvate in the peroxisome (Brisson et al. 1998).
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 Conclusions

50
51 Globally, the results of this study showed that *Eugenia* plants are able to withstand salt stress and
52 can be considered for landscaping project in Mediterranean areas characterized by semiarid climatic
53 conditions. *Eugenia* plants react to avoid leaf ion toxicity, to keep their water status in order to limit water
54 loss and protect the photosynthesis process. Other responses implemented by *Eugenia* plants to adapt to
55 salt stress include increases in the root/canopy ratio and in the chlorophyll content in addition to changes
56 in the leaf anatomy. Finally, *Eugenia* plants cope with the established oxidative stress by activating
57
58
59
60
61
62
63
64
65

certain defence mechanisms (Fig. 3). Nevertheless, irrigation with the same water used on the controls for 16 days (Phase II) seems to be detected by *Eugenia* plants as a new stress situation. This can be due to the fact that *Eugenia* plants implement a plethora of mechanisms that have to be reversed once the saline treatment is finished. In other words, the plants have to retrace their steps to behave as control plants, but it appears that they would need more than 16 days to be able to perform once again as control plants.

Acknowledgements

This work was supported by the Spanish Ministry of Economy and Competitiveness co-financed by FEDER funds (Project CICYT AGL 2011-30022-C02-01-02) and by The Fundación Séneca-Agencia de Ciencia y Tecnología de la Región de Murcia (11883/PI/09 and 15356/PI/10). Authors thank Mrs Ansley Evans for correction of the English.

References

- Acosta-Motos JR, Álvarez S, Hernández JA, Sánchez-Blanco MJ (2014a) Irrigation of *Myrtus communis* L. plants with reclaimed water: morphological and physiological responses to different levels of salinity. *J Hortic Sci Biotechnol* 89:487-494
- Acosta-Motos JR, Álvarez S, Barba-Espín G, Hernández JA, Sánchez-Blanco MJ (2014b) Salts and nutrients present in regenerated waters induce changes in water relations, antioxidative metabolism, ion accumulation and restricted ion uptake in *Myrtus communis* L. plants. *Plant Physiol Biochem* 85:41–50
- Alarcon JJ, Morales MA, Torrecilla A, Sánchez-Blanco MJ (1999) Growth, water relations and accumulation of organic and inorganic solutes in the halophyte *Limonium latifolium* cv. *Avignon* and its interspecific hybrid *Limonium caspia* x *Limonium latifolium* cv. *Beltlaard* during salt stress. *J Plant Physiol* 154:795-801
- Aebi H (1984) Catalase in vitro. *Methods Enzymol* 105:121-126
- Alscher RG, Erturk N, Heath LS (2002) Role of superoxide dismutases (SODs) in controlling oxidative stress in plants. *J Exp Bot* 53:1331–1341

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Álvarez S, Gómez-Bellot MJ, Castillo M, Bañón S, Sánchez-Blanco MJ (2012) Osmotic and saline effect on growth, water relations, and ion uptake and translocation in *Phlomis purpurea* plants. *Environ Exp Bot* 78:138–145
- Álvarez S, Sánchez-Blanco MJ (2014) Long-term effect of salinity on plant quality, water relations, photosynthetic parameters and ion distribution in *Callistemon citrinus*. *Plant Biol* 16:757-764
- Arbona V, Flors V, Jacas J, García-Agustín P, Gómez-Cadenas A (2003) Enzymatic and non-enzymatic antioxidant responses of *Carrizo citrange*, a salt-sensitive citrus rootstock, to different levels of salinity. *Plant Cell Physiol* 44:388–394
- Arrigoni O, Dipierro S, Borraccino G (1981) Ascorbate free radical reductase: a key enzyme of the ascorbic acid system. *FEBS Lett* 125:242-244
- Ashraf M, Foolad MR (2007) Roles of glycine betaine and proline in improving plant abiotic stress resistance. *Environ Exp Bot* 59:206-216
- Ashraf M, Harris PJC (2013) Photosynthesis under stressful environments: An overview. *Photosynthetica* 51:163-190
- Bose J, Rodrigo-Moreno A, Shabala S (2014) ROS homeostasis in halophytes in the context of salinity stress tolerance. *J Exp Bot* 65:1241–1257
- Brisson LF, Zelitch I, Haver EA (1998) Manipulation of catalase levels produces altered photosynthesis in transgenic tobacco plants. *Plant Physiol* 116:259–269
- Cassaniti C, Leonardi C, Flowers T (2009) The effects of sodium chloride on ornamental shrubs. *Sci Hort* 122:586–593
- Cazalé AC, Rouet-Mayer MA, Barbier-Brygoo H, Mathieu Y, Laurière C (1998) Oxidative burst and hypoosmotic stress in tobacco cell suspensions. *Plant Physiol* 116: 659–669
- Chaves MM, Flexas J, Pinheiro C (2009) Photosynthesis under drought and salt stress: Regulation mechanisms from whole plant to cell. *Ann Bot* 103:551-560

- 1 Chaves MM, Costa JM, Madeira Saibo NJ (2011) Recent advances in photosynthesis under drought and
2 salinity. In: Turkan I (ed) Advances in botanical research. Plant responses to drought and salinity
3 stress: Development in a post-genomic era, vol. 57. Elsevier Ltd, San Diego, pp 50-103
4
5
6
7 Colmer TD, Muñiz, R, Flowers TJ (2005) Improving salt tolerance of wheat and barley: future prospects.
8 Aus J Exp Agric 45:1425–1443
9
10
11 Corpas FJ, Gómez M, Hernández JA, del Río LA (1993) Metabolism of activated oxygen in peroxisomes
12 from two *Pisum sativum* L. cultivars with different sensitivity to sodium chloride. J Plant Physiol
13 141:160-165
14
15
16
17
18 Dalton DA, Baird LM, Langeberg L, Taugher CY, Anyan WR, Vance CV, Sarath G (1993) Subcellular
19 localization of oxygen defense enzymes in soybean (*Glycine max* L. Merr.) root nodules. Plant
20 Physiol 102:481-489
21
22
23
24
25
26 Diaz-Vivancos P, Dong YP, Ziegler K, Markovic J, Pallardó FV, Pellny T, Verrier P, Foyer CH (2010)
27 Recruitment of glutathione into the nucleus during cell proliferation adjusts whole cell redox
28 homeostasis in *Arabidopsis thaliana* and lowers the oxidative defence shield. Plant J 64:825–838
29
30
31
32
33 Diaz-Vivancos P, Faize M, Barba-Espin G, Faize L, Petri C, Hernández JA, Burgos L (2013) Ectopic
34 expression of cytosolic superoxide dismutase and ascorbate peroxidase leads to salt stress
35 tolerance in transgenic plums. Plant Biotech J 11:976–985
36
37
38
39
40 Duarte B, Santos D, Marques JC, Caçador I (2013) Ecophysiological adaptations of two halophytes to salt
41 stress: Photosynthesis, PS II photochemistry and anti-oxidant feedback. Implications for resilience
42 in climate change. Plant Physiol Biochem 67:178-188
43
44
45
46
47 Edwards EA, Rawsthorne S, Mullineux PM (1990) Subcellular distribution of multiple forms of
48 glutathione reductase in leaves of pea (*Pisum sativum* L.). Planta 180:278-284
49
50
51
52 Fernández-García N, Olmos E, Bardisi E, García-De la Garma J, López-Berenguer C, Rubio-Asensio JS
53 (2014) Intrinsic water use efficiency controls the adaptation to high salinity in a semi-arid adapted
54 plant, henna (*Lawsonia inermis* L.). J Plant Physiol 171:64-75
55
56
57
58
59 Flower TJ, Yeo AR (1981) Variability in the resistance of sodium chloride salinity within rice (*Oryza*
60 *sativa* L.) varieties. New Phytol 88:363-373
61
62
63
64
65

- 1 Foyer CH, Harbison J (1994) Oxygen metabolism and the regulation of photosynthetic electron transport.
2 In: Foyer CH, Mullineaux P (eds) Causes of photooxidative stresses and amelioration of defense
3 systems in plants. CRC Press, Boca Raton, Florida, USA, pp 1-42
4
5
6
7 Gadallah MAA (1999) Effects of proline and glycinebetaine on *Vicia faba* responses to salt stress. Biol
8 Plant 42:249-257
9
10
11 Gest N, Gautier H, Stevens R (2013) Ascorbate as seen through plant evolution: the rise of a successful
12 molecule? J Exp Bot 64:33-53
13
14
15
16 Gil R, Bautista I, Boscaiu M, Lidón A, Wanklade S, Sánchez H, Llinares J, Vicente O (2014) Responses
17 of five Mediterranean halophytes to seasonal changes in environmental conditions. AoB Plants 6:
18 doi: 10.1093/aobpla/plu049
19
20
21
22
23 Gómez JM, Hernández JA, Jiménez A, del Río LA, Sevilla F (1999) Differential response of
24 antioxidative enzymes of chloroplasts and mitochondria to long-term NaCl stress of pea plants.
25 Free Rad Res 31(suppl.):11-18
26
27
28
29
30 Greenway H, Munns R (1980) Mechanism of salt tolerance in non- halophytes. Annu Rev Plant Phys
31 31:149-190
32
33
34
35 Gueta-Dahan Y, Yaniv Z, Zilinskas BA, Ben-Hayyim G (1997) Salt and oxidative stress: similar and
36 specific responses and their relation to salt tolerance in Citrus. Planta 203:460-469
37
38
39
40 Halliwell B, Gutteridge JMC (2000) Free radicals in biology and medicine. Oxford University Press,
41 London
42
43
44
45 Hardikar SA, Pandey AN (2008) Growth, water status and nutrient accumulation of seedling of *Acacia*
46 *senegal* (L.) Willd. in response to soil salinity. An Biol 30:17-28
47
48
49
50 Hernández JA, Olmos E, Corpas FJ, Sevilla F, del Río LA (1995) Salt-induced oxidative stress in
51 chloroplast of pea plants. Plant Sci 105:151-167
52
53
54
55 Hernández JA, Campillo A, Jiménez A, Alarcón JJ, Sevilla F (1999) Response of antioxidant systems and
56 leaf water relations to NaCl stress in pea plants. New Phytol 141:241-251
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Hernández JA, Jiménez A, Mullineaux PM, Sevilla F (2000) Tolerance of pea (*Pisum sativum* L.) to long-term salt stress is associated with induction of antioxidant defenses. *Plant Cell Environ* 23:853-862
- Hernández JA, Ferrer MA, Jiménez A, Ros-Barceló A, Sevilla F (2001) Antioxidant systems and O₂⁻/H₂O₂ production in the apoplast of *Pisum sativum* L. leaves: its relation with NaCl-induced necrotic lesions in minor veins. *Plant Physiol* 127:817-831
- Hernández JA, Almansa MS (2002) Short-term effects of salt stress on antioxidant systems and leaf water relations of pea plants. *Physiol Plant* 115:251-257
- Hernández JA, Aguilar A, Portillo B, López-Gómez E, Mataix-Beneyto J, García-Legaz MF (2003) The effect of calcium on the antioxidant enzymes from salt-treated loquat and anger plants. *Funct Plant Biol* 30:1127-1137
- Hossain MA, Asada A (1984) Inactivation of ascorbate peroxidase in spinach chloroplasts on dark addition of hydrogen peroxide: its protection by ascorbate. *Plant Cell Physiol* 25:1285-1295
- Ikbāl FE, Hernández JA, Barba-Espín G, Koussa T, Aziz A, Faize M, Diaz-Vivancos P (2014) Enhanced salt-induced antioxidative responses involve a contribution of polyamine biosynthesis in grapevine plants. *J Plant Physiol* 171:779-788
- Jones HG (1983) Estimation of an effective soil water potential at the root surface of transpiring plants. *Plant Cell Environ* 6:671-674
- Koyro HW (2006) Effect of salinity on growth, photosynthesis, water relations and solute composition of the potential cash crop halophyte *Plantago coronopus* (L.). *Environ Exp Bot* 56:136-146
- Lee MH, Cho EJ, Wi SG, Bae H, Kim JE, Cho JY, Lee S, Kim JH, Chung BY (2013) Divergences in morphological changes and antioxidant responses in salt-tolerant and salt-sensitive rice seedlings after salt stress. *Plant Physiol Biochem* 70:325-335
- McCord JM, Fridovich I (1969) Superoxide dismutase. An enzymic function for erythrocyte hemocuprein (hemocuprein). *J Biol Chem* 244:6049-6055
- Marschner H (1995) Mineral nutrition of higher plants, 2nd ed, Academic Press, London
- Maxwell K, Johnson GN (2000) Chlorophyll fluorescence: a practical guide. *J Exp Bot* 51:659-668

- 1 Mehta P, Jajoo A, Mathur S, Bharti S (2010) Chlorophyll a fluorescence study revealing effects of high
2 salt stress on photosystem II in wheat leaves. *Plant Physiol Biochem* 48:16-20
3
4
5 Mittler R, Zilinskas BA (1994) Regulation of pea cytosolic ascorbate peroxidase and other antioxidant
6 enzymes during the progression of drought stress and following recovery from drought. *Plant J*
7
8 5:397-405
9
10
11 Mittal S, Kumari N, Sharma V (2012) Differential response of salt stress on *Brassica juncea*:
12 photosynthetic performance, pigment, proline, D1 and antioxidant enzymes. *Plant Physiol*
13
14 *Biochem* 54:17-26
15
16
17
18 Mittova V, Tal M, Volokita M, Guy M (2003) Up-regulation of the leaf mitochondrial and peroxisomal
19 antioxidative systems in response to salt-induced oxidative stress in the wild salt-tolerant tomato
20 species *Lycopersicon pennellii*. *Plant Cell Environ* 26:845-856
21
22
23
24
25 Moradi F, Ismail AM (2007) Responses of photosynthesis, chlorophyll fluorescence and ROS-scavenging
26 systems to salt stress during seedling and reproductive stages in rice. *Ann Bot* 99:1161-1179
27
28
29
30 Munns R, Weir R (1981) Contribution of sugars to osmotic adjustment in elongating and expanded zones
31 of wheat leaves during moderate water deficits at two light levels. *Aust J Plant Physiol* 8:93-105
32
33
34
35 Munns R, Tester M (2008) Mechanisms of salinity tolerance. *Ann Rev Plant Biol* 59:651-681
36
37
38 Naliwajski MR, Skłodowska M (2014) The oxidative stress and antioxidant systems in cucumber cells
39 during acclimation to salinity. *Biol Plant* 58:47-54
40
41
42
43 Navarro A, Bañón S, Conejero W, Sánchez-Blanco MJ (2008) Ornamental characters, ion accumulation
44 and water status in *Arbutus unedo* seedlings irrigated with saline water and subsequent relief and
45 transplanting. *Environ Exp Bot* 62:364–370
46
47
48
49
50 Nobel PS (1983) *Biophysical plant physiology and ecology*. Freeman and Company, New York, USA, pp
51 61-79
52
53
54
55 Noctor G, Mhamdi A, Chaouch S, Han Y, Neukermans J, Marquez-Garcia B, Queval G, Foyer CH
56 (2012) Glutathione in plants: an integrated overview. *Plant Cell Environ* 35:454–484
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Olmos E, Sánchez-Blanco MJ, Ferrández T, Alarcón JJ (2007) Subcellular effects of drought stress in *Rosmarinus officinalis*. *Plant Biol* 9:77–84
- Osakabe Y, Yamaguchi-Shinozaki K, Shinozaki K, Phan Tran LS (2014) ABA control of plant macroelement membrane transport systems in response to water deficit and high salinity. *New Phytol* 202:35-49
- Ozgun R, Uzilday B, Sekmen AH, Turkan I (2013) Reactive oxygen species regulation and antioxidant defence in halophytes. *Funct Plant Biol* 40:832–847
- Parida AK, Das AB (2005) Salt tolerance and salinity effects on plants: a review. *Ecotoxicol Environ Saf* 60:324-349
- Pellny TK, Locato V, Vivancos PD, Markovic J, De Gara L, Pallardó FV, Foyer CH (2009) Pyridine nucleotide cycling and control of intracellular redox state in relation to poly (ADP-ribose) polymerase activity and nuclear localisation of glutathione during exponential growth of *Arabidopsis* cells in culture. *Mol Plant* 2:442-456
- Pérez-Clemente RM, Montoliu A, Izquierdo-Zandalinas S, De Ollas C, Gómez-Cadenas A (2012) Carrizo citrange plants do not require the presence of roots to modulate the response to osmotic stress. *ScientificWorldJournal* Article ID 795396, 13 pages. doi:10.1100/2012/795396
- Pitman MG (1975) Ion transport in whole plants. In: Baker DA, Hall JL (eds) *Ion transport in plant cells and tissues*. North-Holland Publishing Co, Amsterdam, The Netherlands, pp 267-308
- Planchet E, Verdu I, Delahaie J, Cukier C, Girard C, Morère-Le Paven MC, Limami AM (2014) Abscisic acid-induced nitric oxide and proline accumulation in independent pathways under water-deficit stress during seedling establishment in *Medicago truncatula*. *J Exp Bot* 65:2161-2170
- Romero-Trigueros C, Nortes PA, Pedrero F, Mounzer O, Alarcón JJ, Bayona JM, Nicolás E (2014) Assessment of the viability of using saline reclaimed water in grapefruit in medium to long term. *Span J Agric Res* 12:1137-1148
- Ros-Barceló A (1998) The generation of H₂O₂ in the xylem of *Zinnia elegans* is mediated by an NADPH-oxidase-like enzyme. *Planta* 207:207-216

- 1
2 Roxas VP, Lodhi SA, Garrett DK, Mahan JR, Allen RD (2000) Stress tolerance in transgenic tobacco
3 seedlings that overexpress glutathione S-transferase/glutathione peroxidase. *Plant Cell Environ*
4 41:1229-1234
5
6
7 Sánchez-Blanco MJ, Morales MA, Torrecillas A, Alarcón JJ (1998) Diurnal and seasonal osmotic
8 potential changes in *Lotus creticus* plants grown under saline stress. *Plant Sci* 136:1–10
9
10
11 Shu S, Yuan LY, Guo SR, Sun J, Yuan YH (2013) Effects of exogenous spermine on chlorophyll
12 fluorescence, antioxidant system and ultrastructure of chloroplasts in *Cucumis sativus* L. under salt
13 stress. *Plant Physiol Biochem* 63:209-216
14
15
16
17
18
19 Stepien P, Johnson GN (2009) Contrasting responses of photosynthesis to salt stress in the glycophyte
20 *Arabidopsis* and the halophyte *Thellungiella*: role of the plastid terminal oxidase as an alternative
21 electron sink. *Plant Physiol* 149:1154-1165
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Legend to Figures

Fig. 1 Effect of increased concentrations of NaCl on the uptake rates of Cl⁻ (a), Na⁺ (b), K⁺ (c) and Ca²⁺ (d) ions in *E. myrtifolia* plants at the end of the salinity period (Phase I) and after the recovery period (Phase II). Data represent the mean ± SE from 6 plants. Different letters in the same experimental period indicate significant differences according to Duncan's test ($P \leq 0.05$)

Fig. 2 Concentrations of Cl⁻ (a), Na⁺ (b), K⁺ (c) and Ca²⁺ (d) in different organs of *E. myrtifolia* plants at the end of the salinity period (Phase I) and after the recovery period (Phase II). Data represent the mean ± SE from 6 plants. Different letters in the same experimental period indicate significant differences according to Duncan's test ($P \leq 0.05$)

Fig. 3 Schema showing the effect of long-term salt stress (30 d) on the antioxidative metabolism of *Eugenia* leaves. Under salinity conditions, a decrease in P_N and g_s took place, with an increase in q_P and the electron transport rate and a decrease in q_N . Under these conditions, increases in 1O_2 in PSII and $O_2^{\cdot-}$ in PSI could occur. The recycling of GSH can supply $NADP^+$, which could be considered as an additional response to protect the photosynthetic process in order to minimise ROS generation during the stress period. The increase in SOD activity and the drop in APX activity and ASC content can favour the accumulation of H_2O_2 in different cell compartments as described in other plant species (Corpas et al. 1993; Hernández et al. 1995, 2001; Gómez et al. 1999). In addition, photorespiratory metabolism can be increased and an overproduction of H_2O_2 can occur (Corpas et al. 1993). The H_2O_2 accumulated in chloroplasts, mitochondria and peroxisomes can leak into the cytosol, inducing an oxidative stress. The observed increase in GSH can induce GSH-dependent mechanisms [(Glutathione peroxidase (GPX), Glutathione-S-Transferase (GST)] to control H_2O_2 as well as hydroperoxydes. However, these mechanisms cannot prevent damage to membranes after 30 d of stress

Supporting information

1
2
3 **Suppl. Fig. S1** Contents of Cl⁻ (a), Na⁺ (b), K⁺ (c) and Ca²⁺ (d) in different organs of *E. myrtifolia* plants
4 at the end of the salinity period (Phase I) and after the recovery period (Phase II). Data represent the mean
5 ± SE from 6 plants. Different letters in the same experimental period indicate significant differences
6 according to Duncan's test ($P \leq 0.05$).
7
8
9

10
11
12
13
14 **Suppl. Fig. S2** Influence of the different irrigation treatments on accumulated evapotranspiration (ET) in
15 *E. myrtifolia* plants during stress period (Phase I).
16
17
18
19
20
21

22 **Suppl. Fig. S3** Chlorophyll fluorescence parameters in leaves of *E. myrtifolia* at 15 and 30 days of NaCl
23 stress (Phase I) and after the recovery period (Phase II). Images of the coefficient of photochemical
24 quenching (qP), the effective PSII quantum yield [Y(II)] and the maximal PSII quantum yield (Fv/Fm),
25 the non-photochemical quenching coefficient (qN), non-photochemical quenching (NPQ) and the
26 quantum yield of regulated energy dissipation [Y(NPQ)]
27
28
29
30
31
32
33
34
35

36 **Suppl. Fig. S4** Light microscopy images showing the effect of NaCl on the percentage of area occupied
37 by palisade parenchyma (PP), spongy parenchyma (SP) and intercellular spaces (IS) in leaves from *E.*
38 *myrtifolia* plants at the end of the salinity period. (Phase I: a, control; c, S4; e, S8; g, S12) and after the
39 recovery period (Phase II: b, control; d, S4; f, S8; h, S12)
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig 1











Fig 2



Figure




Table 1 Effect of NaCl on different growth parameters in *E. myrtifolia* plants at the end of the salinity period (Phase I) and after the recovery period (Phase II). Data represent the mean \pm SE from 6 plants. Different letters in the same row indicate significant differences according to Duncan's test ($P \leq 0.05$). Leaf FW, Leaf DW, Leaf Water content, Stem DW and Root DW are given in (g plant^{-1})

Growth parameters	Treatments				
	Control	S4	S8	S12	^a F
(Phase I)					
Total Leaf Area (cm^2)	925 \pm 44b	1105 \pm 63c	775 \pm 10b	549 \pm 55a	27.14***
Leaf FW	23.28 \pm 3.06b	22.27 \pm 2.44b	10.94 \pm 1.74a	10.23 \pm 0.96a	10.32**
Leaf DW	5.05 \pm 0.54bc	6.07 \pm 0.30c	4.28 \pm 0.17b	2.87 \pm 0.22a	15.89***
Leaf Water Content	18.23 \pm 2.64b	16.20 \pm 2.16b	8.45 \pm 1.22a	7.36 \pm 10.80a	8.67**
Stem DW	1.61 \pm 0.32b	1.70 \pm 0.14b	1.33 \pm 0.05ab	0.90 \pm 0.05a	4.18*
Root DW	3.17 \pm 0.56	2.59 \pm 0.10	2.34 \pm 0.18	2.22 \pm 0.09	3.48 n.s.
Root DW / Shoot DW	0.47 \pm 0.03b	0.33 \pm 0.01a	0.48 \pm 0.02b	0.59 \pm 0.04c	17.41***
(Phase II)					
Total Leaf Area (cm^2)	826 \pm 67c	1102 \pm 39d	637 \pm 29b	480 \pm 62a	35.00***
Leaf FW	22.99 \pm 0.74b	27.67 \pm 0.68c	14.59 \pm 1.19a	11.73 \pm 1.54a	45.22***
Leaf DW	6.39 \pm 0.59b	8.98 \pm 0.45c	5.38 \pm 0.06b	3.46 \pm 0.66a	21.30***
Leaf Water Content	16.59 \pm 0.20b	18.69 \pm 0.23c	10.22 \pm 0.75a	8.27 \pm 0.91a	67.37***
Stem DW	2.12 \pm 0.14b	2.97 \pm 0.23c	1.84 \pm 0.10ab	1.27 \pm 0.28a	12.42**
Root DW	3.64 \pm 0.34b	3.73 \pm 0.34b	3.53 \pm 0.26b	2.41 \pm 0.37a	4.48*
Root DW / Shoot DW	0.43 \pm 0.05ab	0.31 \pm 0.01a	0.51 \pm 0.03b	0.52 \pm 0.06b	4.86*

^aF values from one-way ANOVA for the different plant growth parameters analysed. F values were significant at 99.9% (***), 99% (***) or 95% (*) levels of probability. n.s., non-significant values.

Table 2 Effect of increased NaCl levels on soil water potential at the root surface (Ψ_r as MPa), leaf water potential (Ψ_l as MPa), leaf turgor potential (Ψ_t as MPa); leaf osmotic potential at full turgor (Ψ_{100s} as MPa) and proline levels ($\mu\text{mol/g FW}$) after 15 and 30 d of salt treatment (Phase I) and after the recovery period (Phase II) in *E. myrtifolia* plants. Data represent the mean \pm SE from 5 plants. Different letters in the same column indicate significant differences according to Duncan's test ($P \leq 0.05$). For more details, please see Table 1

	Ψ_r	Ψ_l	Ψ_t	Ψ_{100s}	Proline
<i>15 Days(Phase I)</i>					
Control	0d	-0.58 \pm 0.03d	0.64 \pm 0.04	-1.27 \pm 0.03c	8.27 \pm 0.31a
S4	-0.41 \pm 0.06c	-0.73 \pm 0.02c	0.66 \pm 0.09	-1.39 \pm 0.01b	9.17 \pm 0.40ab
S8	-0.58 \pm 0.05b	-0.85 \pm 0.05b	0.60 \pm 0.11	-1.60 \pm 0.03a	9.83 \pm 0.30b
S12	-0.79 \pm 0.04a	-1.00 \pm 0.04a	0.55 \pm 0.09	-1.44 \pm 0.06b	9.77 \pm 0.10b
^a F	52.58***	27.05***	1.37n.s	15.41***	5.81*
<i>30 Days(Phase I)</i>					
Control	0c	-0.63 \pm 0.02d	0.54 \pm 0.02c	-1.03 \pm 0.05b	7.32 \pm 0.24a
S4	-0.22 \pm 0.03b	-0.85 \pm 0.01c	0.52 \pm 0.03c	-1.02 \pm 0.03b	7.67 \pm 0.28a
S8	-0.62 \pm 0.07a	-0.96 \pm 0.02b	0.40 \pm 0.02b	-1.23 \pm 0.07a	9.55 \pm 1.02b
S12	-0.70 \pm 0.10a	-1.12 \pm 0.02a	0.26 \pm 0.03a	-1.13 \pm 0.07ab	9.95 \pm 0.33b
^a F	27.83***	150.91***	29.87***	3.43*	5.42*
<i>Recovery period (Phase II)</i>					
Control	0 \pm b	-0.63 \pm 0.02b	0.32 \pm 0.03	-1.18 \pm 0.11b	6.76 \pm 0.52a
S4	0.16 \pm 0.05b	-0.81 \pm 0.02a	0.32 \pm 0.02	-1.39 \pm 0.01a	7.46 \pm 0.55ab
S8	-0.42 \pm 0.04a	-0.78 \pm 0.03a	0.40 \pm 0.01	-1.55 \pm 0.01a	8.91 \pm 0.24c
S12	-0.43 \pm 0.01a	-0.75 \pm 0.04a	0.29 \pm 0.06	-1.43 \pm 0.01a	8.38 \pm 0.15bc
^a F	18.75**	7.11**	2.35n.s	8.06**	5.48

Table 3 Effect of increased NaCl levels on total chlorophyll content (mg mg^{-1} FW), net photosynthetic rate (P_N as $\mu\text{mol m}^{-2} \text{s}^{-1}$); stomatal conductance (G_s as $\text{mmol m}^{-2} \text{s}^{-1}$); and water use efficiency (WUE as $\mu\text{mol CO}_2 \text{mol}^{-1} \text{H}_2\text{O}$) after 15 and 30 d of salt treatment (Phase I) and after the recovery period (Phase II) in *E. myrtifolia* plants. Data represent the mean \pm SE from 6 plants. Different letters in the same column indicate significant differences according to Duncan's test ($P \leq 0.05$). For more details, please see Table 1

	Total Chlorophyll	P_N	G_s	WUE
<i>15 Days (Phase I)</i>				
Control	1.62 \pm 0.07a	6.76 \pm 0.75b	57.74 \pm 7.07b	123 \pm 14a
S4	1.79 \pm 0.03ab	6.20 \pm 0.77b	39.54 \pm 8.30a	169 \pm 11b
S8	1.81 \pm 0.05ab	5.83 \pm 0.21ab	31.33 \pm 3.65a	197 \pm 17 c
S12	2.04 \pm 0.10b	4.37 \pm 0.22a	23.99 \pm 2.62a	189 \pm 12 bc
^a F	4.19*	3.35*	6.07**	8.82**
<i>30 Days (Phase I)</i>				
Control	1.74 \pm 0.13	5.88 \pm 0.43b	43.60 \pm 2.50b	135 \pm 6 ab
S4	1.69 \pm 0.12	5.58 \pm 0.39 ab	47.62 \pm 4.38b	121 \pm 12a
S8	2.20 \pm 0.06	4.60 \pm 0.44 a	27.42 \pm 2.79 a	170 \pm 11 b
S12	2.07 \pm 0.25	4.48 \pm 0.15 a	31.32 \pm 2.81a	148 \pm 15ab
^a F	1,87n.s	3.66*	9.08**	3.22*
<i>Recovery period (Phase II)</i>				
Control	1.10 \pm 0.03a	6.92 \pm 0.42a	69.64 \pm 11.64a	108 \pm 13 ab
S4	1.37 \pm 0.01ab	9.56 \pm 0.19b	120.16 \pm 11.52b	82 \pm 8a
S8	1.67 \pm 0.04bc	6.06 \pm 0.76a	45.70 \pm 6.37a	139 \pm 16 b
S12	1.61 \pm 0.10c	8.52 \pm 0.35ab	61.66 \pm 4.50a	142 \pm 12 b
^a F	14.32**	10.89***	12.57***	4.99*

Table 4 Effect of increased NaCl levels on fluorescence parameters after 15 and 30 d of salt treatment (Phase I) and after the recovery period (Phase II) in *E. myrtifolia* plants. Data represent the mean from 50 measurements. Different letters in the same column indicate significant differences according to Duncan's test ($P \leq 0.05$). For more details, please see Table 1

	qP	Y(II)	Fv/Fm	qN	NPQ	Y(NPQ)
15 Days (Phase I)						
Control	0.773c	0.473c	0.761b	0.620a	0.273a	0.273a
S4	0.765bc	0.463ab	0.751a	0.639a	0.271a	0.278a
S8	0.754ab	0.469ab	0.765c	0.663b	0.313b	0.302b
S12	0.745a	0.419a	0.758b	0.717c	0.368c	0.344c
^aF	5.54***	19.98***	28.69***	33.44***	36.84***	28.80***
30 Days (Phase I)						
Control	0.754a	0.401a	0.744b	0.743c	0.396c	0.365c
S4	0.829d	0.453b	0.705a	0.644b	0.260a	0.276b
S8	0.769b	0.470c	0.764c	0.633b	0.272b	0.275b
S12	0.805c	0.480c	0.741b	0.606a	0.240a	0.252a
^aF	67.71***	56.59***	86.00***	57.34***	82.38***	69.41***
Recovery period (Phase II)						
Control	0.715b	0.295a	0.682b	0.832c	0.522d	0.476c
S4	0.735b	0.291a	0.647a	0.820c	0.461c	0.458c
S8	0.622a	0.299a	0.706c	0.765b	0.404b	0.431b
S12	0.648a	0.343b	0.730d	0.725a	0.367a	0.391a
^aF	15.32***	9.70***	49.87***	61.21***	43.55***	19.57***

Table 5 Quantitative analysis for morphometric data in leaves from control and NaCl-treated *E. myrtifolia* plants at the end of the salinity period (Phase I) and after the recovery period (Phase II). Data represent the mean \pm SE 10 different sections from each treatment (3 plants of each treatment). Different letters in the same row indicate significant differences according to Duncan's test ($P \leq 0.05$)

For more details, please see Table 1

	Treatments				
	Control	S4	S8	S12	^a F
<i>30 Days (Phase I)</i>					
Palisade parenchyma (%)	36.92 \pm 0.68a	36.16 \pm 1.08a	45.71 \pm 0.74b	36.31 \pm 0.95a	14.64***
Spongy parenchyma (%)	46.57 \pm 0.70b	40.93 \pm 1.80b	31.41 \pm 1.91a	42.92 \pm 1.35b	9.83***
Intercellular space (%)	16.34 \pm 0.83a	21.94 \pm 0.97b	22.89 \pm 1.55b	20.78 \pm 0.75b	6.41**
<i>Recovery period (Phase II)</i>					
Palisade parenchyma (%)	33.85 \pm 0.87a	44.94 \pm 0.83c	43.04 \pm 0.79c	39.27 \pm 1.28b	22.94***
Spongy parenchyma (%)	49.60 \pm 1.42d	30.67 \pm 1.09a	39.79 \pm 1.78b	43.88 \pm 0.19c	40.51***
Intercellular space (%)	16.53 \pm 0.65a	24.43 \pm 0.93b	17.33 \pm 1.49a	16.86 \pm 0.17a	12.34***

Table 6 Effect of increased NaCl levels on oxidative stress parameters in leaves from *E. myrtifolia* plants. Electrolyte leakage (EL) and lipid peroxidation (TBARS) were analysed at the end of the salinity period (Phase I) and after the recovery period (Phase II). Data represent the mean \pm SE from 10 plants. Different letters in the same column indicate significant differences according to Duncan's test ($P \leq 0.05$). For more details, please see Table 1

	EL (%)	TBARS (nmol/g FW)
<i>15 Days (Phase I)</i>		
Control	32.70 \pm 0.34a	2.87 \pm 0.07 a
S4	33.52 \pm 0.45a	3.29 \pm 0.11ab
S8	38.75 \pm 0.69b	3.82 \pm 0.26b
S12	39.45 \pm 0.44b	4.23 \pm 0.38b
^aF	16.49***	6.33*
<i>30 Days (Phase I)</i>		
Control	33.65 \pm 0.64a	3.20 \pm 0.22a
S4	34.33 \pm 0.54a	3.52 \pm 0.19a
S8	39.66 \pm 0.99b	4.12 \pm 0.26b
S12	42.11 \pm 1.22b	4.51 \pm 0.31b
^aF	21.35***	4.92*
<i>Recovery period (Phase II)</i>		
Control	34.07 \pm 1.44a	4.87 \pm 0.16a
S4	34.35 \pm 0.32a	4.45 \pm 0.14a
S8	35.26 \pm 0.59a	5.78 \pm 0.28b
S12	40.48 \pm 1.07b	6.05 \pm 0.35b
^aF	19.85***	10.85**

Table 7 Effect of NaCl on the activity of some antioxidant enzymes in leaves from *E. myrtifolia* plants at the end of the salinity period (Phase I) and after the recovery period (Phase II).Data represent the mean \pm SE from 6 plants. Different letters in the same column indicate significant differences according to Duncan's test ($P \leq 0.05$). For more details, please see Tabl

	CAT $\mu\text{mol min}^{-1} /$ g FW	APX $\text{nmol min}^{-1} /$ g FW	MDHAR $\text{nmol min}^{-1} /$ g FW	GR $\text{nmol min}^{-1} /$ g FW	SOD U/g FW	POX $\mu\text{mol min}^{-1} /$ g FW
<i>15 Days (Phase I)</i>						
Control	8.1 \pm 0.6a	96.1 \pm 0.1b	564.4 \pm 29.2a	34.8 \pm 2.0a	71.4 \pm 3.1a	182.3 \pm 24.1b
S4	19.6 \pm 3.7b	54.2 \pm 6.5a	560.1 \pm 29.3a	37.5 \pm 4.4a	85.1 \pm 2.6a	152.9 \pm 12.4b
S8	14.8 \pm 0.3b	83.1 \pm 4.9ab	646.5 \pm 35.8a	39.5 \pm 8.2a	146.9 \pm 12.1b	56.4 \pm 5.4a
^aF	9.0**	15.4**	2.4n.s	0.1n.s	13.0**	11.9**
<i>30 Days (Phase I)</i>						
Control	13.2 \pm 1.1ab	88.0 \pm 7.2b	274.9 \pm 15.9 a	23.1 \pm 2.4a	100.9 \pm 2.8a	159.0 \pm 15.9b
S4	17.1 \pm 2.3b	47.1 \pm 6.4a	283.0 \pm 19.5 a	39.6 \pm 2.3b	119.0 \pm 6.2b	153.1 \pm 4.6b
S8	11.0 \pm 0.4a	24.4 \pm 4.6a	402.0 \pm 20.5b	26.7 \pm 1.3a	127.3 \pm 8.3b	45.6 \pm 4.6a
^aF	4.1*	20.3***	18.2***	16.2**	5.6*	30.9***
<i>Recovery period (Phase II)</i>						
Control	20.2 \pm 1.0a	40.4 \pm 3.1a	244.7 \pm 21.2b	38.5 \pm 6.3b	168.2 \pm 7.8a	30.7 \pm 1.2a
S4	26.3 \pm 0.54b	42.2 \pm 3.7a	188.1 \pm 5.5a	20.6 \pm 4.4a	232.1 \pm 11.1b	31.4 \pm 2.9ab
S8	27.8 \pm 0.6b	44.7 \pm 3.6a	233.6 \pm 15.7b	32.0 \pm 2.8b	295.8 \pm 5.6c	37.9 \pm 2.4b
^aF	22.1***	3.3*	4.6*	4.19*	41.26***	3.2*

Table 8 Effect of NaCl on the ascorbate and glutathione content in leaves from *E. myrtifolia* plants at the end of the salinity period (Phase I) and after the recovery period (Phase II). Data represent the mean \pm SE from 4 plants. Different letters in the same column indicate significant differences according to Duncan's test ($P \leq 0.05$). For more details, please see Table 1

	GSH nmol/g FW	GSSG nmol/g FW	Redox State GSH/GSHT	Ascorbate μ mol/g FW
<i>15 Days (Phase I)</i>				
Control	1.07 \pm 0.12a	1.62 \pm 0.15a	0.40	6.20 \pm 0.09ba
S4	5.35 \pm 0.13c	1.55 \pm 0.26a	0.79	8.91 \pm 0.46b
S8	2.64 \pm 0.35b	1.12 \pm 0.07a	0.70	7.56 \pm 0.40ab
^aF	41.89***	3.02n.s		6.80*
<i>30 Days (Phase I)</i>				
Control	0.83 \pm 0.25a	3.58 \pm 0.07b	0.20	9.61 \pm 0.28b
S4	1.02 \pm 0.20a	2.73 \pm 0.18a	0.27	6.94 \pm 0.20a
S8	2.29 \pm 0.26b	2.26 \pm 0.21a	0.50	7.50 \pm 0.78a
^aF	10.31**	15.29**		9.32**
<i>Recovery period (Phase II)</i>				
Control	0.86 \pm 0.19a	4.01 \pm 0.16b	0.18	48.04 \pm 8.31b
S4	2.09 \pm 0.28b	2.66 \pm 0.34a	0.44	15.83 \pm 3.93a
S8	2.73 \pm 0.33b	2.55 \pm 0.23a	0.52	11.89 \pm 1.88a
^aF	7.88**	8.00**		12.65**

[Click here to download Supplementary Material: Suppl Fig 1.pdf](#)

[Click here to download Supplementary Material: Suppl Fig 2.pdf](#)

[Click here to download Supplementary Material: Suppl Fig 3.pdf](#)

[Click here to download Supplementary Material: Suppl Fig 4.pdf](#)

[Click here to download Supplementary Material: Suppl. Table S1.docx](#)