

Evidence of low levels of genetic diversity for *Phytophthora austrocedrae* population in Patagonia, Argentina

M.L. Vélez^{a*}, M.P.A. Coetsee^b, M.J. Wingfield^b, M. Rajchenberg^a, A.G. Greslebin^a

^a*Protección Forestal, Centro de Investigación y Extensión Forestal Andino Patagónico (CIEFAP). Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). Universidad Nacional de la Patagonia San Juan Bosco (UNPSJB). Esquel, Argentina. E-mail: mvelez@ciefap.org.ar*

^b*Department of Genetics, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, South Africa.*

*E-mail: mvelez@ciefap.org.ar

ABSTRACT

Phytophthora austrocedrae is a recently discovered pathogen that causes severe mortality of *Austrocedrus chilensis* in Patagonia. The high level of susceptibility of the host tree, together with the distribution pattern of the pathogen in Patagonia, have led to the hypothesis that *P. austrocedrae* was introduced into Argentina. But the genetic diversity of the pathogen that would better inform this question has not been determined. The aim of this study was, therefore, to assess the population structure of *P. austrocedrae* isolates from Argentina in order to gain an understanding of the origin and spread of the pathogen. Genetic diversity was determined based on amplified fragment length polymorphisms (AFLPs). In total, 48 isolates of *P. austrocedrae* were obtained from infected *A. chilensis* trees, and these represented the geographical range of the host. Four primer combinations were used for the AFLP analysis. Of the 332 scored bands, 12% were polymorphic. Gene diversity (h) ranged from 0.01-0.03 and the Shannon index (I) from 0.01-0.04. A high degree of genetic

similarity was observed among the isolates (pairwise S values= 958-1; 0.993 ± 0.009 , mean \pm S.D). A frequency histogram showed that most of the isolate pairs were identical. Principal coordinate analysis using three-dimensional plots did not group any of the isolates based on their geographical origin. The low genetic diversity (within and between sites) and absence of population structure linked to geographic origin, together with the aggressiveness of the pathogen and the disease progression pattern, suggest that *P. austrocedrae* might have been introduced into Argentina.

Keywords: *Austrocedrus chilensis*, “Mal del ciprés”, *Austrocedrus* Root Disease, AFLP analysis, forest Phytophthoras, biological invasions

INTRODUCTION

Austrocedrus chilensis (D. Don) Pic. Serm & Bizarri (ciprés de la cordillera, cypress) is an endemic tree in the Cupressaceae, found in southern Argentina and Chile. It is the most widely distributed species of the small number of conifers found in southern Argentina. *Austrocedrus chilensis* is found across 140,000 ha, in a wide variety of ecological niches, and in different soil types. It grows between 36° 30' and 43° 35' S on the eastern and between 32° 39' and 44° S on the western slopes of the Andes (Veblen et al., 1995). In Argentina, the tree grows in a 60 to 80 km-wide strip along the Andean foothills, across a broad moisture gradient (170 cm/year in the west to 50 cm/year in the east). In the west, *A. chilensis* can be found either in mixed stands with *Nothofagus* spp. or in pure stands on drier sites. In the north, *A. chilensis* trees grow in forests mixed with *Araucaria araucana*. They also grow in open, xeric forests or in isolated clumps at the limit of the Andean forest and the Patagonian steppe, acting as a barrier against the advancing desert. *Austrocedrus chilensis* is important not only because of its ecological functions but because of the high quality of its wood and its aesthetic value.

High levels of *A. chilensis* mortality were first detected in 1948 on Victoria Island (Nahuel Huapi National Park) in Neuquén Province. The disease was later detected during 1953 in a stand located near a forest nursery approximately 150 km from Victoria Island, in Chubut Province. Since then, mortality has extended to almost all of the growth range of *A. chilensis* in Argentina where the disease was referred to as “Mal del Ciprés” (Greslebin & Hansen, 2010).

Phytophthora austrocedrae Gresl. & E.M. Hansen (Pythiales, Peronosporomycetes, Straminipila) was recently described as a new species and is believed to be the primary cause of *A. chilensis* mortality in Patagonia (Greslebin & Hansen, 2010). The disease is generally referred to as *Austrocedrus* Root Disease (ARD) and *P. austrocedrae* is considered to be the main component in the complex pathology that leads to the syndrome known as “Mal del Ciprés” (Greslebin et al., 2007; Greslebin & Hansen, 2010). The pathogen resides in clade 8 of the *Phytophthora* phylogenetic tree of Cooke et al. (2000). Phylogenetic analyses of the nuclear and mitochondrial genes have shown that *P. syringae* and *P. obscura* are the closest relatives of *P. austrocedrae* (Greslebin et al., 2007; Grünwald et al., 2012; Robideau et al., 2011). Among these species, *P. austrocedrae* is homothallic and it is characterized by a combination of very slow growth, semipapillate, non-caducous and non-proliferating sporangia, oogonia with amphigynous antheridia, and low (17.5 °C) optimal temperature for growth (Greslebin et al., 2007).

Phytophthora austrocedrae affects *A. chilensis* trees of all ages and is present across the native growth range of the trees in Argentina (Greslebin & Hansen, 2010). Major symptoms of ARD include chlorosis and a wilting of the foliage and root rot. Trees and especially seedlings can die rapidly, in which case the foliage changes from chlorotic to a red color. Alternatively, infected trees die slowly, first becoming chlorotic followed by a progressive defoliation and eventually tree death after several years (Filip & Rosso, 1999). Studies have shown that the disease originates in the root

systems, is associated with poorly draining and high moisture level soils (Havrylenko et al., 1989; Filip & Rosso, 1999; La Manna & Rajchenberg, 2004). Additional symptoms include necrotic lesions on the roots and stems that can affect the entire breadth of the phloem and the pathogen is able to colonize the xylem, which is evident by superficial discoloration of the woody tissue and the presence of hyphae blocking rays and tracheids in xylem sections (Greslebin & Hansen, 2010, Vélez et al., 2012).

A possible mechanism to explain the decline of *A. chilensis* caused by *P. austrocedrae* has been presented by Vélez et al. (2012). Inoculation studies on 2-year-old saplings with *P. austrocedrae* led to a progressive and significant reduction in net photosynthesis, stomatal conductance and stem-specific hydraulic conductivity. The substantial negative impact on plant physiology was mainly attributed to the extensive death of the bark and cambium tissue, resulting in a disruption of phloem transport. Additionally, blockage of xylem transport by hyphal colonization, presence of resinous plugs, and death of xylem ray parenchyma, leads to a loss of hydraulic conductivity. Involvement of effectors secreted by *P. austrocedrae* or host-derived responses due to the presence of the pathogen was also suggested (Vélez et al., 2012).

The geographic origin of *P. austrocedrae* is unknown although the impact of the pathogen suggests that it has been introduced into Argentina from elsewhere in the world. Until recently, *P. austrocedrae* was not known from any region outside of Argentina, or on any other host than *A. chilensis*. But in 2011, the pathogen was reported as the causal agent of root disease and mortality on *Chamaecyparis nootkatensis* in a public park in East Renfrewshire, Scotland (EPPO Reporting Service N° 6, 2011). This was followed by a report describing a damaging disease in rare native juniper bushes in the Upper Teesdale National Nature Reserve in England (Forestry Commission

Great Britain, 2012). The source of *P. austrocedrae* could not be determined, but it was assumed that the pathogen had been introduced into these areas.

The presence of *P. austrocedrae* outside Argentina has raised concern regarding the origin of the pathogen. Due to its limited geographic range, and the high level of susceptibility of its primary host, it has been hypothesized that *P. austrocedrae* was introduced into Patagonia. However, the population genetic structure of the pathogen has not been established, precluding a more definitive view of this question. The aim of this study was, therefore, to assess the genetic diversity of *P. austrocedrae* in Argentina. The study was also undertaken to gain knowledge that would inform disease management strategies for the pathogen.

MATERIALS AND METHODS

Isolate selection and growth conditions

Isolates of *P. austrocedrae* were obtained from the culture collection of the Centro de Investigación y Extensión Forestal Andino Patagónico (CIEFAP). These isolates were collected during several surveys of declining *A. chilensis* stands in Patagonia, Argentina (**Table 1**). Sampling and isolation methods followed the protocols of Greslebin et al. (2007) and Greslebin & Hansen (2010). Forty-eight isolates from 19 stands were selected for analysis, reflecting the geographical range of *A. chilensis* mortality known at the time of the study, with a distance of approximately 550 km between two most distant locations (**Figure 1**).

In order to verify the identity of the isolates collected, DNA sequence comparisons were made for a sub-set of the isolates using the internally transcribed spacer regions (ITS1 and ITS2) and the cytochrome oxidase I gene (*coxI*). These gene regions were amplified and sequenced for isolates 13-

Phy-309, 15-Phy-271 and 16-Phy-270 (GeneBank accession numbers ITS: JX121857, JX121856, JX121855, *coxI*: JX448319, JX448318, JX448317). DNA was extracted and amplified as outlined below. The ITS region was amplified using primers ITS4 and ITS5 (White et al., 1990) and a region of the *coxI* gene was amplified with primers FM 83 and FM 84 (Martin & Tooley, 2003). PCR reaction mixtures for amplification were the same for both regions. The mixture included dNTPs (0.25 mM of each), MgCl₂ (2.5 mM) PCR buffer, 0.1 μM of each primer, DNA (50 - 100 ng) and *Taq* polymerase produced at the Forestry and Agricultural Biotechnology Institute (2.5 U). The PCR reaction conditions for both regions were the same and followed those described by Martin & Tooley (2003). PCR products were separated on 1 % agarose gel stained with GelRedTM and visualized under UV light. PCR products were purified using a MSB® Spin PCRapace purification kit (Invitex). DNA sequencing reactions were done using a BigDye Terminator v3.1 cycle sequencing kit (ABI) and sequences were determined on an ABI 3100 DNA automated sequencer. The regions were sequenced in both directions using the primers for PCR amplification. Contigs were assembled and edited with the Staden software package (1996). DNA sequences were compared against those available on GenBank using BLASTn.

Isolates of *Phytophthora syringae* were included in this study for comparative purposes. These isolates were obtained from soil samples collected in forests located in different regions of Patagonia using soil baiting. In all, four isolates of *P. syringae* were included as follows: isolate syr-40 was obtained in Corcovado (near site 15, **Figure 1**), syr-157 in Los Alerces National Park (near site 9, **Figure 1**), syr-259 in Lanin National Park (near site 3, **Figure 1**), and syr-319 in Futaleufú, Río Grande valley (near site 13, **Figure 1**).

DNA extraction

Isolates were maintained on clarified tomato juice agar (TA) at 16 °C in the dark (Greslebin et al., 2007). To produce mycelium for DNA extraction, tomato juice broth was inoculated with agar plugs cut from the actively growing margins of fresh cultures. After 30 days of growth at 16 °C, the mycelium was harvested under sterile conditions, washed with sterile DNase/RNase free water and centrifuged to discard the supernatant. Washed mycelium was transferred to Eppendorf tubes and DNA was extracted using the procedure described by Möller et al. (1992) with minor modifications. Mycelium was pulverized in liquid nitrogen and 600 µl of extraction buffer (200 mM Tris-HCl pH 8, 150 mM NaCl, 25 mM EDTA, 50 µg proteinase K, 100 µg RNase A) was added. This mixture was then homogenized at 4 m/s for 20 s using a Fastprep FP120 (QBIogene) homogenizer. The mixture was frozen in liquid nitrogen and incubated for 60 min at 60°C. After incubation, 140 µl 5 M NaCl and 65 µl 10 % CTAB was added, vortexed, and incubated at 60 °C for 10 min. One volume of chloroform/isoamyl alcohol (24:1) was added to the mixture, mixed gently and incubated at 4 °C for 30 min. Samples were centrifuged at 13,800 rpm for 20 min at 4 °C, the supernatant was transferred to new Eppendorf tubes and 0.55 volumes of isopropanol were added followed by centrifugation at 13,800 rpm at room temperature. The supernatant was discarded and the pellet washed with 70 % cold ethanol, at 10,000 rpm for 5 min at 4°C. This step was repeated, the pellet dried and dissolved in 50 µl of DNase/RNase free water. Successful DNA extraction was confirmed by electrophoresis (1 % agarose gel stained with GelRed™ and visualized under UV light). The DNA quality and concentration was determined using a Nanodrop ND1000 spectrophotometer (NanoDrop Technologies Inc.). The final concentration was adjusted to 20 ng/µl.

Amplified Fragment Length Polymorphism (AFLP) analysis

Of the 48 isolates selected for the analysis, five were duplicated in the AFLP analysis to serve as internal controls and to ensure reproducibility. AFLP analysis was performed following the protocol

of Vos et al. (1995) with minor modifications. Restriction digestion of the genomic DNA (100 ng) was done using the endonucleases *EcoRI* and *MseI*. The resulting restriction fragments were ligated to the corresponding enzyme-specific oligonucleotide adapters (Vos et al., 1995). Pre-selective amplifications were performed with zero-base-addition *EcoRI* and *MseI* adapter specific primers using the PCR conditions described in De Vos et al. (2007). Successful restriction, ligation and pre-amplification were confirmed by gel electrophoresis (1 % agarose gel stained with GelRed™ and visualized under UV light). The product of the pre-amplification step was diluted in low TE (10 mM Tris-HCl pH 8, 0.1 mM EDTA; 1:10), and used as template for subsequent amplifications. Final selective amplifications were made using *EcoRI* and *MseI* primers with two-base-additions. Before the final amplification with all the samples, a screening test with five isolates (13-Phy-191, 8-Phy-205, 1-Phy-255, 16-Phy-267, and 5-Phy-290) and 16 primer combinations (E-AC, E-AT, E-AA, and E-TT, in combination of each M-AC, M-CC, M-AG, and M-TG) was done. The *EcoRI* primer was labeled with the infrared dyes, IRDye™ 700 or IRDye™ 800 (LI-COR). PCR conditions were as follows: 13 cycles of 10 s at 94 °C, 30 s at 65 °C with a decrease of 0.7 °C per cycle and 1 min at 72 °C followed by 23 cycles of 10 s at 94 °C, 30 s at 56 °C and 1 min at 72°C with an increase of 1 s per cycle and a final elongation step of 1 min at 72 °C.

AFLP fragment analysis was performed on a model 4200 LI-COR® automated DNA sequencer. Parameters for electrophoresis were set to as follows: 1500 V, 35 mA, 35 W, 45°C, motor speed 3 and signal filter 3. Electrophoresis pre-run time was set to 30 min and run time to 4 h. Digital gel images obtained from the LI-COR system were analyzed and scored manually. Based on the clarity of the resulting fingerprinting profiles (allowing unambiguous scoring), reproducibility and maximal variability obtained between the isolates, four primer combinations (E-AC/M-AC, E-AA/M-TG, E-AC/M-CC, E-CC/M-CC) were selected and used to evaluate the larger group of isolates. The AFLP procedure was replicated from the initial restriction digest step for all study strains.

Each clearly resolved AFLP band was assigned a number based on its migration distance. For each isolate, the presence or absence for all monomorphic and polymorphic bands was scored visually. The presence (1) or absence (0) matrices of each primer combination and a matrix that included all data were analyzed using Popgene version 1.31 (Yeh et al., 1997) to assess the genetic diversity based on Nei's gene diversity index (h) (Nei, 1973) and Shannon's information index (Lewontin, 1972). Values of these indices for each set of primers were compared using One-Way ANOVA. In order to analyze the degree of genetic similarity between isolates, a pairwise distance matrix for each species was generated using the Jaccard, Dice and Simple Matching coefficients of similarity (S) as calculated in Infostat version 2011 (Di Rienzo et al., 2011). Since isolate grouping was shown to be highly similar among the different coefficients, only results from the Dice index are presented in the results section. This coefficient measures the proportion of shared AFLP markers between each pair of isolates, while correcting for the dominance of the AFLP data by disregarding shared absence of bands. Distance dendrograms were then constructed using the Infostat program. *Phytophthora syringae* was used as the outgroup to root the dendrogram. The binary matrix was used as in PAUP, version 4.0b10 (Swofford, 2002) to obtain bootstrap values at the nodes for a tree generated from Nei and Li genetic distances (1000 bootstrap replicates). To test possible grouping of the isolates by provenance and according to variables (band profiles), the binary matrix was also analyzed in a principal coordinate analysis using Infostat software.

RESULTS

In total, after analysis in duplicate, 332 clearly resolved AFLP bands were visually evaluated from the digital images generated using four different primer combinations (**Table 2**). The banding patterns for each of the five duplicated isolates, which served as internal controls, were the same, indicating intra-assay AFLP reproducibility. Of the 332 bands that were scored, only 40 (12 %) were

polymorphic. The total number of bands scored per primer combination ranged from 65 (E-CC/M-CC) to 97 (E-AC/M-AC). The lowest percentage of polymorphic loci (8.3 %) was generated by the E-AC/M-AC primer combination, while the E-AC/M-CC primer combination yielded the highest percentage (16.2 %) of polymorphic loci (**Table 2**). Twenty-four rare alleles (i.e. present in five or less isolates) were identified and none of these could be associated with the region or stands from which the isolates were collected.

Genetic analyses of the bands yielded gene diversities (h) that ranged from 0.01 to 0.03. The Shannon index (I) ranged from 0.01 to 0.04 (**Table 2**). Comparison of the values obtained from the four different primers sets using ANOVA showed that they were not statistically different (h , $P = 0.065$; I , $P = 0.092$).

A high degree of genetic similarity was found for the *P. austrocedrae* isolates. The pairwise similarity (S) values were close to 1, ranging from 0.958 - 1 (0.993 ± 0.009 ; mean \pm S.D.) for the majority of comparisons (**Figure 2**). The largest distances were obtained between isolate 8-Phy-203 and isolates 16-Phy-270 and 15-Phy-271 (0.04), all originating from the Chubut Province. Isolate 8-Phy-203 from Los Alerces National Park, and isolates 16-Phy-270 and 15-Phy-271 were from Corcovado, two sites geographically close to each other (**Figure 1**). Isolates obtained from the most distant locations (Corcovado in the South and San Martín de los Andes in the North) yielded high similarity values (0.969 - 0.998). Isolates originating from the same stand, in general, yielded similarity values (S values: 0.993 ± 0.006) that were comparable with those from different stands (S values: 0.993 ± 0.004).

Dendrograms based on similarity indices from each individual AFLP primer combination were concordant and differed only slightly (data not shown). Data from the four primer combinations

were, therefore, concatenated to generate a dendrogram (**Figure 3**). The isolates resided in one large group that was supported by a 100 % bootstrap value (**Figure 3**). Within this group, 45 isolates grouped in a single cluster (92 % bootstrap support), while three isolates (16-Phy-270, 15-Phy-271 and 13-Phy-309) grouped separately but with no bootstrap support (54 %) (**Figure 3**). DNA sequence comparisons for these 3 isolates with sequences on GenBank showed high similarity (Maximal Identity from 99 to 100 %, E-values $< 6e^{-177}$) to *P. austrocedrae* and therefore was assumed to be *P. austrocedrae*. These isolates derived from stands relatively close to each other in terms of the geographic distribution of the host. Stands in Corcovado (sites 15 and 16) were very close to each other while 'La 106' property in Rio Grande Valley (site 13) is about 60 km from those in Corcovado. Although there seemed to be a geographic pattern in this case, the bootstrap did not support this partition; in addition to this, other isolates obtained in the same stands grouped with isolates collected in other distant stands. This is the case of isolates 13-Phy-191, 13-Phy-195 and 13-Phy-201 from 'La 106' property, and 15-Phy-273 and 16-Phy-267 from the 2 stands in Corcovado (**Figure 3**). Among the 45 isolates that grouped together, a subgroup of 5 isolates from 2 different stands located approximately 90 km from each other could be distinguished (89 % bootstrap support) (**Figure 3**). This subgroup was, in turn, subdivided into two according to isolate provenance (90 % bootstrap support). The subgroup of 5 isolates was obtained from trees that were relatively close to each other (stands with small areas); therefore an effect caused by sampling could have occurred. However, as stated above, it does not seem to be a geographic partitioning since other isolates from the same sites (13-Phy-309 and 8-Phy-209, 8-Phy-211, 8-Phy-213, 8-Phy-215, 8-Phy-338) grouped apart and mixed with isolates obtained in other stands and sites (**Figure 3**). Thus, there was no overall clear partitioning of genetic diversity that corresponded to geographic origin (**Figure 3**). In the principal coordinate analyses, three-dimensional plots did not resolve any clear grouping of isolates on the basis of geographical origin (**Figure 4**).

DISCUSSION

This first study to consider the genetic structure and diversity of *P. austrocedrae* in Argentina. Results from this study revealed high levels of within-species genetic similarity and no evidence of partitioning of genetic diversity among the collection sites. This suggests that the pathogen represents a single population with low heterogeneity in Patagonia and a possible explanation is that *P. austrocedrae* might have been introduced into Argentina. The fact that isolates included in this study were collected from across the broad distribution of *A. chilensis*, during different times of the year and in different years, adds credence to this view.

The fact that the only known host of *P. austrocedrae* in Argentina exhibits high levels of susceptibility is consistent with the behavior of an introduced pathogen, and the results obtained in this study indicate that there is a possibility that *P. austrocedrae* might be exotic to the region. Generally, exotic species, especially near-obligate or obligate pathogens such as *Phytophthora* spp., undergo population “bottlenecks” when they are introduced into new areas (Goodwin, 1997). Therefore, lower genetic variation is expected for these populations in comparison to those where they occur naturally. Moreover, endemic species exhibit population genetic structures that reflect the presence of geographic or environmental barriers, or a combination of both, to gene flow (Goodwin, 1997). In this study, a low genetic variation and absence of geographical differentiation was observed for isolates of *P. austrocedrae*, supporting the notion that it might be introduced into Argentina.

Another plausible explanation for the finding of the low genetic diversity in the pathogen population is that *P. austrocedrae* could recently evolve from another species. With regard to this aspect, *P. syringae* and *P. obscura* are the closest relatives of *P. austrocedrae*, by evaluation of nuclear and

mitochondrial genes (Greslebin et al., 2007; Grünwald et al., 2012; Robideau et al., 2011). Only *P. syringae* was isolated in Patagonia up to the present (Greslebin et al., 2005). More studies are needed to elucidate this possibility.

Previous studies on *Phytophthora* spp. using AFLP analyses, and where the pathogens have been suggested to be introduced into new areas, have reported similar levels of genetic diversity. For example, the percentage of polymorphic loci was slightly lower for *P. austrocedrae* in this study than that reported for *P. nemorosa* in USA forests (Linzer et al., 2009) and for *P. pinifolia* in Chile (Durán et al., 2010). But it was slightly higher than obtained for *P. quercina* (Cooke et al., 2005) in Europe and *P. pseudosyringae* (Linzer et al. 2009) in USA and Europe. These species are homothallic, with the exception of *P. pinifolia* that is sterile, and authors of these studies suggested that they were introduced because of their very low levels of genetic diversity and the lack of geographic structuring that is unexpected for endemic organisms, including those that are able to reproduce homothallically. Although the AFLP primers used in these studies are not exactly the same to those used in this work, the relative number of polymorphic loci can be compared, since it is a parameter that indicate the proportion of variable loci among the total of analyzed loci. Therefore, a similar conclusion that *P. austrocedrae* has been introduced into Patagonia in the present study could be justified.

The first detection of mortality of *A. chilensis* trees occurred during 1948 in Piedras Blancas Bay on Victoria Island in Neuquén Province. The subsequent development of the disease is characteristic of an introduced pathogen that has encountered a highly susceptible host grown over an extended area. Victoria Island is known for the introduction of many exotic woody plants from different continents, especially during the 1920s and 1930s (Simberloff et al., 2002). This, together with the fact that the first appearance of *A. chilensis* mortality occurred in this area, leads us to believe that *P.*

austrocedrae was introduced to the island on infected plants. The results of this study support this view.

The recent discovery that *P. austrocedrae* is causing mortality of *C. nootkatensis* in Scotland and *J. communis* in England (EPPO Reporting Service N° 6, 2011; Forestry Commission Great Britain, 2012) is relevant to the present study. These tree species belong to the same family (Cupressaceae Gray) that *A. chilensis*. *Chamaecyparis nootkatensis* is endemic to the USA and *J. communis* occurs naturally in a wide distribution from the Arctic south to around 30°N latitude in USA, Europe and Asia, in the Northern Hemisphere. The source of *P. austrocedrae* in these areas has not been determined, but it has been suggested that the pathogen was also introduced (EPPO Reporting Service N° 6, 2011). Clearly, it is impossible to determine the origin of *P. austrocedrae* without having isolates from a hypothetical natural host or area of origin. The same situation is true for many *Phytophthora* spp. such as *P. ramorum*, *P. pinifolia*, *P. lateralis*, among others, that have unexpectedly appeared in new areas but for which the likely areas of origin are unknown (Ivors et al., 2004; Durán et al., 2010; Hansen et al., 2000). In this regard, global surveys for *Phytophthora* spp. must continue and, in the case of *P. austrocedrae* that appears to be a conifer-specific pathogen, such surveys should include areas where related conifers occur naturally. For example, the recent discovery of *P. lateralis*, a serious root pathogen of *Chamaecyparis lawsoniana* in western north America, infecting *Chamaecyparis obtusa* var. *formosana*, a native species in Taiwan (Webber et al., 2012), represents a good example of how the area of origin of a *Phytophthora* sp. might be found.

The possibility that *P. austrocedrae* could be native to Patagonia cannot be entirely excluded. The pathogen is a homothallic organism, therefore successful gene flow across the geographic range of the pathogen, together with self-fertile reproduction, could lead also to a population with low genetic diversity. But this hypothesis is not consistent with the high levels of susceptibility of *A. chilensis*

and the disease progression pattern that shows a progression from symptomatic sites to healthy sites (La Manna & Matteucci, 2012).

The high level of genetic similarity found for *P. austrocedrae* isolates in this study could be accounted for if the pathogen had undergone a host shift from another native plant. However, a host jump following anthropogenic introduction (Anderson, 2004; Woolhouse, 2005) would be more plausible when one considers the ecology and epidemiology of other *Phytophthora* spp. (Hansen et al., 2000; Ivors et al., 2004, Brasier et al., 2005; Prospero et al., 2007; Linzer et al., 2009; Durán et al., 2010) and characteristics of emerging infectious diseases of plants in general (Anderson, 2004). It has been shown that host shifts are more likely when the original host and the new host are phylogenetically closely related (Gilbert & Webb, 2007). In this regard, at least nine conifers, phylogenetically related to *A. chilensis* have been introduced into Victoria Island where the disease was first reported and *P. austrocedrae* is found. The areas of origin of these trees would be good targets for surveys aimed at discovering the origin of *P. austrocedrae*.

ACKNOWLEDGEMENTS

This work was supported by Ministerio Nacional de Ciencia, Tecnología e Innovación Productiva (Argentina) and South Africa through an International Cooperation Program (MINCYT_DST SA/10/02.) and Agencia Nacional de Promoción Científica y Técnica (ANPCyT, FONCyT) of Argentina (PICTO 36776, PICT 08-0579). Administración de Parques Nacionales of Argentina and land owners of cypress forests are acknowledged for allowing us to work on their properties. The authors also thank members of the Tree Protection Co-operative Programme (TPCP) and the NRF/DST Centre of Excellence in Tree Health Biotechnology (CTHB), South Africa, for financial support.

REFERENCES

- Anderson PK, Cunningham AA, Patel NG, Morales FJ, Epstein PR, Daszak P, 2004. Emerging infectious diseases of plants: pathogen pollution, climate change and agrotechnology drivers. *Trends in Ecology & Evolution* **19**, 535-544.
- Brasier CM, Beales PA, Kirk SA, Denman S, Rose J, 2005. *Phytophthora kernoviae* sp. nov., an invasive pathogen causing bleeding stem lesions on forest trees and foliar necrosis of ornamentals in the UK. *Mycological Research* **109**, 853-859.
- Cooke DE, Drenth A, Duncan JM, Wagels G, Brasier CM, 2000. A molecular phylogeny of *Phytophthora* and related Oomycetes. *Fungal Genetics and Biology* **30**, 17-32.
- Cooke DEL, Jung T, Williams NA, Schubert R, Oßwald W, Duncan JM, 2005. Genetic diversity of European populations of the oak fine-root pathogen *Phytophthora quercina*. *Forest Pathology* **35**, 57-70.
- De Vos L, Myburg AA, Wingfield MJ, Desjardins AE, Gordon TR, Wingfield BD, 2007. Complete genetic linkage maps from an interspecific cross between *Fusarium circinatum* and *Fusarium subglutinans*. *Fungal Genetics and Biology* **44**, 701-714.
- Di Rienzo JA, Casanoves F, Balzarini MG, Gonzalez L, Tablada M, Robledo CW. InfoStat versión 2011. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina. URL <http://www.infostat.com.ar>
- Durán A, Gryzenhout M, Drenth A, Slippers B, Ahumada R, Wingfield BD, Wingfield MJ, 2010. AFLP analysis reveals a clonal population of *Phytophthora pinifolia* in Chile. *Fungal biology* **114**, 746-752.

European and Mediterranean Plant Protection Organization, 2011. Further findings of *Phytophthora lateralis* and first record of *Phytophthora austrocedrae* in Scotland (GB). Eppo Reporting Service 06-01: 135.

Filip GM, Rosso PH, 1999. Cypress mortality (mal del ciprés) in the Patagonian Andes: comparison with similar forest diseases and declines in North America. *European Journal of Forest Pathology* **29**, 89-96.

Forestry Commission Great Britain, 2012. The Pathology Bulletin Issue 24: News release N°15292. www.forestry.gov.uk

Gilbert GS, Webb CO, 2007. Phylogenetic signal in plant pathogen-host range. *Proceedings of the National Academy of Sciences* **104**, 4979-4983.

Greslebin AG, Hansen EM, Winton L, Rajchenberg M, 2005. *Phytophthora* species from declining *Austrocedrus chilensis* forests in Patagonia, Argentina. *Mycologia* **97**, 218-28.

Greslebin AG, Hansen EM, Sutton W, 2007. *Phytophthora austrocedrae* sp. nov., a new species associated with *Austrocedrus chilensis* mortality in Patagonia (Argentina). *Mycological Research* **111**, 308-316.

Greslebin AG, Hansen EM, 2010. Pathogenicity of *Phytophthora austrocedrae* on *Austrocedrus chilensis* and its relation with *mal del ciprés* in Patagonia. *Plant Pathology* **59**, 604-612.

Goodwin SB, 1997. The population genetics of *Phytophthora*. *Phytopathology* **87**, 462-473.

Grünwald NJ, Werres S, Goss EM, Taylor CR, Fieland VJ, 2012. *Phytophthora obscura* sp. nov., a new species that is a member of the novel *Phytophthora* subclade 8d. *Plant Pathology* **61**, 610-622.

Hansen EM, Goheen DJ, Jules ES, Ullian B, 2000. Managing Port-Orford-Cedar and the introduced pathogen *Phytophthora lateralis*. *Plant Disease* **84**, 4-14.

Havrylenko M, Rosso PH, Fontela SB, 1989. *Austrocedrus chilensis*. Contribución al estudio de su mortalidad en Argentina. *Bosque* **10**, 29-36.

Ivors KL, Hayden KJ, Bonants PJM, Rizzo DM, Garbelotto M, 2004. AFLP and phylogenetic analyses of North American and European populations of *Phytophthora ramorum*. *Mycological Research* **108**, 378-392.

La Manna L, Rajchenberg M, 2004. The decline of *Austrocedrus chilensis* forests in Patagonia, Argentina: soil features as predisposing factors. *Forest Ecology and Management* **190**, 345-357.

La Manna L, Matteucci SD, 2012. Spatial and temporal patterns at small scale in *Austrocedrus chilensis* diseased forests and their effect on disease progression. *European Journal of Forest Research* **131**, 1487-1499.

Lewontin RC, 1972. Testing the theory of natural selection. *Nature* **236**, 181-182.

Linzer RE, Rizzo DM, Cacciola SO, Garbelotto M, 2009. AFLPs detect low genetic diversity for *Phytophthora nemorosa* and *P. pseudosyringae* in the US and Europe. *Mycological Research* **113**, 298-307.

Martin FN, Tooley PW, 2003. Phylogenetic relationships among *Phytophthora* species inferred from sequence analysis of mitochondrially encoded cytochrome oxidase I and II genes. *Mycologia* **95**, 269-284.

Möller EM, Bahnweg G, Sandermann H, Geiger HH, 1992. A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acids Research* **20**, 6115-6116.

Nei M, 1973. Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Sciences* **70**, 3321-3323.

Prospero S, Hansen EM, Grünwald NJ, Winton LM, 2007. Population dynamics of the sudden oak death pathogen *Phytophthora ramorum* in Oregon from 2001 to 2004. *Molecular Ecology* **16**, 2958-2973.

Robideau GP, De Cock AWAM, Coffey MD, Voglmayr H, Brouwer H, Bala K, Chitty DW, Désaulniers N, Eggertson QA, Gachon CMM, Hu C-H, Küpper FC, Rintoul TL, Sarhan E, Verstappen ECP, Zhang Y, Bonants PJM, Ristaino JB, Lévesque CA, 2011. DNA barcoding of oomycetes with cytochrome *c* oxidase subunit I and internal transcribed spacer. *Molecular Ecology Resources* **11**, 1002-1011.

Staden R, 1996. The Staden sequence analysis package. *Molecular Biotechnology* **5**, 233-241.

Simberloff D, Relva MA, Nuñez M, 2002. Gringos en el Bosque: Introduced tree invasion in a *Nothofagus/Austrocedrus* forest. *Biological Invasions* **4**, 35-53.

Swofford DL, 2002. PAUP*. Phylogenetic Analysis Using Parsimony (* and Other Methods), version 4.b10. Sinauer Associates, Sunderland, Massachusetts.

Veblen T, Burns B, Kitzberger T, Lara A, Villalba R, 1995. The ecology of the conifers of southern South America. In: Enright N, Hill R, eds. Ecology of the Southern Conifers. Melbourne, Australia: Melbourne University Press, pp 120-155.

Vélez ML, Silva PV, Troncoso OA, Greslebin AG, 2012. Alteration of physiological parameters of *Austrocedrus chilensis* by the pathogen *Phytophthora austrocedrae*. *Plant Pathology in press*. Doi: 10.1111/j.1365-3059.2011.02585.x

Vos P, Hogers R, Bleeker M, Reijans M, Van der Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M, Zabeau M, 1995. AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research* **23**, 4407-4414.

Webber JF , Vettraino AM, Chang TT, Bellgard SE, Brasier CM, Vannini A, 2012. Isolation of *Phytophthora lateralis* from *Chamaecyparis* foliage in Taiwan. *Forest Pathology* **42**, 136-143.

White TJ, Bruns T, Lee S, Taylor JW, 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MS, Gelfand DH (eds) PCR protocols: a guide to methods and applications, City Publisher, pp 615-322.

Woolhouse MEJ, Haydon DT, Antia R, 2005. Emerging pathogens: the epidemiology and evolution of species jumps. *Trends in Ecology & Evolution* **20**, 238-244.

Yeh F C, Yang R-C, Boyle TBJ, Ye Z-H, Mao JX, 1997. POPGENE, the User-Friendly Shareware for Population Genetic Analysis. University of Alberta, Canada: Molecular Biology and Biotechnology Centre.

FIGURE LEGENDS

Figure 1. Map showing the origin of *Phytophthora austrocedrae* isolates from *Austrocedrus chilensis* in forests of Patagonia. For reference numbers see **Table 1**.

Figure 2. Distribution of Dice coefficients of similarity (Sj) for *Phytophthora austrocedrae*. Frequency histogram showing the distribution of all pairwise Sj for the 48 isolates (n=1128). Frequency indicates percent of comparisons with a given Sj value; Sj values of 1 indicate pairs with 100% similarity.

Figure 3. Dendrogram generated from AFLP distance matrix (1-Sj) for 48 *Phytophthora austrocedrae* isolates obtained along the geographical range of *A. chilensis*. First number of isolates label indicates site of origin (for reference see Table 1). Groups of isolates with zero branch length are comprised of samples with identical AFLP genotypes. Values at the branches are bootstrap values. *Phytophthora syringae* was used as outgroup.

Figure 4. Principal Coordinate Analysis generated in Infostat from the binary matrix obtained from AFLP patterns of 48 *Phytophthora austrocedrae* isolates obtained along the geographical range of *A. chilensis*. First number of isolates label indicates site of origin (for reference see Table 1).




Figure 1




Figure 2

UPGMA (Average linkage)

Distance: (Dice (1-S))




Figure 3




Figure 4