

The influence of volatile semiochemicals from stink bug eggs and oviposition-damaged plants on the foraging behaviour of the egg parasitoid *Telenomus podisi*

M.F.F. Michereff¹, M. Borges¹, M.F.S. Aquino^{1,2},
 R.A. Laumann¹, A.C.M. Mendes Gomes¹ and
 M.C. Blassioli-Moraes^{1*}

¹Embrapa Recursos Genéticos e Biotecnologia (Cenargen), Brasília, Brazil:

²Universidade de Brasília, Brasília, Brazil

Abstract

During host selection, physical and chemical stimuli provide important cues that modify search behaviours of natural enemies. We evaluated the influence of volatiles released by eggs and egg extracts of the stink bug *Euschistus heros* and by soybean plants treated with the eggs and egg extracts on *Telenomus podisi* foraging behaviour. Responses to volatiles were evaluated in Y-tube olfactometers after exposure to (1) one egg cluster for 24 h; (2) plants with eggs laid by the stink bug, tested at 24, 48, and 72 h after treatment; (3) plants with eggs laid artificially, tested at 24, 48, and 72 h after treatment; and (4) plants treated with acetone or hexane extracts of eggs. *Telenomus podisi* was attracted to volatiles emitted by one egg cluster and to acetone extracts of one egg cluster, but not to air or acetone controls. There were no responses to odours of plants treated with eggs or egg extracts. Analysis of acetone extracts of egg clusters by gas chromatography revealed the major components were saturated and unsaturated fatty acids, including hexadecanoic acid, linoleic acid, and (*Z*)-9-octadecenoic acid. Our results suggest that one egg cluster and the acetone extract of one egg cluster contain volatile compounds that can modify *T. podisi* foraging behaviour, and that the amounts of these compounds, probably together with some minor compounds, are important for host recognition by *T. podisi*. Also, the oviposition damage or egg extracts on the plant did not elicit indirect defences that attracted *Telenomus podisi*.

Keywords: Egg extract, *Euschistus heros*, searching behaviour, oviposition damage

(Accepted 7 May 2016; First published online 14 June 2016)

Introduction

The parasitoid host selection process involves two steps, host location and recognition, both of which involve physical and chemical stimuli. Both herbivory-induced plant volatiles (HIPVs) and oviposition-induced plant volatiles (OIPVs) are used by parasitoids to locate their hosts (Vinson, 1998;

Wegener *et al.*, 2001; Büchel *et al.*, 2011). Parasitoids use distance-related cues when searching for hosts. At first they use HIPVs and OIPVs as long-range cues, and then they depend on short-range cues such as chemicals from the host, vibration, and visual signals, when they are closer to a potential host (Borges *et al.*, 1998; Laumann *et al.*, 2009, 2011; Aquino *et al.*, 2012). HIPVs and OIPVs are released by plants as ‘calls for help’—host defences—that attract natural enemies of the herbivore. Plants respond to herbivore injury when the damage occurs and to oviposition prior to damage produced by immature insects. The ability to respond to the presence of eggs is an important plant defence (Hilker *et al.*, 2005; Hilker & Meiners, 2006; Bruce *et al.*, 2010). To accomplish this, plants

*Author for correspondence

Phone: +55 (61) 3448-4932

Fax: +55 (61) 3340-3462

E-mail: carolina.blassioli@embrapa.br

need to recognize cues from the eggs and from the egg–plant interaction, which can be physical, such as weight or a shadow on the leaf surface; or chemical, such as kairomones released by eggs or synomones released by plants with deposited eggs (Hilker & Meiners, 2010, 2011).

Parasitoids can also react to cues from host eggs. *Edovum puttleri* (Hymenoptera: Eulophidae) is attracted to kairomones from eggs of *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae) (Leonard *et al.*, 1987; Hu *et al.*, 1999). Female *Anaphesiole* (Hymenoptera: Mymaridae) recognize chemicals derived from eggs or adults of *Lygus hesperus* (Heteroptera: Miridae) during host recognition (Takasu & Nordlund, 2001); and females of the egg parasitoid *Telenomus euproctidis* (Hymenoptera: Platygasteridae) are attracted to volatiles from egg masses of *Orgyia postica* (Lepidoptera: Lymantriidae) (Arakaki *et al.*, 2011).

Telenomus podisi (Hymenoptera: Platygasteridae) is found in croplands of the Americas and is the most common natural enemy of the stink bug *Euschistus* sp. (Heteroptera: Pentatomidae) (Medeiros *et al.*, 1997, 1998; Pacheco & Corrêa-Ferreira, 2000; Michereff *et al.*, 2015). It is attracted to its primary host, *E. heros*, by chemical cues including sex pheromones, defensive compounds from the methatoracic gland, and cuticular compounds (Borges & Aldrich, 1994; Borges *et al.*, 1998, 1999, 2003; Laumann *et al.*, 2009), HIPVs and OIPVs (Moraes *et al.*, 2005, 2008; Michereff *et al.*, 2011, 2013), vibratory signals (Laumann *et al.*, 2007, 2011), and visual cues (Aquino *et al.*, 2012). *T. podisi* is an egg parasitoid, but it is not attracted to volatiles released by soybean plants with *E. heros* eggs (Moraes *et al.*, 2008; Michereff *et al.*, 2011). Similarly, oviposition by the stink bug *Nezara viridula* (Heteroptera: Pentatomidae) does not induce changes in plant odours that attract the egg parasitoid *Trissolcus basalus* (Hymenoptera: Platygasteridae) (Colazza *et al.*, 2004a). Under natural conditions, stink bugs feed on plants while ovipositing, and egg parasitoids are attracted by the HIPVs that result. The absence of OIPVs might explain the lack of response to volatiles released by plants that are treated only by oviposition by stink bugs. Other studies have shown that stink bug eggs have chemicals that *Platygasteridae* parasitoids use to locate their hosts (Bin *et al.*, 1993), and there is evidence that the eggs themselves have volatiles that attract natural enemies (Sales, 1979; Tognon *et al.*, 2014). Therefore, the aim of this study was to determine whether chemicals from stink bug eggs change the foraging behaviour of *T. podisi* and to investigate what happens to this chemical information when the eggs are laid on leaves, as *T. podisi* do not respond to plants with naturally deposited eggs (Colazza *et al.*, 2004a, b; Moraes *et al.*, 2008; Michereff *et al.*, 2011).

Material and methods

Insect rearing

E. heros individuals were obtained from a laboratory colony started from adults collected in soybean fields near Embrapa Genetic Resources and Biotechnology, Brasília, Brazil (15°47'S, 47°55'W). Bugs were reared in 8-litre plastic containers on a diet of soybeans, sunflower (*Helianthus annuus*) seeds, raw peanuts (*Arachis hypogaea*), fresh green beans (*Phaseolus vulgaris*), and water. The food supply was replenished twice weekly. Females and males were separated after mating and maintained in different cages under the same conditions as described above. Racks covered with tulle fabric and placed

inside the cages provided an oviposition substrate, and they were changed every day to avoid egg contamination.

The egg parasitoid *T. podisi* was obtained from a laboratory colony raised on *E. heros* eggs. The wasps were maintained in acrylic cages (25 cm² angled neck tissue culture flasks; ICN Biomedicals, Irvine, CA, USA) and fed twice weekly with a drop of honey. Following hatching, the parasitoids were kept in acrylic cages for 24 h without host eggs before mating. Two-day-old naive females were used in the experiments. *E. heros* and *T. podisi* were maintained in separate environmental rooms at 27 ± 1°C, 65 ± 10% relative humidity, and a 14-h photoperiod.

Plants

Soybean seeds (Dowling cultivar) were germinated on damp paper for 3 days and then transplanted to pots containing a mixture of soil and organic substrate (1:1 w/w). Plants were kept in a greenhouse and used when in the V3 phenological, i.e., stem elongation, stage. The experimental procedures included (1) plants treated with hexane or acetone egg extracts, (2) plants with eggs laid naturally by stink bugs, (3) plants with eggs laid artificially, and (4) untreated control plants.

Preparation of E. heros egg extracts

Extracts for bioassays were prepared from single clusters of approximately 20 eggs. The eggs were submerged in 40 µl of acetone or *n*-hexane solvent in a 2-ml glass vial, and placed in an ultrasonic bath (Unique, USC – 2800A) for 15 min. The solvent was then collected in a Pasteur pipette, passed through a 4 mm diameter 0.45 µm micropore filter (PFTE-4-4 Iso-Disc™, Supelco, USA) and concentrated under gentle nitrogen flow until total solvent evaporation. After drying, 100 µl of Tween-20 (0.1% v/v) was added, and the solution was vortexed for 30 s. For chemical analysis, extracts of 100 *E. heros* egg clusters (~2000 eggs = 1 g) were prepared using a similar procedure. Eggs were placed in a 4 ml glass vial and immersed in 4 mL of solvent. The procedure required 100 egg clusters because the compounds in one egg cluster are present in very small amounts that are not quantifiable using gas chromatography-flame ionization detection (GC-FID) or GC-mass spectrometry (MS). The extracts were prepared in triplicate for each solvent.

Natural and artificial oviposition on soybean plants

Soybean plants used for natural oviposition bioassays had two egg clusters of 10–12 eggs each. The eggs were deposited by six mated females between 12 and 15 days of age, and their proboscises were removed before they were placed on the soybean plants to avoid herbivory injury. Stink bugs were removed from the plants a few minutes before the bioassays began. For artificial oviposition, single egg clusters with 10–12 eggs were placed on the surfaces of the first and second pairs of leaves of the plant by simple touching and without artificial glue. The natural glue on the base of the egg simulated natural oviposition and avoided stimulating plant defences. Untreated soybean plants were used as controls.

Plant treatment with egg extracts

Plants were treated with three sequential applications of 5 μ l hexane or acetone egg extract. Control plants were treated with 15 μ l of solvent in Tween-20 (0.1% v/v).

Bioassays

Y-tube olfactometer bioassays were conducted to determine whether volatiles emitted from eggs, plants treated with eggs, or egg extracts affected parasitoid searching behaviour. The olfactometers consisted of square acrylic blocks (19 \times 19 cm²) with a 1 cm Y-shaped cavity sandwiched between two glass plates (Morales *et al.*, 2005). The leg of the cavity was 8 cm long, and each arm was 7 cm long. Activated-charcoal filtered, humidified air was pushed through the system at 0.6 litre min⁻¹ and pulled out at 0.2 litre min⁻¹ by a push-pull system. Behaviour of the insects was monitored by a CCD camera (Sony SPT M324CE; Sony, Minato-Ku, Tokyo, Japan) with a 12.5–75.0 mm/F1.8 zoom lens and analysed using software architecture comparison analysis method software (Jorge *et al.*, 2005). A single *T. podisi* female was introduced at the base of the Y-tube and observed for 600 s. The first choice arm, which was the first one that the wasp entered and remained in for at least 20 s, and residence time in each arm were recorded by the software. Residence time was the time that the parasitoid remained in an arm. After every five repetitions, the plants were replaced and the positions of the arms of the olfactometer were changed to avoid bias in the parasitoid responses. Each female was used only one time and forty repetitions were made for each combination tested.

Plants were used in the bioassays at 24, 48, and 72 h after treatment. Treated and control plants were kept in different rooms under the same temperature, humidity, and lighting conditions until used in the experiments to avoid chemical signalling between them. Damaged or undamaged plants were placed in glass chambers and connected to the olfactometer via silicone tubing.

Ten-microliter aliquots of test solution were applied to 1 cm² strips of filter paper (80 g m⁻², 205 μ m thick, 14 μ m average pore size; Qualy J Prolab, Paraná, Brazil), which remained at room temperature for 1 min before being inserted into a syringe connected to the arm of the olfactometer. The solvent was allowed to evaporate. The same procedure was performed for the control filter paper strips containing only the *n*-hexane or acetone solvent. The filter paper and the olfactometer system were exchanged after every five assays.

The bioassay combinations that were used to test whether *E. heros* eggs emit volatile compounds that attract *T. podisi* are shown in figure 1. The attraction of volatiles from one egg cluster was evaluated against both air and the volatiles emitted from the extracts of one egg cluster to determine whether attractant volatiles were present in either the acetone or hexane. To evaluate whether chemicals from the secretions released by *E. heros* females during egg oviposition or by the plants might be involved in the attraction of *T. podisi*, bioassays were conducted using plants with eggs laid naturally and artificially and with plants treated with egg extracts.

Scanning electronic microscopy (SEM)

To evaluate whether *E. heros* oviposition physically damages soybean leaves, they were examined by SEM after oviposition. Approximately 1 cm² pieces cut from abaxial and

adaxial sites on fresh leaves with *E. heros* eggs were fixed with glutaraldehyde 2.5% (v/v) in 0.1 M sodium cacodylate buffer (pH 6.8) for 24 h under low pressure. The leaves were then transferred to fresh solution under low pressure for an additional 30 min. The buffer solution was replaced again and kept at 4°C for 90 min. After fixation, the samples were washed five times for 10 min with 0.1 M sodium cacodylate buffer. The cacodylate buffer was replaced by a 2% osmium tetroxide solution for 60 min at room temperature followed by three sequential washes with sodium cacodylate buffer. The leaf samples were dehydrated in an aqueous series of 30, 50, 70, 90, and 100% methanol (v/v); the samples were kept in each concentration for 20 min. After dehydration, the samples were critical-point dried in liquid CO₂ (Baltec CPD 030, Baltec, Schalksmühle, Germany) between 0 and 5°C at atmospheric pressure. The samples were then sputter coated (Emitech K550) with a 20-nm-thick gold film. The images were obtained using either a Zeiss DMS 962 at 10KV with distance of 13–18 mm or a JSM 840 A at 10 KV with distance of 10–15 mm.

Chemical analysis

Egg extracts were analyzed by GC (Agilent 7890A, DB-5MS) with a 30 m \times 0.25 mm ID column and 0.25 μ m film thickness, (J&W Scientific, Folsom, CA, USA), using a temperature program of 50°C (2 min), 5°C min⁻¹ to 180°C (0.1 min), and 10°C min⁻¹ to 250°C (20 min). The column effluent was analyzed with a FID at 270°C. For GC, 50 μ l of each extract was separated, and 1 μ l of octadecane was added as an internal standard (IS) with a final concentration of 9.8 μ g ml⁻¹. One microliter of each sample was injected using the splitless mode with helium as the carrier gas. The amounts of volatile chemicals released by the plant every 24 h were calculated in relation to the area of the IS. Data were collected with EzChrom Elite software (2008) (Agilent, California, USA) and were recorded Excell software (Microsoft, 2007, EUA).

For qualitative analysis, selected extracts were analyzed using an Agilent 5975MSD instrument equipped with a quadrupole analyzer, a nonpolar DB-5MS column (30 m \times 0.25 mm ID and 0.25 μ m film thickness; J&W Scientific, Folsom, CA, USA), and a splitless injector with helium as the carrier gas. Ionization was by electron impact (70 eV and source temperature 200°C). Data were collected and analyzed with GC-MS ChemStation 2.1 Software (2008) (Agilent, California, USA). Substances in the extracts were identified by comparing spectra with library databases (Software NIST-Wiley database, version 2.0, 2008, USA) or published spectra and confirmed using authentic standards when available.

n-Hexane (95%, suitable for pesticide residue analysis) and acetone (ACS reagent >99.5%) were purchased from Sigma-Aldrich (Steinheim, Germany). Hexanoic acid, octanoic acid, decanoic acid, hexadecanoic acid, hexadecanoic acid methyl ester, hexadecanoic acid ethyl ester, (*Z*)-9-octadecenoic acid, octadecanoic acid, and ethyl stearate were purchased from Sigma-Aldrich (St. Louis, MO or Milwaukee, WI, USA). The ocimene mixture of isomers (90%), (-)-linalool 98%, *n*-tridecane (>99%), *n*-hexadecane (>99%), and *n*-octadecane (>99%) were purchased from Sigma-Aldrich (Steinheim, Germany). Limonene was purchased from TCI America (Tokyo, Japan).




Fig. 1. Experimental scheme of plant treatment. Ten-microliter aliquots of egg extract were applied to filter paper. For plants treated with egg extract or solvent, 15 μ l aliquots were applied to filter paper.

Statistical analysis

For each treatment, the parasitoid first choice data were analyzed using logistic regression to estimate the probability of each choice. The model fitted the side (left or right) on which the test odour was presented as the independent variable. The hypothesis of no preference (i.e., the proportion choosing each odour = 0.5) was tested by the chi-square Wald test. The data for the residence time of the parasitoid in each olfactometer arm, treatment, and control were analyzed by the paired *t*-test. If insects had not moved after 3 min, they were considered nonresponders and were not included in the statistical analysis. The analyses were performed using the R-2.8.0 statistical program (R statistical Development Core Team, 2009) and results considered significant if $P < 0.05$.

Results

Bioassays with volatiles from *E. heros* eggs

T. podisi showed a preference for volatiles emitted from one cluster of *E. heros* eggs when compared with air as the first choice ($\chi^2 = 12.120$, DF = 49, $P < 0.001$) (fig. 2a), but there was no significant difference in residence time ($t = -1.880$, DF = 49, $P = 0.06$, fig. 2b).

Bioassay of volatiles from *E. heros* egg extracts

When the egg extracts were tested against the solvents, the parasitoid was selectively attracted to the volatiles emitted from one-cluster acetone extracts compared with acetone as first choice ($\chi^2 = 4.687$, DF = 39, $P = 0.031$) (fig. 2a); the other treatments did not attract parasitoid females (fig. 2a). The residence time did not differ for any of the treatments when compared with their respective controls (one-cluster acetone extract; $t = -1.494$, DF = 39, $P = 0.143$ and one-cluster hexane extract; $t = -0.231$, DF = 39, $P = 0.818$, fig. 2b).

Bioassay of volatiles from plants with natural and artificial oviposition

T. podisi did not distinguish between the volatiles emitted from undamaged soybean plants and those from plants with natural oviposition as first choice at 24 h ($\chi^2 = 0$, DF = 39, $P = 1$); 48 h ($\chi^2 = 0.893$, DF = 39, $P = 0.344$); and 72 h ($\chi^2 = 0.398$, DF = 39, $P = 0.527$; fig. 3a). There were also no differences in residence time at 24 h ($t = -2.00$, DF = 39, $P = 0.058$); 48 h ($t = -0.954$, DF = 39, $P = 0.345$); or 72 h ($t = -1.20$, DF = 39; $P = 0.235$; fig. 3b). The same pattern was

observed following artificial oviposition for first choice at 24 h ($\chi^2 = 0.398$, DF = 39, $P = 0.527$); 48 h ($\chi^2 = 0$, DF = 39, $P = 1$); and 72 h ($\chi^2 = 0.999$, DF = 39, $P = 0.751$, fig. 3c); and for residence time at 24 h ($t = 0.346$, DF = 39, $P = 0.730$); 48 h ($t = -0.533$, DF = 39, $P = 0.597$); and 72 h ($t = -1.712$, DF = 39, $P = 0.094$); fig. 3d).

T. podisi did not show a preference between the volatiles emitted from the plants with natural oviposition at 24 h compared with either the volatiles from one egg cluster as first choice ($\chi^2 = 2.821$, DF = 39, $P = 0.092$) or residence time ($t = -0.372$, DF = 39, $P = 0.711$, fig. 3e, f). The same pattern was observed for the volatiles emitted from the plants with natural oviposition at 24 h compared with either air as first choice ($\chi^2 = 0.716$, DF = 39, $P = 0.397$) or for residence time ($t = -1.265$, DF = 39, $P = 0.211$, fig. 3e, f).

Bioassay of volatiles from plants treated with egg extracts

The volatiles released from plants treated with one-cluster acetone extracts were not more attractive to *T. podisi* than the volatiles released from control plants treated with acetone as assessed by first choice at 24 h ($\chi^2 = 0.398$, DF = 39, $P = 0.527$); 48 h ($\chi^2 = 0.099$, DF = 39, $P = 0.751$); and 72 h ($\chi^2 = 2.446$, DF = 39, $P = 0.117$, fig. 4a). There were also no differences in residence time at 24 h ($t = -1.825$, DF = 39, $P = 0.075$); 48 h ($t = 0.434$, DF = 39, $P = 0.666$); and 72 h ($t = -1.080$, DF = 39, $P = 0.286$, fig. 4b).

T. podisi did not show a preference for volatiles from plants treated with one-cluster hexane extracts over hexane controls in either first choice at 24 h ($\chi^2 = 0.398$, DF = 39, $P = 0.527$); 48 h ($\chi^2 = 0.398$, DF = 39, $P = 0.527$); or 72 h ($\chi^2 = 0.893$, DF = 39, $P = 0.344$) or residence time at 24 h ($t = 0.276$, DF = 39, $P = 0.783$); 48 h ($t = -1.198$, DF = 39, $P = 0.237$); or 72 h ($t = 0.982$, DF = 39, $P = 0.331$, fig. 4c, d).

SEM of surface structure of plants with *E. heros* eggs

E. heros females laid eggs on soybean plants without significant damage to the leaf tissue (fig. 5a, b). When stink bugs lay their eggs, they secrete an adhesive that fixes their eggs onto the leaf surface (fig. 5c). When the eggs are removed, some of the secretion is removed with them, resulting in damage to leaf tissue (fig. 5c). The SEM images showed that egg deposition did not result in necrosis of leaf tissue and that oviposition did not provoke a hypersensitive response in the plants.




Fig. 2. (a) Response of *T. podisi* females to plant treatments. One egg cluster vs. air; one cluster *E. heros* egg extract vs. acetone or hexane. Asterisks indicate significant differences between treatments using the Wald test with distribution χ^2 , 0.05% probability. Numbers in parentheses indicate the insects that did not respond to any treatment. (b) Residence time (seconds) of *T. podisi* females in the Y-tube olfactometer. One egg cluster vs. air; one cluster extract vs. acetone or hexane. The absence of an asterisk indicates that there is no significant difference between the treatments.




Fig. 3. Responses of *T. podisi* females to odours of plants (a) treated with mated *E. heros* females to evaluate oviposition injury, (c) treated with two *E. heros* egg clusters laid on two different leaves, and (e) with natural oviposition vs. eggs or vs. air. The bioassays were performed 24, 48, and 72 h after treatment. Numbers in parentheses indicate the insects that did not respond to any treatment. Residence time (seconds) of *T. podisi* females in Y-tube olfactometers in response to odours of plants with (b) natural oviposition, (d) artificial oviposition and (f) plants with natural oviposition vs. eggs or vs. air. The absence of an asterisk indicates that there is no significant difference between the treatments.

Chemical analysis of egg extracts

Chemical analysis of 1 h acetone extracts revealed 30 peaks that were consistently present in the chromatograms (Supplementary Material Table 1 and fig. 6). The egg parasitoids responded only to volatiles from acetone extracts;

therefore, only compounds extracted by acetone were quantified. The major compounds identified were C_{16} and C_{18} saturated and unsaturated fatty acids, including: hexadecanoic acid; linoleic acid; (*Z*)-9-octadecenoic acid methyl ester, (*Z*)-9,12-octadecadienoic acid methyl ester; methyl stearate; and (*Z*)-9-octadecenoic acid ethyl ester. Several minor




Fig. 4. Responses of *T. podisi* females to odours of plants (a) treated with an acetone extract of *E. heros* egg clusters (treatment) or (c) with a hexane extract of *E. heros* egg clusters (treatment). The bioassays were performed 24, 48, and 72 h after treatment. Numbers in parentheses represent the insects that did not respond to any treatment. Residence time (in seconds) of *T. podisi* females in Y-tube olfactometers in response to odours of plants (b) treated with an acetone extract of an *E. heros* egg cluster or (d) treated with a hexane extract of an *E. heros* egg cluster (treatment). The absence of an asterisk indicates that there is no significant difference between the treatments.

compounds were identified, including short-chain fatty acids from C₆ to C₁₄ and monoterpenes, specifically limonene, (*E*)- β -ocimene, and linalool (Supplementary Material Table 1).

Discussion

The egg parasitoid *T. podisi* was attracted to volatile chemicals from *E. heros* eggs and had a positive taxis. This parasitoid may use volatiles from stink bug eggs for short-range host searching. Several volatile compounds that might be attractants were extracted from *E. heros* eggs. However, when eggs were laid on soybean plants, these compounds did not appear to be noticed, as the parasitoid was not attracted.

T. podisi was not attracted to volatiles emitted by soybean plants when treated naturally or artificially with one cluster of *E. heros* eggs, as has been previously reported (Moraes *et al.*, 2008; Michereff *et al.*, 2011). Different hypotheses have been proposed to explain this. The SEM images showed that when stink bugs laid eggs on the leaves, there was no visible physical damage to the leaf surface that could lead to localized necrosis and hypersensitive responses (Walling, 2000; Kaloshian & Walling, 2005). Only touching the leaf surface did not elicit an indirect plant defence that might attract a natural enemy (Hilker & Meiners, 2006, 2010). Although some studies have shown that the emission of OIPVs was independent of visible foliar damage or hypersensitive responses (Fatouros *et al.*, 2012; Tamiru *et al.*, 2012), the absence of

response of *T. podisi* to odours released from plants with eggs suggests that there no OIPVs were emitted in this system, which included soybean plants, *E. heros*' eggs and *T. podisi*.

Another explanation is that chemicals in the eggs or ovipositor secretions suppressed the indirect defence of soybean plants, as suggested by Moraes *et al.* (2008). In *Arabidopsis thaliana*, elicitor-treatment suppressed plant defence responses, and the accumulation of salicylic acid at the oviposition site was thought to be involved with the suppression (Bruessow *et al.*, 2010). A similar response was observed following *E. heros* oviposition on soybean plants, which enhanced systemic methyl salicylate production compared with other treatments 96 h after egg laying, but *T. podisi* was not attracted to the volatiles emitted following oviposition damage (Moraes *et al.*, 2008; Michereff *et al.*, 2011). The chemical profile of the plants changed after the *E. heros* eggs were laid, but the change did not result in attracting the egg parasitoid *T. podisi*.

Although OIPVs were not induced in this system, it was expected that the parasitoid would respond to volatiles emitted by the eggs that were present on the plants. The absence of response might be explained by the small quantities of compounds emitted by the eggs that were in contact with the plant foliage. Chemical analysis showed that an average of 5.8 ng of volatile compounds was extracted from each *E. heros* egg. This has been less reported than for other species (Tooker & De Moraes, 2007; Liu *et al.*, 2008). In this study, the amount of hexadecanoic acid was 0.16 ng egg⁻¹, but




Fig. 5. Scanning electronic microscopy images of oviposition damage caused by *E. heros* in soybean leaves. (a) and (b) show oviposition sites in cultivar Dowling with no evidence of a hypersensitive response due to leaf necrosis and (c) shows the glue used to fix the eggs to the sheet. (a) was obtained with a JSM 840A and (b) and (c) were obtained with a Zeiss DMS 962.

222 ng egg⁻¹ has been reported in *Helicoverpa armigera* (Lepidoptera: Noctuidae). The egg volatiles might also have been masked by constitutive volatiles of the soybean plants, which are present in much higher amounts (Michereff *et al.*, 2011). It should be noted that in the olfactometer bioassay there were no visual or other cues to orient the egg parasitoid in addition to the constitutive volatiles from the plants. Normally, egg parasitoid responses to infochemicals include a sequence of steps that involve several cues, such as the distance from the odour source, wind, visual information of plants and eggs, and other environmental stimuli (Fatouros *et al.*, 2008; Hilker & Meiners, 2011).

The saturated and unsaturated fatty acids, including hexadecanoic acid, linoleic acid, and (Z)-9-octadecenoic acid, that were found in acetone egg extracts might be *T. podisi* attractants. Chemical analysis of *Ostrinia nubilalis* (Lepidoptera: Crambidae) and *Mamestra brassicae* (Lepidoptera: Noctuidae) egg extracts revealed the presence of fatty acids, their ethyl esters, and various hydrocarbons, which were found to be kairones that influenced the search behaviour of the egg parasitoid *Trichogramma brassicae* (Hymenoptera: Trichogrammatidae) (Renou *et al.*, 1992). Females of another egg parasitoid, *Telenomus euproctidis* (Hymenoptera: Platygasteridae), were shown to be attracted to *Orgyia postica* (Lepidoptera: Lymantriidae) egg masses, but chemical analysis to identify potential semiochemicals was not done (Arakaki *et al.*, 2011).

Presumably, egg volatiles function at short range when parasitoids are near plants, and compounds with medium and low volatility, such as long-chain fatty acids and fatty acid methyl esters, are the likely signalling molecules. The absence of OIPVs and highly volatile egg compounds that might attract Platygasteridae egg parasitoids can account for the detection of HIPVs, which are reliable indicators of the presence of stink bugs and their eggs (Moraes *et al.*, 2005, 2008; Michereff *et al.*, 2011, 2013; Melo Machado *et al.*, 2014).

HIPVs are important cues for parasitoid host foraging behaviour because stink bugs feed on plants while ovipositing. HIPVs are reliably associated with the presence of eggs on the plant, but the chemical profiles of both HIPVs and OIPVs can be modified by simultaneous infestation of multiple herbivores and/or pathogens that occurs under natural conditions. That can interfere with attraction of natural enemies and has been shown to affect the behaviour of various parasitoids including *Cotesia marginiventris*, *T. basalis*, *T. brassicae* and *Trichogramma evanescens* (Rasmann & Turlings, 2007; Moujahed *et al.*, 2014; Cusumano *et al.*, 2015). Because of this, multitrophic interactions among herbivores that are natural enemies of the soybean should be considered in future studies evaluating parasitoid responses in this model system. It is most likely that egg parasitoids use HIPVs as reliable signals to detect the presence of their host species.

In summary, our study showed that the interaction between soybean plants, *E. heros* eggs and *T. podisi* is different

from several other systems involving plant-herbivore oviposition (with or without physical damage on the leaf surface) and their natural enemies (Hilker & Meiners, 2006; Fatouros *et al.*, 2008, 2012; Tamiru *et al.*, 2011, 2012). We also showed that *E. heros* eggs have compounds that affect *T. podisi* foraging behaviour and that *E. heros* oviposition alone does not induce chemicals that attract *T. podisi* to soybean plants. Additional studies should be conducted to evaluate the effect of the chemical compounds found in *E. heros* eggs on foraging behaviour in Platygasteridae parasitoids. It would be interesting to determine whether increased amounts of these compounds attract natural enemies or change plant responses and whether different soybean cultivars or plant species have similar defences to mask egg volatiles.

SUPPLEMENTARY MATERIAL

The supplementary material for this article can be found at <http://dx.doi.org/10.1017/S0007485316000419>.

Acknowledgements

We thank Daniela Conceição for helping with laboratory rearing of the insects, Guy de Capdeville for helping in image acquisition, and Diego M. Magalhães for comments that greatly improved the manuscript. This work received financial support from the Brazilian Council for Scientific and Technological Development (CNPq) through a grant to M. F. F. Michereff, the Federal District Research Foundation (FAP-DF), and the Brazilian Corporation of Agricultural Research (EMBRAPA).

References

- Aquino, M.F.S., Dias, A.M., Morges, M., Blassioli-Moraes, M.C. & Laumann, R.A. (2012) Influence of visual cues on host-searching and learning behaviour of the parasitoids *Telenomus podisi* and *Trissolcus basal*. *Entomologia Experimentalis et Applicata* **145**, 162–174.
- Arakaki, N., Yamazawa, H. & Wakamura, S. (2011) The egg parasitoid *Telenomus euproctidis* (Hymenoptera: Scelionidae) uses sex pheromone released by immobile female tussock moth *Orgyia postica* (Lepidoptera: Lymantriidae) as kairomone. *Applied Entomology and Zoology* **46**, 195–200.
- Bin, F., Vinson, S.B., Strand, M.R., Colazza, S. & Jones, W.A. (1993) Source of an egg kairomone for *Trissolcus basal*, a parasitoid of *Nezara viridula*. *Physiological Entomology* **18**, 7–15.
- Borges, M. & Aldrich, J.R. (1994) Estudos de semioquímicos para o manejo de Telenominae. *Anais da Sociedade Entomológica do Brasil* **23**, 575–577.
- Borges, M., Schmidt, F.G.V., Sujii, E.R., Medeiros, M.A., Mori, K., Zarbin, P.H.G. & Ferreira, J.T.B. (1998) Field responses of stink bugs to the natural and synthetic pheromone of the Neotropical brown stink bug, *Euschistus heros* (Heteroptera: Pentatomidae). *Physiological Entomology* **23**, 202–207.
- Borges, M., Costa, M.L.M., Sujii, E.R., Cavalcanti, M.G., Redigolo, G.F., Resck, I.S. & Vilela, E.F. (1999) Semiochemical and physical stimuli involved in host recognition by *Telenomus podisi* (Hymenoptera: Scelionidae) toward *Euschistus heros* (Heteroptera: Pentatomidae). *Physiological Entomology* **24**, 227–233.
- Borges, M., Colazza, S., Ramirez-Lucas, P., Chauhan, K.R., Moraes, M.C.B. & Aldrich, J.R. (2003) Kairomonal effect of walking traces from *Euschistus heros* (Heteroptera: Pentatomidae) on two strains of *Telenomus podisi* (Hymenoptera: Scelionidae). *Physiological Entomology* **28**, 349–355.
- Bruce, T.J.A., Midega, C.A.O., Birkett, M.A., Pickett, J.A. & Khan, Z.R. (2010) Is quality more important than quantity? Insect behavioural responses to changes in a volatile blend after stemborer oviposition on an African grass. *Biology Letters* **6**, 314–317.
- Bruessow, F., Gouhier-Darimont, C., Buchala, A., Mettraux, J.P. & Reymond, P. (2010) Insect egg suppress plant defence against chewing herbivores. *Plant Journal* **62**, 876–885.
- Büchel, K., Malskies, S., Mayer, M., Fenning, T.M., Gershenson, J., Hilker, M. & Meiners, T. (2011) How plants give early herbivore alert: volatile terpenoids attract parasitoids to egg-infested elms. *Basic Applied Ecology* **12**, 403–412.
- ChemStation 2.1 Software. (2008) Available online at <http://www.agilent.com/en-us/products/software-informatics/massspec-workstations/gc-msd-chemstation-software>.
- Colazza, S., Mcelfresh, J.S., & Millar, J.G. (2004a) Identification of volatile synomones, induced by *Nezara viridula* feeding and oviposition on bean spp., that attract the egg parasitoid *Trissolcus basal*. *Journal of Chemical Ecology* **30**, 945–964.
- Colazza, S., Fucarino, A., Peri, E., Salerno, G., Conti, E. & Bin, F. (2004b) Insect oviposition induces volatile emission in herbaceous plants that attracts egg parasitoids. *Journal of Experimental Biology* **207**, 47–53.
- Cusumano, A., Weldegergis, B.T., Colazza, S., Dicke, M. & Fatouros, N. (2015) Attraction of egg-killing parasitoids toward induced plant volatiles in a multi-herbivore context. *Oecologia* **179**, 163–174.
- EzChrom Elite software. (2008) Available online at <http://www.agilent.com/en-us/products/liquid-chromatography/gp59442?>
- Fatouros, N.E., Dicke, M., Mumm, R., Meiners, T. & Hilker, M. (2008) Foraging behavior of egg parasitoids exploiting chemical information. *Behavioral Ecology* **19**, 677–689.
- Fatouros, N.E., Lucas-Barbosa, D., Weldegergis, B.T., Pashalidou, F.G., Van Loon, J.J.A., Dicke, M., Harvey, J.A., Gols, R. & Huigens, M.E. (2012) Plant volatiles induced by herbivore egg deposition affect insects of different trophic levels. *PLoS ONE* **7**, 1–14.
- Hilker, M. & Meiners, T. (2006) Early herbivore alert: insect eggs induce plant defense. *Journal of Chemical Ecology* **32**, 1379–1397.
- Hilker, M. & Meiners, T. (2010) How do plants “notice” attack by herbivorous arthropods? *Biological Review* **85**, 267–280.
- Hilker, M. & Meiners, T. (2011) Plants and insect eggs: how do they affect each other? *Phytochemistry* **72**, 1612–1623.
- Hilker, M., Stein, C., Schröder, R., Varama, M. & Mumm, R. (2005) Insect egg deposition induces defence responses in *Pinus sylvestris*: characterisation of the elicitor. *Journal of Experimental Biology* **208**, 1849–1854.
- Hu, J.S., Gelman, D.B. & Bell, R.A. (1999) Effects of selected physical and chemical treatments of Colorado potato beetle eggs on host acceptance and development of the parasitic wasp, *Edovum puttleri*. *Entomologia Experimentalis et Applicata* **90**, 237–245.
- Jorge, L.A.C., Laumann, R.A., Borges, M., Moraes, M.C.B., Cruz, R.A., Milare, B.N. & Palhares, L. (2005) Software para avaliação do comportamento de insetos. *Circular Técnica* n. 30, 7p. São Carlos, SP: Embrapa Instrumentação Agropecuária.
- Kaloshian, I. & Walling, L.L. (2005) Hemipterans as plant pathogens. *Annual Review of Phytopathology* **43**, 491–521.
- Laumann, R.A., Moraes, M.C.B., Cokl, A. & Borges, M. (2007) Eavesdropping on the sexual vibratory communication of

- stink bugs (Hemiptera: Pentatomidae) by the egg parasitoid *Telenomus podisi*. *Animal Behaviour* **73**, 637–649.
- Laumann, R.A., Aquino, M.F.S., Moraes, M.C.B., Pareja, M. & Borges, M.** (2009) Response of the egg parasitoids *Trissolcus basaloides* and *Telenomus podisi* to compounds from defensive secretions of stink bugs. *Journal of Chemical Ecology* **35**, 8–19.
- Laumann, R.A., Cokl, A., Lopes, A.P.S., Ferreira, J.B.C., Moraes, M.C.B. & Borges, M.** (2011) Silent singers are not safe: selective response of a parasitoid to substrate-borne vibratory signals of stink bugs. *Animal Behaviour* **82**, 1175–1183.
- Leonard, D.E., Wu, Z-Xe., Ferro, D.N.** (1987) Responses of parasitoid *Edovum puttleri* to kairomone from eggs of Colorado potato beetle, *Leptinotarsa decemlineata*. *Journal of Chemical Ecology* **13**, 335–344.
- Liu, M., Yu, H. & Li, G.** (2008) Oviposition deterrents from eggs of the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae): chemical identification and analysis by electroantennogram. *Journal of Insect Physiology* **54**, 656–662.
- Medeiros, M.A., Schmidt, F.V.G., Loiacono, M.S., Carvalho, V.F. & Borges, M.** (1997) Parasitismo e predação de ovos de *Euschistus heros* (Fab.) (Heteroptera: Pentatomidae) no Distrito Federal, Brasil. *Anais da Sociedade Entomológica do Brasil* **26**, 397–401.
- Medeiros, M.A., Loiacono, M.S., Borges, M. & Schmidt, F.V.G.** (1998) Incidência natural de parasitoides em ovos de percevejos (Hemiptera: Pentatomidae) encontrados na soja no Distrito Federal. *Pesquisa Agropecuária Brasileira* **33**, 1431–1435.
- Melo Machado, R.C., Sant'Ana, J., Blassioli-Moraes, M.C., Laumann, R.A. & Borges, M.** (2014) Herbivory-induced plant volatiles from *Oryza sativa* and their influence on chemotaxis behaviour of *Tibraca limbativentris* stal. (Hemiptera: Pentatomidae) and egg parasitoids. *Bulletin of Entomological Research* **104**, 347–356.
- Michereff, M.F.F., Laumann, R.A., Borges, M., Michereff Filho, M., Diniz, I.R., Farias Neto, A.L. & Moraes, M.C.B.** (2011) Volatiles mediating plant-herbivory-natural enemy interaction in resistant and susceptible soybean cultivars. *Journal of Chemical Ecology* **37**, 273–285.
- Michereff, M.F.F., Borges, M., Laumann, R.A., Diniz, I.R. & Moraes, M.C.B.** (2013) Influence of volatile compounds from herbivore-damaged soybean plants on searching behavior of the egg parasitoid *Telenomus podisi*. *Entomologia Experimentalis et Applicata* **147**, 9–17.
- Michereff, M.F.F., Michereff Filho, M., Blassioli-Moraes, M.C., Laumann, R.A., Diniz, I.R. & Borges, M.** (2015) Effect of resistant and susceptible soybean cultivars on the attraction of egg parasitoids in field conditions. *Journal of Applied Entomology* **139**, 207–216.
- Moraes, M.C.B., Laumann, R.A., Pires, C.S.S., Sujii, E.R. & Borges, M.** (2005) Induced volatiles in soybean and pigeon pea plants artificially infested with the Neotropical brown stink bug, *Euschistus heros*, and their effect on the egg parasitoid, *Telenomus podisi*. *Entomologia Experimentalis et Applicata* **115**, 227–237.
- Moraes, M.C.B., Pareja, M., Laumann, R.A., Hoffmann-Campo, C.B. & Borges, M.** (2008) Response of the parasitoid *Telenomus podisi* to induced volatiles from soybean damaged by stink bug herbivory and oviposition. *Journal of Plant Interaction* **3**, 1742–1756.
- Moujahed, R., Frati, F., Cusumano, A., Salerno, G., Conti, E., Peri, E. & Colazza, S.** (2014) Egg parasitoid attraction toward induced plant volatile is disrupted by a non-host herbivore attacking above or belowground plant organs. *Frontiers in Plant Science* **5**, 1–10.
- Pacheco, D.J.P. & Corrêa-Ferreira, B.S.** (2000) Parasitismo de *Telenomus podisi* Ashmead (Hymenoptera: Scelionidae) em populações de percevejos pragas da soja. *Anais da Sociedade Entomológica do Brasil* **29**, 295–302.
- Rasmann, S. & Turlings, T.C.J.** (2007) Simultaneous feeding by aboveground and belowground herbivores attenuates plant-mediated attraction of their respective natural enemies. *Ecology Letters* **10**, 926–936.
- R Development Core Team** (2009) *R: A Language and Environment for Statistical Computing*. Vienna, Austria, R Foundation for Statistical Computing. ISBN 3-900051-07-0. Available online at <http://www.R-project.org>
- Renou, M., Nagnan, P., Berthier, A. & Durier, C.** (1992) Identification of compounds from the eggs of *Ostrinia nubilalis* and *Mamestra brassicae* having kairomone activity on *Trichogramma brassicae*. *Entomologia Experimentalis et Applicata* **63**, 291–303.
- Sales, F.M.** (1979) Responsiveness and threshold for host-seeking stimulation of the female, *Trissolcus basaloides* (Wollaston) by the eggs of the host, *Nezara viridula* (L.). *Fitosanidade* **3**, 36–39.
- Takasu, K. & Nordlund, D.A.** (2001) Host recognition kairomones for *Anaphes iole* Girault, an egg parasitoid of the Western tarnished plant bug. *Biological Control* **22**, 60–65.
- Tamiru, A., Bruce, T.J.A., Woodcock, C.M., Caufield, J.C., Midega, C.A.O., Ogol, C.K.P., Mayon, P., Birkett, M.A., Pickett, J.A. & Khan, Z.R.** (2011) Maize landraces recruit egg and larval parasitoids in response to egg deposition by a herbivore. *Ecology Letters* **14**, 1075–1083.
- Tamiru, A., Bruce, T.J.A., Midega, C.A., Woodcock, C.M., Birkett, M.A., Pickett, J.A. & Khan, Z.R.** (2012) Oviposition induced volatile emission from Africa smallholder farmers' maize varieties. *Journal of Chemical Ecology* **38**, 231–234.
- Tognon, R., Sant'Ana, J. & Jahnke, S.M.** (2014) Influence of original host on chemotactic behaviour and parasitism in *Telenomus podisi* Ashmead (Hymenoptera: Platygasteridae). *Bulletin of Entomological Research* **104**, 781–787.
- Tooker, J.F. & De Moraes, C.M.** (2007) Jasmonate, salicylate and benzoate in insect eggs. *Journal of Chemical Ecology* **33**, 331–343.
- Vinson, S.B.** (1998) The general host selection behavior of parasitoid Hymenoptera and a comparison of initial strategies utilized by larviphagous and oophagous species. *Biological Control* **11**, 79–96.
- Walling, L.L.** (2000) The myriad plant responses to herbivores. *Journal of Plant Growth Regulation* **19**, 195–216.
- Wegener, R., Schulz, S., Meiners, T., Hadwich, K. & Hilker, M.** (2001) Analysis of volatiles induced by oviposition of elm leaf beetle *Xanthogaleruca luteola* on *Ulmus minor*. *Journal of Chemical Ecology* **27**, 499–515.