



Pyricularia graminis-tritici, a new *Pyricularia* species causing wheat blast

V.L. Castroagudín¹, S.I. Moreira², D.A.S. Pereira^{1,3}, S.S. Moreira¹, P.C. Brunner³, J.L.N. Maciel⁴, P.W. Crous^{5,6,7}, B.A. McDonald³, E. Alves², P.C. Ceresini¹

Key words

cryptic species
host adaptation
phylogenetics
systematics
Triticum aestivum

Abstract *Pyricularia oryzae* is a species complex that causes blast disease on more than 50 species of poaceous plants. *Pyricularia oryzae* has a worldwide distribution as a rice pathogen and in the last 30 years emerged as an important wheat pathogen in southern Brazil. We conducted phylogenetic analyses using 10 housekeeping loci for 128 isolates of *P. oryzae* sampled from sympatric populations of wheat, rice, and grasses growing in or near wheat fields. Phylogenetic analyses grouped the isolates into three major clades. Clade 1 comprised isolates associated only with rice and corresponds to the previously described rice blast pathogen *P. oryzae* pathotype *Oryza* (PoO). Clade 2 comprised isolates associated almost exclusively with wheat and corresponds to the previously described wheat blast pathogen *P. oryzae* pathotype *Triticum* (PoT). Clade 3 contained isolates obtained from wheat as well as other *Poaceae* hosts. We found that Clade 3 is distinct from *P. oryzae* and represents a new species, *Pyricularia graminis-tritici* (Pgt). No morphological differences were observed among these species, but a distinctive pathogenicity spectrum was observed. Pgt and PoT were pathogenic and highly aggressive on *Triticum aestivum* (wheat), *Hordeum vulgare* (barley), *Urochloa brizantha* (signal grass), and *Avena sativa* (oats). PoO was highly virulent on the original rice host (*Oryza sativa*), and also on wheat, barley, and oats, but not on signal grass. We conclude that blast disease on wheat and its associated *Poaceae* hosts in Brazil is caused by multiple *Pyricularia* species. *Pyricularia graminis-tritici* was recently found causing wheat blast in Bangladesh. This indicates that *P. graminis-tritici* represents a serious threat to wheat cultivation globally.

Article info Received: 29 April 2016; Accepted: 8 June 2016; Published: 24 June 2016.

INTRODUCTION

Pyricularia oryzae is a species complex (Couch & Kohn 2002) that causes blast disease on more than 50 species of poaceous plants, including important crops such as rice, wheat, barley, millet, and oats (Urashima & Kato 1998, Couch & Kohn 2002, Takabayashi et al. 2002, Murakami et al. 2003, Couch et al. 2005). On the basis of host specificity, mating ability, and genetic relatedness, *P. oryzae* isolates were classified into several subgroups with restricted host ranges, including: the *Oryza* pathotype, pathogenic on rice (*Oryza sativa*); the *Setaria* pathotype, pathogenic on foxtail millet (*Setaria italica*); the *Panicum* pathotype, pathogenic on common millet (*Panicum miliaceum*); the *Eleusine* pathotype, pathogenic on finger millet (*Eleusine coracana*); the *Triticum* pathotype, pathogenic on wheat (*Triticum aestivum*); the *Avena* pathotype, pathogenic on oats (*Avena sativa*); and the *Lolium* pathotype, pathogenic

on perennial ryegrass (*Lolium perenne*) (Urashima et al. 1993, Kato et al. 2000, Tosa et al. 2004, Tosa & Chuma 2014). Kato and collaborators (Kato et al. 2000) reported that isolates of *P. oryzae* recovered from *Eleusine*, *Panicum*, *Oryza*, *Setaria*, and *Triticum* spp. form a highly related group that is partially inter-fertile with the *Oryza* subgroup (i.e. the rice blast pathogen). In addition, the *Oryza* and *Setaria* pathotypes contain physiological races that show distinct patterns of virulence on cultivars within their host species (Tosa & Chuma 2014). Both host species-specificity and cultivar-specificity can be governed by gene-for-gene interactions (Silue et al. 1992, Takabayashi et al. 2002, Tosa et al. 2006, Valent & Khang 2010).

The *P. oryzae* pathotype *Triticum* is considered the causal agent of wheat blast in South America and has also been associated with blast disease on barley, rye, triticale, and signal grass (*Urochloa* sp., ex *Brachiaria* sp.) in central-western and southern Brazil (Lima & Minella 2003, Verzignassi et al. 2012). Wheat blast was first reported in Paraná State, Brazil in 1985 (Igarashi et al. 1986, Anjos et al. 1996). Due to the lack of resistant cultivars and effective fungicides for disease management, wheat blast is widely distributed across all the wheat-cropping areas in Brazil, causing crop losses from 40–100 % (Silva et al. 2009, Maciel 2011, Castroagudín et al. 2015). Wheat blast also occurs in Bolivia, Argentina, and Paraguay (Duveiller et al. 2010). The disease was not found outside South America (Maciel 2011) until a recent outbreak reported in Bangladesh (Callaway 2016), though wheat blast is considered a major quarantine disease and a threat to wheat crops in the United States (Duveiller et al. 2007, Kohli et al. 2011).

As wheat blast emerged in an area of southern Brazil where rice blast is prevalent, it was originally proposed that the rice

¹ Department of Phytopathology, Rural Engineering, and Soil Science (Departamento de Fitossanidade, Engenharia Rural e Solos), UNESP- University of São Paulo State, Ilha Solteira, São Paulo, Brazil; corresponding author e-mail: paulo.ceresini@bio.feis.unesp.br.

² Department of Phytopathology, Federal University of Lavras, Lavras, Minas Gerais, Brazil.

³ Plant Pathology Group, Institute of Integrative Biology, ETH Zürich, Zürich, Switzerland.

⁴ Brazilian Agriculture Research Corporation-Wheat (EMBRAPA-Trigo), Passo Fundo, Rio Grande do Sul, Brazil.

⁵ CBS-KNAW Fungal Biodiversity Centre, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands.

⁶ Department of Microbiology and Plant Pathology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria 0002, South Africa.

⁷ Microbiology, Department of Biology, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands.

pathogen had evolved to parasitize wheat (Igarashi et al. 1986). Urashima et al. (1993) provided evidence based on pathogenicity, reproductive isolation, and genetic data that indicated the existence of two distinct groups of *P. oryzae* causing wheat blast in Brazil: one that infects rice and wheat, and one that infects only wheat. In that study, wheat-derived isolates were reported to infect grass plants from six different tribes within *Poaceae*. In addition, crosses of wheat-derived isolates with strains from *Eleusine coracana*, *Urochloa plantaginea* (ex *Brachiaria plantaginea*), and *Setaria italica* produced mature perithecia with viable ascospores, i.e. evidence of fertile crosses (Urashima et al. 1993). On the contrary, progeny from the crosses between wheat- and rice-derived isolates were infertile (Urashima et al. 1993). In the same study, crosses between wheat-derived isolates and isolates obtained from *Cenchrus echinatus*, *Setaria geniculata*, and *Echinochloa colonum* produced no perithecia (Urashima et al. 1993). The work of Urashima and his collaborators indicated that two distinct pyricularia-like pathogens cause wheat blast disease in Brazil. However, it is not clear whether a population of *P. oryzae* able to infect both rice and wheat coexists with a population that infects only wheat.

Several studies suggested that the wheat-adapted *P. oryzae* population was derived *de novo* from a non-rice host. DNA fingerprinting with the repetitive DNA probes MGR563 and MGR586 found a high level of differentiation between *P. oryzae* pathotype *Oryza* (PoO) and *P. oryzae* pathotype *Triticum* (PoT) from Brazil (Farman 2002). In fact, the fingerprints from wheat-derived isolates resembled those from isolates non-pathogenic to rice (Hamer 1991, Valent & Chumley 1991, Urashima et al. 1999, Farman 2002). Maciel et al. (2014) showed that the Brazilian wheat-adapted population of *P. oryzae* was highly differentiated ($F_{CT} = 0.896$, $P \leq 0.001$) from the local rice-adapted population. Analyses of the current pathotype diversity of *P. oryzae* showed that none of the 69 wheat-derived isolates were able to infect rice (Maciel et al. 2014).

Phylogenetic analyses demonstrated that *Pyricularia* is a species-rich genus in which different species evolved through repeated radiation events from a common ancestor (Hirata et al. 2007, Choi et al. 2013, Klaubauf et al. 2014). Multi-locus phylogenetic analyses revealed that *P. oryzae* and *P. grisea* are independent phylogenetic species (Taylor et al. 2000, Couch & Kohn 2002) and showed that the contemporary rice-infecting pathogen (*P. oryzae* pathotype *Oryza*) originated via a host shift from millet onto rice ~7 000 years ago during rice domestication in China (Couch et al. 2005). More recent phylogenetic analyses combined pre-existing biological and morphological data to re-examine the relationships among pyricularia-like species. These comprehensive studies favoured the classification of new cryptic species that were recently identified within *Pyricularia* and other relevant changes within the order *Magnaporthales* (Hirata et al. 2007, Choi et al. 2013, Luo & Zhang 2013, Klaubauf et al. 2014, Murata et al. 2014). Most relevant for agricultural scientists is that despite the extensively reported differentiation between *P. oryzae* pathotypes *Oryzae* and *Triticum*, these two pathotypes have been kept under the same species name *P. oryzae*. Therefore, we sought to determine whether the pathotypes *Oryza* and *Triticum* of *P. oryzae* are distinct species that should be given different names. We conducted phylogenetic analyses based on 10 housekeeping genes using sympatric populations of *Pyricularia* sampled from rice, wheat, and other poaceous hosts in Brazil. We also conducted cultural, morphological, and pathogenic characterisation of the *Pyricularia* isolates to provide a complete description for each species. Our phylogenetic analyses revealed a new *Pyricularia* species causing blast on wheat and other poaceous hosts in Brazil. We name and describe *Pyricularia graminis-tritici* in this report.

MATERIALS AND METHODS

Fungal isolates and DNA extraction

A unique collection of 128 monoconidial isolates of *Pyricularia* spp. obtained in sympatry from the Brazilian wheat agro-ecosystem was analysed in this study (Table 1). *Pyricularia* spp. isolates were obtained from *Triticum aestivum* ($N = 79$), *Oryza sativa* ($N = 23$), *Avena sativa* ($N = 5$), *Cenchrus echinatus* ($N = 3$), *Cynodon* sp. ($N = 1$), *Digitaria sanguinalis* ($N = 4$), *Elionurus candidus* ($N = 2$), *Echinochloa crusgalli* ($N = 1$), *Eleusine indica* ($N = 1$), *Rhynchelytrum repens* ($N = 3$), and *Urochloa brizantha* (ex *Brachiaria brizantha*) ($N = 6$). Isolates recovered from wheat and other poaceous hosts located within or adjacent to sampled wheat plots were obtained from symptomatic head and leaf tissue in commercial wheat fields located in seven states in Brazil during 2012. A detailed description of wheat field sampling strategies was provided earlier (Castroagudín et al. 2015). The rice-derived isolates of *P. oryzae* were recovered from rice leaves, necks and panicles exhibiting typical rice blast symptoms, comprising a representative group including all races of *P. oryzae* pathotype *Oryza* prevalent in the major Brazilian rice growing areas (Maciel et al. 2014). The rice-derived isolates were provided by EMBRAPA-Rice and Beans, Santo Antônio de Goiás, Goiás, Brazil. The isolate collection is maintained at the Laboratory of Phytopathology, UNESP-DEFERS Campus Ilha Solteira, São Paulo, Brazil. A duplicate of the collection is hosted at the Laboratory of Phytopathology, EMBRAPA-Wheat, Passo Fundo, Brazil. Specimens were deposited at Culture Collection Mycobank Prof. Maria Auxiliadora Cavalcanti, Federal University of Pernambuco, Recife, Brazil, and at the Coleção de Culturas da Microbiologia Agrícola (Agriculture Microbiology Culture Collection) of the Federal University of Lavras, Lavras, Minas Gerais, Brazil. Holotype specimen was deposited at INCT-HISA Herbário Virtual da Flora e dos Fungos at UNESP – Campus Ilha Solteira (Virtual Herbarium of Flora and Fungi, University of São Paulo State – Campus Ilha Solteira, Ilha Solteira, São Paulo, Brazil).

DNA extraction, amplification, and sequencing

Genomic DNA was extracted from freeze-dried mycelia with the GenElute Plant Genomic DNA Miniprep Kit (Sigma-Aldrich, St. Louis, MO, USA), according to the specifications of the manufacturer. Partial sequences of 10 nuclear housekeeping loci previously used to characterise *Pyricularia* species (Carbone & Kohn 1999, Couch & Kohn 2002, Couch et al. 2005, Zhang et al. 2011) were included in the analyses. The loci amplified were: *ACT* (actin), *BAC6* (putative vacuolar import and degradation protein), *βT-1* (beta-tubulin), *CAL* (calmodulin), *CH7-BAC7* (hypothetical protein), *CH7-BAC9* (anonymous sequence), *CHS1* (chitin synthase 1), *EF-1α* (translation elongation factor 1-alpha), *MPG1* (hydrophobin), and *NUT1* (nitrogen regulatory protein 1). The loci were amplified using PCR cycling conditions described previously (Carbone & Kohn 1999, Couch et al. 2005). The PCR primers and the annealing temperatures used to amplify each locus are described in Table 2. The PCR products were purified and sequenced by Macrogen Inc. (Seoul, Korea) using the ABI Prism BigDye Terminator v.3.1 Cycle Sequencing Ready Reaction Kit in an ABI 3730xl automated sequencer (Applied Biosystems, Foster City, CA). Newly generated DNA sequences were deposited in NCBI's GenBank nucleotide database (Table 1).

Phylogenetic analyses

The complete set of sequence data was obtained from 125 isolates of *Pyricularia* spp., including two identified as *P. pennisetigena* (URM7372 = CML3524, isolate 12.0.100) and *P. grisea* (URM7371 = CML3525, isolate 12.0.082) from Brazil, which

Table 1 (cont.)

Species, isolate	Race	Host	Origin	Sampling year	ACT	BAC6	β T-1	CAL	CH7-BAC7	CH7-BAC9	CHS	EF-1 α	MPG1	NUT1
<i>Triticum aestivum</i>	-		São Paulo	2012	KU952187	KU952313	KU953067	KU952941	KU952439	KU952564	KU953192	KU953317	KU952690	KU952816
<i>Triticum aestivum</i>	-		São Paulo	2012	KU952188	KU952314	KU953068	KU952942	KU952440	KU952565	KU953193	KU953318	KU952691	KU952817
<i>Triticum aestivum</i>	-		Minas Gerais	2012	KU952189	KU952315	KU953069	KU952943	KU952441	KU952566	KU953194	KU953319	KU952692	KU952818
<i>Triticum aestivum</i>	-		São Paulo	2012	KU952190	KU952316	KU953070	KU952944	KU952442	KU952567	KU953195	KU953320	KU952693	KU952819
<i>Triticum aestivum</i>	-		Goias	2012	KU952191	KU952317	KU953071	KU952945	KU952443	KU952568	KU953196	KU953321	KU952694	KU952820
<i>Triticum aestivum</i>	-		Goias	2012	KU952192	KU952318	KU953072	KU952946	KU952444	KU952569	KU953197	KU953322	KU952695	KU952821
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952193	KU952319	KU953073	KU952947	KU952445	KU952570	KU953198	KU953323	KU952696	KU952822
<i>Triticum aestivum</i>	-		Minas Gerais	2012	KU952194	KU952320	KU953074	KU952948	KU952446	KU952571	KU953199	KU953324	KU952697	KU952823
<i>Triticum aestivum</i>	-		Minas Gerais	2012	KU952195	KU952321	KU953075	KU952949	KU952447	KU952572	KU953200	KU953325	KU952698	KU952824
<i>Triticum aestivum</i>	-		Minas Gerais	2012	KU952196	KU952322	KU953076	KU952950	KU952448	KU952573	KU953201	KU953326	KU952699	KU952825
<i>Triticum aestivum</i>	-		Minas Gerais	2012	KU952197	KU952323	KU953077	KU952951	KU952449	KU952574	KU953202	KU953327	KU952700	KU952826
<i>Triticum aestivum</i>	-		Goias	2012	KU952198	KU952324	KU953078	KU952952	KU952450	KU952575	KU953203	KU953328	KU952701	KU952827
<i>Triticum aestivum</i>	-		Federal District	2012	KU952199	KU952325	KU953079	KU952953	KU952451	KU952576	KU953204	KU953329	KU952702	KU952828
<i>Triticum aestivum</i>	-		Federal District	2012	KU952200	KU952326	KU953080	KU952954	KU952452	KU952577	KU953205	KU953330	KU952703	KU952829
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952201	KU952327	KU953081	KU952955	KU952453	KU952578	KU953206	KU953331	KU952704	KU952830
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952202	KU952328	KU953082	KU952956	KU952454	KU952579	KU953207	KU953332	KU952705	KU952831
<i>Triticum aestivum</i>	-		Minas Gerais	2012	KU952203	KU952329	KU953083	KU952957	KU952455	KU952580	KU953208	KU953333	KU952706	KU952832
<i>Triticum aestivum</i>	-		Minas Gerais	2012	KU952204	KU952330	KU953084	KU952958	KU952456	KU952581	KU953209	KU953334	KU952707	KU952833
<i>Triticum aestivum</i>	-		Minas Gerais	2012	KU952205	KU952331	KU953085	KU952959	KU952457	KU952582	KU953210	KU953335	KU952708	KU952834
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952206	KU952332	KU953086	KU952960	KU952458	KU952583	KU953211	KU953336	KU952709	KU952835
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952207	KU952333	KU953087	KU952961	KU952459	KU952584	KU953212	KU953337	KU952710	KU952836
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952208	KU952334	KU953088	KU952962	KU952460	KU952585	KU953213	KU953338	KU952711	KU952837
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952209	KU952335	KU953089	KU952963	KU952461	KU952586	KU953214	KU953339	KU952712	KU952838
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952210	KU952336	KU953090	KU952964	KU952462	KU952587	KU953215	KU953340	KU952713	KU952839
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952211	KU952337	KU953091	KU952965	KU952463	KU952588	KU953216	KU953341	KU952714	KU952840
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952212	KU952338	KU953092	KU952966	KU952464	KU952589	KU953217	KU953342	KU952715	KU952841
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952213	KU952339	KU953093	KU952967	KU952465	KU952590	KU953218	KU953343	KU952716	KU952842
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952214	KU952340	KU953094	KU952968	KU952466	KU952591	KU953219	KU953344	KU952717	KU952843
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952215	KU952341	KU953095	KU952969	KU952467	KU952592	KU953220	KU953345	KU952718	KU952844
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952216	KU952342	KU953096	KU952970	KU952468	KU952593	KU953221	KU953346	KU952719	KU952845
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952217	KU952343	KU953097	KU952971	KU952469	KU952594	KU953222	KU953347	KU952720	KU952846
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952218	KU952344	KU953098	KU952972	KU952470	KU952595	KU953223	KU953348	KU952721	KU952847
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952219	KU952345	KU953099	KU952973	KU952471	KU952596	KU953224	KU953349	KU952722	KU952848
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952220	KU952346	KU953100	KU952974	KU952472	KU952597	KU953225	KU953350	KU952723	KU952849
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952221	KU952347	KU953101	KU952975	KU952473	KU952598	KU953226	KU953351	KU952724	KU952850
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952222	KU952348	KU953102	KU952976	KU952474	KU952599	KU953227	KU953352	KU952725	KU952851
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952223	KU952349	KU953103	KU952977	KU952475	KU952600	KU953228	KU953353	KU952726	KU952852
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952224	KU952350	KU953104	KU952978	KU952476	KU952601	KU953229	KU953354	KU952727	KU952853
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952225	KU952351	KU953105	KU952979	KU952477	KU952602	KU953230	KU953355	KU952728	KU952854
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952226	KU952352	KU953106	KU952980	KU952478	KU952603	KU953231	KU953356	KU952729	KU952855
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952227	KU952353	KU953107	KU952981	KU952479	KU952604	KU953232	KU953357	KU952730	KU952856
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952228	KU952354	KU953108	KU952982	KU952480	KU952605	KU953233	KU953358	KU952731	KU952857
<i>Triticum aestivum</i>	-		Rio Grande do Sul	2012	KU952229	KU952355	KU953109	KU952983	KU952481	KU952606	KU953234	KU953359	KU952732	KU952858
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952230	KU952356	KU953110	KU952984	KU952482	KU952607	KU953235	KU953360	KU952733	KU952859
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952231	KU952357	KU953111	KU952985	KU952483	KU952608	KU953236	KU953361	KU952734	KU952860
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952232	KU952358	KU953112	KU952986	KU952484	KU952609	KU953237	KU953362	KU952735	KU952861
<i>Triticum aestivum</i>	-		Mato Grosso do Sul	2012	KU952233	KU952359	KU953113	KU952987	KU952485	KU952610	KU953238	KU953363	KU952736	KU952862
<i>Triticum aestivum</i>	-		Paraná	2012	KU952234	KU952360	KU953114	KU952988	KU952486	KU952611	KU953239	KU953364	KU952737	KU952863
<i>Triticum aestivum</i>	-		Paraná	2012	KU952235	KU952361	KU953115	KU952989	KU952487	KU952612	KU953240	KU953365	KU952738	KU952864
<i>Triticum aestivum</i>	-		Paraná	2012	KU952236	KU952362	KU953116	KU952990	KU952488	KU952613	KU953241	KU953366	KU952739	KU952865
<i>Triticum aestivum</i>	-		Paraná	2012	KU952237	KU952363	KU953117	KU952991	KU952489	KU952614	KU953242	KU953367	KU952740	KU952866
<i>Oryza sativa</i>	97		Tocantins	2007	KU952175	KU952301	KU953055	KU952929	KU952427	KU952552	KU953180	KU953305	KU952678	KU952804
<i>Oryza sativa</i>	284		Tocantins	2007	KU952158	KU952284	KU953038	KU952912	KU952410	KU952535	KU953163	KU953288	KU952661	KU952787
<i>Oryza sativa</i>	323		Tocantins	2006	KU952159	KU952285	KU953039	KU952913	KU952411	KU952536	KU953164	KU953289	KU952662	KU952788

P. oryzae pathotype Oryza

364	IC-17	<i>Oryza sativa</i>	Tocantins	2007	KU952160	KU952286	KU953040	KU952914	KU952412	KU952537	KU953165	KU953290	KU952663	KU952789
421	ID-2	<i>Oryza sativa</i>	Tocantins	2007	KU952161	KU952287	KU953041	KU952915	KU952413	KU952538	KU953166	KU953291	KU952664	KU952790
611	IA-65	<i>Oryza sativa</i>	Tocantins	2007	KU952162	KU952288	KU953042	KU952916	KU952414	KU952539	KU953167	KU953292	KU952665	KU952791
641	IB-41	<i>Oryza sativa</i>	Goiás	2007	KU952163	KU952289	KU953043	KU952917	KU952415	KU952540	KU953168	KU953293	KU952666	KU952792
658	IB-9	<i>Oryza sativa</i>	Goiás	2006	KU952164	KU952290	KU953044	KU952918	KU952416	KU952541	KU953169	KU953294	KU952667	KU952793
674	IB-33	<i>Oryza sativa</i>	Goiás	2007	KU952165	KU952291	KU953045	KU952919	KU952417	KU952542	KU953170	KU953295	KU952668	KU952794
678 ^{abc}	IA-33	<i>Oryza sativa</i>	Goiás	2006	KU952166	KU952292	KU953046	KU952920	KU952418	KU952543	KU953171	KU953296	KU952669	KU952795
695	IA-41	<i>Oryza sativa</i>	Tocantins	2007	KU952167	KU952293	KU953047	KU952921	KU952419	KU952544	KU953172	KU953297	KU952670	KU952796
704 ^{ac}	IA-1	<i>Oryza sativa</i>	Tocantins	2007	KU952168	KU952294	KU953048	KU952922	KU952420	KU952545	KU953173	KU953298	KU952671	KU952797
706	IA-25	<i>Oryza sativa</i>	Tocantins	2007	KU952169	KU952295	KU953049	KU952923	KU952421	KU952546	KU953174	KU953299	KU952672	KU952798
8762 ^{abc}	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952170	KU952296	KU953050	KU952924	KU952422	KU952547	KU953175	KU953300	KU952673	KU952799
8763	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952171	KU952297	KU953051	KU952925	KU952423	KU952548	KU953176	KU953301	KU952674	KU952800
8772	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952172	KU952298	KU953052	KU952926	KU952424	KU952549	KU953177	KU953302	KU952675	KU952801
8844	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952173	KU952299	KU953053	KU952927	KU952425	KU952550	KU953178	KU953303	KU952676	KU952802
8847	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952174	KU952300	KU953054	KU952928	KU952426	KU952551	KU953179	KU953304	KU952677	KU952803
10659 ^b	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952153	KU952279	KU953033	KU952907	KU952405	KU952530	KU953158	KU953283	KU952656	KU952782
10783	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952154	KU952280	KU953034	KU952908	KU952406	KU952531	KU953159	KU953284	KU952657	KU952783
10877	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952155	KU952281	KU953035	KU952909	KU952407	KU952532	KU953160	KU953285	KU952658	KU952784
10879	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952156	KU952282	KU953036	KU952910	KU952408	KU952533	KU953161	KU953286	KU952659	KU952785
10880 ^{abc}	-	<i>Oryza sativa</i>	Central Brazil	2013	KU952157	KU952283	KU953037	KU952911	KU952409	KU952534	KU953162	KU953287	KU952660	KU952786
Outgroup isolates														
<i>P. pennsylvanica</i> , 12.0.100		<i>Cenchrus echinatus</i>	Mato Grosso do Sul	2012	KU963214	KU963216	KU953118	KU963218	KU952490	KU963220	KU953243	KU953368	KU963222	KU952867
<i>P. grisea</i> , 12.0.082		<i>Digitaria sanguinalis</i>	Mato Grosso do Sul	2012	KU963215	KU963217	KU953119	KU963219	KU952491	KU963221	KU953244	KU953369	KU963223	KU952868

^a Isolates included in the cultural and morphological characterization assays.

^b Isolates included in the pathogenicity spectra assays.

^c Isolates listed in the Taxonomy section as specimens examined.

^d „-“ indicates no data available.

were used as outgroups. Sequence data from the 10 loci were assembled, aligned, and concatenated using Geneious R v. 9.0.5 (Biomatters, Auckland, New Zealand) for further phylogenetic analyses.

The phylogeny for the *Pyricularia* species was reconstructed through Bayesian inference using BEAST v. 1.8.2 and in-files created with the help of BEAUti (Drummond et al. 2012). The 10-locus dataset was partitioned and the best substitution model for each locus was determined using JModelTest2 (Darriba et al. 2012). Exploratory BEAST runs were conducted to determine the optimal clock- and tree-models. Model comparisons were based on the likelihoods using the Akaike information criterion (AICM) as implemented in the program Tracer v. 1.6 (Rambaut et al. 2014). The selected nucleotide substitution model was GTR for all loci, the strict clock model and the birth-death speciation process as the tree model.

Four independent final runs were conducted with MCMC length set to 10^8 generations with sampling intervals every 1 000 generations. Runs were assessed for convergence and combined using LogCombiner v. 1.8.0, which is part of the BEAST package. Posterior sampled trees were extracted using TreeAnnotator v. 1.8.2. (Drummond et al. 2012) with the following parameters: burn-in 10 %, 0.50 posterior probability limit, maximum clade credibility target tree type, and mean node height. The final tree was visualised with FigTree v. 1.4.2 (Institute of Evolutionary Biology, University of Edinburgh, <http://tree.bio.ed.ac.uk/software/figtree>). A phylogenetic tree was reconstructed for *MPG1* using the same settings as described for the combined tree. The resulting trees and respective alignments were deposited into TreeBASE (submission 19365). Based on the phylogenetic results, non-fixed and fixed nucleotide differences across all loci among the major clades were calculated using DnaSP (Librado & Rozas 2009).

Cultural characterisation

To examine macroscopic features, a representative subgroup of 30 isolates (Table 1) were grown on Corn Meal Agar (CMA), Malt Extract Agar (MEA), Oatmeal Agar (OA), Potato Dextrose Agar (PDA), and Synthetic Nutrient-poor Agar (SNA). All media were prepared as previously described (Crous et al. 2009) and amended with streptomycin sulphate (INLAB, São Paulo, Brazil) 0.05 g/L, and chloramphenicol (INLAB, São Paulo, Brazil) 0.05 g/L.

Stored isolates were re-activated on PDA. For this assay, a 6-mm-diam disk of colonized PDA from a 7-d-old re-activated culture was transferred to the centre of a Petri plate containing one of the media described above. Colony diameter and cultural features were assessed after 7 d of incubation at 25 °C under a 12 h dark / 12 h fluorescent light regime, following the procedures described by Klaubauf et al. (2014). Three replicates were made for each isolate and the assay was conducted twice. For colony descriptions, isolates were grouped according to their clustering in the phylogenetic analyses. A general description representing the colony morphology of each group of isolates was recorded. In addition, one isolate from each group was chosen as representative of the group.

Morphological characterisation

The same subgroup of 30 isolates selected for the description of colony morphology was examined using bright field and electron microscopy to characterise fungal structures. Isolates were re-activated on CMA and incubated for 7 d at 25 °C in darkness. They were subsequently transferred to SNA with sterile barley seeds to induce sporulation and incubated for 3 wk at 25 °C under a 12 h dark / 12 h fluorescent light regime. Samples were prepared following methods described previously (Bozzola & Russell 1999).

Table 2 Primers used in this study.

Locus	Forward primer (5' - 3')	Reverse primer (5' - 3')	AT (°C) ^a	Expected PCR product (bp)	Reference
<i>ACT</i>	ACT-34F: CGTCTTCCGTAAGTGCCC	ACT-322R: GCCCATACCAATCATGATAC	58	279	This study
<i>BAC6</i>	BAC6-F: ACATCATTGTCTCCTCGTC	BAC6-R: GTTCCTGTCTATTTCATTTCAA	54	283	Couch et al. 2005
<i>βT-1</i>	BT-26F: CCAGCTCAACTCTGATCTCC	BT-630R: GGTACTCGGAAACAAGATCG	56–58 ^b	604	This study
<i>CAL</i>	CAL-35F: CTTACCGAAGAGCAAGTTTCCG	CAL-607R: TYTTCCTGGCCATCATGGTS	55	648	This study
<i>CH7-BAC7</i>	CH7-BAC7-F: AAGACACGAGAGCAAAGAAAGAAG	CH7-BAC7-R: CGATACATTACAGTGCCTACGAA	55	313	Couch et al. 2005
<i>CH7-BAC9</i>	CH7-BAC9-F: TGTAAGAAGCTCGGTGACTGAT	CH7-BAC7-R: AGTGTGCTTGAACGGCTAA	59	296	Couch et al. 2005
<i>CHS1</i>	CHS-79F: TGGGGCAAGGATGCTTGGAAAGAAG	CHS-354R: TGGAAGAACCATCTGTGAGAGTTG	55	300	Carbone & Kohn 1999
<i>EF-1α</i>	EF-98F: CTYGGTGTAGGCAGCTCA	EF-820R: GAAMTTCAGGCRATGTGGG	55	722	This study
<i>MPG1</i>	MPG1-F: AGATCCCATCGAGCTTCTC	MPG1-R: TCCTCACAGAACTCCAAC	55	368	Couch et al. 2005
<i>NUT1</i>	NUT1-F: AAGTATGGCGCTTCTTCCAGC	NUT1-R: GCGCATTGGTCTTTAGTGGT	55	268	Couch et al. 2005

^a AT: Annealing temperature.

^b AT of 56 °C was used with DNA from isolates obtained from wheat and rice, and annealing temperature of 58 °C was used with DNA of isolates obtained from other poaceous hosts.

Observations were made with a Nikon SMZ25 stereo-microscope, and with a Zeiss Axio Imager 2 light microscope using differential interference contrast (DIC) illumination and a Nikon DS-Ri2 camera and software. The bright field images were taken with a Nikon SMZ1500 stereoscope microscope using NIS Elements D 3.2 software. Scanning electron microscope (SEM) images and measurements were acquired on a Zeiss LEOEVO 40 microscope using SmartSem Zeiss software (Oberkochen, Germany) operating at 10 kV and 10 to 30 mm work distance. When possible, biometric data were obtained from 30 observations per fungal structure per isolate. The photo plates were created on Corel Draw X7 software (Corel Corporation, Ottawa, Canada).

Pathogenicity spectrum

A subgroup of 18 isolates was tested for pathogenicity spectra in greenhouse assays on barley (*Hordeum vulgare*) cvs. BRS Korb, signal grass (*Urochloa brizantha*, ex *Brachiaria brizantha*) cvs. Piatã and Marandú, oats (*Avena sativa*) cvs. EMBRAPA 29 and IAPAR 61, rice (*Oryza sativa*) cv. IRGA 409, and wheat cv. Anahuac 75. Seeds of the different hosts were planted in 10-cm-diam plastic pots filled with Tropstrato HT potting mix (Vida Verde, Mogi Mirim, São Paulo, Brazil). Fifteen seeds were planted per pot. Fifteen d after seedling emergence, pots were thinned to eight seedlings per pot for barley, signal grass, oats, and rice; and to five seedlings per pot for wheat. Pots were kept in the greenhouse under natural conditions until inoculation and watered daily from the top. Plants were fertilised with NPK 10 : 10 : 10 granular fertiliser (N : P₂O₅ : K₂O, Vida Verde, Mogi Mirim, São Paulo, Brazil). A forty gram dose of NPK granular fertiliser was sprinkled across every 100 pots 1 d after emergence. Fertilisation was repeated every 15 d until inoculation. In addition, rice plants were fertilised with a solution of 4 g/L FeSO₄·7H₂O (Dinâmica, Diadema, São Paulo, Brazil) once after emergence, with 1 L of solution applied to every 100 pots.

Isolates were recovered from long-term storage and re-activated on PDA plates and then transferred either to OA plates (rice-derived isolates) or PDA plates (wheat and other isolates originating from poaceous hosts). Fifteen plates were prepared for each isolate. Plates were incubated for 15 d at 25 °C under a 12 h dark/12 h fluorescent light regime. Mycelium was gently scraped and washed with 3–5 mL of sterile distilled water amended with Tween 80 (two drops/L) to release the spores. Conidia concentration was quantified using a Neubauer counting chamber and adjusted to 1 × 10⁵ spores/mL for inoculation. Pathogenicity assays were conducted on seedlings, 1-mo-old plants at growth stage 14 (Zadocks et al. 1974) on all hosts, and on immature heads of 2-mo-old wheat plants at the be-

ginning of anthesis in growth stage 60 (Zadocks et al. 1974). Spore suspensions (1 × 10⁵ spores/mL) were uniformly applied either onto the adaxial leaf surfaces or onto wheat heads until runoff. Fifty millilitres of spore suspension was used for every 20 inoculated pots.

Inoculated pots were placed onto plastic trays and incubated in a plant growth chamber for 7 d at 26 °C (barley, oats, rice, and wheat) or 30 °C (signal grass). Plants were kept in the dark for the first 24 h, followed by a 12 h dark/12 h fluorescent light regime. Plants were watered every other day from the bottom to avoid cross-contamination. Humidifiers were used to insure that relative humidity would stay above 85 % within the chamber during the entire experiment. Temperature and relative humidity were recorded in the chamber using an ITLOG80 Datalogger (Instrutemp, Belenzinho, São Paulo, Brazil). As negative controls, five pots of each host were mock-inoculated with sterile deionised water amended with Tween 80 (two drops/L) in each experimental replication.

Plants were examined for lesions 7 d after inoculation. For the seedling inoculation tests, the disease severity index was calculated using an ordinal scale from 0 to 5 as previously described (Urashima et al. 2005). The disease severity index (DI) was scored as follows: lesion type 0 = no visible reaction; 1 = minute, pinhead-sized spots; 2 = small brown to dark brown lesions with no distinguishable centres; 3 = small eyespot shaped lesions with grey centres; 4 = typical elliptical blast lesions with grey centres; 5 = completely dead plant. Index values 0, 1, and 2 were considered non-compatible and index values 3, 4 and 5 were considered compatible. When different types of lesions were found on a single leaf, the most abundant lesions were considered.

Disease severity on wheat heads was assessed following the procedure described by Maciel et al. (2014), calculating the percentage of each wheat head affected by blast using Assess v. 2.0 image analysis software (APS, St. Paul, Minnesota). Wheat head tissue was considered affected by blast when it was chlorotic and/or it was covered with pathogen spores. For each head, a picture from each side of the head was taken, and the percentage of affected area in the two pictures was averaged.

Seedling and head inoculation experiments were conducted using a one-factor completely randomized unbalanced design. Five pots containing five (wheat) or eight (barley, signal grass, oats, and rice) plants in the seedling tests, or five non-detached heads in the wheat-head tests were inoculated with each of the 18 isolates. The seedling inoculation experiments were conducted twice. The head inoculation experiment was conducted six times, but only two randomly chosen replicates were used for further statistical analyses. For statistical analyses,

isolates were grouped according to their phylogenetic clustering (i.e. based on the species clades identified using the 10 loci sequences).

Analyses of variance (ANOVA) were performed to evaluate the effects of experiment's replicates, *Pyricularia* species, and their interactions in the different inoculation tests. Analyses were performed independently for each host species. For non-parametric data (seedlings inoculation tests) ANOVAs were conducted using the PROC NPAR1WAY procedure computed with the Wilcoxon rank-sum test and by using Monte Carlo estimations for the exact *p*-values (*P*) with the EXACT/MC statement, at $\alpha = 0.01$. A Dunn all Pairs for Joint Ranks test was used for non-parametric means comparisons. In the seedlings inoculation experiment, replicates were not significantly different (exact $P \geq 0.05$), thus the two replicates were combined for these analyses. For parametric data (wheat heads inoculation tests) ANOVAs were conducted with the PROC GLM procedure, considering species as fixed factors and isolates as random factors nested inside species factors. Fisher's protected Least Significant Difference (LSD) test was used for comparison of disease severity means for species, at $\alpha = 0.05$. Since the experiment was unbalanced, the harmonic cell size was used to calculate the average LSD. The experiment effect was statistically significant ($P = 0.02$), therefore the two replicates of the experiment were analysed independently. All statistical analyses were performed with Statistical Analysis System program, v. 9.4 (SAS Institute, Cary, North Carolina)

RESULTS

Phylogenetic analyses

The final alignment for partial sequences of the 10 genes had a total length of 3 381 bases (3 301 un-gapped bases) from 125 isolates, including sequences retrieved from Brazilian isolates of *P. grisea* and *P. pennisetigena* used as outgroups. A total of 471 polymorphic sites were found, equivalent to 14.3 % of the un-gapped alignment total length, and 168 of these sites (5.1 %) were phylogenetically informative (Table 3). This resulted in 109 multilocus haplotypes, i.e. 87.2 % of isolates had a unique multilocus haplotype.

The Bayesian analyses grouped the isolates into three major phylogenetic clades (Fig. 1, 2). In the 10-locus phylogeny, Clade 1 (Bayesian posterior probability, BPP = 1) comprised isolates exclusively associated with rice and corresponds to the previously described *P. oryzae* pathotype *Oryza* (PoO). Clade 2 (BPP = 0.99) comprised isolates almost exclusively associated with wheat. A single isolate (12.0.009i) collected from signal grass plants invading a wheat field in Paraná state also clustered within this clade. This clade corresponds to the previously described *P. oryzae* pathotype *Triticum* (PoT). Clade 3 (BPP = 0.99) contained isolates obtained from wheat as well as other *Poaceae* hosts. Based on the combined evidence presented in this study, we propose that this clade is distinct from *P. oryzae* and represents a new species, *Pyricularia graminis-tritici* (Pgt).

Non-fixed and fixed nucleotide differences among the three identified phylogenetic clades were examined for each locus, excluding the outgroups (Table 3, 4). A total of 242 polymorphic sites were found, corresponding to 7.3 % of the un-gapped alignment total length. Of those sites, 120 (3.6 %) were phylogenetically informative. Four of the 10 loci (*βT-1*, *CH7-BAC9*, *EF-1α*, and *MPG1*) showed a total of 18 (0.6 %) fixed differences across the three clades (Table 4, 5). *Pyricularia graminis-tritici* could be distinguished from PoT by 14 differences at *MPG1*. These fixed differences were at the following positions:

Table 3 Number of polymorphic sites in ten loci across *Pyricularia* species examined in this study.

Locus	Alignment length (bp)	Un-gapped sequence mean length (bp)	Polymorphic sites ^a	
			including outgroups ^b	excluding outgroups ^c
<i>ACT</i>	184	179	16 (2) ^d	0 (0)
<i>BAC6</i>	254	253	18 (0)	0 (0)
<i>βT-1</i>	501	500	28 (9)	19 (9)
<i>CAL</i>	524	520	92 (33)	12 (5)
<i>CH7-BAC7</i>	285	285	54 (34)	54 (34)
<i>CH7-BAC9</i>	293	268	40 (20)	38 (20)
<i>CHS</i>	229	224	78 (8)	26 (2)
<i>EF-1α</i>	658	643	83 (31)	66 (30)
<i>MPG1</i>	229	205	55 (26)	22 (16)
<i>NUT1</i>	224	224	7 (5)	5 (4)
Total	3381	3301	471 (168)	242 (120)

^a Sequences of isolates 12.0.100 (*P. pennisetigena*, URM7372) and 12.0.082 (*P. grisea*, URM7371) were used as outgroups.

^b *N* = 125.

^c *N* = 123.

^d The number of phylogenetically informative sites is indicated between parenthesis.

10 (C), 13–14 (TC), 20 (A), 22–25 (CCAG), 27 (C), 33–34 (CA), 41–42 (AG), and 87 (C). Likewise, Pgt could be distinguished from PoO by 18 fixed differences. These mutations are: one fixed difference at *βT-1*: 338 (A), one at *CH7-BAC9*: 20 (C), one at *EF-1α*: 325 (T), and 15 fixed differences at *MPG1*, as follows: 4 (T), 10 (C), 13–14 (TC), 20 (A), 22–25 (CCAG), 27 (C), 33–34 (CA), 41–42 (AG), and 87 (C). PoT was differentiated from PoO only by fixed differences: one difference at *CH7-BAC9*: 20 (C) and one at *EF-1α*: 325 (T) (Table 4, 5).

Sequences for only six genes were obtained for three isolates; therefore these isolates were not included in the phylogenetic analyses. However, by analysing variation in the diagnostic genes *CH7-BAC9* and *MPG1*, we were able to assign isolate 12.0.642i to Pgt, and isolates 12.0.007i and 12.0.012i to PoT.

Cultural and morphological characterisation

For description of cultural and morphological characteristics, *Pyricularia* isolates were grouped according to their phylogenetic placement, following the assignments *P. graminis-tritici* (Pgt), *P. oryzae* pathotype *Triticum* (PoT) and *P. oryzae* pathotype *Oryza* (PoO).

In general, similar colony morphologies were observed for isolates of Pgt, PoT, and PoO on the five media tested. No morphological differences were observed among the *Pyricularia* species. Cultural and morphological characteristics observed for *Pyricularia graminis-tritici* and *Pyricularia oryzae* pathotypes *Triticum* and *Oryza* (Fig. 6–8, a–j) are described in the Taxonomy section.

Pathogenicity spectrum of *Pyricularia* spp. on wheat, barley, signal grass, oats, and rice

The replicates of the seedlings inoculation tests were combined due to the lack of experiment effect (Table 6). *Pyricularia* species caused symptoms ranging from hypersensitive response lesions composed of diminutive, 1-mm-diam brown spots (mean disease index (DI) = 1), to typical elliptical blast lesions with grey centres (> 5 mm diam), usually coalescing and causing plant death on all hosts (DI ≥ 3) (Kato et al. 2000, Cruz et al. 2016) (Fig. 3–5). This virulence variation was observed even among isolates of the same *Pyricularia* species and pathotypes, indicating the presence of host-physiological race interactions. For all tests, host seedlings or wheat heads used as negative controls showed no blast lesions on their leaves (DI = 0.00).

Table 4 Number of fixed polymorphic sites in ten loci across *Pyricularia* species.

Locus	ACT	BAC6	β T-1	CAL	CH7-BAC7	CH7-BAC9	CHS	EF-1 α	MPG1	NUT1	Total	% ^a	
Species, clade	Alignment length (bp)	184	254	501	524	285	293	229	658	229	224	3381	
	Ungapped sequence mean length (bp)	179	253	500	520	285	268	224	643	205	224	3301	
<i>P. graminis-tritici</i> vs. <i>P. oryzae</i> pathotype <i>Triticum</i>		0	0	0	0	0	0	0	14	0	14	0.42	
<i>P. graminis-tritici</i> vs. <i>P. oryzae</i> pathotype <i>Oryza</i>		0	0	1	0	0	1	0	15	0	18	0.55	
<i>P. oryzae</i> pathotype <i>Triticum</i> vs. <i>P. oryzae</i> pathotype <i>Oryza</i>		0	0	0	0	0	1	0	1	0	2	0.06	
Total		0	0	1	0	0	1	0	15	0	18	0.55	

^a Percentage of fixed mutation with reference to the total number of 3301 nucleotides in the ungapped alignment.




Fig. 1 Phylogeny inferred by Bayesian Inference from the combined sequences of 10 partial loci (actin, *BAC6*, β -tubulin, calmodulin, *CH7-BAC7*, *CH7-BAC9*, chitin synthase 1, translation elongation factor 1- α , *MPG1* hydrophobin, and nitrogen regulatory protein 1) from isolates of *Pyricularia* spp. The 50 % majority-rule consensus tree is shown. The numbers above the branches are the Bayesian posterior probabilities (BPP) for node support with BPP > 0.95. *Pyricularia grisea* and *P. pennisetigena* were used as outgroups. The original host of the isolate can be distinguished by the colour of the isolate number: black = wheat; green = other poaceous hosts; and orange = rice. The asterisk (*) indicates the isolates listed in the Taxonomy section as specimens examined.

Table 5 Fixed polymorphic sites in four loci across *Pyricularia* spp.

Species, clade	Locus	$\beta T-1$	CH7-BAC9	EF-1 α	MPG1															
	Alignment position	776	1771	2597	2934	2940	2943	2944	2950	2952	2953	2954	2955	2957	2964	2965	2973	2974	3019	
	Locus position	338	20	325	4	10	13	14	20	22	23	24	25	27	33	34	41	42	87	
<i>Pyricularia graminis-tritici</i>	A	C	T	T	C	T	C	A	C	C	A	G	C	C	A	A	G	C		
<i>P. oryzae</i> pathotype <i>Triticum</i>	A/C	C	T	T/C	T	C	G	C	T	T	C	-	T	T	C	-	-	A		
<i>P. oryzae</i> pathotype <i>Oryza</i>	C	A	C	C	T	C	G	C	T	T	C	-	T	T	C	-	-	A		
<i>P. pennisetigena</i>	A	C	C	T	A	A	T	T	A	T	C	A	T	T	C	-	G	A		
<i>P. grisea</i>	C	C	C	A	T	T	T	C	A	T	G	G	C	C	G	A	-	A		




Fig. 2 Phylogeny inferred by Bayesian Inference from the sequences of the *MPG1* hydrophobin locus from isolates of *Pyricularia* spp. The 50 % majority-rule consensus tree is shown. The numbers above the branches are the Bayesian posterior probabilities (BPP) for node support with BPP > 0.95. *Pyricularia grisea* and *P. pennisetigena* were used as outgroups. The original host of the isolate can be distinguished by the colour of the isolate number: black = wheat; green = other poaceous hosts; and orange = rice. The asterisk (*) indicates the isolates listed in the Taxonomy section as specimens examined.




Fig. 3 Boxplot distribution of leaf blast severity of seedlings of five poaceous hosts in response to inoculations with isolates of *P. graminis-tritici* (Pgt, $N = 7$), *P. oryzae* pathotype *Triticum* (PoT, $N = 7$), and *P. oryzae* pathotype *Oryza* (PoO, $N = 4$). Boxplots represent blast severity as mean disease index assessed 7 d after inoculation using an ordinal scale from 0 to 5, and based on lesion type (Urashima et al. 2005). Disease index means with the same letter are not significantly different according to Dunn's All Pairs for Joint Ranks non-parametric test ($P > \chi^2 \leq 0.05$). a. Inoculation tests on seedlings of wheat (*Triticum aestivum*); b. barley (*Hordeum vulgare*) cv. BRS Korbell; c. signal grass (*Urochloa brizantha*, ex *Brachiaria brizanta*) cv. Marandú; d. signal grass cv. Piatã; e. oats (*Avena sativa*) cv. EMBRAPA 29; f. oats cv. IAPAR 61; g. rice (*Oryza sativa*) cv. IRGA 409.

Table 6 Pathogenicity of isolates of *Pyricularia* spp. on seedlings of five poaceous hosts.

Species	Host Cultivar	Mean scores for disease index ^a						
		Wheat	Barley	Signal grass		Oat		Rice
		Anahuac 75	BRS Korbell	Marandú	Piatã	EMBRAPA 29	IAPAR 61	IRGA 409
<i>Pyricularia graminis-tritici</i> (N = 7)		4.0882 a	3.8286 a	1.7612 a	0.3857 ab	3.4328 a	3.4627 a	0.0000 b
<i>P. oryzae</i> pathotype <i>Triticum</i> (N = 7)		4.4857 a	3.8986 a	2.0882 a	0.4714 a	2.7121 a	3.0145 a	0.0143 b
<i>P. oryzae</i> pathotype <i>Oryza</i> (N = 4)		2.0000 b	3.9143 a	0.1750 b	0.2051 b	1.2750 b	0.8500 b	1.8000 a
Species effect								
χ^2		80.6093	0.5303	48.8753	2.9844	56.0390	81.2610	92.7152
$P > \chi^2$		< 0.0001	0.7671	< 0.0001	0.2249	< 0.0001	< 0.0001	< 0.0001
Experiment effect								
χ^2		1.8216	3.9535	0.5244	2.9081	2.3851	0.3639	0.7286
$P > \chi^2$		0.1771	0.0500	0.4690	0.0881	0.1225	0.5463	0.3934

^a Mean disease index was averaged over five repetitions per test, and two test replicates were conducted. Each repetition (pot) had five seedlings for wheat, and eight seedlings for the other hosts. Disease index was assessed 7 d after inoculation using an ordinal scale from 0 to 5, and based on lesion type (Urashima et al. 2005). In this scale, 0 = no visible reaction; 1 = minute, pinhead-sized spots; 2 = small brown to dark brown lesions with no distinguishable centers; 3 = small eyespot shaped lesions; with grey centers; 4 = typical elliptical blast lesions with grey centers; 5 = complete dead plant. Disease index means with the same letter are not significantly different according to Dunn's All Pairs for Joint Ranks non-parametric test ($P > \chi^2 \leq 0.05$).

Table 7 Pathogenicity of isolates of *Pyricularia* spp. on non-detached heads of wheat (*Triticum aestivum*) cv. Anahuac 75.

Species, clade	Disease index (% head affected area) ^a			
	Experiment 1		Experiment 2	
	Least Mean Square	Standard Error	Least Mean Square	Standard Error
<i>Pyricularia graminis-tritici</i> (N = 7)	57.0364 a	1.6566	47.9202 a	2.3065
<i>P. oryzae</i> pathotype <i>Triticum</i> (N = 7)	39.7740 b	1.6996	43.6509 a	2.3065
<i>P. oryzae</i> pathotype <i>Oryza</i> (N = 4)	2.1330 c	2.1241	8.3485 b	2.8691
Species effect				
F	209.0400		65.2000	
P	< 0.0001		< 0.0001	
LSD	5.123		7.016	

^a Disease index was calculated as the percentage of the wheat head affected by blast using Assess v. 2.0 Image Analysis software. Head tissue was considered diseased when it was chlorotic and/or covered in pathogen spores. Disease was assessed 7 d after inoculation. Mean disease index was averaged over five repetitions (wheat heads) for each test replicate. The inoculation experiment was conducted twice, and replicates were analyzed independently due to significant experiment effect ($P = 0.0170$). Disease index means with the same letter are not significantly different according to Fisher's protected Least Significant Difference (LSD) test at $P \leq 0.05$.




Fig. 4 Boxplot distribution of blast severity observed on heads of wheat (*Triticum aestivum*) cv. Anahuac after inoculations with isolates of *P. graminis-tritici* (Pgt, N = 7), *P. oryzae* pathotype *Triticum* (PoT, N = 7), and *P. oryzae* pathotype *Oryza* (PoO, N = 4). Heads were not detached from the plant. Boxplots represent blast severity as mean disease index assessed 7 d after inoculation as percentage wheat head affected by blast using Assess v. 2.0 Image Analysis software. Head tissue was considered diseased when it was chlorotic and/or covered in pathogen spores. The test was conducted twice, and replicates (experiment 1 and 2) were analysed independently (a, b). Disease index means with the same letter are not significantly different according to Fisher's protected Least Significant Difference test at $P \leq 0.05$.




Fig. 5 Blast symptoms on leaves and heads of poaceous host after inoculation with *Pyricularia* species. Inoculated hosts: a and f. wheat (*Triticum aestivum*); b. barley (*Hordeum vulgare*); c. signal grass (*Urochloa brizantha*, ex *Brachiaria brizantha*); d. oats (*Avena sativa*); e. rice (*Oryza sativa*). *Pyricularia* species: *Pyricularia graminis-tritici* (Pgt), *P. oryzae* pathotype *Triticum* (PoT), and *P. oryzae* pathotype *Oryza* (PoO). Control plants (Ctr) were inoculated with sterile deionized water amended with Tween 80 (2 drops/L). Plants were assessed for disease symptoms 7 d after inoculation.

Inoculation tests on seedlings of wheat cv. Anahuac 75 showed significant differences among *Pyricularia* species in pathogenicity ($P > \chi^2 < 0.0001$). Seedlings were highly susceptible to isolates of PoT and Pgt (DIs of 4.48 and 4.09, respectively). In addition, isolates of PoO caused lesions on wheat seedlings (DI = 2.00); however, conspicuous differences were observed in the levels of virulence of isolates of this group. Isolates 8762 and 10659 sporadically produced lesions that ranged from minute, pinhead-sized spots (type 1 lesion) to small eyespot shaped lesions with grey centres (type 3 lesions). On the other hand, isolates 678 and 10880 consistently produced typical elliptical blast lesions with grey centres (type 4 lesions) (Fig. 3a, 5a).

Seedlings of barley cv. BRS Korbell did not show significant differences in their susceptible response to the inoculated *Pyricularia* species ($P > \chi^2 = 0.7671$). All species were highly virulent on this host (DIs ≥ 3.82), showing that barley is very susceptible to both wheat and rice blast pathogens (Fig. 3b, 5b).

Inoculations on signal grass seedlings showed that cv. Marandú was more susceptible to *Pyricularia* species than cv. Piatã. On cv. Marandú, PoT (DI = 2.08) showed the highest level of virulence, but it was not significantly different from Pgt (DI = 1.76). PoO was not pathogenic on this cultivar (DI = 0.18). None of the species were pathogenic on signal grass cv. Piatã (DIs

ranged from 0.21 to 0.47, and were not significantly different at $P > \chi^2 = 0.2249$) (Fig. 3c, d, 5c).

Inoculation tests on oats showed similar seedling reactions for cvs. EMBRAPA 29 and IAPAR 61. Both Pgt and PoT had similar, high average levels of aggressiveness with DIs > 2.71 for cv. EMBRAPA 29 and DI > 3.01 for cv. IAPAR 61. Furthermore, significant differences in the level of aggressiveness of individual isolates of these species were observed. The most aggressive isolates on oats cv. EMBRAPA 29 were 12.0.534i (Pgt), 12.1.169 and 12.1.119 (both PoT), and the least aggressive isolates were 12.0.607i (Pgt), 12.1.032i and 12.1.291 (both PoT). Likewise, on cv. IAPAR 61 the most aggressive isolates were 12.0.607i (Pgt), 12.1.158 and 12.1.119 (both PoT), and the least aggressive isolates were 12.0.642i (Pgt), 12.0.009i and 12.1.291 (both PoT). Isolates of PoO showed the lowest level of aggressiveness on oats (DI = 1.28 on cv. EMBRAPA 29, and 0.85 on cv. IAPAR 61), significantly lower ($P > \chi^2 < 0.0001$) compared to PoT and Pgt. Differences in virulence among isolates of PoO were significant only on cv. IAPAR 61, on which isolate 10659 was the most aggressive while isolate 8762 was not pathogenic (Fig. 3e, f, 5d).

Inoculation tests on rice seedlings showed generally low levels of disease severity. On cultivar IRGA 409, PoO was pathogenic

with a mean DI = 1.80 which was significantly different from the DI of the other two species ($P > \chi^2 < 0.0001$). Pgt and PoT were not pathogenic on rice (DI = 0.00 and DI = 0.01, respectively). PoO isolates showed a wide range of aggressiveness. Whereas isolates 8762 and 10880 consistently produced small eyespot-shaped lesions with grey centres (type 3 lesions) and sporadically typical elliptical blast lesions (type 4 lesions), isolate 678 produced small dark brown lesions with no distinguishable centres (type 2 lesions) and isolate 10659 sporadically produced type 2 lesions or no lesions at all on cv. IRGA 409 (Fig. 3h, 5e). This variation in virulence among the isolates is consistent with race-cultivar interactions.

A significant experiment effect was observed in the wheat head inoculation tests ($P = 0.02$). Therefore, statistical analyses of the two test replicates were conducted independently (Table 7, Fig. 4, 5f). The mean disease indexes obtained for PoT and PoO were higher in the second experiment; nevertheless, results from both experiments were congruent. All species tested were pathogenic on heads of wheat cv. Anahuac 75 and significant differences were found in their levels of aggressiveness ($P < 0.0001$ for both experiment 1 and experiment 2). Pgt was the most aggressive species, followed by PoT (Table 7). Isolates of PoO were able to infect wheat heads, but the disease did not progress to more than 10 % of the head of cv. Anahuac 75. However, similar to the seedling inoculation tests, PoO isolate 10880 was very aggressive on wheat heads, infecting 20–60 % of the inoculated heads (mean DI = 33.39 %; Fig. 4, 5f).

TAXONOMY

Pyricularia graminis-tritici V.L. Castroagudin, S.I. Moreira, J.L.N. Maciel, B.A. McDonald, Crous & P.C. Ceresini, sp. nov. — MycoBank MB816086; Fig. 6

Etymology. Referring to the major association of this fungal species with multiple grasses, and to the most common cultivated species this fungal species infects causing blast, *Triticum aestivum*.

Typus. BRAZIL, Goiás, isolated from head of *Triticum aestivum*, 2012, J.L.N. Maciel (holotype HISA 10298, culture ex-type URM7380 = CML 3547 = isolate 12.1.037).

On SNA on sterile barley seeds — *Mycelium* consisting of smooth, hyaline, branched, septate hyphae, 2–3 µm diam. *Conidiophores* solitary, erect, straight or curved, unbranched, 1–5-septate, medium brown, smooth, (14–)125(–255) × (1–)3.5(–6) µm. Abundant conidiogenesis observed on the top half of the conidiophore. *Conidiogenous cells* 50–80(–170) × 3–5 µm, terminal and intercalary, pale brown, smooth, forming a rachis with sympodial proliferation, with several protruding denticles, 1–2 µm long, 1.5–2 µm diam. *Conidia* solitary, pyriform to obclavate, pale brown, finely verruculose, granular to guttulate, 2-septate, (23–)25–29(–32) × (8–)9(–10) µm; apical cell 10–13 µm height, basal cell 6–9 µm long; frill hilum, protruding, 1–1.5 µm long, 1.5–2 µm diam, unthickened, not darkened; central cell turning dark brown with age. *Chlamydospores* and *microconidia* not observed.

Culture characteristics — Colonies on CMA with moderate dark grey aerial mycelium, irregular margins, reaching up to 6.5 cm diam after 1 wk; reverse dark grey. Colonies on MEA with abundant white aerial mycelium, and pale grey sporulation at the centre; reaching up to 7.6 cm diam after 1 wk; reverse dark grey; sometimes, fewer colonies (5.1 cm diam) with dark grey sporulation at centre and abundant white aerial mycelium at margins. Colonies on OA with dark grey sporulation in concentric circles, with sparse margins, up to 5.8 cm; reverse pale grey; sometimes, larger growth with abundant white aerial mycelium, pale grey at the centre. Colonies on PDA with abundant white aerial mycelium, olivaceous at centre, growth in concentric

circles, up to 6.5 cm diam; reverse black in centre with white margins. Colonies on SNA with sparse olivaceous mycelium irregular margins, up to 5.2 cm diam; reverse sparse olivaceous.

Specimens examined. BRAZIL, Goiás, isolated from head of *Triticum aestivum*, 2012, J.L.N. Maciel (URM7380, isolate 12.1.037); Mato Grosso do Sul, isolated from leaves of *Avena sativa*, 2012, J.L.N. Maciel (URM7366 = CML3516, isolate 12.0.345); Mato Grosso do Sul, isolated from leaves of *Echinochloa crusgalli*, 2012, J.L.N. Maciel (URM7381, isolate 12.0.326); Mato Grosso do Sul, isolated from leaves of *Elionorus candidus*, 2012, J.L.N. Maciel (URM7377, isolate 12.0.194); Mato Grosso do Sul, isolated from leaves of *Urochloa brizantha*, 2012, J.L.N. Maciel (URM7367 = CML3517, isolate 12.0.366); Paraná, isolated from leaves of *Cenchrus equinatus*, 2012, J.L.N. Maciel (URM7378, isolate 12.0.642i); Paraná, isolated from leaves of *Cynodon* spp., 2012, J.L.N. Maciel (URM7375, isolate 12.0.578i); Paraná, isolated from leaves of *Digitaria sanguinalis*, 2012, J.L.N. Maciel (URM7376, isolate 12.0.555i); Paraná, isolated from leaves of *Eleusine indica*, 2012, J.L.N. Maciel (URM7365 = CML3518, isolate 12.0.534i); Paraná, isolated from leaves of *Rhynchelytrum repens*, 2012, J.L.N. Maciel (URM7384, isolate 12.0.607i); Rio Grande do Sul, isolated from head of *T. aestivum*, 2012, J.L.N. Maciel (URM7387, isolate 12.1.191).

Notes — *Pyricularia graminis-tritici* causes blast disease on *Triticum aestivum*, *Avena sativa*, *Hordeum vulgare*, and *Urochloa brizantha* but not on *Oryza sativa*.

Based on morphological and cultural comparisons, isolates of *P. graminis-tritici* are indistinguishable from those of *P. oryzae* pathotypes *Oryza* and *Triticum*. However, these taxa are readily distinguished based on their DNA phylogeny, host range and pathogenicity spectra. Sequencing of the *MPG1* gene is a diagnostic tool to distinguish *P. graminis-tritici* from *P. oryzae*.

Pyricularia oryzae Cavara, Fungi Longobard. Exsicc. 1: no. 49. 1891

= *Magnaporthe oryzae* B.C. Couch, Mycologia 94: 692. 2002.

Pyricularia oryzae* pathotype *Triticum (Kato et al. 2000) — Fig. 7

On SNA on sterile barley seeds — *Mycelium* consisting of smooth, hyaline, branched, septate hyphae, 1.5–2 µm diam. *Conidiophores* solitary, erect, straight or curved, unbranched, medium brown, smooth, 60–150 × 4–6 µm, 2–3-septate; base arising from hyphae, not swollen, lacking rhizoids. *Conidiogenous cells* 40–95 × 3–5 µm, integrated, terminal and intercalary, pale brown, smooth, forming a rachis with several protruding denticles, 0.5–1 µm long, 1.5–2 µm diam. *Conidia* solitary, pyriform to obclavate, pale brown, smooth, granular to guttulate, 2-septate, (25–)27–29(–32) × (8–)9(–10) µm; apical cell 10–13 µm long, basal cell 6–9 µm long; hilum truncate, protruding, 1–1.5 µm long, 1.5–2 µm diam, unthickened, not darkened. *Chlamydospores* and *microconidia* not observed (based on isolate CPC 26580 = 12.1.132).

Culture characteristics — On CMA colonies with moderate dark grey aerial mycelium with irregular margins, sometimes with black aerial mycelium with sporulation in concentric circles, or sparse white mycelial colonies, reaching up to 5.9 cm diam after 1 wk; reverse dark grey with brown margins. On MEA, colonies presented different forms: cottony white aerial mycelia within concentric growth rings, sometimes with a grey sporulation at the centre, reaching up to 6.9 cm diam after 1 wk; reverse dark grey. Colonies on OA with grey aerial mycelium and sporulation in concentric circles; sometimes surface mycelia were white or cream, showing concentric growth, up to 7.9 cm diam; reverse dark grey; sometimes, larger growth with abundant white aerial mycelium, pale grey at the centre. PDA colonies exhibited many variations in culture, often with concentric growth: abundant white aerial mycelia and pale grey sporulation at centre; abundant white aerial mycelia; or




Fig. 6 *Pyricularia graminis-tritici*. a–j. Cultures of isolate 12.1.037 grown for 7 d at 12 h photoperiod and 25 °C in CMA (a, f), MEA (b, g), OA (c, h), PDA (d, i), and SNA (e, j) media; k–l. sporulation on SNA on sterile barley seeds; m–o. scanning electron micrographs of conidiophores and conidia; p–x. bright field microscopy images of conidiophores and conidia. — Scale bars = 10 μm.




Fig. 7 *Pyricularia oryzae* pathotype *Triticum*. a–j. Cultures of isolate 12.1.291 grown for 7 d at 12 h photoperiod and 25 °C in CMA (a, f), MEA (b, g), OA (c, h), PDA (d, i), and SNA (e, j) media; k–l. sporulation on SNA on sterile barley seeds; m–o. scanning electron micrographs of conidiophores and conidia; p–v. bright field microscopy images of conidiophores and conidia. — Scale bars = 10 µm.

dark grey mycelia at the bottom, with white aerial mycelia up to 7 cm diam; reverse, concentric growth, black in centre with olivaceous margins. On SNA the colonies with dark green centres with sparse pale brown margins; or pale grey at the centre and sparse pale brown margins; reverse dark green to black at the centre and with pale brown margins.

Specimens examined. BRAZIL, Mato Grosso do Sul, isolated from head of *Triticum aestivum*, 2012, J.L.N. Maciel (URM7388, isolate 12.1.132); Mato Grosso do Sul, isolated from head of *T. aestivum*, 2012, J.L.N. Maciel (URM7368 = CML3521, isolate 12.1.158); Mato Grosso do Sul, isolated from head of *T. aestivum*, 2012, J.L.N. Maciel (URM7386, isolate 12.1.169); Paraná, isolated from head of *T. aestivum*, 2012, J.L.N. Maciel (URM7369 =

CML3522, isolate 12.1.291); Paraná, isolated from leaves of *Urochloa brizantha*, 2012, J.L.N. Maciel (URM7385, isolate 12.0.0091); Rio Grande do Sul, isolated from head of *T. aestivum*, 2012, J.L.N. Maciel (URM7389, isolate 12.1.205).

Pyricularia oryzae pathotype *Oryza* (Kato et al. 2000) — Fig. 8

On SNA on sterile barley seeds — *Mycelium* consisting of smooth, hyaline, branched, septate hyphae, 2–3 µm diam. *Conidiophores* were (70.5–)146.5(–247) × (3.5–)4.5(–5.5) µm, solitary, erect, straight or curved, septate, hyaline, sometimes light brown. Sometimes, the conidiophores branched. Conidio-

genous cells apical and intercalary, sporulating frequently at the apical part, with protruding denticles 0.9–1.1 μm long. *Conidia* pyriform to obclavate, narrowed towards the tip, rounded at the base, 2-septate, hyaline to pale olivaceous, (18–)24–28(–32) \times (8–)9(–10) μm ; apical cell 7–14 μm long, basal cell 7–12 μm long; hilum 1.5–2 μm diam. *Chlamydospores* and *microconidia* not observed.

Culture characteristics — On CMA the predominant colony morphology was the moderate pale grey aerial mycelium with irregular margins reaching up to 5.6 cm diam after 1 wk; reverse dark grey centre and grey edges; fewer colonies with regular

margin formed by sparse white aerial mycelia; sometimes, moderate dark grey aerial mycelium with irregular margins; or white aerial mycelium. Colonies on MEA were often pale grey, sporulation in concentric circles, with dark grey margins; sometimes dark grey at the bottom with sparse white aerial mycelia; or white colonies with regular margins, dark grey at the centre, reaching up to 7.6 cm diam after 1 wk; reverse dark grey. On OA colonies with dark grey sporulation at centre and regular margins of white aerial mycelia up to 7.3 cm. PDA colonies were variable, with grey growth in concentric circles, sometimes pale grey or olivaceous; in some cases, with regular




Fig. 8 *Pyricularia oryzae* pathotype *Oryza*. a–j. Cultures of isolate 10880 grown for 7 d at 12 h photoperiod and 25 °C in CMA (a, f), MEA (b, g), OA (c, h), PDA (d, i), and SNA (e, j) media; k–l. sporulation on SNA on sterile barley seeds; m–o. scanning electron micrographs of conidiophores and conidia; p–t. bright field microscopy images of conidiophores and conidia. — Scale bars = 10 μm .

margins of white mycelia, reaching up to 6.4 cm; reverse dark grey. On SNA colonies with pale green or dark green mycelia, with sparse margins; in rare cases with abundant pale grey aerial mycelia at centre and white mycelia in regular margins, up to 3.1 cm; reverse dark green in centre and olivaceous at the borders.

Specimens examined. BRAZIL, Central Brazil, isolated from leaves of *Oryza sativa*, 2013, *Unknown* (URM7382, isolate 8762); Central Brazil, isolated from leaves of *O. sativa*, 2013, *Unknown* (URM7370 = CML3523, isolate 10880); Goiás, isolated from leaves of *O. sativa*, 2006, *Unknown* (URM7379, isolate 678); Tocantins, isolated from leaves of *O. sativa*, 2007, *Unknown* (URM7383, isolate 704).

DISCUSSION

We conducted comprehensive phylogenetic, morphological, and pathogenicity analyses to characterise *Pyricularia* isolates associated with the blast disease on rice, wheat and other poaceous hosts from the Brazilian agro-ecosystem. Urashima, Igarashi & Kato (1993) demonstrated that the blast pathogens infecting wheat and rice were distinct. These authors also reported that isolates recovered from wheat did not infect rice and that most isolates recovered from rice did not infect wheat, except for a few isolates capable of producing small leaf lesions. Although Urashima & Kato (1998), and several follow-up studies demonstrated that the wheat and rice pathogens were phenotypically and genetically different, they have been treated as subgroups of the same species: *Pyricularia oryzae* (Urashima & Kato 1998, Kato et al. 2000, Murakami et al. 2000, Couch & Kohn 2002, Farman 2002, Klaubauf et al. 2014, Chiapello et al. 2015).

The results of our phylogenetic analyses indicate that wheat blast is caused by *Pyricularia* strains assigned to Clade 2, previously described as *P. oryzae* pathotype *Triticum*, and to Clade 3 (Fig. 1, Table 5). Here, we propose that Clade 3 is distinct from *P. oryzae* and represents a new species, *Pyricularia graminis-tritici* (Pgt).

We confirmed that the two host-associated clades *P. oryzae* pathotype *Triticum* and *P. oryzae* pathotype *Oryza* correspond to different pathotypes. This distinction is supported by the combined phylogenetic reconstruction that clearly separates the two taxa. Interestingly, the combined tree (Fig. 2) does not suggest that PoO and PoT are sister taxa. Instead, PoT forms a sister group with Pgt that includes all isolates collected from wheat and other poaceous hosts. This combined group is the sister group to the rice-associated PoO. However, we postulate that this pattern should be interpreted with caution as explained below.

Among the *Pyricularia* species examined in this study, non-fixed polymorphic sites and phylogenetically informative sites were found in nine of the ten loci examined (locus *BAC6* was monomorphic). Fixed nucleotide differences that are diagnostic for the three taxa were located in four loci: *βT-1*, *CH7-BAC9*, *EF-1α*, and *MPG1*. Among these, *MPG1* was the most diagnostic locus with 15 fixed differences. Hence, sequencing the *MPG1* locus could provide a simple and informative tool to establish the identity of *Pyricularia* isolates at the species level.

Fig. 2 shows the phylogenetic tree reconstructed for *MPG1* using the same settings as described for the combined tree. Significant differences in tree topology are visible compared to the combined tree. Variation at the *MPG1* locus can distinguish Pgt and PoO with high confidence. However, this analysis splits PoT into two sub-clades. Furthermore, PoO and PoT now join together to form the sister-group, as opposed to Pgt. The observation that single loci can produce different phylogenetic patterns has been referred to as 'phylogenetic incongruence'. The concept of genealogical concordance of different sequence loci (genealogical concordance phylogenetic species recognition, GCPSR) was proposed as a possible solution for phylogenetic

species recognition (Taylor et al. 2000, Dettman et al. 2003). In the GCPSR approach, concordant grouping of species based on several sequences is regarded as evidence for restricted exchange of genetic material and, thus, for the reproductive isolation of taxonomic units, indicating speciation. However, in an extensive analysis Grünig et al. (2007) showed that this combined phylogenetic approach also has its limits. The authors concluded that in ambiguous cases (such as cryptic species complexes) phylogenetic approaches should be complemented with population genetic analyses that more easily detect reproductive isolation between taxa. Until additional evidence emerges, likely based on comparative population genomics analyses that include entire genome sequences, we suggest a conservative interpretation and propose to maintain the pathotype-based denomination system of *P. oryzae* pathotype *Oryza* and *P. oryzae Triticum* (Kato et al. 2000), recognizing that PoT and Pgt may eventually be fused into a single, highly diverse species.

Under our experimental conditions, *P. graminis-tritici* and *P. oryzae* pathotypes *Oryza* and *Triticum* did not present consistent cultural or morphological differences. However, distinctive pathogenicity spectra were observed. *Pyricularia graminis-tritici* and *P. oryzae* pathotypes *Triticum* and *Oryza* caused blast symptoms on wheat, barley, and oats with different levels of aggressiveness. These findings agree with Urashima's pioneering observation that two different pyricularia-like pathogens caused wheat blast disease in Brazil (Urashima et al. 2005). Furthermore, our results confirmed that isolates of *P. oryzae* pathotype *Oryza* can cause blast on seedlings and heads of wheat under greenhouse conditions that favour infection, as previously reported (Urashima et al. 1993, Urashima & Kato 1998). An important question that remains to be answered is whether compatible interactions also occur under natural field conditions. Our observation that none of the wheat-derived isolates was genetically assigned to PoO suggests that PoO infections on wheat are very rare or absent under natural field conditions.

In conclusion, our study suggests that blast disease on wheat and other *Poaceae* in Brazil represents a disease complex caused by more than one species of *Pyricularia*. A recent population genomics analysis performed by D. Croll showed that the Bangladeshi wheat blast strains responsible for the 2016 outbreak were closely related to strains of *Pyricularia graminis-tritici* collected in Brazilian wheat fields (Callaway 2016). Given these findings, recognising and properly naming the causal agents of wheat blast will not only increase our understanding of the biology and epidemiology of the disease, but will also enable the establishment of proper quarantine regulations to limit the spread of these pathogens into disease-free areas that grow susceptible wheat cultivars, including Asia, Europe, and North America (McTaggart et al. 2016).

Acknowledgements This work was funded by FAPESP (São Paulo Research Foundation, Brazil) research grants to P.C. Ceresini (2013/10655-4 and 2015/10453-8), EMBRAPA/Monsanto research grant (Macroprogram II) to J.L.N. Maciel, and research grants from FINEP (Funding Authority for Studies and Projects, Brazil) and FAPEMIG (Minas Gerais Research Foundation, Brazil) to E. Alves (CAG-APQ-01975-5). P.C. Ceresini and E. Alves were supported by research fellowships from Brazilian National Council for Scientific and Technological Development - CNPq (Pq-2 307361/2012-8 and 307295/2015-0). S.I. Moreira was supported by Doctorate research fellowship from CAPES (Higher Education Personnel Improvement Coordination, Brazil). V. L. Castroagudin was supported by Post-Doctorate research fellowships from CNPq (PDJ 150490/2013-5, from 2012–2014), and FAPESP/CAPES (PDJ 2014/25904-2, from 2015–2016). We thank CAPES for sponsoring the establishment of the 'Centro de Diversidade Genética no Agroecossistema' (Pro-equipamentos 775202/2012). Authorization for scientific activities # 39131-3 from the Brazilian Ministry of Environment (MMA) / 'Chico Mendes' Institute for Conservation of Biodiversity (ICMBIO) / System for Authorization and Information in Biodiversity (ICMBIO).

REFERENCES

- Anjos JRND, Silva DBD, Charchar MJD, et al. 1996. Occurrence of blast fungus (*Pyricularia grisea*) on wheat and rye in the savanna region of Central Brazil. *Pesquisa Agropecuária Brasileira* 31: 79–82.
- Bozzola JJ, Russell LD. 1999. *Electron microscopy: principles and techniques for biologists*: 670. Boston, Jones & Bartlett Publishers.
- Callaway E. 2016. Devastating wheat fungus appears in Asia for first time. *Nature* 532: 421–422.
- Carbone I, Kohn LM. 1999. A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 91: 553–556.
- Castroagudín VL, Ceresini PC, Oliveira SC, et al. 2015. Resistance to Qol fungicides is widespread in Brazilian populations of the wheat blast pathogen *Magnaporthe oryzae*. *Phytopathology* 104: 284–294.
- Chiappello H, Mallet L, Guérin C, et al. 2015. Deciphering genome content and evolutionary relationships of isolates from the fungus *Magnaporthe oryzae* attacking different hosts plants. *Genome Biology and Evolution* 7: 2896–2912.
- Choi J, Park S-Y, Kim B-R, et al. 2013. Comparative analysis of pathogenicity and phylogenetic relationship in *Magnaporthe grisea* species complex. *PLoS ONE*: 8, 2: e57196. doi:57110.51371/journal.pone.0057196.
- Couch BC, Fudal I, Lebrun MH, et al. 2005. Origins of host-specific populations of the blast pathogen *Magnaporthe oryzae* in crop domestication with subsequent expansion of pandemic clones on rice and weeds of rice. *Genetics* 170: 613–630.
- Couch BC, Kohn LM. 2002. A multilocus gene genealogy concordant with host preference indicates segregation of a new species, *Magnaporthe oryzae*, from *M. grisea*. *Mycologia* 94: 683–693.
- Crous PW, Verkley GJM, Groenwald JZ, et al. 2009. *Fungal Biodiversity*. Utrecht, The Netherlands: CBS-KNAW Fungal Biodiversity Centre.
- Cruz MFA, Rios JA, Araujo L, et al. 2016. Infection process of *Pyricularia oryzae* on the leaves of wheat seedling. *Tropical Plant Pathology* 41: 123–127.
- Darriba D, Taboada GL, Doallo R, et al. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9: 772.
- Dettman JR, Jacobson DJ, Turner E, et al. 2003. Reproductive isolation and phylogenetic divergence in *Neurospora*: comparing methods of species recognition in a model eukaryote. *Evolution* 57: 2721–2741.
- Drummond AJ, Suchard MA, Xie D, et al. 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution* 29: 1969–1973.
- Duveiller E, Hodson D, Tiedmann A. 2010. Wheat blast caused by *Magnaporthe grisea*: a reality and new challenge for wheat research. *International Wheat Conference*, 8: 247–248.
- Duveiller E, Singh RP, Nicol JM. 2007. The challenges of maintaining wheat productivity: pests, diseases, and potential epidemics. *Euphytica* 157: 417–430.
- Farman ML. 2002. *Pyricularia grisea* isolates causing gray leaf spot on perennial ryegrass (*Lolium perenne*) in the United States: relationship to *P. grisea* isolates from other host plants. *Phytopathology* 92: 245–254.
- Grünig CR, Brunner PC, Duò A, et al. 2007. Suitability of methods for species recognition in the *Phialocephala fortinii*-*Acephala applanata* species complex using DNA analysis. *Fungal Genetics and Biology* 44: 773–788.
- Hamer JE. 1991. Molecular probes for rice blast disease. *Science* 252: 632–633.
- Hirata K, Kusaba M, Chuma I, et al. 2007. Speciation in *Pyricularia* inferred from multilocus phylogenetic analysis. *Mycological Research* 111: 799–808.
- Igarashi S, Utimada CM, Igarashi LC, et al. 1986. *Pyricularia* em trigo. 1. Ocorrência de *Pyricularia* spp. no estado do Paraná. *Fitopatologia Brasileira* 11: 351–352.
- Kato H, Yamamoto M, Yamaguchi-Ozaki T, et al. 2000. Pathogenicity, mating ability and DNA restriction fragment length polymorphisms of *Pyricularia* populations isolated from Gramineae, Bambusoideae and Zingiberaceae plants. *Journal of General Plant Pathology* 66: 30–47.
- Klaubauf S, Tharreau D, Fournier E, et al. 2014. Resolving the polyphyletic nature of *Pyricularia* (*Pyriculariaceae*). *Studies in Mycology* 79: 85–120.
- Kohli MM, Mehta YR, Guzman E, et al. 2011. *Pyricularia* blast – a threat to wheat cultivation. *Czech Journal of Genetics and Plant Breeding* 47: S130–S134.
- Librado P, Rozas J. 2009. DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25: 1451–1452.
- Lima MIP, Minella E. 2003. Occurrence of head blast in barley. *Fitopatologia Brasileira* 28: 207.
- Luo J, Zhang N. 2013. *Magnaportheiopsis*, a new genus in *Magnaportheaceae* (*Ascomycota*). *Mycologia* 105: 1019–1029.
- Maciel JLN. 2011. *Magnaporthe oryzae*, the blast pathogen: current status and options for its control. *Plant Science Reviews* 2011: 233–240.
- Maciel JLN, Ceresini PC, Castroagudín VL, et al. 2014. Population structure and pathotype diversity of the wheat blast pathogen *Magnaporthe oryzae* 25 years after its emergence in Brazil. *Phytopathology* 104: 95–107.
- McTaggart AR, Van der Nest MA, Steenkamp ET, et al. 2016. Fungal genomics challenges the dogma of name-based biosecurity. *PLoS Pathogens* 12: e1005475. doi: 10.1371/journal.ppat.1005475.
- Murakami J, Tomita R, Kataoka T, et al. 2003. Analysis of host species specificity of *Magnaporthe grisea* toward foxtail millet using a genetic cross between isolates from wheat and foxtail millet. *Phytopathology* 93: 42–45.
- Murakami J, Tosa Y, Kataoka T, et al. 2000. Analysis of host species specificity of *Magnaporthe grisea* toward wheat using a genetic cross between isolates from wheat and foxtail millet. *Phytopathology* 90: 1060–1067.
- Murata N, Aoki T, Kusaba M, et al. 2014. Various species of *Pyricularia* constitute a robust clade distinct from *Magnaporthe salvinii* and its relatives in *Magnaportheaceae*. *Journal of General Plant Pathology* 80: 66–72.
- Rambaut A, Suchard MA, Xie D, et al. 2014. Tracer v1.6, available from <http://beast.bio.ed.ac.uk/Tracer>.
- Silue DJ, Nottéghem JL, Tharreau D. 1992. Evidence for a gene-for-gene relationship in the *Oryza sativa*-*Magnaporthe grisea* pathosystem. *Phytopathology* 82: 577–580.
- Silva CP, Nomura E, Freitas EG, et al. 2009. Efficiency of alternative treatments in the control of *Pyricularia grisea* on wheat seeds. *Tropical Plant Pathology* 34: 127–131.
- Takabayashi N, Tosa Y, Oh HS, et al. 2002. A gene-for-gene relationship underlying the species-specific parasitism of *Avena/Triticum* isolates of *Magnaporthe grisea* on wheat cultivars. *Phytopathology* 92: 1182–1188.
- Taylor JW, Jacobson DJ, Kroken S, et al. 2000. Phylogenetic species recognition and species concepts in fungi. *Fungal Genetics and Biology* 31: 21–32.
- Tosa Y, Chuma I. 2014. Classification and parasitic specialization of blast fungi. *Journal of General Plant Pathology* 80: 202–209.
- Tosa Y, Hirata K, Tamba H, et al. 2004. Genetic constitution and pathogenicity of *Lolium* isolates of *Magnaporthe oryzae* in comparison with host species-specific pathotypes of the blast fungus. *Phytopathology* 94: 454–462.
- Tosa Y, Tamba H, Tanaka K, et al. 2006. Genetic analysis of host species specificity of *Magnaporthe oryzae* isolates from rice and wheat. *Phytopathology* 96: 480–484.
- Urashima AS, Galbieri R, Stabili A. 2005. DNA fingerprinting and sexual characterization revealed two distinct populations of *Magnaporthe grisea* in wheat blast from Brazil. *Czech Journal of Genetics and Plant Breeding* 41: 238–245.
- Urashima AS, Hashimoto Y, Don LD, et al. 1999. Molecular analysis of the wheat blast population in Brazil with a homolog of retrotransposon MGR583. *Annals of the Phytopathological Society of Japan* 65: 429–436.
- Urashima AS, Igarashi S, Kato H. 1993. Host range, mating type, and fertility of *Pyricularia grisea* from wheat in Brazil. *Plant Disease* 77: 1211–1216.
- Urashima AS, Kato H. 1998. Pathogenic relationship between isolates of *Pyricularia grisea* of wheat and other hosts at different host developmental stages. *Fitopatologia Brasileira* 23: 30–35.
- Valent B, Chumley FG. 1991. Molecular genetic analysis of the rice blast fungus, *Magnaporthe grisea*. *Annual Review of Phytopathology* 29: 443–467.
- Valent B, Khang CH. 2010. Recent advances in rice blast effector research. *Current Opinion in Plant Biology* 13: 434–441.
- Verzignassi RS, Poltronieri LS, Benchimol RL, et al. 2012. *Pyricularia grisea*: new pathogen on *Brachiaria brizantha* cv. Marandu in Pará. *Summa Phytopathologica* 38: 254.
- Zadocks JC, Chang TT, Konzak CF. 1974. A decimal code for the growth stages of cereals. *Weed Research* 14: 415–421.
- Zhang N, Zhao S, Shen Q. 2011. A six-gene phylogeny reveals the evolution of mode of infection in the rice blast fungus and allied species. *Mycologia* 103: 1267–1276.