




<b>Title</b>	<b>The first fossil Perilestidae (Odonata: Zygoptera) from mid-Cretaceous Burmese amber</b>
<b>Author(s)</b>	<b>Zheng, D; Wang, B; Jarzembowski, E; Chang, S; Nel, A</b>
<b>Citation</b>	<b>Cretaceous Research, 2016, v. 65, p. 199-205</b>
<b>Issued Date</b>	<b>2016</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/231813">http://hdl.handle.net/10722/231813</a></b>
<b>Rights</b>	<b>© &lt;2016&gt;. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>; This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.</b>

# Accepted Manuscript

The first fossil Perilestidae (Odonata: Zygoptera) from mid-Cretaceous Burmese amber

Daran Zheng, Bo Wang, Edmund A. Jarzembowski, Su-Chin Chang, André Nel



PII: S0195-6671(16)30092-1

DOI: [10.1016/j.cretres.2016.05.002](https://doi.org/10.1016/j.cretres.2016.05.002)

Reference: YCRES 3400

To appear in: *Cretaceous Research*

Received Date: 25 March 2016

Revised Date: 5 May 2016

Accepted Date: 8 May 2016

Please cite this article as: Zheng, D., Wang, B., Jarzembowski, E.A., Chang, S.-C., Nel, A., The first fossil Perilestidae (Odonata: Zygoptera) from mid-Cretaceous Burmese amber, *Cretaceous Research* (2016), doi: 10.1016/j.cretres.2016.05.002.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 The first fossil Perilestidae (Odonata: Zygoptera) from mid-Cretaceous Burmese  
2 amber

3

4 Daran Zheng <sup>a, b, \*</sup>, Bo Wang <sup>a, c</sup>, Edmund A. Jarzembowski <sup>a, d</sup>, Su-Chin Chang <sup>b, \*</sup>,  
5 André Nel <sup>c</sup>

6

7 <sup>a</sup> *State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of*  
8 *Geology and Palaeontology, Chinese Academy of Sciences, 39 East Beijing Road,*  
9 *Nanjing 210008, China*

10 <sup>b</sup> *Department of Earth Sciences, The University of Hong Kong, Hong Kong Special*  
11 *Administrative Region, China*

12 <sup>c</sup> *Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology,*  
13 *Chinese Academy of Sciences, 1, Beichen West Road, Beijing 100101, China*

14 <sup>d</sup> *Department of Earth Sciences, The Natural History Museum, London SW7 5BD, UK*

15 <sup>e</sup> *Institut de Systématique, Évolution, Biodiversité, ISYEB-UMR 7205-CNRS, MNHN,*  
16 *UPMC, EPHE, Muséum national d'Histoire naturelle, Sorbonne Universités, 57 rue*  
17 *Cuvier, CP 50, Entomologie, F-75005, Paris, France (anel@mnhn.fr)*

18

19 \* Corresponding author. Department of Earth Sciences, The University of Hong Kong,  
20 Hong Kong Special Administrative Region, China.

21 E-mail address: dranzheng@gmail.com (D. Zheng), suchin@hku.hk (S.-C. Chang).

22

## 23 A B S T R A C T

24 *Palaeoperilestes electronicus* gen. et sp. nov. is the first perilestid damselfly  
25 described from mid-Cretaceous Burmese amber. This new damselfly can be attributed  
26 to the family Perilestidae by the midfork being distal of the subnodus and the base of  
27 IR2 quite near to the base of RP2, both features found in the extant genera *Perilestes*  
28 and *Perissolestes*. *Palaeoperilestes electronicus* gen. et sp. nov. has a strongly  
29 zigzagged IR1, however, differing from *Perilestes* and *Perissolestes* which have a  
30 straight IR1. The discovery not only adds to the diversity of damselflies in Burmese  
31 amber, but also puts the origin of Perilestidae back to at least the mid-Cretaceous.

32  
33 Key words: Perilestidae, Zygoptera, Odonata, Cenomanian, Cretaceous, Burmese  
34 amber

35  
36 **1. Introduction**

37 The Perilestidae Kennedy, 1920 is a family of small damselflies, often called  
38 shortwings or twigtails, characterized by short wings (20–25 mm) and very long,  
39 slender and colour-banded abdomens (40–56 mm) (Williamson and Williamson 1924;  
40 Haber and Wagner, 2014). The adults often perch on plant stems or dead twigs near  
41 streams and frequent the understory and glades within dense forest, thus making them  
42 easily overlooked in the field. Perilestidae now consists of two Neotropical genera:  
43 *Perilestes* Hagen in Selys-Longchamps, 1862 and *Perissolestes* Kennedy, 1941, with  
44 the former consisting of eight species while the later comprises 11 species (Dijkstra et

45 al., 2011, 2014; Schorr and Paulson, 2015; Machado, 2015). The African endemic  
46 genus *Nubiolestes* Fraser, 1944 was previously attributed to this family (Bechly, 1996;  
47 Neiss and Neusa, 2010); however, it is the sister genus of the Perilestidae sensu stricto  
48 in the phylogenetic analysis of Dijkstra et al. (2014). In the present paper, a new  
49 damselfly, *Palaeoperilestes electronicus* gen. et sp. nov., is described from the  
50 mid-Cretaceous Burmese amber. This is the first fossil representative of the extant  
51 family Perilestidae. The new discovery increases our knowledge about these unique  
52 damselflies.

53

## 54 **2. Material and methods**

55 The specimen described herein was collected from the Hukawng Valley of  
56 Kachin Province, Myanmar (locality in Kania et al., 2015: fig. 1). The age of Burmese  
57 amber is radiometrically dated at  $98.79 \pm 0.62$  Ma (earliest Cenomanian; Cohen et al.,  
58 2013) based on U–Pb zircon dating of the volcanoclastic matrix (Shi et al., 2012).

59 The amber containing the damselfly is yellow and transparent. The damselfly is  
60 preserved together with a big horsefly. The damselfly wings are close to the amber  
61 surface and are slightly curved towards the horsefly. This makes it difficult for further  
62 polishing and obtaining clear photomicrographs. Photomicrographs were taken using  
63 a Zeiss Stereo Discovery V16 microscope system and Zen software. In most instances,  
64 incident and transmitted light were used simultaneously. All images are digitally  
65 stacked photomicrographic composites of approximately 40 individual focal planes  
66 obtained using the free software Combine ZP for a better illustration of the 3D

67 structures. The line drawings were prepared from photographs using image-editing  
68 software (CorelDraw X7 and Adobe Photoshop CS6). The specimen is housed in the  
69 Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences  
70 (NIGPAS). All taxonomic acts established in the present work have been registered in  
71 ZooBank (see below), together with the electronic publication LSID:  
72 urn:lsid:zoobank.org:pub:28289195-14CB-4434-92CD-6D6F5865ABBF.

73 The nomenclature of the odonatan wing venation used in this paper is based on  
74 the interpretations of Riek (1976) and Riek and Kukalová-Peck (1984), as modified  
75 by Nel et al. (1993) and Bechly (1996). The higher classification of fossil and extant  
76 Odonatoptera, as well as family and generic characters followed in the present work,  
77 are based on the phylogenetic system proposed by Bechly (1996) and Dijkstra et al.  
78 (2014) for the phylogeny of extant Zygoptera. Wing abbreviations are as follows:  
79 CuA, cubitus anterior; IR, intercalary radial veins; MA, median anterior; MP, median  
80 posterior; N, nodus; Pt, pterostigma; RA, radius anterior; RP, radius posterior; Sn,  
81 subnodal crossvein. All measurements are given in mm.

### 83 **3. Systematic palaeontology**

84 Order: Odonata Fabricius, 1793

85 Suborder: Zygoptera Selys-Longchamps, 1854

86 Family: Perilestidae Kennedy, 1920

87 Type genus. *Perilestes* Hagen in Selys-Longchamps, 1862

88 New genus. *Palaeoperilestes* gen. nov.

89 (urn:lsid:zoobank.org:act:08EA6631-83A4-4EBA-8BEC-A6932257A86B)

90 Type species. *Palaeoperilestes electronicus* sp. nov.

91 *Etymology*. Named from the Greek word παλαιός for ‘old’ and the type genus

92 *Perilestes*. Gender unknown.

93 *Diagnosis*. Wing characters: hindwing midfork (base of RP3/4) shifted distal of N;

94 IR1 strongly zigzagged and shortened, originating basal of Pt base; IR2 distinctly

95 shortened, arising on base of RP2 in forewing but one cell basal of that in hindwing;

96 IR2 two cells distal of midfork in hindwing; hindwing MP long and reaching posterior

97 wing margin slightly distal of base of IR1; CuA short and ending on posterior wing

98 margin near base of IR2; postnodal and postsubnodal crossveins somewhat aligned;

99 all intercalary veins (except IR1 and IR2) suppressed; longitudinal veins RA, IR1,

100 RP1, IR2, and RP2 strongly converging to wing apex; Pt one cell long and well

101 braced.

102

103 *Palaeoperilestes electronicus* sp. nov.

104 (urn:lsid:zoobank.org:act:15000F2A-E86C-451F-8857-45BA86B873F6)

105 Figs. 1–5

106

107 *Etymology*. Named after the Greek word electron for ‘amber’.

108 *Holotype*. NIGP163955. The distal two third of two wings, probably a forewing and a

109 hindwing because of their respective positions; deposited in the Nanjing Institute of

110 Geology and Palaeontology, Chinese Academy of Sciences, Nanjing, China.

111 *Locality and Horizon.* Hukawng Valley, Kachin Province, Myanmar; lowermost

112 Cenomanian, Upper Cretaceous.

113 *Diagnosis.* As for genus.

114 *Description.* Forewing incomplete (Figs 2A, 3A). Preserved wing length 6.81 mm,  
115 maximum width 2.36 mm, length from base of RP2 to Pt 3.8 mm, from Pt to wing  
116 apex 1.87 mm. Five postnodal crossveins and six postsubnodal crossveins present  
117 distal of N and basal of Pt, with two basal rows aligned but three distal rows not  
118 aligned. Three postnodal and postsubnodal crossveins present distal of Pt, not aligned.  
119 IR2 slightly zigzagged, arising from RP2 and slightly distal of base of RP2, lying 0.25  
120 mm distally. IR1 strongly zigzagged, three cells and 2.45 mm distal of base of RP2,  
121 and two cells basal of Pt base. RP1 with a slight angle below Pt brace. MA distally  
122 zigzagged and long. CuA ending on posterior wing margin just below base of IR2. Pt  
123 one cell long (Fig. 4), 0.6 mm long and 0.35 mm wide, well braced; star- or  
124 pyramid-like microstructures distributed on Pt surface; Pt brace in same orientation to  
125 base of Pt. All intercalary veins (except IR1 and IR2) suppressed. Longitudinal veins  
126 RA, IR1, RP1, IR2, and RP2 strongly converging to wing apex.

127 Hindwing incomplete (Figs 2B–C, 3B), resembling forewing except for  
128 following differences. Preserved wing length 9.01 mm, maximum width 2.37 mm,  
129 length from base of RP3/4 to base of RP2 2.7 mm, from base of RP2 to Pt 3.32 mm,  
130 from Pt to wing apex 1.57 mm. Seven postnodal crossveins and eight postsubnodal  
131 crossveins present before Pt, somewhat aligned. Five postnodal crossveins and five  
132 postsubnodal crossveins present distal of Pt, non-aligned. Midfork present distal of N.



133 Base of IR2 two cells and 1.99 mm distal of midfork. Base of RP2 one cell distal of  
134 base of IR2, lying 0.7 mm distally. IR1 strongly zigzagged, two cells and 1.38 mm  
135 distal of base of RP2, and two cells basal of Pt base. MA long, basally straight but  
136 strongly zigzagged distally, ending on posterior wing margin slightly basal of Pt brace.  
137 MP long, basally straight but slightly zigzagged distally, ending on posterior wing  
138 margin slightly distal of base of IR1. CuA ending on posterior wing margin slightly  
139 basal of base of IR2.

140 Three fragmentary legs preserved (Fig. 5), paired long spines present on tibia and  
141 tarsi; tibia armed with about seven or eight pairs of spines; tarsi three segmented with  
142 third tarsomere length equal to first two tarsomeres, and armed with about five or six  
143 pairs of spines; apical claws symmetrical.

#### 145 **4. Discussion**

146 The absence of the wing bases makes the attribution of this damselfly rather  
147 difficult. However, *Palaeoperilestes* has a star like microstructures distributed on the  
148 surface of the pterostigma (Fig. 4), a derived type for the superfamily Lestoidea  
149 Calvert, 1901 (Bechly, 2016), indicating the strong relationship between  
150 *Palaeoperilestes* and Lestoidea. Besides, *Palaeoperilestes* has the base of IR2 shifted  
151 several cells distal of the midfork. This unique character is only shared by a few  
152 zygopteran taxa, i.e., Chorismagrionidae Tillyard and Fraser, 1938, Perilestidae  
153 Tillyard and Fraser, 1938, Nubiolestinae Bechly, 1996 and the "megapodagrionid"  
154 genus *Arrhenocnemis* Lieftinck, 1933 (according to Bechly, 2016).

155 *Arrhenocnemis* was reassigned to Megapodagrionidae by Lieftinck, (1971);  
156 however, it was later attributed to the Paltynemididae: Calicnemiinae (see Gassmann,  
157 2005; Orr and Kalkman, 2010). *Arrhenocnemis* consists of three species, viz., *A.*  
158 *sinuatipennis* Lieftinck, 1933, *A. amphidactylis* Lieftinck, 1949, and *A. parvibullis*  
159 Orr and Kalkman, 2010, all from New Guinea. *Palaeoperilestes* resembles  
160 *Arrhenocnemis* in having the base of RP2 one cell distal of the base of IR2, and the  
161 base of IR1 being three cells distal of the base of RP2. However, any affinity of  
162 *Palaeoperilestes* with *Arrhenocnemis* can be excluded by the presence of crenulated  
163 distal wing margins, the midfork being aligned with Sn, and a non-zigzagged IR1 in  
164 *Arrhenocnemis*.

165 Chorismagrionidae comprise the relict species *Chorismagrion risi* Morton, 1914  
166 (Fig. 6), only recorded in Australia. Fraser (1957) considered the Chorismagrionidae  
167 as ‘an annectant between the families Perilestidae and Chlorolestidae’, but Dijkstra et  
168 al. (2014) placed *Chorismagrion* in the Synlestidae. The open discoidal cell in  
169 *Chorismagrion* cannot indicate affinities between these two groups, since this  
170 character has evolved several times within Zygoptera and Epiproctophora (Bechly,  
171 2016). The absence of wing base characters makes it more difficult to distinguish  
172 *Palaeoperilestes* from *Chorismagrion*. The new specimen shares with *Chorismagrion*  
173 a strongly zigzagged IR1. However, *Chorismagrion* has a midfork aligned with Sn,  
174 the base of IR2 more cells basal of the base of RP1, the base of IR1 nearer to the base  
175 of RP2 than to the base of Pt, and Pt covering two cells, obviously differing from the  
176 new specimen.

177 Nubiolestinae Bechly, 1996 comprises the relict genus *Nubiolestes* (type species:  
178 *Nubiolestes diotima* (Schmidt, 1943, Fig. 6), recorded in tropical Africa and restricted  
179 to Cameroon (Dijkstra and Vick, 2004). *Palaeoperilestes* resembles *Nubiolestes* in the  
180 base of IR2 being near to the base of RP2 and the base of IR1 being basal of Pt. The  
181 differences of *Palaeoperilestes* from *Nubiolestes* are: the midfork is distal of the  
182 subnodus instead of being aligned as in *Nubiolestes*, IR1 is quite zigzagged instead of  
183 being straight, and the base of IR2 is two cells distal of the midfork instead of six cells  
184 (Schmidt, 1943; Fraser, 1944).

185 *Palaeoperilestes* has IR2 distinctly shortened, arising on RP2 in the forewing and  
186 one cell basal of base of RP2 in the hindwing. These structures resemble the situation  
187 in the extant family Perilestidae. *Palaeoperilestes* shares with *Perilestes* and  
188 *Perissolestes* (Fig. 6) the midfork distal of the subnodus, the base of IR2 quite near to  
189 the base of RP2, but differs from them in having a quite zigzagged IR1 (straight IR1  
190 in *Perilestes* and *Perissolestes*).

191 In conclusion, *Palaeoperilestes electronicus* cannot be attributed to any known  
192 genus. However, *Palaeoperilestes* resembles *Perilestes* and *Perissolestes* more than  
193 any of the other genera discussed above. Thus we suggest a new genus provisionally  
194 attributed to the family Perilestidae (Fig. 6).

195 It should be noted that the family Austroperilestidae Petrulevičius and Nel, 2005  
196 (*Austroperilestes hunco* Petrulevičius and Nel, 2005), described from the Lower  
197 Eocene of Argentina, was considered to be related to Perilestidae (Petrulevičius and  
198 Nel, 2005). However, *A. hunco* can be easily differentiated from *P. electronicus* by the

199 presence of a 'lestine' oblique vein, IR2 being opposite the subnodus, RP3/4 being  
200 basal of the nodus, a long pterostigma and a very dense wing venation.

201

## 202 **5. Conclusions**

203 The first fossil representative of the extant family Perilestidae, *Palaeoperilestes*  
204 *electronicus* gen. et sp. nov., is described from mid-Cretaceous Burmese amber. The  
205 new discovery adds to the diversity of damselflies in the mid-Cretaceous amber.  
206 Gondwana was considered to be the ancestral area of the Lestoidea (van Tol et al.,  
207 2009), and the recent Perilestidae only occurs in the Neotropical region. The new  
208 discovery puts the appearance of perilestid damselfly back to at least the  
209 mid-Cretaceous in India.

210

## 211 **Acknowledgements**

212 This research was supported by the National Basic Research Program of China  
213 (2012CB821900), National Natural Science Foundation of China (41572010,  
214 J1210006), Youth Innovation Promotion Association of Chinese Academy of Sciences  
215 (No. 2011224), the Chinese Academy of Sciences President's International Fellowship  
216 Initiative (2011T2Z04), Open projects of Key Laboratory of Economic Stratigraphy  
217 and Palaeogeography, Chinese Academy of Science (Grant no. 2016KF05), and the  
218 HKU Seed Funding Program for Basic Research (Grant no. 201210159058). We offer  
219 our sincere gratitude to Dr. Eduardo Koutsoukos for editing, and two anonymous  
220 reviewers for the very useful comments on the earlier version of the manuscript.

221

222 **References**

223 Bechly, G., 1996. Morphologische Untersuchungen am Flügelgeäder der rezenten

224 Libellen und deren Stammgruppenvertreter (Insecta; Pterygota; Odonata), unter

225 besonderer Berücksichtigung der Phylogenetischen Systematik und des

226 Grundplanes der Odonata. *Petalura*, 2: 1–402.

227 Bechly, G., 2016. Phylogenetic systematics of Odonata. Homepage on the Internet at:

228 <https://dl.dropboxusercontent.com/u/13756162/Website/odonata/system.htm>

229 (accessed 5 May 2016).

230 Calvert, P. P., 1901. Odonata. In: *Biologia Centrali Americana: Insecta Neuroptera*. R.

231 H. Porter, London, pp. 17–72.

232 Cohen, K. M., Finney, S. C., Gibbard, P. L., Fan, J., 2013. The ICS international

233 chronostratigraphic chart. *Episodes*, 36: 199–204.

234 Dijkstra, K.-D. B., Vick, G. S., 2004. Critical species of Odonata in western Africa. In:

235 Clausnitzer, V. & Jodicke, R., (Eds.). *Guardians of the watershed. Global status*236 *of dragonflies: critical species, threat and conservation. International Journal of*237 *Odonatology*, 7: 229–238.

238 Dijkstra, K.-D. B., Boudot, J.-P., Clausnitzer, V., Kipping, J., Kisakye, J. J., Ogbogu, S.

239 S., Samraoui, B., Samways, M. J., Schütte, K., Simaika, J. P., Suhling, F.,

240 Tchiboza, S., 2011. Dragonflies and damselflies of Africa (Odonata): history,

241 diversity, distribution, and conservation. In: Darwall, W. R. T., Smith, K. G.,

242 Allen, D. J., Holland, R. A., Harrison, I. J. & Brooks, E. G. E. (Eds.). *The*

- 243 Diversity of life in African Freshwaters: under water, under threat. An analysis of  
244 the status and distribution of freshwater species throughout mainland Africa.  
245 IUCN, pp. 128–177.
- 246 Dijkstra, K.-D. B., Kalkman, V. J., Dow, R. A., Stokvis, F. R., van Tol, J., 2014.  
247 Redefining the damselfly families: a comprehensive molecular phylogeny of  
248 Zygoptera (Odonata). *Systematic Entomology*, 39: 68–96.
- 249 Fabricius, J. C., 1793. *Entomologia systematica emendata et aucta, secundum classes,*  
250 *ordines, genera, species, adjectis synonymis, locis, observationibus,*  
251 *descriptionibus.* C.G. Proft, Hafniae (= Copenhagen), 3: 1–487 and 1–349.
- 252 Fraser, F. C., 1944. The significance of vestigial oblique veins in the evolution of  
253 intercalated veins in the odonate wing, with the description of a new genus.  
254 *Proceedings of the Royal Entomological Society, London, (B), 13 (5–6): 58-67.*
- 255 Fraser, F. C. 1957. A reclassification of the order Odonata, based on some new  
256 interpretations of the venation of the dragonfly wing. *Royal Zoological Society*  
257 *of New South Wales, Sydney, 12: 1–133.*
- 258 Fraser, F. C., 1960. *The Odonata of Australia, Tasmania, and New Zealand. A*  
259 *handbook of the dragonflies of Australasia with keys for the identification of all*  
260 *species, Royal Zoological Society of New South Wales, Sydney, 1–61.*
- 261 Fabricius, J. C., 1793. *Entomologia systematica emendata et aucta, secundum classes,*  
262 *ordines, genera, species, adjectis synonymis, locis, observationibus,*  
263 *descriptionibus.* C.G. Proft, Hafniae (= Copenhagen), 3: 1–487 and 1–349.
- 264 Gassmann, D., 2005. The phylogeny of Southeast Asian and Indo-Pacific

- 265 Calicnemiinae (Odonata:Platycnemididae). Bonner Zoologische Beiträge, 53  
266 (2004): 37–80.
- 267 Haber, W., Wagner, D., 2014. Perilestidae: shortwings. Dragonflies and damselflies of  
268 Ecuador (Odonata). Available on the Internet at:  
269 [http://bdei2.cs.umb.edu/~whaber/Odonata\\_of\\_Ecuador/index.html](http://bdei2.cs.umb.edu/~whaber/Odonata_of_Ecuador/index.html) (accessed 16  
270 March 2016).
- 271 Kania, I., Wang, B., Szwed, J., 2015. *Dicranoptycha* Osten Sacken, 1860 (Diptera,  
272 Limoniidae) from the earliest Upper Cretaceous Burmese amber. Cretaceous  
273 Research, 52: 522–530.
- 274 Kennedy, C. H., 1920. The phylogeny of the zygopterous dragonflies as based on the  
275 evidence of the penes. Ohio Journal of Science, 21: 19–29.
- 276 Kennedy, C. H., 1941. Perilestinae in Ecuador and Peru, revisional notes and  
277 descriptions (Lestidae: Odonata). Annals of the Entomological Society of  
278 America, 34 (3): 658–688.
- 279 Lieftinck, M. A., 1933. The dragonflies of New Guinea and neighbouring islands. Part  
280 II. Descriptions of a new genus and species of Platycneminae (Agrionidae) and  
281 of new Libellulidae. Nova Guinea, 17: 1–66.
- 282 Lieftinck, M. A., 1949. The dragonflies (Odonata) of New Guinea and neighbouring  
283 islands. Part VII. Results of the third Archbold expedition 1938–1939 and of the  
284 Le Roux expedition 1939 to the Netherlands, New Guinea (II. Zygoptera). Nova  
285 Guinea (N.S.), 5: 1–271.
- 286 Lieftinck, M. A., 1971. A catalogue of type specimens of Odonata preserved in the

- 287 Netherlands, with a supplementary list of the Odonata types described by Dutch  
288 scientists deposited in foreign institutional collections. Tijdschrift voor  
289 Entomologie, 114: 65–139.
- 290 Machado, A. B. M., 2015. *Perilestes eustaquioi* sp. nov. and new distributional  
291 records of Perilestidae (Odonata) in Brazil. Zoologia, 32(5): 428–430.
- 292 Morton, K. J., 1914. A remarkable new genus and new species of Odonata, of the  
293 legion Podagrion, Selys, from North Queensland. Transactions of the  
294 Entomological Society of London, 62: 169–172.
- 295 Nel, A., Martinez-Delclòs, X., Paicheler, J. C., Henrotay, M., 1993. Les  
296 ‘Anisozygoptera’ fossiles. Phylogénie et classification (Odonata). Martinia Hors  
297 Série, 3: 1–311.
- 298 Neiss, U. G., Neusa, H., 2010. The larva of *Perilestes attenuatus* Selys, 1886  
299 (Odonata: Perilestidae) from Amazonas, Brazil. Zootaxa, 2614: 53–58.
- 300 Orr, A. G., Kalkman, V. J., 2010. *Arrhenocnemis parvibullis* sp. nov. (Odonata:  
301 Platycnemididae), a new calicnemiine damselfly from Papua New Guinea, with a  
302 description of the female of *A. amphidactylis* Lieftinck, 1949. Australian  
303 Entomologist, 37 (4): 137–146.
- 304 Petrulevičius, J. F., Nel, A., 2005. Austroperilestidae, a new family of damselflies  
305 from the earliest Eocene of Argentina (Insecta: Odonata). Phylogenetic  
306 relationships within Odonata. Journal of Paleontology, 79 (4): 658–662.
- 307 Riek, E. F., 1976. A new collection of insects from the Upper Triassic of South Africa.  
308 Annals of the Natal Museum, 22: 791–820.



- 309 Riek, E. F., Kukalová-Peck, J., 1984. A new interpretation of dragonfly wing venation  
310 based upon Early Carboniferous fossils from Argentina (Insecta: Odonatoidea)  
311 and basic character states in pterygote wings. *Canadian Journal of Zoology*, 62:  
312 1150–1166.
- 313 Schmidt, E., 1942. Bemerkungen über Lestiden. 2. Eine neue Gattung und Art aus  
314 Kamerun (*Eolestes diotima*). *Mitteilungen der Deutsche Entomologische*  
315 *Gesellschaft*, 11: 102–111.
- 316 Schorr, M., Paulson, D., 2015. World Odonata list. Available on the Internet at:  
317 <http://www.pugetsound.edu/> (accessed 16 March 2016).
- 318 Selys-longchamps, E. de, 1854. Monographie des Calopterygines. *Mémoires de la*  
319 *Société royale des Sciences de Liège* 9, xi + 291 pp.
- 320 Selys-longchamps, E. de, 1862. Synopsis des agrionines. Troisième légion: Podagrion.  
321 *Bulletins de l'Académie Royale des Sciences, des Lettres et des Beaux-Arts de*  
322 *Belgique* (2me série), 14: 5–44.
- 323 Shi, G., Grimaldi, D. A., Harlow, G. E., Wang, J., Wang, J., Wang, M., Lei, W., Li, Q.,  
324 Li, X., 2012. Age constraint on Burmese amber based on U-Pb dating of zircons.  
325 *Cretaceous Research*, 37: 155–163.
- 326 Tillyard, R. J., Fraser, F. C., 1938–1940. A reclassification of the order Odonata.  
327 *Australian Zoologist* 9, 125–169, 195–221, and 359–390.
- 328 van Tol, J., Reijnen, B. T., Thomassen, H. A., 2009. Phylogeny and biogeography of  
329 the Platystictidae (Odonata). In: van Tol, J., *Phylogeny and biogeography of the*  
330 *Platystictidae (Odonata)*. PhD Thesis, Leiden University, Leiden, pp. 3–70.

331 Williamson, E. B., Williamson, J. H., 1924. The genus *Perilestes* (Odonata).  
332 Miscellaneous Publications of the Museum of Zoology University of Michigan,  
333 14: 1–39.

334

335 **Figure captions**

336 **Fig 1.** *Palaeoperilestes electronicus* gen. et sp. nov., holotype, NIGP163955,  
337 photomicrograph of specimen.

338 **Fig 2.** *Palaeoperilestes electronicus* gen. et sp. nov., holotype, NIGP163955. A,  
339 photomicrograph of forewing; B, photomicrograph of mid hindwing; C,  
340 photomicrograph of distal part of hindwing.

341 **Fig 3.** *Palaeoperilestes electronicus* gen. et sp. nov., holotype, NIGP163955, line  
342 drawing showing wing venation.


343 **Fig 4.** *Palaeoperilestes electronicus* gen. et sp. nov., holotype, NIGP163955,  
344 photomicrograph showing details of Pt.

345 **Fig 5.** *Palaeoperilestes electronicus* gen. et sp. nov., holotype, NIGP163955,  
346 photomicrograph showing leg details.


347 **Fig 6.** Putative position of *Palaeoperilestes* gen. nov. in phylogenetic tree of  
348 Zygoptera. All line drawings are based on forewings except for *Palaeoperilestes* gen.  
349 nov. from a more complete hindwing (Cladogram based on Dijkstra et al. 2014, line  
350 drawing of *Chorismagrion risi* Morton, 1914 after Fraser, 1960; line drawing of  
351 *Nubiolestes diotima* Schmidt, 1943 after Fraser, 1944; line drawings of *Perilestes*  
352 *gracillimus* Kennedy, 1941 and *Perissolestes remotus* Williamson & Williamson,


353 1924 after Kennedy, 1941).


ACCEPTED MANUSCRIPT



ACCEPTED







ACCEPTED MANUSCRIPT

