



Title	New DSH planetary nebulae and candidates from optical and infrared surveys
Author(s)	Kronberger, M; Parker, QA; Jacoby, GH; Acker, A; Alves, F; Bojicic, I; Eigenthaler, P; Frew, DJ; Harmer, D; Patchick, D; Reid, W; Schedler, J
Citation	The 11th Pacific Rim Conference on Stellar Astrophysics, Hong Kong, 14-17 December 2015. In Journal of Physics: Conference Series, 2016, v. 728 n. 7, p. 072012:1- 072012:2
Issued Date	2016
URL	http://hdl.handle.net/10722/237687
Rights	Journal of Physics: Conference Series (Online Edition). Copyright © Institute of Physics Publishing.; This is an author-created, un-copied version of an article published in [insert name of journal]. IOP Publishing Ltd is not responsible for any errors or omissions in this version of the manuscript or any version derived from it. The Version of Record is available online at http://dx.doi.org/[insert DOI].; This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

New DSH planetary nebulae and candidates from optical and infrared surveys

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 J. Phys.: Conf. Ser. 728 072012

(<http://iopscience.iop.org/1742-6596/728/7/072012>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 147.8.230.84

This content was downloaded on 19/01/2017 at 02:43

Please note that [terms and conditions apply](#).

You may also be interested in:

[On the applicability of Sato's equation to capacitative radiofrequency sheaths](#)

J Balakrishnan and G R Nagabhushana

[The invariance of DSHs and DVHs under bladder filling changes](#)

Mischa S Hoogeman, Stephanie T H Peeters, Josien de Bois et al.

[Simulation of ion transport characteristics in a collisionless rf sheath](#)

Zu-li Liu, Xing-bin Jing and Kai-lun Yao

[Experimental investigation on linewidth characteristics of a single-frequency phosphate fiber laser at 1.0 \$\mu\text{m}\$](#)

Can Li, Shanhui Xu, Yu Xiao et al.

[Injection laser with a diffraction grating in its resonator](#)

D Akkerman, Alexandr P Bogatov, P G Eliseev et al.

[The Dirac equation with a Coulomb potential in D dimensions](#)

Shi-Hai Dong

[Gas temperature measurement in thermal radiating environments using a suction thermocouple apparatus](#)

A Z'Graggen, H Friess and A Steinfeld

[A Preliminary Survey of the Paramagnetic Resonance Phenomena observed in Rare Earth Ethyl Sulphates](#)

R J Elliott and K W H Stevens

[Magnetic symmetry in quasicrystals](#)

K Rama Mohana Rao and P Hemagiri Rao

New DSH planetary nebulae and candidates from optical and infrared surveys

Matthias Kronberger¹, Quentin A Parker², George H Jacoby^{1,3},
Agnes Acker⁴, Filipe Alves¹, Ivan Bojicic², Paul Eigenthaler⁵, David
J Frew², Dianne Harmer⁶, Dana Patchick¹, Warren Reid^{7,8} and
Johannes Schedler⁹

¹ Deep Sky Hunters collaboration

² Department of Physics, The University of Hong Kong, Hong Kong

³ Giant Magellan Telescope/Carnegie Observatories, Pasadena, CA 91101, USA

⁴ Observatoire astronomique de Strasbourg, 11 rue de l'Université, F-67000 Strasbourg, France

⁵ Instituto de Astrofísica, Pontificia Universidad Católica de Chile, Santiago, Chile

⁶ National Optical Astronomy Observatory, 950 N. Cherry Avenue, Tucson, AZ 85719, USA

⁷ Department of Physics and Astronomy, Macquarie University, Sydney, NSW 2109, Australia

⁸ University of Western Sydney, Locked Bag 1797, Penrith South DC, NSW 1797, Australia

⁹ CHART32 Team

E-mail: ¹ matthias.kronberger@gmx.at

Abstract. To date, the planetary nebula (PN) survey of the Deep Sky Hunters collaboration has led to the detection of more than 250 previously unknown candidate planetary nebulae (PNe). About 60% of them were found during the past two years and are expected to be true, likely or possible PNe because careful vetting has already thrown out more doubtful objects. The majority of the new PN candidates are located within the boundaries of the SHS and IPHAS H α surveys and were discovered by combining MIR data from the WideField Infrared Survey Explorer (WISE) with optical data from the IPHAS, SHS and DSS surveys, and UV data from the Galaxy Evolution Explorer (GALEX).

Since the discovery of M 27 by the French astronomer Charles Messier about 250 years ago, ≈ 3500 more Galactic PNe have been discovered [1]. However, this number is still short of the total number of Galactic PNe estimated from local volume counts ($N_{tot=11,000}$ [2]). To a great extent, this difference can be attributed to the circumstance that, until recently, PN surveys were conducted mainly at optical wavelengths and over a relatively limited latitude range of the Galactic plane. Consequently, a considerable gain in the number of Galactic PNe can be expected from surveys in the MIR range, allowing for the discovery of highly obscured dust-rich and high excitation PNe that are invisible at optical wavelengths.

The Deep Sky Hunters [3] (DSH) have been scanning the Milky Way for new clusters and nebulae for more than decade now. Until fairly recently, our survey for PN candidates was restricted to medium Galactic latitudes ($5^\circ \leq |b| \leq 20^\circ$), and based on broadband Digitized Sky Survey (DSS) images [2,4,5] as well as on [S II]/H α /[O III] narrowband images of the Milky Way taken with modestly-sized amateur telescopes equipped with modern CCD imagers [6]. Since then, the focus has shifted to fields closer to the Galactic Equator, and from optical to MIR and UV wavelengths, thanks to the availability of high resolution imagery from the WISE and GALEX missions. This has allowed us to detect 162 true, likely and possible PNe during the past






Figure 1. Aitoff-Hammer projection of the Galactic Plane, showing the distribution of all True, Likely and Possible PNe of the DSH sample.

two years, increasing the census of DSH PNe and candidates by more than 150%. It is worth noting that many of the new PN candidates, although originally discovered in the MIR, are clearly visible on SHS [7] and IPHAS [8] survey images but were overlooked thus far, most likely because of their faintness, because of insufficient contrast between $H\alpha$ and their comparative broadband imagery, or because of their proximity to bright sources in the field. Object sizes of the new PN candidates found at optical wavelengths range from essentially unresolved to several arcmin. All good quality DSH candidates and all those confirmed spectroscopically will be ingested into the new HASH database [9].

Figure 1 shows the distribution of all True, Likely and Possible PNe of the DSH sample in an Aitoff-Hammer projection of the Galactic Plane. Objects listed as True, Likely and Possible PNe in the HASH database are shown for comparison. Gaps in the DSH candidate distribution correspond to unsurveyed Milky Way regions. We plan to obtain confirmatory spectroscopy of the Possible and Likely PNe during the next years, and $H\alpha + [O III]$ images of another ≈ 130 objects that we consider to be PN candidates based alone on their MIR characteristics.

References

- [1] Parker Q A, Bojicic I, Frew D J, Acker A, Ochsenbein F and MASP Database Team 2015 *American Astronomy Society Meeting* **225** 108.06
- [2] Jacoby G H *et al* 2010 *Publ. Astron. Soc. Australia* **27** 156
- [3] Kronberger M *et al* 2006 *Astron. Astrophys.* **447** 921
- [4] Kronberger M *et al* 2012 *Proc. Int. Astron. Union (S283)* **7** 414
- [5] Kronberger M *et al* 2014 *Asymmetrical Planetary Nebulae VI* 48
- [6] Acker A, Boffin H M J, Outters N, Miszalski B, Sabin L, Le Dû P and Alves F 2012 *Revista Mexicana de Astronomia y Astrofisica* **38** 223
- [7] Parker Q A *et al* 2005 *Mon. Not. R. Astron. Soc* **362** 689
- [8] Drew J E *et al* 2005 *Mon. Not. R. Astron. Soc* **362** 753
- [9] Parker Q A *et al these proceedings*