

Title	Entropy for gravitational Chern-Simons terms by squashed cone method
Author(s)	Guo, Wu Zhong; Miao, Rong Xin
Citation	Journal of High Energy Physics (2016), 2016
Issue Date	2016-04
URL	http://hdl.handle.net/2433/218569
Right	© The Author(s) 2016; JHEP is an open-access journal funded by SCOAP3 and licensed under CC BY 4.0
Type	Journal Article
Textversion	publisher

Entropy for gravitational Chern-Simons terms by squashed cone method

Wu-Zhong Guo^{a,b} and Rong-Xin Miao^c

^aState Key Laboratory of Theoretical Physics,
 Institute of Theoretical Physics, Chinese Academy of Science,
 Zhong Guan Cun East Street 55 #, P.O. Box 2735, Beijing 100190, P.R. China

^bYukawa Institute for Theoretical Physics (YITP), Kyoto University,
 Kitashirakawa Oiwakecho, Sakyo-ku, Kyoto 606-8502, Japan

^cMax Planck Institute for Gravitational Physics (Albert Einstein Institute),
 Am Mühlenberg 1, 14476 Golm, Germany

E-mail: wuzhong@itp.ac.cn, rong-xin.miao@aei.mpg.de

ABSTRACT: In this paper we investigate the entropy of gravitational Chern-Simons terms for the horizon with non-vanishing extrinsic curvatures, or the holographic entanglement entropy for arbitrary entangling surface. In 3D there is no anomaly of entropy. But the original squashed cone method can not be used directly to get the correct result. For higher dimensions the anomaly of entropy would appear, still, we can not use the squashed cone method directly. That is because the Chern-Simons action is not gauge invariant. To get a reasonable result we suggest two methods. One is by adding a boundary term to recover the gauge invariance. This boundary term can be derived from the variation of the Chern-Simons action. The other one is by using the Chern-Simons relation $d\Omega_{4n-1} = \text{tr}(\mathbf{R}^{2n})$. We notice that the entropy of $\text{tr}(\mathbf{R}^{2n})$ is a total derivative locally, i.e. $S = ds_{CS}$. We propose to identify s_{CS} with the entropy of gravitational Chern-Simons terms Ω_{4n-1} . In the first method we could get the correct result for Wald entropy in arbitrary dimension. In the second approach, in addition to Wald entropy, we can also obtain the anomaly of entropy with non-zero extrinsic curvatures. Our results imply that the entropy of a topological invariant, such as the Pontryagin term $\text{tr}(\mathbf{R}^{2n})$ and the Euler density, is a topological invariant on the entangling surface.

KEYWORDS: AdS-CFT Correspondence, Classical Theories of Gravity

ARXIV EPRINT: [1506.08397](https://arxiv.org/abs/1506.08397)

Contents

1	Introduction	1
2	Review of generalized gravitational entropy	3
3	Gravitational Chern-Simons term	4
4	Entropy of gravitational Chern-Simons terms in 3D	5
4.1	Result by using the squashed cone method directly	6
4.2	A solution	6
4.3	The surface	8
5	Entropy of gravitational Chern-Simons terms in 7D	9
5.1	Approach to 7D case by a topological method	9
6	Wald entropy in arbitrary dimension	12
6.1	The topological method	12
6.2	Another approach	13
7	Conclusion and discussion	15
A	Useful components of Γ and R	15
B	Non-covariant part of Ω	16
C	Details of the calculation in section 5	17
D	Comparison with [38]	17

1 Introduction

The entropy is often used as a quality to reflect the degree of freedoms of a system. In the gravitational field black hole entropy, i.e., Bekenstein-Hawking entropy [1, 2], is related to geometry of the spacetime, and performs as a thermal quality. On the other hand in the gauge field theory with gravity dual, the entanglement entropy for a subsystem could also have a geometry description in the gravity side, which is known as the Ryu-Takayanagi formula [3]. In general the geometry description of the entropy is closely connected with the detail of the theory. Wald formula [4] provides the connection between the action and entropy for general covariant gravitational theory.

The recent idea concerning about the generalized gravitational entropy [5] gives a strong evidence for the Ryu-Takayanagi formula on general entangling surfaces. Generalization to theory other than Einstein gravity is not so straightforward. Instead, one has to deal with the squashed conical singularity carefully and take into account the possible contribution from the extrinsic curvatures. The first breakthrough was made by [6], in which the authors develop a nice approach of regularization to study the entropy of curvature-squared gravity. However, recently it is found that this method produces wrong entropy for general higher derivative gravity [7]. The correct approach to derive the entropy of general higher derivative gravity was developed by [8], see also [9]. In this paper we use the method of [8] and denote it by the squashed cone method. For other interesting developments of the holographic entanglement entropy, please refer to [10–15].

The squashed cone method by Lewkowycz-Maldacena [5] and Dong [8] works well for covariant gravitational theories. However, it may break down or need to be modified for non-covariant gravitational theories. There are two kind of non-covariant gravitational theories. The first one is that neither the action nor the equations of motion are gauge invariant. The black hole thermodynamics is not well-defined for this kind of gravitational theories [16–18]. The second one is that the action is gauge invariant up to some boundary terms. Theory with gravitational Chern-Simons term is one of this kind of non-covariant theories. The Chern-Simons(CS) term as a possible correction to Einstein gravity is motivated by the low-energy effective action from superstring theories. In 3D the modification of the black hole entropy by CS term is studied in many literatures, see [19–22]. In higher dimensions the answer is also found in [26, 27] by generalizing the covariant phase formalism. There are also some studies on the contribution to thermodynamics and transport in hydrodynamics from the gravitational anomalies [28, 29], which is related to the gravitational CS term by AdS/CFT [30–32]. In this paper we would like to study the problem for generalized gravitational entropy. Specially, we would use the regularization process developed in [8]. The method works well for covariant theory, but for CS theory, we find that the method should be modified significantly to get a consistent result. The modification is related with the local gauge transformation of theory with gravitational Chern-Simons term, for this transformation would produce a total derivative terms. Actually the regularized process would ignore the possible effect caused by the gauge transformation. The modification is based on the consideration to fix the gauge freedom. In the following we refer to entropy either to generalized gravity entropy or to holographic entanglement entropy for they are the same thing in a sense.

The paper is organized as follows. In the next section we briefly introduce the regularization process developed in [8]. In section 3 we briefly review the gravitational Chern-Simons terms in arbitrary dimension. We will calculate the entropy in 3D in this section 4. In section 5, we propose an approach to derive the entropy of gravitational Chern-Simons terms. We work out the entropy exactly in 7D space-time. In section 6, we use this method to get the Wald entropy in arbitrary dimension. We will also discuss other approach to get the correct Wald entropy. We will conclude and discuss some related problems in section 7. Some useful formula and detail calculation can be found in appendix.

2 Review of generalized gravitational entropy

The generalized gravitational entropy is based on the “replica trick” in Euclidean spacetime. The un-normalized density matrix of the gravitational field is related to the partition function $Z(1)$ by $\text{tr} \rho = Z(1)$. We could normalize the density matrix and define $\hat{\rho} = \rho / \text{tr} \rho$. The n -th replica spacetime B_n would admit $\text{tr} \hat{\rho}^n = Z(n) / Z(1)^n$, where $Z(n)$ is the partition function of B_n . In the semi-classical limit we would have $\text{tr} \hat{\rho}^n \simeq \exp(-I(n) + nI(1))$, where $I(n)$ is the classical on-shell action of B_n , $I(1)$ is the action of the original spacetime. One could consider the orbifold $\hat{B}_n \equiv B_n / Z_n$. This leads to $I_n = n\hat{I}(n)$, where $\hat{I}(n)$ means the action with the solution \hat{B}_n without counting the contribution from the conical defect in \hat{B}_n . The entropy in $O(G^{-1})$ can be expressed as

$$S = \partial_\epsilon \hat{I}(\epsilon), \tag{2.1}$$

where we denote $n = 1 + \epsilon$. Now one fills the singular cone, the calculation becomes

$$S = -\partial_\epsilon I(\epsilon), \tag{2.2}$$

where $I(\epsilon)$ is the action of the regularized squashed cone. This equation is the starting point to calculate the entropy. We refer the readers to [5] and [8] for more argument and explanation to the generalized gravitational entropy and holographic entanglement entropy.

But in the theory that is not-covariant this statement can not be used directly. Such as the gravity with Chern-Simons term we will discuss below, the local gauge transformation or the non-covariant part of the diffeomorphism will lead to a boundary term, which also contributes to the $O(\epsilon)$. For 3D gravity theory with Chern-Simons terms this problem is first noticed in papers [19] and [35], in which the authors obtain additional contributions to $I(\epsilon)$ by making integration by parts. In this paper we provide alternative methods to solve this problem. Our main discussion below is about how to eliminate the ambiguity in the non-covariant theory when using the squashed cone method by Lewkowycz-Maldacena [5] and Dong [8]. Our approach is consistent with the results of [19, 35, 38].

Regardless of the difference that we mention above, one have to find a way to regularize the squashed cone. We would follow the regularization process in [8]. According to [8], the metric of regularized cone is

$$ds^2 = e^{2A}[dzd\bar{z} + e^{2A}T(\bar{z}dz - zd\bar{z})^2] + (g_{ij} + 2K_{aij}x^a + Q_{abij}x^ax^b)dy^i dy^j + 2ie^{2A}(U_i + V_{ai}x^a)(\bar{z}dz - zd\bar{z})dy^i + \dots \tag{2.3}$$

where $T, g_{ij}, K_{aij}, Q_{abij}, U_i, V_{ai}$ are independent of z and \bar{z} , with the exception that $Q_{z\bar{z}ij} = Q_{\bar{z}zij}$ contains a factor e^{2A} . The warp factor A is regularized by a thickness parameter a as $A = -\frac{\epsilon}{2} \log(z\bar{z} + a^2)$. The result is independent of the choice of regularization.

The contribution from the Wald entropy is related with the fact

$$\int dzd\bar{z} e^{-\beta A} \partial_z \partial_{\bar{z}} A = -\pi\epsilon. \tag{2.4}$$

The key observation of [8] is that

$$\int \rho d\rho \partial_z A \partial_{\bar{z}} A e^{-\beta A} = -\frac{\epsilon}{4\beta}, \tag{2.5}$$

where $z = \rho e^{i\tau}$. The would-be logarithmic divergence gains a $\frac{1}{\epsilon}$ enhancement:

$$\int \rho d\rho \frac{1}{\rho^2} e^{\beta\epsilon} \sim \frac{1}{\beta\epsilon}. \tag{2.6}$$

This will give the anomaly contribution to the entropy. One is suggested to refer the recent paper [10] in which we discuss more possible terms that may contribute to entropy. For our purpose in this paper (2.4)(2.5) are enough.

3 Gravitational Chern-Simons term

In the this section we would like to introduce some definitions and properties of CS terms. The gravitational CS terms can be constructed in two different ways, one is by the one-form of Christoffel symbol Γ , the other one is the spin connection ω .

By using ω the $(2n+1)$ -dimensional gravitational CS terms Ω_{2n+1} are formally defined as

$$d\Omega_{2n+1}(\omega) = Tr \mathbf{R}^{n+1}, \tag{3.1}$$

where $\mathbf{R} = d\omega + \omega\omega$ is the two-form curvature, and we suppress the wedge between the forms. Ω_{2n+1} can be expressed as

$$\Omega_{2n+1} = (n+1) \int_0^1 t^n str(\omega \mathbf{R}_t^n), \tag{3.2}$$

where $\mathbf{R}_t^n \equiv \mathbf{R} + (t-1)\omega^2$, and ‘‘str’’ is defined by

$$str(A_1, A_2, \dots, A_n) \equiv \frac{1}{n!} \sum_{\pi} Tr(A_{\pi(1)} A_{\pi(2)} \dots A_{\pi(n)}), \tag{3.3}$$

π denotes the permutations of $\{1, 2, \dots, n\}$. The CS action is

$$I_{CS} = \frac{\lambda}{32\pi G} \int_{M_{2n+1}} \frac{2^n}{n+1} \Omega_{2n+1}, \tag{3.4}$$

λ is the coupling constant. The full action is

$$I = \frac{1}{16\pi G} \int d^{2n+1}x \sqrt{-g} \left(R + \frac{n(2n+1)}{l^2} \right) + I_{CS}. \tag{3.5}$$

The spin connection ω can be construct by vielbeins $E = \{e^{a\nu}\}$, which is defined by $G_{\mu\nu} = e_{\nu}^{a\kappa} e_{\mu}^{a\sigma} \delta_{a\kappa a\sigma}$. As an example we could choose the vielbeins of (2.3) up to $O(\rho)$ as follows.

$$\begin{aligned} e_{a_1} &= e_{\mu a_1} dx^{\mu} = \frac{e^A}{2} (dz + d\bar{z}) + e^A (\bar{z} - z) U_i dy^i, \\ e_{a_2} &= -i \frac{e^A}{2} (dz - d\bar{z}) - e^A (\bar{z} + z) U_i dy^i, \\ e_{a_i} &= \bar{e}_{j a_i} dy^j + x^a K_{a j a_i} dy^j, \end{aligned} \tag{3.6}$$

where $\bar{e}_{ja_i}\bar{e}_{ka_i} = g_{jk}$ and $K_{aja_i} = K_{aji}e_{a_i}^i$. Here a_1, a_2 denotes the local Lorentz indices with respect to z, \bar{z} . One can check that the above vielbeins can yield the correct metric in order $O(\rho)$:

$$e_{\mu a_1}e_{\nu a_1} + e_{\mu a_2}e_{\nu a_2} + e_{\mu a_i}e_{\nu a_i} = G_{\mu\nu} + O(\rho^2). \quad (3.7)$$

The choose of vielbeins are not unique, different ones are related by performing local Lorentz transformations,

$$e'_{a_\mu} = \Lambda(x)_{a_\mu}^{b_\nu} e_{b_\nu}. \quad (3.8)$$

We can calculate the spin connection ω by the following formula,

$$\begin{aligned} \omega_{\mu, a_\nu a_\kappa} &= \frac{1}{2}(C_{a_\nu \sigma \mu} e_{a_\kappa}^\sigma + C_{a_\kappa \mu \sigma} e_{a_\nu}^\sigma - C_{a_\sigma \alpha \beta} e_{a_\nu}^\alpha e_{a_\kappa}^\beta e_\mu^{a_\sigma}) \\ C_{\mu\nu}^{a\sigma} &= \partial_\mu e_\nu^{a\sigma} - \partial_\nu e_\mu^{a\sigma}. \end{aligned} \quad (3.9)$$

The spin connection ω can also be related with the one-form Christoffel symbols Γ by the vielbeins,

$$\omega_\mu^x{}_y = \Gamma_{\mu y}^x - e^{ax} \partial_\mu e_{ya}, \quad (3.10)$$

where we have mapped all the index into spacetime. We list some useful components of the Christoffel symbols and spin connections in appendix A.

By varying the action (3.2) new terms would contribute to the equation of motion. The result is derived in [26, 33],

$$R_{\mu\nu} - \frac{1}{2}G_{\mu\nu}R - \frac{n(2n+1)}{l^2}G_{\mu\nu} + \lambda C_{\mu\nu} = 0 \quad (3.11)$$

where

$$C^{\mu\nu} = \nabla_\alpha S^{\mu\nu\alpha} \quad (3.12)$$

with

$$S^{\mu\nu\alpha} = -\frac{1}{2}\epsilon^{\lambda_1\lambda_2\dots\lambda_{2n}\mu} R^\nu{}_{\kappa_1\lambda_1\lambda_2} R^{\kappa_1}{}_{\kappa_2\lambda_3\lambda_4} \dots R^{\kappa_{2n-2}\alpha}{}_{\lambda_{2n-1}\lambda_{2n}}. \quad (3.13)$$

The CS term expressed by Γ is similar to ω , with ω replaced by Γ in (3.2).

4 Entropy of gravitational Chern-Simons terms in 3D

3D theory with CS term is also known as topologically massive gravity [34]. The black hole entropy in this theory has been discussed in many authors, see e.g. [19–25]. Non-trivial correction appears for rotating BTZ black. We would like to use the squashed cone method to study the correction of CS term to black hole entropy (also the HEE formula) in 3D spacetime.

For the Euclidean theory, the action of 3D CS term is

$$I_{CS} = \frac{-i\lambda}{64\pi G} \int d^3x \sqrt{G} \epsilon^{\mu\nu\sigma} \left(\Gamma_{\beta\mu}^\alpha R_{\alpha\nu\sigma}^\beta - \frac{2}{3} \Gamma_{\beta\mu}^\alpha \Gamma_{\gamma\nu}^\beta \Gamma_{\alpha\sigma}^\gamma \right), \quad (4.1)$$

where we have integrated t , $-i$ appears because of the Wick rotation, see [27].

4.1 Result by using the squashed cone method directly

Firstly we use the squashed cone method by Lewkowycz-Maldacena [5] and Dong [8] directly. To get the contribution to HEE from the CS action, we need work in metric (2.3), and find the $O(\epsilon)$. We calculate the components of Γ and \mathbf{R} in the appendix A. It's easy to see that only the Wald entropy appears, and the final result of the entropy is

$$S = \frac{i\lambda}{16G} \int_{\Sigma} dy (\Gamma_{yz}^z \epsilon^z{}_z + \Gamma_{y\bar{z}}^{\bar{z}} \epsilon^{\bar{z}}{}_{\bar{z}}) = \frac{i\lambda}{16G} \int_{\Sigma} \Gamma_{\sigma\mu}^{\nu} \epsilon^{\mu}{}_{\nu} dx^{\sigma} \tag{4.2}$$

where Σ is codimension-2 surface in the bulk.

It's also interesting to check the result by using the spin connection ω . The action would be

$$I_{CS} = \frac{-i\lambda}{64\pi G} \int d^3x \sqrt{G} \epsilon^{\mu\nu\sigma} \left(\omega_{\mu}{}^{\alpha}{}_{\beta} R^{\beta}{}_{\alpha\nu\sigma} - \frac{2}{3} \omega_{\mu}{}^{\alpha}{}_{\beta} \omega_{\nu}{}^{\beta}{}_{\gamma} \omega_{\sigma}{}^{\gamma}{}_{\alpha} \right). \tag{4.3}$$

We also calculate all the components of ω in appendix A. Besides the Wald entropy there are also contributions from the second term in (4.3). But the contribution finally vanishes. The result is also

$$S = \frac{i\lambda}{16G} \int_{\Sigma} \Gamma_{\sigma\mu}^{\nu} \epsilon^{\mu}{}_{\nu} dx^{\sigma}, \tag{4.4}$$

where we use $\omega_{y^z}{}^z = \Gamma_{yz}^z + O(z)$. It is natural because Γ is related with ω by local gauge transformation, and the result is gauge-invariant.

But (4.2), (4.4) is not consistent with the result given in literature [19–22]. The inconsistency warn us to be careful when dealing with the non-covariant theory. Just like the Wald method [4] to calculate the entropy of CS term, some modification is expected to get the correct result [25]. We would give a solution to this problem in the next subsection for the squashed cone method.

4.2 A solution

The vielbeins (3.6) still have a gauge freedom (3.8). An arbitrary local Lorentz transformation would produce an additional total derivative term for the action. This term contributes to entropy if we use the squashed cone method before integrating out the total derivation term. The freedom should be eliminated if we want a reasonable result.

We denote the action of CS term after regularization as $I(\epsilon)$, the entropy $S = -\partial_{\epsilon} I(\epsilon)$. As we can see from metric (2.3) the vielbeins E_{μ}^a and the spin connection $\omega_{\mu}{}^a{}_b$ would also depend on ϵ . Under an infinitesimal local Lorentz transformation parameterized by $\theta^a{}_b$,

$$\begin{aligned} \delta_{\theta} e^b &= -\theta^b{}_a e^a, \\ \delta_{\theta} \omega^a{}_b &= d\theta^a{}_b + [\omega, \theta]^a{}_b. \end{aligned} \tag{4.5}$$

The 3D CS action

$$I_{CS}^{(3)} = \frac{-i\lambda}{32\pi G} \int_{M_3} Tr \left(\omega d\omega + \frac{2}{3} \omega^3 \right) \tag{4.6}$$

would have a variation that is total derivative term,

$$\delta_{\theta} I_{CS}^{(3)} = \frac{-i\lambda}{32\pi G} \int_{M_3} dTr(\theta d\omega(\epsilon)). \tag{4.7}$$

When we work in the metric (2.3), the above term (4.7) contains ϵ , which contributes to entropy. This suggests new terms should be added to the action to eliminate the ambiguity. We find the following one satisfies the requirement,

$$\Delta I_{CS}^{(3)} = \frac{i\lambda}{32\pi G} \int_{M_3} dTr(\omega(0)\omega(\epsilon)). \tag{4.8}$$

The total derivative term does not modify the equation of motion. In the limit $\epsilon \rightarrow 0$ it vanishes, we get the original action. Under the infinitesimal local Lorentz transformation,

$$\begin{aligned} \delta_\theta \Delta I_{CS}^{(3)} &= \frac{i\lambda}{32\pi G} \int_{M_3} dTr(\delta_\theta \omega(0)\omega(\epsilon) + \omega(0)\delta_\theta \omega(\epsilon)) \\ &= \frac{i\lambda}{32\pi G} \int_{M_3} \left(dTr\theta d\omega(\epsilon) - dTr\theta d\omega(0) \right) \end{aligned} \tag{4.9}$$

The variation of the modified action,

$$\delta_\theta \tilde{I}_{CS}^{(3)} = \delta_\theta \left(I_{CS}^{(3)} + \Delta I_{CS}^{(3)} \right) = \frac{i\lambda}{32\pi G} \int_{M_3} dTr\theta d\omega(0). \tag{4.10}$$

This is what we expected. Now the contribution from (4.8) is

$$\Delta S = \frac{i\lambda}{16G} \int_{\Sigma} \Gamma_{\sigma\mu}^\nu \epsilon^\mu{}_\nu dx^\sigma. \tag{4.11}$$

Including this contribution we recover previous result for black hole in 3D gravity with CS term. Then the HEE formula for 3D gravity theory with CS term is

$$\frac{1}{4G} \int dy \sqrt{g_{yy}} + \frac{i\lambda}{8G} \int \Gamma_{\sigma\mu}^\nu \epsilon^\mu{}_\nu dx^\sigma. \tag{4.12}$$

The HEE formula of theory with a gravitational CS term was first derived in [35]. One could check (4.12) is consistent with the result of [35] and [38]. Before we move on let's comment more on the squashed cone method for non-covariant theory. We show that the result obtained by using the squashed cone method by Lewkowycz-Maldacena [5] and Dong [8] is not correct. To get the correct result one could add a suitable boundary term, e.g., (4.8). Actually this method is consistent with the one used in [19] or [35]. In [19] and [35] the authors also used the squashed cone method. To get the correct result of the action in order $O(\epsilon)$, they make integration by parts and throw away a boundary term, see the formula (B.6) and (B.7) in [35]. One could also understand this as the statement of [19], $\omega = \omega_{\text{sing}} + \omega_{\text{reg}}$, so $R = d\omega_{\text{sing}} + d\omega_{\text{reg}} + \dots$, $\int \omega R = \int \omega_{\text{reg}} R_{\text{sing}} + \omega_{\text{sing}} d\omega_{\text{reg}} + \dots = 2 \int \omega_{\text{reg}} R_{\text{sing}} \dots$. A factor 2 also appears in this approach. If one just counts the terms that is related to $\partial_z \partial_{\bar{z}} A$ and $\partial_z A \partial_{\bar{z}} A$, the result is not correct. In fact the author in [36] just considered the contribution from terms related to $\partial_z \partial_{\bar{z}} A$, one could check the result in [36] is not consistent with (4.12). In higher dimension the anomaly term of the entropy (2.5) will appear. The generalization to higher dimension is straightforward but much more complex. We will comment on this problem later.

4.3 The surface

If we know the solution of a black hole in 3D, the result (4.2) can be directly used to calculate the entropy. But for HEE we have to find the surface Σ firstly. We would use boundary condition method to determine where the Σ should be. We will follow the same strategy as [5, 8].

We could parameterize the coordinate y in 3D, the metric is

$$ds^2 = e^{2A}[dzd\bar{z} + e^{2A}T(\bar{z}dz - zd\bar{z})^2] + (1 + 2K_ax^a + Q_{ab}x^ax^b)dy^2 + 2ie^{2A}(U + V_ax^a)(\bar{z}dz - zd\bar{z})dy + \dots \quad (4.13)$$

In three dimensions

$$C^{\mu\nu} = \epsilon^{\mu\kappa\sigma} \nabla_\kappa \left(R_\sigma^\nu - \frac{1}{4} \delta_\sigma^\nu R \right) \quad (4.14)$$

The equation of motion is

$$E_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}G_{\mu\nu}R - \frac{1}{l^2}G_{\mu\nu} - i\lambda C_{\mu\nu} = 0 \quad (4.15)$$

Let's check the divergent terms in E_{zz} . The result is

$$E_{zz} = \frac{\epsilon}{z} \left(-\frac{1}{2}K_z(y) + \lambda(U(y) - 3V_z)K_z(y) - \frac{i\lambda}{2}K'_z(y) \right) + \dots, \quad (4.16)$$

where the “...” means terms less divergent. We should set the divergent term in (4.16) to zero. We get the constraint,

$$-\frac{1}{2}K_z(y) + \lambda(U(y) - 3V_z)K_z(y) - \frac{i\lambda}{2}K'_z(y) = 0, \quad (4.17)$$

when $\lambda = 0$ we get the conditions for Einstein gravity ([5]). For $E_{\bar{z}\bar{z}}$ we would get a constraint on $K_{\bar{z}}$ with $z \leftrightarrow \bar{z}$ and $U(y) \leftrightarrow -U(y)$, $V_z \rightarrow -V_{\bar{z}}$ in (4.17). The CS term would give a non-trivial correction to constraint on the bulk surface Σ .

The question is considered in [35], they conclude that the minimization of (4.12) results in the Mathisson-Papapetrou-Dixon(MPD) equations for a spinning particle in 3D, which is exactly the equation (4.17). To get the correction to HEE by CS term, one need to solve the MPD equation firstly.¹

Actually without knowing the equation of motion of the surface, one also could get the leading contribution of the correction to HEE. According to method of [37], the coordinate y could be parameterized as $y = \rho$ near the boundary, where ρ is the coordinate of the bulk direction in the FG gauge, which states that any spacetime asymptotical to AdS admit the expansion

$$ds^2 = \frac{l^2}{4} \frac{d\rho^2}{\rho^2} + \frac{1}{\rho}(g_{(0)ij} + \rho g_{(1)ij} + \rho^2 g_{(2)ij} + \dots), \quad (4.18)$$

where $g_{(0)ij}$ is the boundary metric. Now we have two coordinates to describe the bulk metric, i.e., $X^\mu \in \{\rho, x_1, x_2\}$ and $\{z, \bar{z}, y\}$, where x_1 and x_2 are the boundary coordinates.

¹We would like to thank Prof. Takayanagi for reminding the paper [35] when preparing the draft.

To find the leading contribution of the entropy one needs to know the transformation between the two coordinates. It is possible to get the coordinate transformation near the boundary ($\rho \rightarrow 0$), as

$$\begin{aligned}\rho &= y + Bzy^{3/2} + C\bar{z}y^{3/2} + \dots \\ x^i &= x_0^i(y) + A^iy^{1/2}z + \bar{A}^iy^{1/2}\bar{z} + \dots,\end{aligned}\tag{4.19}$$

with $x_0^i = x_0^i(0) + Dy\dots$, where B, C, A^i, \bar{A}^i, D are constants, there are some relation among these parameters, which is not important for our purpose. This transformation can be obtained by considering the following constraints.

$$\frac{\partial X^\mu}{\partial z} \frac{\partial X^\nu}{\partial y} G_{\mu\nu}|_{z=0, \bar{z}=0} = 0, \text{ and } \frac{\partial X^\mu}{\partial z} \frac{\partial X^\nu}{\partial \bar{z}} G_{\mu\nu}|_{z=0, \bar{z}=0} = \frac{1}{2}.\tag{4.20}$$

We know from (4.12) that the additional terms for the HEE formula is proportional to

$$S_{\text{addition}} \propto \int dy U_y.\tag{4.21}$$

U_y can be written in the coordinate $\{\rho, x_1, x_2\}$ as

$$U_y \propto \frac{\partial X^\mu}{\partial y} \frac{\partial X^\nu}{\partial \bar{z}} \nabla_\mu n_{z\nu},\tag{4.22}$$

where $n_z^\nu \equiv \frac{\partial x^\nu}{\partial z}$, the derivative ∇ is defined in the coordinate $\{\rho, x_1, x_2\}$. One could take the coordinate transformation (4.19) into (4.22), and find the ρ^{-1} term is vanishing, thus $U_y \propto O(\rho^0)$. As we know the first term in (4.12) would contribute a log divergence term for the HEE. So the additional term in the theory with CS term would not contribute to the leading divergence. In [35] the authors calculate some examples in which the bulk are asymptotically to AdS_3 , the result is consistent with conclusion above.

5 Entropy of gravitational Chern-Simons terms in 7D

We find the contribution from the 3D CS term for HEE or black hole entropy. In 3D the possible correction related to extrinsic curvature do not appear. It's also interesting to investigate this property in higher dimensional theory. It is well known that gravitational CS term only exists in $(4n - 1)$ dimensional spacetime. We will discuss 7D theory in the following. Like the 3 dimensional case, one can't obtain the result directly by using (2.2). This is because of the fact that the action is not covariant. In this section we will use a "topological method" to get the result. We argue that the result is correct. As an important check this method could produce the correct result for Wald entropy in arbitrary dimension.

5.1 Approach to 7D case by a topological method

In this section we would like to use a "topological method" to derive the entropy for 7D theory with gravitational CS term. This method is based on the observation that the entropy of

a topological Invariant is a local total derivative.² This is indeed the case for Euler densities, or equivalently, the Lovelock gravity in critical dimensions [8], see also [39–41]. As we shall prove below, this is also the case for the Pontryagin density $tr(\mathbf{R}^4)$. Recall that we have

$$d\Omega_7(\omega) = Tr\mathbf{R}^4. \tag{5.1}$$

We propose to derive the entropy of 7D CS term s_7 from the following identity

$$ds_7 = \text{Entropy of } d\Omega_7(\omega) = \text{Entropy of } Tr\mathbf{R}^4. \tag{5.2}$$

Since the right hand side of (5.1) is invariant under the local Lorentz transformation, s_7 (up to an exact form) would also be free of the ambiguity. To make s_7 really be the entropy of 7d CS term, we have assumed

$$\text{Entropy of } d\Omega_7(\omega) = d(\text{Entropy of } \Omega_7(\omega)). \tag{5.3}$$

An evidence for the above approach is that we derive the correct entropy with zero extrinsic curvature, which is obtained by using the generalized covariant phase formalism in [26, 27].

Now let us start to derive the entropy of 7D CS term. We use the spin connection formalism in this section. Let's take a theory in 8D with the action,

$$I_8 = \int_{M_8} tr(\mathbf{R}^4). \tag{5.4}$$

By using the relation (3.1) one have

$$I_7 = \int_{M_7} \Omega_7, \tag{5.5}$$

where M_7 is a 7D manifold as the boundary of M_8 . The action (5.4) is invariant under the local Lorentz transformation. We would firstly get the entropy for the theory with such an action (5.4). The details of the calculation can be found in appendix C, we list the result as follows.

$$\begin{aligned} S_8 = -i\pi \int_{\Sigma_6} \sqrt{\det(g)} \hat{\epsilon}^{z\bar{z}i_1i_2i_3\dots i_6} & \left[-6K_{zi_1j_1} K_{\bar{z}i_2}^{j_1} R_{zj_2i_3i_4} R_{\bar{z}i_5i_6}^{j_2} \right. \\ & +64\partial_{i_1} U_{i_2} \partial_{i_3} U_{i_4} K_{z i_5}^{j_1} K_{\bar{z} i_6 j_1} + 48i K_{z i_1 j_1} K_{\bar{z} i_2}^{j_1} \partial_{i_3} U_{i_4} K_{z i_5 j_2} K_{\bar{z} i_6}^{j_2} \\ & -6K_{\bar{z} i_1}^{j_1} R_{z j_1 i_2 i_3} K_{z i_4 j_2} R_{\bar{z} i_5 i_6}^{j_2} + 2K_{z i_1 j_1} r_{i_2 i_3}^{j_2 j_1} K_{\bar{z} i_4 j_3} r_{j_2 i_5 i_6}^{j_3} \\ & -12K_{z i_1 j_1} r_{j_2 i_2 i_3}^{j_1} K_{\bar{z} i_4}^{j_2} K_{z i_5}^{j_3} K_{\bar{z} j_3 i_6} + 8i K_{z i_1 j_1} r_{j_2 i_2 i_3}^{j_1} K_{\bar{z} i_4}^{j_2} \partial_{i_5} U_{i_6} \\ & -32i \partial_{i_1} U_{i_2} \partial_{i_3} U_{i_4} \partial_{i_5} U_{i_6} + 8i \partial_{i_1} U_{i_2} R_{z j_1 i_3 i_4} R_{\bar{z} i_5 i_6}^{j_1} \\ & \left. + R_{z j_1 i_1 i_2} r_{j_2 i_3 i_4}^{j_1} R_{\bar{z} i_5 i_6}^{j_2} \right] + (z \leftrightarrow \bar{z}). \tag{5.6} \end{aligned}$$

where the integration is on the codimension-2 surface Σ_6 , “-i” appears because we are using Euclidean version. The result is still quite complex, to see it more clear, we would

²After our paper we notice that a recent paper also uses the same method to deal with the entropy CS term [38].

rewrite the result by forms. On the surface Σ_6 , where $z = \bar{z} = 0$, K_{aij} and U_i are one-form, R_{aijk} and r_{ijkl} are two forms. One could map the spacetime index into the local Lorenz coordinate by the vielbeins (3.6). From (3.6) we get

$$e_z^{a_1} = e_{\bar{z}}^{a_2} = \frac{1}{2}, \quad e_z^{a_2} = -e_{\bar{z}}^{a_2} = -\frac{i}{2}, \quad e_j^{a_i} = \bar{e}_j^{a_i}, \quad (5.7)$$

on the codimension-2 surface Σ_6 . In the following the index $a, a', a'' \dots$ denote the element of the set $\{a_1, a_2\}$, $b, b', b'' \dots$ denote the set $\{a_i\}$, $c, c' \dots$ denote the set $\{a_1, a_2, a_i\}$. By using (5.7) and (A.2) we have the following relation³

$$K_{zij} = e_z^a e_i^b \omega_{j,ba}, \quad -2iU_i = e_z^{a'} e_z^{a''} \omega_{i,a'a''}, \quad r_{ijkl} = e_i^{b'} e_j^{b''} r_{b'b''kl}, \quad (5.8)$$

one could rewrite (5.6) as⁴

$$\begin{aligned} S_8 = -i\pi \int_{\Sigma_6} & \left[-6\epsilon^{aa'} \omega_{ba} \omega_{a'}^b \mathbf{R}_{a''b'} \mathbf{R}^{b'a''} + 64dU dU \epsilon^{aa'} \omega_{ba} \omega_{a'}^b \right. \\ & + 24idU (\epsilon^{aa'} \omega_{ba} \omega_{a'}^b)^2 + 8\omega_{ba} \mathbf{R}^{ba} \epsilon^{a'a''} \omega_{b'a'} \mathbf{R}^{b'a''} \\ & + 8\epsilon^{aa'} \omega_{ba} r_{b'}^b r_{b''}^{b'} \omega_{a'}^{b''} - 6\omega_{ba} r_{b'}^b \omega_{b''}^{b'} \epsilon^{a'a''} \omega_{b''}^{b'} \omega_{a'}^{b''} \\ & + 32\omega_{ba} r_{b'}^b \omega_{b''}^{b'} dU - 128dU dU dU + 32dU \mathbf{R}_{ab} \mathbf{R}^{ba} \\ & \left. + 8\epsilon^{aa'} \mathbf{R}_{ab} r_{b'}^b \mathbf{R}^{b'a'} \right], \quad (5.9) \end{aligned}$$

where $\epsilon^{aa'} \equiv \epsilon^{z\bar{z}} e_z^a e_{\bar{z}}^{a'} + \epsilon^{\bar{z}z} e_{\bar{z}}^a e_z^{a'}$, $\epsilon^{z\bar{z}} = -\epsilon^{\bar{z}z} = -ig^{z\bar{z}}$. To simplify result we need the relation

$$\mathbf{R}_{ab} \equiv d\omega_{ab} + \omega_{ac} \omega_b^c = d\omega_{ab} + \omega_{aa'} \omega_b^{a'} + \omega_{ab'} \omega_b^{b'}, \quad (5.10)$$

$$r_{bb'} = d\omega_{bb'} + \omega_{bb''} \omega_{b''}^{b'}. \quad (5.11)$$

The result is

$$S_8 = \int_{\Sigma_6} ds_7, \quad (5.12)$$

with

$$\begin{aligned} s_7 = -i\pi & \left[-128U dU dU - 6\epsilon^{aa'} \omega_{ba} \omega_{a'}^b \omega_{a''b''} R^{b''a''} \right. \\ & - 64U dU \epsilon^{aa'} \omega_{ab} \omega_{a'}^b + 32U d\omega_{ab} d\omega^{ba} - 32dU \omega_{ab} \omega_{b'}^b \omega^{b'a} \\ & - 8\epsilon^{aa'} \omega_{ba} \omega_{b'}^b d\omega_{b''}^{b'} \omega_{a'}^{b''} - 8\epsilon^{aa'} d\omega_{ba} \omega_{b'}^b \omega_{b''}^{b'} \omega_{a'}^{b''} + 16U \omega_{ab} d\omega_{b'}^b \omega^{b'a} \\ & \left. + 8\epsilon^{a''a} \omega_{ab} \omega_{b'}^b \omega_{b''}^{b'} \omega_c^{b''} \omega_{a''}^c + 8\epsilon^{aa'} \omega_{ab} d\omega_{b'}^b d\omega_{a'}^{b'} \right]. \quad (5.13) \end{aligned}$$

One could define the “density” of the entropy in 8D s_8 , and the relation $s_8 = ds_7$. This is the expected relation that we mentioned in the beginning of this subsection. s_8 is the entropy that we obtain from the Pontryagin density, it has the similar relation (3.1) which must be satisfied by the Pontryagin class. The entropy for the CS term in 7D is

$$S_7 = \int_{\Sigma_5} (s_7 + ds'), \quad (5.14)$$

³For example, $K_{zij} = \omega_{j,iz} = \omega_{j,cc'} e_i^c e_z^{c'} = \omega_{j,ba} e_i^b e_z^a$, since $e_z^b = e_i^a = 0$, where $a \in \{a_1, a_2\}$, $b \in \{a_i\}$.

⁴We note that $dx^{\nu_1} \wedge dx^{\nu_2} \dots \wedge dx^{\nu_n} = \epsilon^{\nu_1 \nu_2 \dots \nu_n} \sqrt{\det(g)} dx^n$.

where the surface Σ_5 is the codimension-2 surface in 7D, which is also a boundary of Σ_6 ,⁵ \mathbf{s}' is arbitrary. Our above approach actually do not use the vielbeins in the regularized spacetime (3.6), it is not expected the result is effected by the local gauge transformation. We also show (5.13) is same as the result in [38] up to a boundary term in appendix D.

Here we only calculate the 7D result. We expect it can be generalized to $(4n - 1)$ D without any difficulty in principle. Conversely, we could say our result above provides another evidence to support our proposal that the entropy of a topological Invariant is also a topological Invariant.

6 Wald entropy in arbitrary dimension

The result (5.13) is new, and now we don't have other methods to compare with. When setting extrinsic curvature to zero, the entropy is just the Wald entropy. One could derive the Wald entropy by other method, see [26, 27]. So the Wald entropy is a good example to check the topological method. As we have discussed in section 4.2 for 3D case, one can not obtain the correct result by using the squashed cone method directly. We also mentioned the trick used in [19] and [35] is equal to add a boundary term in the action. We would like to use the topological method and boundary term method in this section to calculate the Wald entropy.

6.1 The topological method

Here we use a topological approach to this problem by considering the relation (3.1). For 3 dimension

$$d\Omega_3 = Tr \mathbf{R}^2, \tag{6.1}$$

here \mathbf{R} lives in 4 dimension spacetime M_4 , Ω_3 lives in M_3 . The above terms are invariant under local Lorentz transformation. Let's define

$$\tilde{I}_4 = \int_{M_4} Tr \mathbf{R}^2 = \int_{M_3} \Omega_3. \tag{6.2}$$

It's easy to check the contribution from $(\partial_z A, \partial_{\bar{z}} A)$ vanishes. The total contribution of (6.2) to the entropy is

$$\tilde{S}_4 = 8i\pi \int_{\Sigma_2} \mathbf{R}_N, \tag{6.3}$$

where the integration is on the a dimension-2 surface Σ_2 , \mathbf{R}_N is defined as

$$\mathbf{R}_N \equiv \frac{1}{2} tr \epsilon \mathbf{R}. \tag{6.4}$$

Note that $\mathbf{R}_N = d\Gamma_N$, where “ d ” is the defined on Σ_2 ,

$$\Gamma_N \equiv \frac{1}{2} tr(\epsilon \Gamma). \tag{6.5}$$

⁵Actually the action in 8D depends on one more coordinate, the result of the corresponding entropy is dependent on the coordinate. But one can assume to choose a suitable Mainfold for M_8 , on which the boundary is M_7 , and the boundary of the surface Σ_6 is Σ_5 .

One could rewrite (6.3) as

$$\tilde{S}_4 = 8i\pi \int_{\Sigma_1} \Gamma_N, \quad (6.6)$$

where Σ_1 is a codimension-3 surface in M_4 , as well as a codimension-2 surface in M_3 . Formally considering (6.2) we have the entropy

$$S_3 = 8i\pi \int_{\Sigma_1} \Gamma_N, \quad (6.7)$$

if one uses the trick to find suitable M_4 such that M_3 is a boundary of M_4 , and Σ_1 as a boundary of Σ_2 .

The result is same as (4.12), and also [19, 35]. Now it is easy to generalize the method to higher dimension. Actually the generalization is quite trivial. With the relation

$$d\Omega_{2n+1} = \text{tr} \mathbf{R}^{n+1}, \quad (6.8)$$

one could have

$$\tilde{I}_{2n+2} = \int_{M_{2n+2}} \text{tr} \mathbf{R}^{n+1} = \int_{M_{2n+1}} \Omega_{2n+1}, \quad (6.9)$$

The entropy from \tilde{I}_n is

$$\tilde{S}_{2n+2} = 4i\pi(n+1) \int_{\Sigma_{2n}} \mathbf{R}_N^n. \quad (6.10)$$

With the relation $\mathbf{R}_N = d\Gamma_N$, one have

$$S_{2n+1} = 4i\pi(n+1) \int_{\Sigma_{2n-1}} \Gamma_N \mathbf{R}_N^{n-1}. \quad (6.11)$$

The result is consistent with the one that is obtained by the covariant phase formalism [26, 27].

6.2 Another approach

Actually we also could use the trick in 3 dimension to get the correct result for Wald entropy in arbitrary dimension. We will briefly state the process in the following. As the appendix B shows, a local gauge transformation would lead to a boundary term which is related with ϵ in the regularized spacetime. Even if we only consider the Wald entropy, i.e., the extrinsic curvature is vanishing, this ambiguity still exist. The following term is suitable to add to the action,⁶

$$\Delta I_n = -n(n+1) \int_0^1 dt (t-1)t^{n-1} \int_M \text{dstr}(\omega(0), \omega(\epsilon), \mathbf{R}_t^{n-1}), \quad (6.13)$$

where $\omega(\epsilon)$ is the spin connection in the regularized metric (2.3), \mathbf{R}_t is constructed by $\omega(\epsilon)$. After complex calculation one could get the variance of the action with the additional term,

$$\delta_\theta(I_n + \Delta I_n) = -n(n+1) \int_0^1 dt (t-1)t^{n-1} \int_M \left[-\text{str}(\mathbf{R}_1, d\theta, \mathbf{R}_t^{n-1}) \right]$$

⁶The CS action is normalized as

$$I_n = (n+1) \int_M \Omega_{2n+1}. \quad (6.12)$$

$$\begin{aligned}
 & +str(\omega(0), \mathbf{R}_2, \mathbf{R}_t^{n-1})] \\
 & +2n(n+1) \int_0^1 dt (t-1)^2 t^{n-1} \int_M \left[str(\mathbf{R}_1, \omega, \mathbf{R}_2, \mathbf{R}_t^{n-2}) \right. \\
 & \left. -str(\omega(0), D_t \omega, \mathbf{R}_2, \mathbf{R}_t^{n-2}) + str(\omega(0), \omega, \mathbf{R}_3, \mathbf{R}_t^{n-2}) \right], \quad (6.14)
 \end{aligned}$$

with the following definitions,

$$\mathbf{R}_1 = D_t \omega(0), \quad \mathbf{R}_2 = D_t d\theta, \quad \mathbf{R}_3 = D_t D_t d\theta = [\mathbf{R}_t, d\theta], \quad (6.15)$$

see the definition of “ D_t ” in appendix B. One could extract the $O(\epsilon)$ terms in (6.14), and finally get

$$\Delta S = -4i\pi(n+1) \int_{\Sigma} df \mathbf{R}_N^{n-1}, \quad (6.16)$$

where Σ is the codimension-2 surface, f is defined as

$$f \equiv \frac{n-1}{n+1} tr(\theta \epsilon). \quad (6.17)$$

R_N is defined as

$$\mathbf{R}_N \equiv \frac{1}{2} tr(\epsilon \mathbf{R}) \quad (6.18)$$

where ϵ is 2-dimension Levi-Civita symbol with $\epsilon_{12} = -\epsilon_{21} = 1$. (6.17) is actually a integration of total derivative term. Thus the local gauge transformation will not effect the final result of the entropy, which is what we expect. With the additional term one could get the entropy,

$$S = 4i\pi(n+1) \int_{\Sigma} \omega_N \mathbf{R}_N^{n-1}, \quad (6.19)$$

where \mathbf{R}_N is defined by (6.18), and ω_N is defined as

$$\omega_N = \frac{1}{2} tr(\epsilon \omega), \quad (6.20)$$

the result is consistent with [26, 27] and (6.11). (6.18) and (6.20) are related with each other by

$$d\omega_N = \mathbf{R}_N, \quad (6.21)$$

for $tr(\epsilon \omega \omega) = 0$. Then it's obvious (6.16) can be seen as the gauge transformation

$$\omega_N \rightarrow \omega_N - df. \quad (6.22)$$

Note that in 3D $f = 0$, so the result in 3D is gauge invariant.

We use the spin connection formulism of CS action in the above discussion. But it's easy to generalize the result to Christoffel symbols formulism. The non-covariant part of the diffeomorphism δ_{ξ} of Γ is,

$$\hat{\delta}_{\xi} \Gamma = d\Lambda, \quad (6.23)$$

where $\Lambda^a_b = \partial_b \xi^a$. If one replaces the parameter θ with Λ , as well as Γ with ω , all the result for Christoffel symbols formulism can be obtained.

We show that the two methods both could get the correct result for Wald entropy. The topological method is more reasonable and simpler. This method depends on the special property of the gravitational CS terms. For other non-covariant theories, one could not generalize the topological method. As we have mentioned adding a boundary term is equal to making integration by parts [38]. In [38] the authors make more details discussion on how to obtain the correct result by making integration by parts. In principle there is no difficulty to write down a suitable boundary term to get the result (5.13).

7 Conclusion and discussion

In this paper we analyze the entropy when gravitational Chern-Simons terms are added into the action. We find it is necessary to modify the squashed cone method by Lewkowycz-Maldacena [5] and Dong [8] for theory with gravitational Chern-Simons terms. The necessity is related to the non-covariant part transformation of diffeomorphism since a total derivative term would appear under such transformation. The covariant theory is free of this ambiguity. One possible solution to this problem is to add a total derivative term into the original Lagrangian, which does not affect the equation of motion, at the same time eliminates the ambiguity caused by the diffeomorphism. This term is also vanishing in the limit $\epsilon \rightarrow 0$. We show how to add a boundary term to get the correct Wald entropy. In principle it is also straightforward to find a suitable boundary term to get correct result for general case, i.e., the extrinsic curvature is non-vanishing, by following [38].

On the other hand we propose a ‘topological approach’ to calculate the entropy of gravitational Chern-Simons terms when the extrinsic curvature K_{aij} is non-vanishing. It yields the correct Wald entropy in arbitrary dimension and gives non-trivial results when the extrinsic curvature does not vanish. Our results imply that the entropy of a topological invariant seems to also be a topological invariant. There may exist some mathematical interpretations or correspondence for this nice property. We hope someone could clarify this problem in future.

Acknowledgments

R. X. Miao is supported by Sino-Germann (CSC-DAAD) Postdoc Scholarship Program. W. Z. Guo is supported by Postgraduate Scholarship Program of China Scholarship Council. We thank Prof. Miao Li for his encouragement and support. We are grateful to School of Astronomy and Space Science at Sun Yat-Sen University for hospitality where part of work was done. We thank Tadashi Takayanagi, Song He, Noburo Shiba for discussions.

A Useful components of Γ and R

$$\Gamma_{\bar{z}\bar{z}}^{\bar{z}} = 2\partial_{\bar{z}}A, \quad \Gamma_{i\bar{z}}^{\bar{z}} = 2iU_i, \quad \Gamma_{\bar{z}i}^j = g^{jk}K_{\bar{z}ki}, \quad \Gamma_{ij}^z = -2e^{-2A}K_{\bar{z}ji}, \quad \Gamma_{li}^m = \gamma_{li}^m, \quad (\text{A.1})$$

$$\omega_z^z{}_z = -\omega_z^{\bar{z}}{}_{\bar{z}} = \partial_z A, \quad \omega_i^z{}_z = -2iU_i, \quad \omega_i^j{}_z = K_z^j{}_i, \quad \omega_i^z{}_j = -2e^{-2A}K_{\bar{z}}^j{}_i, \quad \omega_i^k{}_j = \bar{\omega}_i^k{}_j, \quad (\text{A.2})$$

$$\begin{aligned}
 R_{z\bar{z}z\bar{z}} &= e^{2A} \left(\partial_z \partial_{\bar{z}} A - 3T e^{2A} \right), \\
 R_{z\bar{z}zi} &= \frac{1}{2} e^{2A} \left[2i U_i (\bar{z} \partial_z \partial_{\bar{z}} A + \partial_z A + z \partial_z \partial_z A) + 3i V_{zi} \right] \\
 R_{zizj} &= \frac{1}{2} \left[4K_{zij} \partial_z A - 2Q_{zzij} + 2g^{lk} K_{zlj} K_{zik} \right] \\
 R_{z\bar{z}ij} &= \frac{1}{2} e^{2A} \left[2i \partial_i U_j - 2e^{-2A} g^{ln} K_{\bar{z}nj} K_{zli} \right] - (i \leftrightarrow j), \\
 R_{zi\bar{z}j} &= \frac{1}{2} e^{2A} \left[i(\partial_i U_j - \partial_j U_i) + 2e^{-2A} g^{mn} K_{\bar{z}ni} K_{z mj} + 4U_i U_j - 2e^{-2A} Q_{z\bar{z}ij} \right] \\
 R_{zij k} &= \frac{1}{2} e^{2A} \left[-2e^{-2A} \partial_j K_{zki} - 4ie^{-2A} U_j K_{zik} - 2e^{-2A} K_{zlj} \gamma_{ik}^l \right] - (j \leftrightarrow k) \\
 R_{ikjl} &= r_{ikjl} - \left[2e^{-2A} (K_{zij} K_{\bar{z}kl} + K_{\bar{z}ij} K_{zkl}) \right] - (j \leftrightarrow k).
 \end{aligned} \tag{A.3}$$

B Non-covariant part of Ω

Following the step as [26], we define the ‘‘covariant’’ derivative

$$D = d + [\omega, \], \quad D_t = d + [t\omega, \], \tag{B.1}$$

note that $\frac{d}{dt}(t\mathbf{R}_t) = D_t \omega$ one can get

$$\begin{aligned}
 \hat{\delta}\Omega_{2n+1} &= (n+1) \int_0^1 dt t^n \left(\text{str}(\hat{\delta}\omega, \mathbf{R}_t^n) + n \text{str}(\omega, \hat{\delta}\mathbf{R}_t, \mathbf{R}_t^{n-1}) \right) \\
 &= (n+1) \int_0^1 dt \left(t^n \text{str}(d\theta, \mathbf{R}_t^n) + n t^{n-1} \text{str}(\omega, (t-1) D_t d\theta, \mathbf{R}_t^{n-1}) \right) \\
 &= (n+1) \int_0^1 dt \left[t^n \text{str}(d\theta, \mathbf{R}_t^n) - n t^{n-1} d \text{str}(\omega, (t-1) d\theta, \mathbf{R}_t^{n-1}) \right. \\
 &\quad \left. + n \text{str} \left(\frac{d}{dt}(t\mathbf{R}_t), (t-1) d\theta, (t\mathbf{R}_t)^{n-1} \right) \right],
 \end{aligned} \tag{B.2}$$

where the last step we use

$$d \text{str}(\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_n) = \sum_{i=1}^n (-)^{a_1+a_2+\dots+a_{i-1}} \text{str}(\mathbf{A}_1, \dots, D(\mathbf{A}_i), \dots), \tag{B.3}$$

\mathbf{A}_i denotes the a_i -form, the covariant derivative $D \equiv d + [\Theta, \], \Theta$ is 1-form. Then

$$\begin{aligned}
 \int_0^1 dt n \text{str} \left(\frac{d}{dt}(t\mathbf{R}_t), (t-1) d\theta, (t\mathbf{R}_t)^{n-1} \right) &= \\
 &= \int_0^1 dt \left(\frac{d}{dt} \text{str}((t-1) d\theta, t^n \mathbf{R}_t^n) - t^n \text{str}(d\theta, \mathbf{R}_t^n) \right),
 \end{aligned} \tag{B.4}$$

the first term vanishes. We get

$$\hat{\delta}\Omega_{2n+1} = -n(n+1) d \int_0^1 dt t^{n-1} (t-1) \text{str}(\omega, d\theta, \mathbf{R}_t^{n-1}). \tag{B.5}$$

C Details of the calculation in section 5

The action (5.4) can be written by the spacetime components as

$$I_8 = -\frac{i}{16} \int_{M_8} \sqrt{\det(G)} \epsilon^{\beta_1 \beta_2 \dots \beta_8} R^{\alpha_1}_{\alpha_2 \beta_1 \beta_2} R^{\alpha_2}_{\alpha_3 \beta_3 \beta_4} R^{\alpha_3}_{\alpha_4 \beta_5 \beta_6} R^{\alpha_4}_{\alpha_1 \beta_7 \beta_8}, \quad (\text{C.1})$$

where “-i” appears because we work in the Euclidean spacetime. The contribution to the entropy from $R_{z\bar{z}z\bar{z}}$ is

$$\begin{aligned} I^{(1)} = -2i \int_{M_8} \sqrt{\det(G)} \epsilon^{z\bar{z}i_1 i_2 i_3 i_4 i_5 i_6} R_{z\bar{z}z\bar{z}} & \left[4R_{z\bar{z}i_1 i_2} R_{z\bar{z}i_3 i_4} R_{z\bar{z}i_5 i_6} \right. \\ & + 2R_{z\bar{z}i_1 i_2} R_{zj_1 i_3 i_4} R_{\bar{z}i_5 i_6}^{j_1} + 2R_{zj_1 i_1 i_2} R_{\bar{z}i_3 i_4}^{j_1} R_{z\bar{z}i_5 i_6} \\ & \left. + R_{zj_1 i_1 i_2} R_{j_2 i_3 i_4}^{j_1} R_{\bar{z}i_5 i_6}^{j_2} \right] \end{aligned} \quad (\text{C.2})$$

The contribution from $(R_{zizj}, R_{\bar{z}i\bar{z}j})$ is

$$\begin{aligned} I^{(2)} = -4i \int_{M_8} \sqrt{\det(G)} \epsilon^{z\bar{z}i_1 i_2 i_3 i_4 i_5 i_6} \partial_z A \partial_{\bar{z}} A & \left[R_{zj_1 i_1 i_2} R_{\bar{z}i_3 i_4}^{j_1} K_{zi_5 j_2} K_{\bar{z}i_6}^{j_2} (G^{z\bar{z}})^2 \right. \\ & + K_{zi_1 j_1} K_{\bar{z}i_2}^{j_2} R_{z\bar{z}i_3 i_4} R_{z\bar{z}i_5 i_6} (G^{z\bar{z}})^3 + K_{zi_1}^{j_1} K_{\bar{z}i_2 j_2} R_{zi_3 i_4}^{j_2} R_{\bar{z}j_1 i_5 i_6} (G^{z\bar{z}})^2 \\ & + K_{zi_1}^{j_1} K_{\bar{z}i_2 j_2} R_{\bar{z}i_3 i_4}^{j_2} R_{zj_1 i_5 i_6} (G^{z\bar{z}})^2 + K_{zi_1}^{j_1} K_{\bar{z}i_1 j_2} R_{j_4 i_3 i_4}^{j_3} R_{j_1 i_5 i_6}^{j_3} G^{z\bar{z}} \\ & \left. - K_{zi_1 j_1} R_{zi_2 i_3}^{j_1} K_{\bar{z}i_4 j_2} R_{\bar{z}i_5 i_6}^{j_2} + K_{zi_1 j_1} R_{j_2 i_2 i_3}^{j_1} K_{\bar{z}i_4}^{j_2} R_{z\bar{z}i_5 i_6} \right] + (z \leftrightarrow \bar{z}). \end{aligned} \quad (\text{C.3})$$

One could get the $O(\epsilon)$ term of the above terms after complex calculation,

$$\begin{aligned} S_8 = -i\pi \int_{\Sigma_6} \sqrt{\det(g)} \hat{\epsilon}^{z\bar{z}i_1 i_2 i_3 \dots i_6} & \left[-6K_{zi_1 j_1} K_{\bar{z}i_2}^{j_1} R_{zj_2 i_3 i_4} R_{\bar{z}i_5 i_6}^{j_2} \right. \\ & + 64\partial_{i_1} U_{i_2} \partial_{i_3} U_{i_4} K_{zi_5}^{j_1} K_{\bar{z}i_6 j_1} + 48iK_{zi_1 j_1} K_{\bar{z}i_2}^{j_1} \partial_{i_3} U_{i_4} K_{zi_5 j_2} K_{\bar{z}i_6}^{j_2} \\ & - 6K_{\bar{z}i_1}^{j_1} R_{zj_1 i_2 i_3} K_{zi_4 j_2} R_{\bar{z}i_5 i_6}^{j_2} + 2K_{zi_1 j_1} r_{i_2 i_3}^{j_2 j_1} K_{\bar{z}i_4 j_3} r_{j_2 i_5 i_6}^{j_3} \\ & - 12K_{zi_1 j_1} r_{j_2 i_2 i_3}^{j_1} K_{\bar{z}i_4}^{j_2} K_{zi_5}^{j_3} K_{\bar{z}j_3 i_6} + 8iK_{zi_1 j_1} r_{j_2 i_2 i_3}^{j_1} K_{\bar{z}i_4}^{j_2} \partial_{i_5} U_{i_6} \\ & - 32i\partial_{i_1} U_{i_2} \partial_{i_3} U_{i_4} \partial_{i_5} U_{i_6} + 8i\partial_{i_1} U_{i_2} R_{zj_1 i_3 i_4} R_{\bar{z}i_5 i_6}^{j_1} \\ & \left. + R_{zj_1 i_1 i_2} r_{j_2 i_3 i_4}^{j_1} R_{\bar{z}i_5 i_6}^{j_2} \right] + (z \leftrightarrow \bar{z}). \end{aligned} \quad (\text{C.4})$$

D Comparison with [38]

We would like to show our result (5.13) is different from the one in the recent paper [38] only by a boundary term, in which the author calculate the same quantity. One could see the paper [38] for more details. Since the notations in [38] are not all same as ones we use here, we rewrite the result in [38] by our notations as

$$\begin{aligned} s_{ent,CS}^{7d, single} = 16\pi & \left[8iU dU^2 + [4idU + 3\omega_{\bar{z}} \cdot \omega_z] \wedge [\omega_z \cdot \hat{D}\omega_{\bar{z}} + \omega_{\bar{z}} \cdot \hat{D}\omega_z] \right. \\ & \left. + \hat{D}\omega_z \cdot r \cdot \omega_{\bar{z}} - \hat{D}\omega_{\bar{z}} \cdot r \cdot \omega_z \right], \end{aligned} \quad (\text{D.1})$$

where ω_z denotes ω_{bz} , $\hat{D}\omega_{bz} \equiv D\omega_{bz} + 2iU\omega_{bz}$. One could get the following results

$$16\pi \times 4idU \wedge [\omega_z \cdot \hat{D}\omega_{\bar{z}} + \omega_{\bar{z}} \cdot \hat{D}\omega_z] \tag{D.2}$$

$$= 64i\pi dUU\epsilon^{aa'}\omega_{ab}\omega_{a'}^b - 32i\pi U d\omega_{ab}d\omega^{ba} + 32i\pi dU\omega_{ab}\omega_{b'}^b\omega^{b'a} - 32i\pi d[U\omega_{ab}d\omega^{ba}],$$

$$16\pi \times 3\omega_{\bar{z}} \cdot \omega_z \wedge [\omega_z \cdot \hat{D}\omega_{\bar{z}} + \omega_{\bar{z}} \cdot \hat{D}\omega_z] \tag{D.3}$$

$$= 6i\pi\epsilon^{aa'}\omega_{ba}\omega_{a'}^b\omega_{a''}^{b''}R^{b''a''}$$

$$16\pi \times [\hat{D}\omega_z \cdot r \cdot \omega_{\bar{z}} - \hat{D}\omega_{\bar{z}} \cdot r \cdot \omega_z] \tag{D.4}$$

$$= 8i\pi\epsilon^{aa'}\omega_{ba}\omega_{b'}^b d\omega_{b''}^{b''}\omega_{a'}^{b''} + 8i\pi\epsilon^{aa'}d\omega_{ba}\omega_{b'}^b\omega_{b''}^{b''}\omega_{a'}^{b''} - 16i\pi U\omega_{ab}d\omega_{b'}^b\omega^{b'a} \\ - 8i\pi\epsilon^{aa''}\omega_{ba}\omega_{b'}^b\omega_{b''}^{b''}\omega_c^{c a''} - 8i\pi\epsilon^{aa'}\omega_{ab}d\omega_{b'}^b d\omega_{a'}^{b'}$$

Now we could see the difference between (D.1) and (5.13) is just a boundary term $-32i\pi d[U\omega_{ab}d\omega^{ba}]$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J.D. Bekenstein, *Black holes and entropy*, *Phys. Rev. D* **7** (1973) 2333 [[INSPIRE](#)].
- [2] S.W. Hawking, *Particle creation by black holes*, *Commun. Math. Phys.* **43** (1975) 199 [[Erratum ibid.](#) **46** (1976) 206] [[INSPIRE](#)].
- [3] S. Ryu and T. Takayanagi, *Holographic derivation of entanglement entropy from AdS/CFT*, *Phys. Rev. Lett.* **96** (2006) 181602 [[hep-th/0603001](#)] [[INSPIRE](#)].
- [4] R.M. Wald, *Black hole entropy is the Noether charge*, *Phys. Rev. D* **48** (1993) 3427 [[gr-qc/9307038](#)] [[INSPIRE](#)].
- [5] A. Lewkowycz and J. Maldacena, *Generalized gravitational entropy*, *JHEP* **08** (2013) 090 [[arXiv:1304.4926](#)] [[INSPIRE](#)].
- [6] D.V. Fursaev, A. Patrushev and S.N. Solodukhin, *Distributional geometry of squashed cones*, *Phys. Rev. D* **88** (2013) 044054 [[arXiv:1306.4000](#)] [[INSPIRE](#)].
- [7] X. Dong and R.-X. Miao, *Generalized gravitational entropy from total derivative action*, *JHEP* **12** (2015) 100 [[arXiv:1510.04273](#)] [[INSPIRE](#)].
- [8] X. Dong, *Holographic entanglement entropy for general higher derivative gravity*, *JHEP* **01** (2014) 044 [[arXiv:1310.5713](#)] [[INSPIRE](#)].
- [9] J. Camps, *Generalized entropy and higher derivative gravity*, *JHEP* **03** (2014) 070 [[arXiv:1310.6659](#)] [[INSPIRE](#)].
- [10] R.-X. Miao and W.-Z. Guo, *Holographic entanglement entropy for the most general higher derivative gravity*, *JHEP* **08** (2015) 031 [[arXiv:1411.5579](#)] [[INSPIRE](#)].
- [11] R.-X. Miao, *Universal terms of entanglement entropy for 6d CFTs*, *JHEP* **10** (2015) 049 [[arXiv:1503.05538](#)] [[INSPIRE](#)].
- [12] Y. Huang and R.-X. Miao, *A note on the resolution of the entropy discrepancy*, *Phys. Lett. B* **749** (2015) 489 [[arXiv:1504.02301](#)] [[INSPIRE](#)].

- [13] A. Bhattacharyya, A. Kaviraj and A. Sinha, *Entanglement entropy in higher derivative holography*, *JHEP* **08** (2013) 012 [[arXiv:1305.6694](#)] [[INSPIRE](#)].
- [14] A. Bhattacharyya, M. Sharma and A. Sinha, *On generalized gravitational entropy, squashed cones and holography*, *JHEP* **01** (2014) 021 [[arXiv:1308.5748](#)] [[INSPIRE](#)].
- [15] A. Bhattacharyya and M. Sharma, *On entanglement entropy functionals in higher derivative gravity theories*, *JHEP* **10** (2014) 130 [[arXiv:1405.3511](#)] [[INSPIRE](#)].
- [16] S.L. Dubovsky and S.M. Sibiryakov, *Spontaneous breaking of Lorentz invariance, black holes and perpetuum mobile of the 2nd kind*, *Phys. Lett. B* **638** (2006) 509 [[hep-th/0603158](#)] [[INSPIRE](#)].
- [17] T. Jacobson and A.C. Wall, *Black hole thermodynamics and Lorentz symmetry*, *Found. Phys.* **40** (2010) 1076 [[arXiv:0804.2720](#)] [[INSPIRE](#)].
- [18] R.-X. Miao, M. Li and Y.-G. Miao, *Violation of the first law of black hole thermodynamics in $f(T)$ gravity*, *JCAP* **11** (2011) 033 [[arXiv:1107.0515](#)] [[INSPIRE](#)].
- [19] S.N. Solodukhin, *Holography with gravitational Chern-Simons*, *Phys. Rev. D* **74** (2006) 024015 [[hep-th/0509148](#)] [[INSPIRE](#)].
- [20] R.F. Perez, *Conserved current for the Cotton tensor, black hole entropy and equivariant Pontryagin forms*, *Class. Quant. Grav.* **27** (2010) 135015 [[arXiv:1004.3161](#)] [[INSPIRE](#)].
- [21] P. Kraus and F. Larsen, *Holographic gravitational anomalies*, *JHEP* **01** (2006) 022 [[hep-th/0508218](#)] [[INSPIRE](#)].
- [22] M.-I. Park, *BTZ black hole with gravitational Chern-Simons: thermodynamics and statistical entropy*, *Phys. Rev. D* **77** (2008) 026011 [[hep-th/0608165](#)] [[INSPIRE](#)].
- [23] O. Mišković and R. Olea, *Background-independent charges in topologically massive gravity*, *JHEP* **12** (2009) 046 [[arXiv:0909.2275](#)] [[INSPIRE](#)].
- [24] P. Kraus and F. Larsen, *Microscopic black hole entropy in theories with higher derivatives*, *JHEP* **09** (2005) 034 [[hep-th/0506176](#)] [[INSPIRE](#)].
- [25] Y. Tachikawa, *Black hole entropy in the presence of Chern-Simons terms*, *Class. Quant. Grav.* **24** (2007) 737 [[hep-th/0611141](#)] [[INSPIRE](#)].
- [26] L. Bonora, M. Cvitan, P. Dominis Prester, S. Pallua and I. Smolic, *Gravitational Chern-Simons Lagrangians and black hole entropy*, *JHEP* **07** (2011) 085 [[arXiv:1104.2523](#)] [[INSPIRE](#)].
- [27] L. Bonora, M. Cvitan, P.D. Prester, S. Pallua and I. Smolic, *Gravitational Chern-Simons terms and black hole entropy. Global aspects*, *JHEP* **10** (2012) 077 [[arXiv:1207.6969](#)] [[INSPIRE](#)].
- [28] K. Jensen, R. Loganayagam and A. Yarom, *Anomaly inflow and thermal equilibrium*, *JHEP* **05** (2014) 134 [[arXiv:1310.7024](#)] [[INSPIRE](#)].
- [29] K. Jensen, R. Loganayagam and A. Yarom, *Chern-Simons terms from thermal circles and anomalies*, *JHEP* **05** (2014) 110 [[arXiv:1311.2935](#)] [[INSPIRE](#)].
- [30] T. Azeyanagi, R. Loganayagam, G.S. Ng and M.J. Rodriguez, *Holographic thermal helicity*, *JHEP* **08** (2014) 040 [[arXiv:1311.2940](#)] [[INSPIRE](#)].
- [31] T. Azeyanagi, R. Loganayagam, G.S. Ng and M.J. Rodriguez, *Covariant Noether charge for higher dimensional Chern-Simons terms*, *JHEP* **05** (2015) 041 [[arXiv:1407.6364](#)] [[INSPIRE](#)].

- [32] T. Azeyanagi, R. Loganayagam and G.S. Ng, *Anomalies, Chern-Simons terms and black hole entropy*, *JHEP* **09** (2015) 121 [[arXiv:1505.02816](#)] [[INSPIRE](#)].
- [33] S.N. Solodukhin, *Holographic description of gravitational anomalies*, *JHEP* **07** (2006) 003 [[hep-th/0512216](#)] [[INSPIRE](#)].
- [34] S. Deser, R. Jackiw and S. Templeton, *Topologically massive gauge theories*, *Annals Phys.* **140** (1982) 372 [*Erratum ibid.* **185** (1988) 406] [*Erratum ibid.* **281** (2000) 409] [[INSPIRE](#)].
- [35] A. Castro, S. Detournay, N. Iqbal and E. Perlmutter, *Holographic entanglement entropy and gravitational anomalies*, *JHEP* **07** (2014) 114 [[arXiv:1405.2792](#)] [[INSPIRE](#)].
- [36] J.-R. Sun, *Note on Chern-Simons term correction to holographic entanglement entropy*, *JHEP* **05** (2009) 061 [[arXiv:0810.0967](#)] [[INSPIRE](#)].
- [37] A. Schwimmer and S. Theisen, *Entanglement entropy, trace anomalies and holography*, *Nucl. Phys. B* **801** (2008) 1 [[arXiv:0802.1017](#)] [[INSPIRE](#)].
- [38] T. Azeyanagi, R. Loganayagam and G.S. Ng, *Holographic entanglement for Chern-Simons terms*, [arXiv:1507.02298](#) [[INSPIRE](#)].
- [39] L.-Y. Hung, R.C. Myers and M. Smolkin, *On holographic entanglement entropy and higher curvature gravity*, *JHEP* **04** (2011) 025 [[arXiv:1101.5813](#)] [[INSPIRE](#)].
- [40] J. de Boer, M. Kulaxizi and A. Parnachev, *Holographic entanglement entropy in Lovelock gravities*, *JHEP* **07** (2011) 109 [[arXiv:1101.5781](#)] [[INSPIRE](#)].
- [41] W.-Z. Guo, S. He and J. Tao, *Note on entanglement temperature for low thermal excited states in higher derivative gravity*, *JHEP* **08** (2013) 050 [[arXiv:1305.2682](#)] [[INSPIRE](#)].