

Title	ONE-STEP EXTENSIONS OF SUBNORMAL 2-VARIABLE WEIGHTED SHIFTS : BASED ON JOINT WORK WITH R. CURTO AND J. YOON. (Theory of operator means and related topics)
Author(s)	LEE, SANG HOON
Citation	数理解析研究所講究録 (2015), 1935: 25-30
Issue Date	2015-04
URL	http://hdl.handle.net/2433/223691
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

ONE-STEP EXTENSIONS OF SUBNORMAL 2-VARIABLE WEIGHTED SHIFTS

SANG HOON LEE
 (BASED ON JOINT WORK WITH R. CURTO AND J. YOON.)

1. INTRODUCTION

Consider the following reconstruction-of-the-measure problem:

Problem 1.1 (A). Given two probability measures μ_1 and μ_2 on \mathbb{R}_+^2 , find necessary and sufficient conditions for the existence of a probability measure μ on \mathbb{R}_+^2 such that

$$(1.1) \quad \frac{s \, d\mu(s, t)}{\int s \, d\mu(s, t)} = d\mu_1(s, t) \quad \text{and} \quad \frac{t \, d\mu(s, t)}{\int t \, d\mu(s, t)} = d\mu_2(s, t).$$

Note that (1.1) implies that $t d\mu_1(s, t) = \lambda s d\mu_2(s, t)$ for some $\lambda > 0$.

In this talk, we solve this interpolation problem using techniques from multivariable operator theory, namely the theory of 2-variable weighted shifts.

Definition 1.2. $T \in \mathcal{B}(\mathcal{H})$: normal if $T^*T = TT^*$,

subnormal if $T = N|_{\mathcal{H}}$, where N normal and $N(\mathcal{H}) \subseteq \mathcal{H}$,

hyponormal if $[T^*, T] := T^*T - TT^* \geq 0$.

Definition 1.3. $\mathbf{T} \equiv (T_1, \dots, T_n)$: hyponormal if

$$[\mathbf{T}^*, \mathbf{T}] := \left([T_j^*, T_i]_{i,j=1}^n \right) \geq 0.$$

$$= \begin{pmatrix} [T_1^*, T_1] & [T_2^*, T_1] & \cdots & [T_n^*, T_1] \\ [T_1^*, T_2] & [T_2^*, T_2] & \cdots & [T_n^*, T_2] \\ \vdots & \vdots & \ddots & \vdots \\ [T_1^*, T_n] & [T_2^*, T_n] & \cdots & [T_n^*, T_n] \end{pmatrix} \geq 0.$$

Definition 1.4. The n -tuple $\mathbf{T} \equiv (T_1, T_2, \dots, T_n)$ is said to be normal if \mathbf{T} is commuting and each T_i is normal, and \mathbf{T} is subnormal if \mathbf{T} is the restriction of a normal n -tuple to a common invariant subspace.

- Clearly, normal \implies subnormal \implies hyponormal.
- Normality(sub-, hypo-) of \mathbf{T} is not affected by permuting of the operators T_i .
- If (T_1, \dots, T_n) is normal(sub-, hypo-) then so is $(k_1 T_1, \dots, k_n T_n)$ for any $k_1, \dots, k_n \in \mathbb{C}$.
- If (T_1, \dots, T_n) is normal(sub-, hypo-) then any operator in $LS\{T_1, \dots, T_n\}$ is normal(sub-, hypo-).

Problem 1.5 (Lifting Problem for Commuting Subnormals). Find necessary and sufficient conditions for a pair of subnormal operators on a Hilbert space to admit commuting normal extensions i.e., to be subnormal.

Necessary Conditions: Commuting

Sufficient Conditions: Doubly commuting, either T_1 or T_2 is normal, either T_1 or T_2 is isometry,...

2000 Mathematics Subject Classification. Primary 47B20, 47B37, 47A13, 28A50; Secondary 44A60, 47A20.
 Key words and phrases. one-step extension, 2-variable weighted shifts, subnormal pair, Berger measure.

Besides their relevance for the construction of examples and counterexamples in Hilbert space operator theory, weighted shifts can also be used to detect properties such as subnormality, via the Lambert-Lubin Criterion([15, 17]):

Theorem 1.6 ([15]). *If $T \in \mathcal{B}(\mathcal{H})$ is one-one, then T is subnormal if and only if T_x is subnormal for all $x(\neq 0) \in \mathcal{H}$ where T_x is the weighted shift with weights $\{\frac{\|T^{n+1}x\|}{\|T^n x\|}\}_{n=0}^\infty$.*

Theorem 1.7 ([17]). *If $T_1, T_2 \in \mathcal{B}(\mathcal{H})$ are commuting and one-one, then $\mathbf{T} \equiv (T_1, T_2)$ is subnormal if and only if \mathbf{T}_x is subnormal for all $x(\neq 0) \in \mathcal{H}$ where \mathbf{T}_x is the 2-variable weighted shift with weights*

$$\alpha_{m,n} := \frac{\|T_1^{m+1}T_2^n x\|}{\|T_1^m T_2^n x\|} \text{ and } \beta_{m,n} := \frac{\|T_1^m T_2^{n+1} x\|}{\|T_1^m T_2^n x\|}.$$

Thus, to study the subnormality of commuting pairs, we focus on weighted shifts in the sequel.

Example 1.8 (1-variable weighted shift). For a bounded sequence $a \equiv \{a_n\}_{n=0}^\infty$ of positive real numbers (called *weights*), let $W_a : \ell^2(\mathbb{Z}_+) \rightarrow \ell^2(\mathbb{Z}_+)$ be the associated unilateral weighted shift, defined by $W_a e_n := a_n e_{n+1}$ (all $n \geq 0$), where $\{e_n\}_{n=0}^\infty$ is the canonical orthonormal basis in $\ell^2(\mathbb{Z}_+)$.

For a weighted shift W_a , the moments of a are given as

$$\gamma_k \equiv \gamma_k(a) := \begin{cases} 1, & \text{if } k = 0 \\ a_0^2 \cdots a_{k-1}^2, & \text{if } k \geq 1. \end{cases}$$

It is easy to see that W_a is never normal, and that it is hyponormal if and only if $a_0 \leq a_1 \leq \cdots$.

We shall often write $\text{shift}(a_0, a_1, \dots)$ to denote the weighted shift W_a .

Example 1.9 (2-variable weighted shift). For $\alpha \equiv \{\alpha_{\mathbf{k}}\}, \beta \equiv \{\beta_{\mathbf{k}}\} \in \ell^\infty(\mathbb{Z}_+^2)$, we define the 2-variable weighted shift $W_{(\alpha, \beta)} \equiv (W_\alpha, W_\beta)$ on $\ell^2(\mathbb{Z}_+^2)$ by

$$W_\alpha e_{\mathbf{k}} := \alpha_{\mathbf{k}} e_{\mathbf{k} + \varepsilon_1} \text{ and } W_\beta e_{\mathbf{k}} := \beta_{\mathbf{k}} e_{\mathbf{k} + \varepsilon_2},$$

where $\varepsilon_1 := (1, 0), \varepsilon_2 := (0, 1)$ and $\{e_{\mathbf{k}} : \mathbf{k} \in \mathbb{Z}_+^2\}$ is the canonical orthonormal basis of $\ell^2(\mathbb{Z}_+^2)$.

In an entirely similar way one can define multivariable weighted shifts.





FIGURE 1. Weight diagram for 2-variable weighted shift $W_{(\alpha, \beta)}$

Clearly,

$$(1.2) \quad W_\alpha W_\beta = W_\beta W_\alpha \iff \alpha_{\mathbf{k}} \beta_{\mathbf{k} + \varepsilon_1} = \beta_{\mathbf{k}} \alpha_{\mathbf{k} + \varepsilon_2} \quad (\forall \mathbf{k} \in \mathbb{Z}_+^2).$$



In the sequel, we assume that all 2-variable weighted shifts $W_{(\alpha, \beta)}$ are commuting, i.e., it satisfies condition (1.2).

Given $\mathbf{k} \in \mathbb{Z}_+^2$, the moments $\gamma_{\mathbf{k}} \equiv \gamma_{\mathbf{k}}(\alpha, \beta)$ of (α, β) of order \mathbf{k} is defined by

$$\begin{cases} 1 & \text{if } \mathbf{k} \equiv (k_1, k_2) = (0, 0) \\ \alpha_{(0,0)}^2 \cdots \alpha_{(k_1-1,0)}^2 & \text{if } k_1 \geq 1 \text{ and } k_2 = 0 \\ \beta_{(0,0)}^2 \cdots \beta_{(0,k_2-1)}^2 & \text{if } k_1 = 0 \text{ and } k_2 \geq 1 \\ \alpha_{(0,0)}^2 \cdots \alpha_{(k_1-1,0)}^2 \cdot \beta_{(k_1,0)}^2 \cdots \beta_{(k_1,k_2-1)}^2 & \text{if } k_1 \geq 1 \text{ and } k_2 \geq 1. \end{cases}$$

We remark that, due to the commutativity condition (1.2), $\gamma_{\mathbf{k}}$ can be computed using any nondecreasing path from $(0, 0)$ to \mathbf{k} .

Question 1.10. Which weighted shifts are subnormal?

Theorem 1.11 (Berger's Theorem(1-variable)). W_a is subnormal if and only if there exists a probability measure ξ (called the Berger measure of W_a) supported in $[0, \|W_a\|^2]$ such that $\gamma_{\mathbf{k}}(a) = \int s^{\mathbf{k}} d\xi(s)$ ($\mathbf{k} \geq 0$).

Theorem 1.12 (Berger's Theorem(2-variable)([14])). $W_{(\alpha, \beta)}$ is subnormal if and only if there is a probability measure μ (called the Berger measure of $W_{(\alpha, \beta)}$) supported in the 2-dimensional rectangle $R = [0, \|W_\alpha\|^2] \times [0, \|W_\beta\|^2]$ such that

$$\gamma_{\mathbf{k}}(\alpha, \beta) = \int_R s^{k_1} t^{k_2} d\mu(s, t), \forall \mathbf{k} \equiv (k_1, k_2) \in \mathbb{Z}_+^2.$$

2. AUXILIARY LEMMAS

For a 2-variable weighted shift $W_{(\alpha, \beta)}$, we let \mathcal{M} (resp. \mathcal{N}) be the invariant subspace of $\ell^2(\mathbb{Z}_+^2)$ spanned by the canonical orthonormal basis vectors associated to indices $\mathbf{k} = (k_1, k_2)$ with $k_1 \geq 0$ and $k_2 \geq 1$ (resp. $k_1 \geq 1$ and $k_2 \geq 0$).

We consider the following problem:

Problem 2.1 (B). Assume that $W_{(\alpha, \beta)}|_{\mathcal{M}}$ and $W_{(\alpha, \beta)}|_{\mathcal{N}}$ are subnormal with the Berger measures $\mu_{\mathcal{M}}$ and $\mu_{\mathcal{N}}$, respectively. Find necessary and sufficient conditions on $\mu_{\mathcal{M}}$, $\mu_{\mathcal{N}}$ and β_{00} for the subnormality of $W_{(\alpha, \beta)}$.

Note that Problem (B) is equivalent to Problem (A).


If W_a is subnormal with Berger measure ξ , and if we let for fixed $i \geq 1$,

$$\mathcal{L}_i := \bigvee \{e_n : n \geq i\}$$

then the Berger measure ξ_i of $W_a|_{\mathcal{L}_i}$ is $\frac{s^i}{\gamma_i} d\xi(s)$.

Lemma 2.2 (1-variable subnormal backward extension ([5])). If $W_a|_{\mathcal{L}_1}$ is subnormal with Berger measure ξ_1 then W_a is subnormal if and only if

$$(2.1) \quad \frac{1}{s} \in L^1(\xi_1) \quad \text{and} \quad a_0^2 \left\| \frac{1}{s} \right\|_{L^1(\xi_1)} \leq 1.$$

FIGURE 2. 2-variable weighted shift $W_{(\alpha,\beta)}$ in Problem (B)

In this case, the Berger measure ξ of W_a is $d\xi(s) = \frac{a_0^2}{s} d\xi_1(s) + (1 - a_0^2) \left\| \frac{1}{s} \right\|_{L^1(\xi_1)} d\delta_0(s)$.

- Let μ and ν be two positive measures on \mathbb{R}_+ . We say that $\mu \leq \nu$ if $\mu(E) \leq \nu(E)$ for each Borel subset $E \subseteq \mathbb{R}_+$.
- Let μ be a probability measure on $\mathbb{R}_+ \times \mathbb{R}_+$ and assume that $\frac{1}{t} \in L^1(\mu)$. The *extremal measure* μ_{ext} (which is also a probability measure) on $\mathbb{R}_+ \times \mathbb{R}_+$ is given by

$$d\mu_{ext}(s, t) := (1 - \delta_0(t)) \frac{1}{t \left\| \frac{1}{t} \right\|_{L^1(\mu)}} d\mu(s, t).$$

Here δ_0 denotes Dirac measure at 0.

- Given a measure μ on $X \times Y$, the *marginal measure* μ^X is a measure on X given by

$$\mu^X := \mu \circ \pi_X^{-1},$$

where $\pi_X : X \times Y \rightarrow X$ is the canonical projection onto X .

Lemma 2.3 (2-variable subnormal backward extension ([10])). *Assume that $W_{(\alpha,\beta)}|_{\mathcal{M}}$ is subnormal with the Berger measure $\mu_{\mathcal{M}}$ and that shift $(\alpha_{00}, \alpha_{10}, \dots)$ is subnormal with Berger measure ξ . Then $W_{(\alpha,\beta)}$ is subnormal if and only if the following conditions hold:*

- $\frac{1}{t} \in L^1(\mu_{\mathcal{M}})$;
- $\beta_{00}^2 \leq \left(\left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}})} \right)^{-1}$;
- $\beta_{00}^2 \left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}})} (\mu_{\mathcal{M}})_{ext}^X \leq \xi$.


Moreover, if $\beta_{00}^2 \left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}_1})} = 1$, then $(\mu_{\mathcal{M}})_{ext}^X = \xi$.

Lemma 2.4 ([11]). *Let μ be the Berger measure of a subnormal 2-variable weighted shift $W_{(\alpha,\beta)}$, and let ξ be the Berger measure of the associated 0-th horizontal 1-variable shift $(\alpha_{00}, \alpha_{10}, \dots)$. Then $\xi = \mu^X$.*

3. MAIN RESULT AND APPLICATION

We provides a concrete solution of Problem (B) in terms of $\mu_{\mathcal{M}}$, $\mu_{\mathcal{N}}$ and β_{00} .

Theorem 3.1 (Main Theorem). *Assume that $W_{(\alpha,\beta)}|_{\mathcal{M}}$ and $W_{(\alpha,\beta)}|_{\mathcal{N}}$ are subnormal with associated Berger measures $\mu_{\mathcal{M}}$ and $\mu_{\mathcal{N}}$, respectively, and let $c := \frac{\int s d\mu_{\mathcal{M}}}{\int t d\mu_{\mathcal{N}}} \equiv \frac{\alpha_{01}^2}{\beta_{10}^2}$. Then $W_{(\alpha,\beta)}$ is subnormal if and only if the following*

FIGURE 3. 2-variable weighted shift $W_{(\alpha,\beta)}$ in Lemma 2.4

conditions hold:

(i) $\frac{1}{t} \in L^1(\mu_{\mathcal{M}})$ and $\frac{1}{s} \in L^1(\mu_{\mathcal{N}})$;


(ii) $\beta_{00}^2 \left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}})} \leq 1$;

(iii) $\beta_{00}^2 \left\{ \left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}})} (\mu_{\mathcal{M}})_{ext}^X + c \left\| \frac{1}{s} \right\|_{L^1(\mu_{\mathcal{N}})} \delta_0 - \frac{c}{s} (\mu_{\mathcal{N}})^X \right\} \leq \delta_0$.

For a measure μ with $\frac{1}{s} \in L^1(\mu)$, we write $d\tilde{\mu}(s) := \frac{1}{s \left\| \frac{1}{s} \right\|_{L^1(\mu)}} d\mu(s)$.

Lemma 3.2 ([8]). *Let $W_{(\alpha,\beta)}$ be the 2-variable weighted shift given in Figure 4. Then $W_{(\alpha,\beta)}|_{\mathcal{M}}$ is subnormal if and only if $\psi := \eta_1 - \alpha_{01}^2 \left\| \frac{1}{s} \right\|_{L^1(\sigma)} \tau$ is a positive measure. In this case, the Berger measure of $W_{(\alpha,\beta)}|_{\mathcal{M}}$ is*

$$\mu_{\mathcal{M}} = \alpha_{01}^2 \left\| \frac{1}{s} \right\|_{L^1(\sigma)} \tilde{\sigma} \times \tau + \delta_0 \times \psi.$$

FIGURE 4. 2-variable weighted shift $W_{(\alpha,\beta)}$ in Lemma 3.2

As a special case of Main Theorem, we have:

Theorem 3.3 (The case when $W_{(\alpha,\beta)}$ has a core of tensor form). *Assume that $W_{(\alpha,\beta)}|_{\mathcal{M}}$ and $W_{(\alpha,\beta)}|_{\mathcal{N}}$ are subnormal with associated Berger measures $\mu_{\mathcal{M}}$ and $\mu_{\mathcal{N}}$, respectively, and let $\rho := \mu_{\mathcal{M}}^X$. Also assume that $\mu_{\mathcal{M} \cap \mathcal{N}} = \sigma \times \tau$ for some 1-variable probability measures σ and τ . Then $\rho = \mu_{\mathcal{M}}^X = (\mu_{\mathcal{M}})_{ext}^X$, and hence $W_{(\alpha,\beta)}$ is subnormal if and only if the following conditions hold:*

(i) $\frac{1}{t} \in L^1(\mu_{\mathcal{M}})$ and $\frac{1}{s} \in L^1(\mu_{\mathcal{N}})$;

- (ii) $\beta_{00}^2 \left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}})} \leq 1$;
 (iii) $\left(\beta_{00}^2 \left\| \frac{1}{t} \right\|_{L^1(\mu_{\mathcal{M}})} \right) \rho \leq \xi$, where ξ is the Berger measure of shift $(\alpha_{00}, \alpha_{10}, \dots)$.




FIGURE 5. 2-variable weighted shift $W_{(\alpha,\beta)}$ in Theorem 3.3

REFERENCES

- [1] A. Athavale, On joint hyponormality of operators, *Proc. Amer. Math. Soc.* 103(1988), 417-423.
- [2] J. Bram, Subnormal operators, *Duke Math. J.* 22(1955), 75-94.
- [3] J. Conway, *The Theory of Subnormal Operators*, Mathematical Surveys and Monographs, vol. 36, Amer. Math. Soc., Providence, 1991.
- [4] R. Curto, Joint hyponormality: A bridge between hyponormality and subnormality, *Proc. Symposia Pure Math.* 51(1990), 69-91.
- [5] R. Curto, Quadratically hyponormal weighted shifts, *Integral Equations Operator Theory* 13(1990), 49-66.
- [6] R. Curto, S.H. Lee and J. Yoon, k -hyponormality of multivariable weighted shifts, *J. Funct. Anal.*, 229(2005), 462-480.
- [7] R. Curto, S.H. Lee and J. Yoon, Hyponormality and subnormality for powers of commuting pairs of subnormal operators, *J. Funct. Anal.* 245(2007), 390-412.
- [8] R. Curto, S.H. Lee and J. Yoon, Reconstruction of the Berger measure when the core is of tensor form, *Actas del XVI Coloquio Latinoamericano de Álgebra, Bibl. Rev. Mat. Iberoamericana*, (2007), 317-331.
- [9] R. Curto, P. Muhly and J. Xia, Hyponormal pairs of commuting operators, *Operator Theory: Adv. Appl.* 35(1988), 1-22.
- [10] R. Curto, J. Yoon, Jointly hyponormal pairs of subnormal operators need not be jointly subnormal, *Trans. Amer. Math. Soc.* 358(2006), 5139-5159.
- [11] R. Curto, J. Yoon, Disintegration-of-measure techniques for commuting multivariable weighted shifts, *Proc. London Math. Soc.*, 92(2006), 321-402.
- [12] Gellar and Wallen, Subnormal weighted shifts and the Halmos-Bram criterion, *Proc. Japan Acad.*, 46(1970), 375-378.
- [13] P.R. Halmos, Ten problems in Hilbert space, *Bull. Amer. Math. Soc.* 76(1970), 887-933.
- [14] N.P. Jewell and A.R. Lubin, Commuting weighted shifts and analytic function theory in several variables, *J. Operator Theory* 1(1979), 207-223.
- [15] A. Lambert, Subnormality and weighted shifts, *J. London Math. Soc.* 14(3)(1976), 476-480.
- [16] A. Lubin, Weighted shifts and product of subnormal operators, *Indiana Univ. Math. J.*, 26(1977), 839-845.
- [17] A.R. Lubin, Weighted shifts and commuting normal extension, *J. Austral Math. Soc. Ser. A* 27(1)(1979), 17-26.
- [18] J. Yoon, Disintegration of measures and contractive 2-variable weighted shifts, *Integral Equations Operator Theory*, 59(2007), 281-298.

DEPARTMENT OF MATHEMATICS, CHUNGNAM NATIONAL UNIVERSITY, DAEJEON, 305-764, REPUBLIC OF KOREA
 E-mail address: s1ee@cnu.ac.kr