

Randomized, Double-Blind, Placebo-Controlled Phase III Study of Tasquinimod in Men With Metastatic Castration-Resistant Prostate Cancer

Cora Sternberg, Andrew Armstrong, Roberto Pili, Siobhan Ng, Robert Huddart, Neeraj Agarwal, Denis Khvorostenko, Olexiy Lyulko, Arija Brize, Nicholas Vogelzang, Rémy Delva, Mihai Harza, Anastasios Thanos, Nicholas James, Patrick Werbrouck, Martin Bögemann, Thomas Hutson, Piotr Milecki, Simon Chowdhury, Enrique Gallardo, Gilberto Schwartzmann, Jean-Christophe Pouget, Frédérique Baton, Thore Nederman, Helen Tuvesson, and Michael Carducci

ABSTRACT

Purpose

Tasquinimod, a novel oral therapy targeting the tumor microenvironment, significantly improved progression-free survival (PFS) in a randomized, placebo-controlled phase II trial in men with metastatic castration-resistant prostate cancer (mCRPC). This phase III study was conducted to confirm the phase II results and to detect an overall survival (OS) benefit.

Patients and Methods

Men with chemotherapy-naïve mCRPC and evidence of bone metastases were assigned (2:1) to receive tasquinimod once per day or placebo until progression or toxicity. The primary end point was radiographic PFS (rPFS; time from random assignment to radiologic progression or death) per Prostate Cancer Working Group 2 criteria and RECIST 1.1. The study had 99.9% power to detect an rPFS hazard ratio (HR) of 0.6 with a two-sided alpha error of .05 and 80% power to detect a target HR of 0.8 for OS, the key secondary end point.

Results

In all, 1,245 patients were randomly assigned to either tasquinimod ($n = 832$) or placebo ($n = 413$) between March 2011 and December 2012 at 241 sites in 37 countries. Baseline characteristics were balanced between groups: median age, 71 years; Karnofsky performance status $\geq 90\%$, 77.3%; and visceral metastases, 21.1%. Estimated median rPFS by central review was 7.0 months (95% CI, 5.8 to 8.2 months) with tasquinimod and 4.4 months (95% CI, 3.5 to 5.5 months) with placebo (HR, 0.64; 95% CI, 0.54 to 0.75; $P < .001$). Median OS was 21.3 months (95% CI, 19.5 to 23.0 months) with tasquinimod and 24.0 months (95% CI, 21.4 to 26.9 months) with placebo (HR, 1.10; 95% CI, 0.94 to 1.28; $P = .25$). Grade ≥ 3 adverse events were more frequent with tasquinimod (42.8% v 33.6%), the most common being anemia, fatigue, and cancer pain.

Conclusion

In chemotherapy-naïve men with mCRPC, tasquinimod significantly improved rPFS compared with placebo. However, no OS benefit was observed.

J Clin Oncol 34:2636-2643. © 2016 by American Society of Clinical Oncology

INTRODUCTION

Treatment options for metastatic castration-resistant prostate cancer (mCRPC) have expanded with the introduction of several new agents that delay disease progression and improve overall survival (OS). These include second-generation androgen-directed therapies, radium-223, sipuleucel-T, and the taxanes cabazitaxel and docetaxel. Despite these advances,

mCRPC remains incurable, and survival benefits typically achieved with newer agents are modest while resistance remains common.¹⁻⁷ New agents with alternative mechanisms of action that further improve survival while minimizing toxicity are needed.

The tumor microenvironment is increasingly recognized as playing a major role in the formation and growth of metastases.⁸ In addition, the host microenvironment has been shown to promote prostate cancer invasion, systemic spread, bone

Author affiliations appear at the end of this article.

Published online ahead of print at www.jco.org on June 13, 2016.

Supported by Active Biotech.

C.S. and A.A. contributed equally to this work as first co-authors.

Presented in part at the European Cancer Congress 2015, Vienna, Austria, September 25-29, 2015.

Authors' disclosures of potential conflicts of interest are found in the article online at www.jco.org. Contributions are found at the end of this article.

Clinical trial information: NCT01234311.

Corresponding author: Michael Carducci, MD, The Johns Hopkins University School of Medicine, 1650 Orleans St, CRB1 1M59, Baltimore, MD 21287-1000; e-mail: carducci@jhmi.edu

© 2016 by American Society of Clinical Oncology

0732-183X/16/3422w-2636w/\$20.00

DOI: 10.1200/JCO.2016.66.9697

colonization, and osteoblastic metastasis.⁹ Drugs that target the tumor microenvironment therefore offer a potentially new approach in the treatment of advanced prostate cancer.¹⁰ Tasquinimod (ABR-215050; Active Biotech, Lund, Sweden) is an oral immunotherapy with demonstrated effects on the tumor microenvironment that counteract tumor growth.^{11,12} One molecular target of tasquinimod is the immunomodulatory protein S100A9, which plays a role in the accumulation and function of innate immune cells, specifically regulatory myeloid cells.¹¹⁻¹³ Targeting regulatory myeloid cells within the tumor microenvironment leads to decreased immune suppression and angiogenesis and prevention of metastasis development. Tasquinimod may also reduce angiogenesis by downregulation of HIF1-controlled genes via interaction with histone deacetylases.¹⁴

In a randomized, placebo-controlled phase II study in men with mCRPC, tasquinimod significantly improved progression-free survival (PFS; median, 7.6 v 3.3 months; hazard ratio [HR], 0.57; $P < .01$).¹⁵ In long-term follow-up, multivariate analysis indicated that the PFS improvement may be associated with improved OS, particularly in patients with bone metastases.¹⁶ The objective of this phase III study was to confirm the benefit of tasquinimod in delaying disease progression and improving OS in men with mCRPC.

PATIENTS AND METHODS

Patients

Eligible patients had histologically confirmed prostate adenocarcinoma with evidence of bone metastases, serum testosterone ≤ 50 ng/dL, disease progression (increasing serum prostate-specific antigen [PSA] as defined by the Prostate Cancer Working Group 2 [PCWG2],¹⁷ progression of soft tissue metastasis, or bone disease progression), and Karnofsky performance status $\geq 70\%$. Concurrent use of luteinizing hormone-releasing hormone agonists or antagonists and bone agents (denosumab or bisphosphonates) was permitted.

No cytotoxic chemotherapy within 2 years or previous anticancer therapy within 4 weeks (2 weeks for sipuleucel-T) of random assignment was allowed. Prior enzalutamide or abiraterone was permitted. Other exclusion criteria included presence of prostate cancer pain requiring opiate analgesics, systemic exposure to ketoconazole, and ongoing corticosteroid treatment equivalent to a prednisolone or prednisone dose of > 10 mg/day.

Study Design

This multinational, randomized, double-blind, placebo-controlled phase III study was conducted at 241 sites in 37 countries (Appendix Table A1, online only). Patients were randomly assigned in a 2:1 ratio to receive tasquinimod or placebo by using an interactive voice response system. Random assignment was stratified by Karnofsky performance status ($\geq 90\%$ v $< 90\%$), presence or absence of visceral disease (all metastatic soft tissue except lymph nodes and local recurrence), and geographic region (North America, Europe, the Middle East, Africa, Asia-Pacific, and Latin America). Tasquinimod or placebo was administered orally at a starting dose of 0.25 mg/day for at least 2 weeks.¹⁸ If tolerability was established, the dose was escalated to 0.5 mg/day for 2 weeks and then to 1 mg/day. Patients unable to tolerate the escalated doses could continue in the study at their maximum tolerated dose. Treatment continued until symptomatic disease progressed so that it required alternative antitumor therapy or until poor tolerability occurred. After the end of treatment, patients continued follow-up with visits every 3 months until death or until 727 patients had reached the survival end point.

The study was approved by the institutional review boards or ethics committees at each participating center and was conducted in accordance with the Declaration of Helsinki. All patients provided written informed consent before study entry.

End Points

The primary end point was radiographic PFS (rPFS), the time from random assignment to radiologic progression or death, whatever the cause. Radiographic progression was defined as soft tissue progression (RECIST 1.1),¹⁹ bone progression detected with confirmatory bone scans (PCWG2),⁷ or radiographically confirmed spinal cord compression or fracture as a result of malignant progression. Soft tissue lesions were evaluated by computed tomography or magnetic resonance imaging scans by the investigator. All scans underwent independent central review, with reviewers blinded to study treatment and investigator assessments.

The key secondary end point was OS, defined as time from random assignment to death. Other prespecified secondary end points included time to radiologic progression, time to symptomatic progression, time to PSA progression, time to initiation of further cytotoxic therapy, time to opiate use, and time to deterioration of quality-of-life (QoL) measure (Functional Assessment of Cancer Therapy-Prostate [FACT-P]). Safety was assessed on the basis of physical examination, vital signs measurements, clinical laboratory analyses, and adverse events (AEs; coded using Medical Dictionary for Regulatory Activities [MedDRA]; graded using Common Terminology Criteria for Adverse Events [CTCAE] version 4.0).

Statistical Analysis

The planned sample size of 1,200 patients (800 in the tasquinimod arm and 400 in the placebo arm) provided 99.9% power at a two-sided significance level of 0.05 to detect an HR of 0.6 for the primary end point of rPFS, corresponding to an increase in median PFS from 3.4 to 5.7 months. The study was also designed to detect an HR of 0.8 for the key secondary end point of OS, corresponding to an increase in median OS from 22 to 27.5 months. Specifying a two-sided significance level of 0.05, the study had 80% power to detect the OS difference after 727 deaths had been observed. The OS end point comparisons incorporated group sequential design involving two interim analyses (at 473 and 582 events) and a final analysis at 727 events using O'Brien-Fleming stopping boundaries²⁰: first interim analysis, $P \leq .0109$; second interim analysis, $P \leq .0212$; and final analysis, $P \leq .0422$. rPFS was analyzed at the first planned interim analysis for OS (after 473 events). If the comparison of rPFS reached statistical significance ($P \leq .05$), the first comparison of OS was performed; however, the results were not reported until the final analysis.

A stratified log-rank test by factors at random assignment was used to compare rPFS, OS, and the time-to-event secondary end points for tasquinimod versus placebo (analysis of PSA doubling time was not stratified). To describe time-to-event variables, Kaplan-Meier curves and life tables by treatment group were generated, and CIs were calculated.²¹ Patients who did not experience an event were censored at the date of their last adequate assessment, previous assessment, last visit, or death, depending on the end point and analysis. Treatment effect was estimated by calculating the HR and its 95% CI from a Cox proportional hazards model stratified by factors at random assignment. For rPFS and OS, Cox proportional hazards models were performed for predefined subgroups and multivariate analyses. In the latter analyses, after testing each prespecified prognostic factor with a univariate analysis, a backward selection approach was used. Treatment was always included in the models.

All efficacy end points were analyzed by planned treatment in the intent-to-treat population (all randomly assigned patients, regardless of whether any study treatment dosing was completed). The safety analysis population comprised all patients who received at least one dose of study treatment. Safety was analyzed according to treatment received. All statistical analyses were performed by using SAS version 9.2 (SAS Institute, Cary, NC).

RESULTS

Patient Disposition

Of 1,645 patients screened, 1,245 were randomly assigned to receive tasquinimod ($n = 832$) or placebo ($n = 413$) between March 29, 2011, and December 7, 2012 (Appendix Table A1). Two patients from each group did not receive treatment after random assignment (Fig 1). Baseline characteristics were generally well balanced between the groups (Table 1). A greater proportion of patients in the tasquinimod group had higher levels of baseline tumor pain (Visual Analog Scale > 4 : 18.6% ν 14.5%). Median time since diagnosis was shorter in the tasquinimod group than in the placebo group (45.7 ν 57.7 months).

At final analysis (cutoff date: February 13, 2015), median follow-up was 30.0 months in the tasquinimod arm and 30.7 months in the placebo arm, and 96.1% of patients had discontinued treatment. The most common reasons for discontinuation (tasquinimod ν placebo) were radiographic progression (23.8% ν 36.5%), symptomatic progression requiring new anti-cancer therapy (21.3% ν 18.8%), and poor tolerability or AEs (17.9% ν 8.8%; Fig 1).

Efficacy

The final analysis of the primary end point of rPFS was performed at the time of the first interim analysis of OS. Radiographic progression by central review, or death, occurred in 396 patients (48%) in the tasquinimod group and in 258 patients (62%) in the placebo group. Estimated median rPFS was 7.0 months (95% CI, 5.8 to 8.2 months) for tasquinimod and 4.4 months (95% CI, 3.5 to 5.5 months) for placebo, corresponding to a 36% reduction in the risk of radiographic progression or death with tasquinimod versus placebo (HR, 0.64; 95% CI, 0.54 to 0.75; $P < .001$; Fig 2A). Similar results were seen in

the assessment by local review: estimated median rPFS was 5.7 months (95% CI, 5.5 to 6.2 months) and 4.1 months (95% CI, 3.1 to 5.1 months), respectively (HR 0.69; 95% CI, 0.60 to 0.80; $P < .001$).

OS results were not significant at either of the two interim analyses and, because no safety concerns were raised, the Data and Safety Monitoring Board recommended continuation of the study according to the protocol. At final analysis of OS, 492 deaths (59.1%) had occurred in the tasquinimod group and 238 deaths (57.6%) had occurred in the placebo group. Tasquinimod did not improve OS compared with placebo (median OS, 21.3 months [95% CI, 19.5 to 23.0 months] with tasquinimod and 24.0 months [95% CI, 21.4 to 26.9 months] with placebo; HR, 1.10; 95% CI, 0.94 to 1.28; $P = .25$; Fig 2B). The rPFS and OS results were consistent when examined across predefined patient subgroups without evidence of significant heterogeneity (Fig 3).

In general, secondary end points that favored tasquinimod over placebo included the radiographic- and PSA-based outcomes (Table 2 and Appendix Table A2, online only). In contrast, symptomatically assessed end points, such as time to symptomatic progression, time to opiate use, and deterioration in QoL, favored placebo. Time to initiation of salvage therapy was longer with tasquinimod than with placebo (11.4 ν 8.1 months; $P = .001$), as was time to initiation of further cytotoxic therapy (25.8 ν 16.0 months; $P = .021$).

One quarter of patients (315 [25.3%] of 1,245) had undergone orchiectomy, and most patients (1,178 [94.6%] of 1,245) had received hormonal therapy pre-enrollment (mostly bicalutamide, flutamide, and luteinizing hormone-releasing hormone analogs). In contrast, only a few patients had received prior abiraterone (five patients [0.6%] in the tasquinimod group ν seven patients [1.7%] in the placebo group) or enzalutamide (zero ν one [0.2%]). These treatments were more commonly




Fig 1. CONSORT diagram. TASQ, tasquinimod.

Table 1. Baseline Demographic and Disease Characteristics				
Characteristic	Tasquinimod (n = 832)		Placebo (n = 413)	
	No.	%	No.	%
Median age, years (range)	71.0 (43-92)		71.0 (48-92)	
Age group (years)				
≤ 65	214	25.7	106	25.7
66-75	371	44.6	186	45.0
76-80	144	17.3	64	15.5
> 80	103	12.4	57	13.8
Race*				
White	729	87.6	359	86.9
Black	20	2.4	8	1.9
Asian	46	5.5	27	6.5
Other	37	4.4	18	4.4
Ethnicity				
Hispanic/Latino	97	11.7	42	10.2
Non-Hispanic/Latino	735	88.3	371	89.8
Median time since diagnosis, months (range)	45.7 (0.1-299.6)		57.7 (0.3-319.9)	
Karnofsky performance status†				
< 90%	187	22.5	95	23.0
≥ 90%	645	77.5	318	77.0
Geographic region of enrollment‡				
North America	143	17.2	72	17.4
Europe/Middle East/Africa	505	60.7	254	61.5
Asia-Pacific	94	11.3	46	11.1
Latin America	90	10.8	41	9.9
Tumor pain (VAS)‡				
0	371	44.6	195	47.2
1-3	286	34.4	157	38.0
4-10	155	18.6	60	14.5
Median PSA, μg/L (range)	54.3 (0.6-8,710.7)		50.1 (0.2-5,679.5)	
Gleason score of 8 to 10 at diagnosis	398	47.8	190	46.0
Visceral disease present†	176	21.2	87	21.1
Location of metastases				
Visceral§	161	19.4	76	18.4
Bone	824	99.0	409	99.0
Node	297	35.7	179	43.3
No. of bone metastases				
< 10	377	45.3	194	47.0
≥ 10	447	53.7	215	52.1
Previous second-generation hormonal therapy¶	65	7.8	48	11.6

Abbreviations: PSA, prostate-specific antigen; VAS, Visual Analog Scale.

*Data missing for one patient in the placebo group.

†According to interactive voice response system data, except for Europe, Middle East, and Asia-Pacific subcategories for geographic region.

‡Data missing for 20 patients in the tasquinimod group and one patient in the placebo group.

§According to electronic case report form data. Indicated location does not exclude other sites.

¶Abiraterone, enzalutamide, ketoconazole, or any other second-generation hormonal treatment.

available during the follow-up period after withdrawal from study treatment and were used more in the placebo group (abiraterone, 209 [25%] *v* 127 [31%]; enzalutamide, 66 [8%] *v* 48 [12%]). More than one third of patients received docetaxel after the study (281 [34%] *v* 166 [40%]).




Fig 2. Kaplan-Meier analysis of (A) radiologic progression-free survival (rPFS; central review) and (B) overall survival (OS). HR, hazard ratio; TASQ, tasquinimod.

Drug Exposure and Safety

Overall median treatment duration was 137 days (range, 1 to 1,377 days) for tasquinimod and 133 days (range, 8 to 1,179 days) for placebo, and most patients (82% and 92%, respectively) escalated to the maximum dose of 1 mg/day.

The proportion of patients with at least one dose reduction from maximum dose was higher in the tasquinimod group than in the placebo group (17.5% *v* 5.6% for the 1 mg/day dose and 1.4% *v* 0% for the 0.5 mg/day dose). The majority of patients in both treatment groups experienced at least one treatment-emergent AE (Table 3). A greater proportion of patients in the tasquinimod group discontinued treatment because of AEs (17.7% *v* 10.2%), mainly as a result of decreased appetite, fatigue, asthenia, or nausea.




Fig 3. Radiologic progression-free survival (rPFS) and overall survival (OS) outcomes in patient subgroups. *Number of patients included in analyses of rPFS/OS. ALP, alkaline phosphatase; Hb, hemoglobin; ITT, intent to treat; LDH, lactate dehydrogenase; PSA, prostate-specific antigen; TASQ, tasquinimod; VAS, Visual Analog Scale.

The most common reported AEs were GI disorders (60.2% for the tasquinimod group v 47.9% for the placebo group), general disorders and administration site conditions (55.1% v 39.9%), and musculoskeletal and connective tissue disorders (48.2% v 36.7%). The most frequently reported AEs are summarized in Table 3.

A total of 229 patients (27.6%) in the tasquinimod group and 97 patients (23.6%) in the placebo group experienced at least one serious AE, the most common being renal and urinary disorders (7.3% v 7.3%), infections and infestations (5.1% v 4.1%), and blood and lymphatic system disorders (4.3% v 4.1%).

Table 2. Secondary Efficacy End Points

Progression	Tasquinimod (n = 832)		Placebo (n = 413)		HR	95% CI	P
	Median (months)	95% CI	Median (months)	95% CI			
Radiologic progression							
Local	8.0	5.8 to 8.3	4.6	3.2 to 5.5	0.683	0.591 to 0.789	< .001
Central	8.4	8.1 to 9.2	5.5	4.5 to 5.6	0.628	0.534 to 0.739	< .001
Soft tissue progression (RECIST 1.1)							
Local	16.6	13.6 to 19.4	8.3	5.9 to 10.9	0.586	0.483 to 0.711	< .001
Central	16.6	14.6 to 20.5	11.1	8.2 to 14.0	0.621	0.504 to 0.765	< .001
Symptomatic progression*	9.5	7.8 to 11.1	11.9	8.9 to 14.1	1.171	1.014 to 1.353	.031
Initiation of salvage therapy†	11.4	9.1 to 13.1	8.1	6.7 to 9.7	0.778	0.667 to 0.907	.001
Initiation of further cytotoxic therapy	25.8	22.1 to 35.9	16.0	13.6 to 23.2	0.809	0.675 to 0.969	.021
Opiate use for cancer pain	29.5	25.1 to NR	35.9	29.4 to NR	1.328	1.060 to 1.664	.013
FACT-P deterioration (criterion 1)‡	3.0	2.9 to 3.3	5.8	5.6 to 6.5	1.447	1.265 to 1.655	< .001
PSA progression	2.9	2.8 to 2.9	2.8	2.8 to 2.8	0.826	0.723 to 0.945	.003

NOTE. Time to skeletal-related events and time to symptomatic progression as a result of skeletal-related events could not be calculated because of the low number of events.

Abbreviations: FACT-P, Functional Assessment of Cancer Therapy-Prostate; HR, hazard ratio; NR, not reached; PSA, prostate-specific antigen.

*Including death as a result of prostate cancer.

†Including radionuclide, chemotherapy, or radiation therapy.

‡Deterioration event was classified as the first of (1) death as a result of prostate cancer, (2) significant and meaningful decline in FACT-P total score, or (3) disease progression, defined as radiologic progression and a missing FACT-P at the same scheduled visit.

Table 3. Most Common AEs Occurring in at Least 5% of Patients in Either Treatment Group

AE	Tasquinimod (n = 830)				Placebo (n = 411)			
	All Grades		Grades 3 to 5		All Grades		Grades 3 to 5	
	No.	%	No.	%	No.	%	No.	%
All AEs	791	95.3	355	42.8	381	92.7	138	33.6
Cancer pain	264	31.8	27	3.3	129	31.4	10	2.4
Decreased appetite	250	30.1	15	1.8	67	16.3	4	1.0
Nausea	222	26.7	7	0.8	89	21.7	3	0.7
Fatigue	217	26.1	28	3.4	72	17.5	9	2.2
Constipation	194	23.4	8	1.0	67	16.3	2	0.5
Anemia	179	21.6	69	8.3	67	16.3	31	7.5
Asthenia	140	16.9	23	2.8	51	12.4	8	1.9
Decreased weight	125	15.1	15	1.8	35	8.5	3	0.7
Back pain	105	12.7	10	1.2	38	9.2	1	0.2
Pain in extremity	104	12.5	10	1.2	31	7.5	1	0.2
Arthralgia	101	12.2	8	1.0	52	12.7	0	
Diarrhea	94	11.3	3	0.4	42	10.2	3	0.7
Insomnia	87	10.5	2	0.2	30	7.3	0	
Vomiting	87	10.5	3	0.4	28	6.8	3	0.7
Peripheral edema	85	10.2	3	0.4	28	6.8	1	0.2

Abbreviation: AE, adverse event.

The incidence of vascular disorders was similar for the tasquinimod and placebo groups (12.4% *v* 13.1%), as was the incidence of deep vein thrombosis (0.7% *v* 1.5%). Cardiac disorders were more frequent with tasquinimod (all grades, 10% *v* 6.8%; grades 3 to 5, 3.4% *v* 1.6%; serious AEs, 3.9% *v* 1.9%). The frequencies of specific cardiac events for tasquinimod and placebo groups, respectively, were atrial fibrillation (2.8% *v* 0.7%), angina pectoris (1.2% *v* 0.7%), cardiac failure (1.2% *v* 0.2%), pericardial effusion (0.8% *v* 0%), pericarditis (0.4% *v* 0%), coronary artery disease (0.4% *v* 0%), and myocardial infarction (0.5% *v* 0.2%). The incidence of death as a result of AEs was similar between the groups: 27 patients (3.3%) in the tasquinimod group and 15 patients (3.6%) in the placebo group. There were four (0.5%) cardiac AE-related deaths in the tasquinimod group and one (0.2%) in the placebo group.

DISCUSSION

Tasquinimod was shown in a randomized phase II study to improve PFS in patients with mCRPC, and it was further indicated that this effect might be associated with an OS benefit.^{15,16} The primary objective of this phase III study was to confirm the phase II findings, and therefore a similar design was used with rPFS as the primary end point. However, the study was designed with sufficient statistical power to detect a potential OS benefit, and OS was the main secondary end point. The results showed that rPFS was significantly delayed by tasquinimod (36% reduced risk of radiographic progression or death *v* placebo, by central review; HR, 0.64), thereby confirming the phase II findings. There was good agreement between independent radiologists and local investigator assessment, suggesting that rPFS can be reliably ascertained, and

recent data suggest that delays in rPFS may be associated with prolonged survival.²²

However, the significant rPFS benefit with tasquinimod did not translate into improved survival over time. Subgroup analyses demonstrated consistent results for rPFS and OS and did not highlight any clear heterogeneity for an OS benefit among any of the subgroups. Tasquinimod seemed to provide clinical benefit over placebo with respect to a number of other objective radiology-based measures as well as for time to PSA progression. Time to initiation of further cytotoxic therapy was prolonged by 9.8 months likely because of the delayed progression with tasquinimod treatment. However, this was not the case for more subjective outcomes such as time to opiate use for cancer pain, time to tumor-related pain progression, and time to QoL deterioration, all of which were better in the placebo group. The most common AEs over-represented in the tasquinimod group included the types of events that are also commonly seen as signs of cancer progression and general health deterioration and thus may have contributed to the unfavorable outcome of symptomatically assessed end points.

Assessing clinical benefit in mCRPC is challenging, given the heterogeneous nature of the disease and differential effects of subsequent therapy on traditional end points, such as OS and postprogression time-to-event end points.²³ PCWG2 guidelines on defining disease progression⁷ have been adopted as the standard primary efficacy measure in most recent clinical trials in mCRPC, and there is widespread interest in the use of PCWG2-defined rPFS as a surrogate end point of survival benefit. A recent analysis of the phase III COU-AA-302 (Abiraterone Acetate in Asymptomatic or Mildly Symptomatic Patients With Metastatic Castration-Resistant Prostate Cancer) trial demonstrated a significant correlation between rPFS and OS.²² However, the lack of correlation between rPFS and OS in this study and in other phase III studies in mCRPC²⁴⁻²⁶ illustrates that significant improvements in rPFS may not always translate into longer-term survival benefit.

Among several possible explanations for the lack of OS benefit in this study, one contributory factor may be the availability of more effective salvage therapies that prolong OS treatment after the study,²⁷ many of which were not widely available at the time of the phase II study. The current availability of such agents (eg, abiraterone and enzalutamide) may have had an impact on the course of disease because patients in the placebo group gained access before those in the tasquinimod group on account of their earlier withdrawal from study treatment. Indeed, post-treatment use of abiraterone and enzalutamide was more common among patients in the placebo group. Furthermore, baseline characteristics suggest a more aggressive cancer population in the tasquinimod arm as indicated by an imbalance in median time since diagnosis and baseline Visual Analog Scale score for tumor-related pain. It may also be that the survival results were influenced by a combination of the relatively modest effect on rPFS and other confounding factors, suggesting that tasquinimod may not have sufficient efficacy as a single agent to have an impact on long-term OS.

Further study of predictive biomarkers of tasquinimod efficacy may be warranted to determine whether certain subgroups will derive an OS advantage. Data from the phase II trial suggested that men with low baseline thrombospondin-1 levels derived the greatest benefit from tasquinimod.¹⁶ Because tasquinimod is

known to increase this antiangiogenic marker in preclinical tumor models,²⁸ there may be a mechanistic basis for further examination of predictive biomarkers identified in this study. Preclinical evidence also suggests that tasquinimod has immunomodulatory activity, shown as an inhibitory effect on myeloid-derived suppressive cells and M2-polarized tumor-associated macrophages.¹³ Identification of a potential immunologic biomarker will help with patient selection and determination of the most rational combination strategy for developing S100A9 inhibitors.

The tolerability of tasquinimod was good overall, and the vast majority of patients were able to escalate to the maximum 1-mg/day dose according to the predefined schedule. Dose interruptions or reductions were infrequent, and the overall safety profile was consistent with that observed in the phase II study. Tasquinimod was associated with a higher rate of withdrawals as a result of AEs. GI and musculoskeletal disorders occurred at a slightly higher frequency with tasquinimod, as seen in the phase II study. The overall incidence of cardiovascular events was low but, as observed previously,^{15,16} was slightly higher with tasquinimod. This higher rate of cardiovascular events may have contributed to the lack of survival benefit due to early drug discontinuation. However, treatment-related deaths were not increased with tasquinimod, suggesting lack of efficacy rather than toxicity as the main contributing factor.

In conclusion, this phase III study confirmed that tasquinimod improved rPFS in patients with mCRPC compared with placebo. This benefit did not translate into an improvement in OS. The tolerability profile of tasquinimod was consistent with that in previous studies. On the basis of the lack of OS benefit observed in

this study, further clinical development of tasquinimod in this patient population was not pursued.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at www.jco.org.

AUTHOR CONTRIBUTIONS

Conception and design: Cora Sternberg, Andrew Armstrong, Roberto Pili, Thore Nederman, Helen Tuvesson, Michael Carducci

Provision of study materials or patients: Andrew Armstrong, Siobhan Ng, Robert Huddart, Neeraj Agarwal, Rémy Delva, Nicholas James, Thomas Hutson, Enrique Gallardo, Thore Nederman, Helen Tuvesson, Michael Carducci

Collection and assembly of data: Cora Sternberg, Andrew Armstrong, Roberto Pili, Robert Huddart, Neeraj Agarwal, Denis Khvorostenko, Olexiy Lyulko, Arija Brize, Nicholas Vogelzang, Rémy Delva, Mihai Harza, Anastasios Thanos, Nicholas James, Patrick Werbrouck, Martin Bögemann, Thomas Hutson, Piotr Milecki, Enrique Gallardo, Gilberto Schwartzmann, Michael Carducci

Data analysis and interpretation: Cora Sternberg, Andrew Armstrong, Roberto Pili, Siobhan Ng, Robert Huddart, Neeraj Agarwal, Nicholas James, Thomas Hutson, Piotr Milecki, Simon Chowdhury, Jean-Christophe Pouget, Frédérique Baton, Thore Nederman, Helen Tuvesson, Michael Carducci

Manuscript writing: All authors

Final approval of manuscript: All authors

REFERENCES

- Beer TM, Armstrong AJ, Rathkopf DE, et al: Enzalutamide in metastatic prostate cancer before chemotherapy. *N Engl J Med* 371:424-433, 2014
- de Bono JS, Oudard S, Ozguroglu M, et al: Prednisone plus cabazitaxel or mitoxantrone for metastatic castration-resistant prostate cancer progressing after docetaxel treatment: A randomised open-label trial. *Lancet* 376:1147-1154, 2010
- de Bono JS, Logothetis CJ, Molina A, et al: Abiraterone and increased survival in metastatic prostate cancer. *N Engl J Med* 364:1995-2005, 2011
- Kantoff PW, Higano CS, Shore ND, et al: Sipuleucel-T immunotherapy for castration-resistant prostate cancer. *N Engl J Med* 363:411-422, 2010
- Parker C, Nilsson S, Heinrich D, et al: Alpha emitter radium-223 and survival in metastatic prostate cancer. *N Engl J Med* 369:213-223, 2013
- Ryan CJ, Smith MR, de Bono JS, et al: Abiraterone in metastatic prostate cancer without previous chemotherapy. *N Engl J Med* 368:138-148, 2013
- Scher HI, Halabi S, Tannock I, et al: Design and end points of clinical trials for patients with progressive prostate cancer and castrate levels of testosterone: Recommendations of the Prostate Cancer Clinical Trials Working Group. *J Clin Oncol* 26:1148-1159, 2008
- Sounni NE, Noel A: Targeting the tumor microenvironment for cancer therapy. *Clin Chem* 59:85-93, 2013
- Ganguly SS, Li X, Miranti CK: The host microenvironment influences prostate cancer invasion, systemic spread, bone colonization, and osteoblastic metastasis. *Front Oncol* 4:364, 2014
- Clarke JM, Armstrong AJ: Novel therapies for the treatment of advanced prostate cancer. *Curr Treat Options Oncol* 14:109-126, 2013
- Gupta N, Al Ustvani O, Shen L, et al: Mechanism of action and clinical activity of tasquinimod in castrate-resistant prostate cancer. *Onco Targets Ther* 7:223-234, 2014
- Raymond E, Dalgleish A, Damber JE, et al: Mechanisms of action of tasquinimod on the tumour microenvironment. *Cancer Chemother Pharmacol* 73:1-8, 2014
- Shen L, Sundstedt A, Ciesielski M, et al: Tasquinimod modulates suppressive myeloid cells and enhances cancer immunotherapies in murine models. *Cancer Immunol Res* 3:136-148, 2015
- Isaacs JT, Antony L, Dalrymple SL, et al: Tasquinimod is an allosteric modulator of HDAC4 survival signaling within the compromised cancer microenvironment. *Cancer Res* 73:1386-1399, 2013
- Pili R, Häggman M, Stadler WM, et al: Phase II randomized, double-blind, placebo-controlled study of tasquinimod in men with minimally symptomatic metastatic castrate-resistant prostate cancer. *J Clin Oncol* 29:4022-4028, 2011
- Armstrong AJ, Häggman M, Stadler WM, et al: Long-term survival and biomarker correlates of tasquinimod efficacy in a multicenter randomized study of men with minimally symptomatic metastatic castration-resistant prostate cancer. *Clin Cancer Res* 19:6891-6901, 2013
- Scher HI, Fizazi K, Saad F, et al: Increased survival with enzalutamide in prostate cancer after chemotherapy. *N Engl J Med* 367:1187-1197, 2012
- Bratt O, Häggman M, Ahlgren G, et al: Open-label, clinical phase I studies of tasquinimod in patients with castration-resistant prostate cancer. *Br J Cancer* 101:1233-1240, 2009
- Eisenhauer EA, Therasse P, Bogaerts J, et al: New response evaluation criteria in solid tumours: Revised RECIST guideline (version 1.1). *Eur J Cancer* 45:228-247, 2009
- Lan KKG, Demets DL: Discrete sequential boundaries for clinical trials. *Biometrika* 70:659-663, 1983
- Brookmeyer R, Crowley J: A confidence interval for the median survival time. *Biometrics* 38:29-41, 1982
- Morris MJ, Molina A, Small EJ, et al: Radiographic progression-free survival as a response biomarker in metastatic castration-resistant prostate cancer: COU-AA-302 results. *J Clin Oncol* 33:1356-1363, 2015
- Armstrong AJ, Halabi S: Making progress on progression in metastatic prostate cancer. *J Clin Oncol* 33:1322-1324, 2015
- Fizazi K, Jones R, Oudard S, et al: Phase III, randomized, double-blind, multicenter trial comparing orteronel (TAK-700) plus prednisone with placebo plus prednisone in patients with metastatic castration-resistant prostate cancer that has progressed during or after docetaxel-based therapy: ELM-PC 5. *J Clin Oncol* 33:723-731, 2015
- Kelly WK, Halabi S, Carducci M, et al: Randomized, double-blind, placebo-controlled phase III trial comparing docetaxel and prednisone with or without bevacizumab in men with metastatic castration-resistant prostate cancer: CALGB 90401. *J Clin Oncol* 30:1534-1540, 2012

26. Michaelson MD, Oudard S, Ou YC, et al: Randomized, placebo-controlled, phase III trial of sunitinib plus prednisone versus prednisone alone in progressive, metastatic, castration-resistant prostate cancer. *J Clin Oncol* 32:76-82, 2014

27. Crawford ED, Higano CS, Shore ND, et al: Treating patients with metastatic castration resistant prostate cancer: A comprehensive review of available therapies. *J Urol* 194:1537-1547, 2015

28. Olsson A, Björk A, Vallon-Christersson J, et al: Tasquinimod (ABR-215050), a quinoline-3-carboxamide anti-angiogenic agent, modulates the expression of thrombospondin-1 in human prostate tumors. *Mol Cancer* 9:107, 2010

Affiliations

Cora Sternberg, San Camillo Forlanini Hospitals, Rome, Italy; Andrew Armstrong, Duke Cancer Institute, Duke University, Durham, NC; Roberto Pili, Indiana University School of Medicine, Indianapolis, IN; Siobhan Ng, St John of God Medical Centre, Subiaco, Western Australia, Australia; Robert Huddart, Royal Marsden Hospital, Sutton; Nicholas James, Queen Elizabeth Hospital, Birmingham; Simon Chowdhury, Guy's Hospital and Sarah Cannon Research UK, London, United Kingdom; Neeraj Agarwal, University of Utah, Salt Lake City, UT; Denis Khvorostenko, Leningrad Regional Oncology Dispensary, St Petersburg, Russia; Olexiy Lyulko, Zaporizhzhya Regional Clinical Hospital, Zaporizhzhya, Ukraine; Arija Brize, Riga Eastern Clinical University Hospital, Riga, Latvia; Nicholas Vogelzang, Comprehensive Cancer Centers of Nevada, Las Vegas, NV; Rémy Delva, Centre Régional de Lutte Contre le Cancer Paul Papin, Angers; Jean-Christophe Pouget and Frédérique Baton, Ipsen Innovation, Les Ulis, France; Mihai Harza, Fundeni Clinical Institute, Bucharest, Romania; Anastasios Thanos, Agios Savas Anticancer Oncology Hospital of Athens, Athens, Greece; Patrick Werbrouck, Algemeen Ziekenhuis Groeninge, Kortrijk, Belgium; Martin Bögemann, Universitätsklinikum Münster, Münster, Germany; Thomas Hutson, Texas Oncology, Dallas, TX; Piotr Milecki, Poznan University of Medical Sciences, Poznan, Poland; Enrique Gallardo, Corporació Sanitaria Parc Taulí, Sabadell, Spain; Gilberto Schwartzmann, Hospital De Clinicas De Porto Alegre, Porto Alegre, Brazil; Thore Nederman and Helen Tuveesson, Active Biotech, Lund, Sweden; and Michael Carducci, Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins University, Baltimore, MD.



Participate in ASCO's Practice Guidelines Implementation Network and Influence Cancer Care

ASCO members are invited to serve in the society's Practice Guidelines Implementation Network (PGIN), a network of oncology professionals who raise awareness of ASCO's evidence-based recommendations on cancer care.

Participation in PGIN provides an opportunity for members to positively influence the way that clinical oncology is delivered now and in the future. PGIN members have the opportunity to:

- Participate in Guideline Panels and Advisory Groups
- Aid in developing and reviewing Guidelines and Guideline Clinical Tools and Resources
- Serve as an "ambassador" to state societies
- Better implement Guidelines

To learn how you can participate, visit asco.org/guidelines, or contact PGIN@asco.org.



AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Randomized, Double-Blind, Placebo-Controlled Phase III Study of Tasquinimod in Men With Metastatic Castration-Resistant Prostate Cancer

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or jco.ascopubs.org/site/ifc.

Cora Sternberg

Honoraria: Pfizer, Novartis, Janssen Pharmaceuticals, Sanofi, GlaxoSmithKline

Consulting or Advisory Role: Bristol-Myers Squibb, Bayer HealthCare Pharmaceuticals, Astellas Pharma, Eisai

Research Funding: Sanofi (Inst), Bayer (Inst), Pfizer (Inst), Exelixis (Inst), Astellas Pharma (Inst), Medivation (Inst), Active Biotech (Inst), Genentech (Inst)

Andrew Armstrong

Honoraria: Dendreon, Sanofi

Consulting or Advisory Role: Bayer HealthCare Pharmaceuticals, Sanofi, Novartis, Dendreon, Medivation, Janssen Biotech

Speakers' Bureau: Dendreon, Sanofi

Research Funding: Dendreon (Inst), Sanofi (Inst), Bayer HealthCare Pharmaceuticals (Inst), Pfizer (Inst), Novartis (Inst), Bristol-Myers Squibb (Inst), KangLaiTe (Inst), Janssen Oncology (Inst), Medivation (Inst), Astellas Pharma (Inst), ImClone Systems (Inst), Active Biotech (Inst)

Patents, Royalties, Other Intellectual Property: Circulating tumor cell novel capture technology (Inst)

Travel, Accommodations, Expenses: Dendreon, Medivation, Janssen Biotech

Roberto Pili

Consulting or Advisory Role: Ipsen

Research Funding: Active Biotech

Siobhan Ng

Consulting or Advisory Role: Janssen-Cilag, Astellas Pharma, Pfizer, Novartis, GlaxoSmithKline, Sanofi

Expert Testimony: Janssen-Cilag, Astellas Pharma

Travel, Accommodations, Expenses: Sanofi, Novartis, Astellas

Robert Huddart

Leadership: Cancer Centre London

Consulting or Advisory Role: Astellas Pharma, Merck Sharp & Dohme, Janssen Oncology

Speakers' Bureau: Pierre Fabre

Research Funding: Merck Sharp & Dohme

Travel, Accommodations, Expenses: Janssen Oncology, Merck Sharp & Dohme

Neeraj Agarwal

Consulting or Advisory Role: Pfizer, Medivation/Astellas Pharma, Exelixis, Argos Therapeutics, Cerulean Pharma

Denis Khvorostenko

No relationship to disclose

Olexiy Lyulko

No relationship to disclose

Arija Brize

No relationship to disclose

Nicholas Vogelzang

Stock or Other Ownership: Caris Life Sciences

Honoraria: UpToDate, Endocyte

Consulting or Advisory Role: Janssen Biotech, Amgen, Cerulean Pharma, Pfizer

Speakers' Bureau: Medivation, Dendreon, Bayer AG, Sanofi, GlaxoSmithKline, Genentech

Research Funding: PAREXEL International (Inst), Progenics Pharmaceuticals (Inst), US Oncology (Inst), Viamet Pharmaceuticals (Inst), Endocyte (Inst), GlaxoSmithKline (Inst)

Travel, Accommodations, Expenses: Genentech, Celgene, US Oncology, Dendreon, Novartis, Pfizer, Bayer AG, Exelixis

Rémy Delva

No relationship to disclose

Mihai Harza

No relationship to disclose

Anastasios Thanos

No relationship to disclose

Nicholas James

Honoraria: Sanofi, Bayer HealthCare Pharmaceuticals, OncoGenex, Janssen Pharmaceuticals, Astellas Pharma, Pierre Fabre

Consulting or Advisory Role: Sanofi, Bayer HealthCare Pharmaceuticals, Merck, Astellas Pharma, Janssen Pharmaceuticals

Speakers' Bureau: Pierre Fabre, Ferring Pharmaceuticals, Sanofi, Astellas Pharma, Janssen Pharmaceuticals, Ipsen

Research Funding: Janssen Pharmaceuticals (Inst), Astellas Pharma (Inst), Pfizer (Inst), Sanofi (Inst), Novartis (Inst), Active Biotech (Inst), Bayer HealthCare Pharmaceuticals (Inst), Merck (Inst), Genentech (Inst)

Patrick Werbrouck

No relationship to disclose

Martin Bögemann

Honoraria: Janssen-Cilag, Astellas Pharma, Amgen, Sanofi, Pfizer, Bristol-Myers Squibb

Consulting or Advisory Role: Janssen-Cilag, Astellas Pharma, Amgen, Sanofi, Pfizer, Bristol-Myers Squibb

Travel, Accommodations, Expenses: Bayer Vital GmbH, Janssen-Cilag, Astellas Pharma, TEVA Pharmaceuticals

Thomas Hutson

Employment: Texas Oncology

Honoraria: Bayer/Onyx, Pfizer, Medivation, Novartis, Johnson & Johnson, GlaxoSmithKline, Dendreon

Consulting or Advisory Role: Bayer/Onyx, Pfizer, Novartis, Astellas Pharma, GlaxoSmithKline

Speakers' Bureau: Dendreon, Pfizer, GlaxoSmithKline, Bayer/Onyx, Johnson & Johnson, Medivation

Research Funding: Pfizer (Inst), GlaxoSmithKline (Inst), Johnson & Johnson (Inst), Medivation (Inst)

Piotr Milecki

Honoraria: Astellas Pharma, Ipsen, AstraZeneca

Travel, Accommodations, Expenses: Astellas Pharma, Ipsen Pharmaceuticals

Simon Chowdhury

Honoraria: Pfizer, Novartis, Janssen-Cilag, Astellas Pharma

Consulting or Advisory Role: Novartis, Sanofi, Johnson & Johnson, Astellas Pharma, Pfizer, Bayer AG

Speakers' Bureau: Astellas Pharma, Sanofi, Johnson & Johnson, Bayer AG

Research Funding: Sanofi

Travel, Accommodations, Expenses: Astellas Pharma, Johnson & Johnson, Bayer

Enrique Gallardo

Honoraria: Novartis, Pfizer, Ipsen

Consulting or Advisory Role: Pfizer, GlaxoSmithKline, Novartis, Astellas Pharma, Janssen Oncology, Bayer Schering Pharma

Speakers' Bureau: Pfizer, Novartis, GlaxoSmithKline, Astellas Pharma, LEO Pharma, Sanofi

Travel, Accommodations, Expenses: Pfizer, Novartis, Astellas Pharma, Janssen Oncology, Pierre Fabre, Sanofi

Gilberto Schwartzmann

No relationship to disclose

Jean-Christophe Pouget

Employment: Ipsen

Frédérique Baton

Employment: Ipsen

Stock or Other Ownership: Ipsen

Thore Nederman

Employment: Active Biotech

Stock or Other Ownership: Active Biotech

Helen Tuvevson

Employment: Active Biotech

Leadership: Active Biotech

Stock or Other Ownership: Active Biotech

Michael Carducci

Consulting or Advisory Role: Medivation, Astellas Pharma, AstraZeneca, Clovis Oncology, Merck

Acknowledgment

We thank Neil Anderson from Adelphi Communications who provided medical writing support that was funded by Ipsen.

Appendix

Table A1. Phase III Study Sites (241) in 37 Countries

Site	Principal Investigator
Australia	
The Tweed Hospital, Tweed Heads	Ehtesham Abdi, MBBS
Coffs Harbor Health Campus, Coffs Harbor	Karen Briscoe, MBBS
Australian Prostate Cancer Research Centre, Richmond	Anthony Costello, MD, MBBS
St John of God Subiaco Hospital, Subiaco	Siobhan Ng, MBBS
St Vincent's Hospital, Darlinghurst	Richard Epstein, MD, PhD, MBBS
Royal Prince Alfred Hospital, Camperdown	Lisa Horvath, MBBS, PhD
India	
Jehangir Clinical Development Centre, Pune	Bhalchandra Kashyapi, MBBS, MCh, MS
New Zealand	
Tauranga Urology Research, Tauranga	Peter Gilling, MD
Canterbury Urology Research Trust, Christchurch	Frank Kueppers, MD, PhD
Roundhay Medical Centre and Nelson Public Hospital, Nelson; Wairau Public Hospital, Blenheim	Patrick Meffan, MBChB, FRACS
Palmerston North Hospital, Palmerston North	Quinten King, MBBCh, FRCS
Korea	
Gangnam Severance Hospital, Seoul	Byung Ha Chung, MD, PhD
Chonnam National University Hospital, Gwangju	Taek Won Kang, MD, PhD
Seoul St Mary's Hospital, Seoul	Sae Woong Kim, MD, PhD
Asan Medical Center, Seoul	Choung-Soo Kim, MD
Severance Hospital, Seoul	Sung Joon Hong, MD, PhD, MS
Samsung Medical Center, Seoul	Hyun Moo Lee, MD, PhD
Taiwan	
Chang Gung Medical Foundation, Taoyuan	Cheng-Keng Chuang, MD
Taichung Veterans General Hospital, Taichung	Yen-Chuan Ou, MD, PhD
National Taiwan University Hospital, Taipei	Yu-Chieh Tsai, MD
Kaohsiung Veterans General Hospital, Kaohsiung	Tong-Lin Wu, MD, EMBA
China	
Urology Surgery Department, Beijing	Lijun Chen, MD
Fudan University Shanghai Cancer Center, Shanghai	Dingwei Ye, MD
Chengdu Military General Hospital, Chengdu	Liang Wang, MD
Urology Surgery Department, Shantou	Junhong Zheng, MD
Huashan Hospital, Shanghai	Qiang Ding, MD
Zhongnan Hospital of Wuhan University, Wuhan	Fuxiang Zhou, MD
Argentina	
Centro Oncológico "Ágave," Santa Fe	Natalia Broglio Sicco, MD
Centro de Diagnóstico Urológico, Buenos Aires	Luis Fernando Montes de Oca, MD
Centro Oncológico Fundacion Korla, St Rosa	Pablo Picon, MD
Hospital Italiano de Buenos Aires, Buenos Aires	María Pallotta, MD
Brazil	
Hospital Evangelico de Cachoeiro de Itapemirim, Cachoeiro de Itapemirim	Sabina Aleixo, MD
Granbery Juiz de Fora Hospital Universitario da Universidade Federal de Juiz de Fora Avenue Eugenio do Nascimento, Juiz de Fora	Christiane Alves, MD
Rio de Janeiro Hospital St Maria Madalena Estrada do Dende, Rio de Janeiro	Iane Cardoso, MD
Natal Liga Norte Riograndense Contra o Cancer Unidade I Hospital Luiz Antonio, Natal	Danielli Matias, MD
Florianopolis Hospital, Bala Sul Medical Center, Florianopolis	Yeni Neron, MD
Salvador Hospital da Bahia, Salvador	José Nogueira, MD
Sao Paulo Centro de Pesquisa Clinical, Sao Paulo	Roberto Rocha, MD
Centro Oncologico Mogi das Cruzes, Sao Paulo	Daniel Grabarz, MD
Passo Fundo Hospital Sao Vicente de Paulo, Passo Fundo	Luis Schlittler, MD
Hospital de Clinicas de Porto Alegre, Porto Alegre	Gilberto Schwartzmann, MD
Rio Sul Centro de Atencao e Saude Humana, Rio de Janeiro Hospital, Rio de Janeiro	Leonardo Osorio, MD; Heloisa M. Resende, MD
Universidade Estadual Paulista, Sao Paulo	Guareide Carelli, MD
Liga Paranaense de Combate ao Cancer, Hospital Erasto Gaertner, Curitiba	Flavio Tomasich, MD

(continued on following page)

Table A1. Phase III Study Sites (241) in 37 Countries (continued)

Site	Principal Investigator
Chile	
Hospital Clínico Vina de Mar Limache, Vina de Mar	Alejandro Acevedo Gaete, MD
Clinicia Alemana de Temuco, Temuco	Mario Gorena, MD
Uromed Ave Salvador 351, Santiago	Anibal Salazar Huerta, MD
Hospital Dipreca, Santiago	Luis Soto Diaz, MD
Hospital de Carabineros General Humberto Arriagada Valdivieso, Santiago	Nelson Orellana Salinas, MD
Colombia	
Fundacion Clinica Valle del Lili, Cali	Manuel Duque Galan, MD
Fundacion St Fe de Bogota, Bogota	Carlos Vargas, MD
Mexico	
Hospital Angeles Puebla, Puebla	José Arroyo Kuribreña, MD
Hospital Aranda de la Parra, Guanajuato	Marco Badillo Santoyo, MD
Christus Muguerza del Parque, Chihuahua	Roberto Hidalgo-Silva, MD
Hospital Angeles Culiacan, Culiacan	Gustavo Gaxiola Meza, MD
Consultorio Medico, Zapopan	Jose Rodriguez Rivera, MD
Panama	
Centro Hemato Oncologico Paitilla Consultorios Medicos Royal Center, Panama City	Juan Bares Weeden, MD
Clinica Hospital San Fernando Centro Especializado, Panama City	Javier Del Rosario Gibbs, MD
Medical Research Center Edificio Consultotios America, Panama City	Roberto Lopez Sanchez, MD
Clinica Hospital San Fernando, Panama City	Alejandro Manduley, MD
Peru	
Instituto Régional de Enfermedades Neoplasicas del Sur Ave de la Salud Arequipa, Arequipa	Ernesto Vargas Quezada, MD
Hospital Nacional Carlos Alberto Seguin Escobedo Calle Peral y Ayacucho, Arequipa	Hernan Moron Escobar, MD
Belgium	
Centre Hospitalier Universitaire Sart Tilman, Liege	Brieuc Sautois, MD
Algemeen Ziekenhuis Groeninge Burgemeester, Kortijk	Patrick Werbrouck, MD
Erasme Hospital, Université Libre de Bruxelles, Brussels	Thierry Roumeguère, MD
Algemeen Ziekenhuis Maria Middelaes, Gent	Filip Ameye, MD
Bulgaria	
Hospital for Treatment of Oncology Disease, Sofia	Borislav Dimitrov, MD
Oncology Center Plovdiv, Plovdiv	Petar Petrov, MD
Hospital Oncology Disease Medical Oncotherapy and Palliative Care Base II, Varna	Violina Taskova, MD
University Multiprofile Hospital Chemotherapy Department, Pleven	Rumyana Micheva, MD
Czech Republic	
Fakultni Nemocnice v Motole, Urology Clinic, Prague	Josef Stolz, MD
Urocentrum Praha, Prague	Michaela Matouskova, MD
Krajiska Zdravotni – Masarykova Nemocnice, Clinic of Urology and Robotic Surgery, Usti nad Labem	Jan Schraml, MD
Nemocnice Novy Jicin, Urology Department, Novy Jicin	Miroslav Stursa, MD
Fakultni Nemocnice Olomouc, Urology Clinic, Olomouc	Vladimir Student, MD
Estonia	
Tartu University Hospital, Tartu	Jaanus Kahu, MD
France	
Central Hospital Cannes, Cannes	Regis Kaphan, MD
Beau Soleil Clinic, Montpellier	Xavier Rebillard, MD
Foch Hospital, Suresnes	Christine Theodore, MD
Regional Center de Lutte, Angers	Rémy Delva, MD
Centre Hospital Lyon-Sud, Pierre-Benite	Alain Ruffion, MD
Centre Régional de Lutte Contre le Cancer Val d'Aurelle, Montpellier	David Azria, MD
Centre Hospitalier Universitaire Rennes Hospital Pontchaillou, Rennes	Sebastien Vincendeau, MD
Curie Institute, Paris	Philippe Beuzeboc, MD
Hopital Saint-Louis Service d'Oncologie, Paris	Stéphane Culine, MD
Centre Hospitalier Universitaire La Timone Adultes-Service d'Oncologie Medicale, Marseille	Jean-Laurent Deville, MD
Germany	
Eberhard-Karls University Tübingen, Clinic for Urology, Tübingen	Arnulf Stenzl, MD
Urologic Study Office, Nürtingen	Susan Feyerabend, MD
University Hospital Münster, Urology, Münster	Martin Bögemann, MD
Hospitals Nordoberpfalz, Urology, Weiden	Theodor Klotz, MD
University Hospital Gießen and Marburg, Clinic for Urology and Pediatric Urology, Marburg	Alex Hegele, MD
University Hospital Cologne, Clinic for Urology, Cologne	Sebastian Wille, MD
University Hospital Munich, Clinic for Urology, Munich	Phillip Nuhn, MD; Claudius Füllhase, MD; Patrick Bastian, MD

(continued on following page)

Tasquinimod in Metastatic Castration-Resistant Prostate Cancer

Table A1. Phase III Study Sites (241) in 37 Countries (continued)

Site	Principal Investigator
University Hospital Carl Gustav Carus Dresden, Dresden Martini-Clinic at the University Hospital Hamburg-Eppendorf, Hamburg University Hospital Mannheim, Clinic for Urology, Mannheim	Manfred Wirth, MD Thomas Steuber, MD, PD Christian Bolenz, MD, PD
Greece Alexandra Hospital Department of Clinical Therapeutics, Athens Athens Oncology Hospital Urology Clinic, Athens Thessaloniki General Hospital, Thessaloniki Patras University General Hospital, Rion Patras	Eleni Efstathiou, MD, PhD Anastasios Thanos, MD Athanasios Papatthasiou, MD Petros Perimenis, MD
Israel Oncology Institute, The Chaim Sheba Medical Center, Tel Hashomer The Lady Davis Carmel Medical Center, Haifa Assaf Harofe Medical Center Oncology Department, Zerifin Tel Aviv Sourasky Medical Center, Oncology Department, Tel Aviv Oncology Institute Rambam Health Care Campus, Haifa Institute of Oncology Davidoff Cancer Center, Rabin Medical Center, Tikva Bnai Zion Medical Center, Haifa Sharett Institute of Oncology, Hadassah University Hospital, Jerusalem Soroka University Medical Center, Be'er Sheva	Raanan Berger, MD, PhD Avi Stein, MD Avishay Sella, MD Eliahu Gez, MD Avivit Peer, MD Eli Rosenbaum, MD Ofer Nativ, MD Stephen Frank, MD Wilmosh Mermershtain, MD
Italy San Camillo Forlanini Hospital, Rome Scientific Institute Romagnolo Via Piero Maroncelli, Meldola Oncology Institute Veneto, Padova Hospital di Lecco, Lecco Oncologia Falck Hospital, Niguarda Ca Granda Piazzale Hospital, Milan Azienda Ospedaliero Universitaria Giovanni Battista di Torino Molinette, Turin Istitute di Cremona, Cremona Ospedale degli Infermi di Faenza Unità di Oncologia di Oncologia Medica, Faenza Hospital San Carlo Borromeo, Milan	Cora Sternberg, MD Cecilia Menna, MD Umberto Basso, MD Antonio Ardizzoia, MD Salvatore Siena, MD Liberio Ciuffreda, MD Rodolfo Passalacqua, MD Francesco Carrozza, MD; Giorgio Cruciani, MD Maria Locatelli, MD
Latvia Riga Eastern Clinical University Hospital, Latvian Oncology Center, Riga Private practice of Dzintra Litavniece, Liepaja P. Stradins Clinical University Hospital, Riga	Arija Brize, MD Dzintra Litavniece, MD Egils Vjaters, MD
Lebanon Middle East Institute of Health, Bsalim El Meten American University of Beirut Medical Center, Beirut	Abi Gerges Dany, MD Ali Shamseddine, MD
Lithuania Vilnius University Hospital, Vilnius Institute of Oncology Vilnius University, Vilnius Lithuanian University Health Sciences Kaunas Clinics, Kaunas	Feliksas Jankevicius, MD Albertas Ulys, MD, PhD Daimantas Milonas, MD
The Netherlands St Elizabeth Hospital, Tilburg Martini Ziekenhuis, Groningen Canisius Wilhelmina Hospital, Nijmegen Leiden University Medical Center, Leiden Vrije Universiteit Medical Center, Amsterdam University Medical Center St Radboud, Nijmegen	P. Kil, MD, PhD L.F.A. Wymenga, MD, PhD H. Vergunst, MD A.J. Gelderblom, MD R.J.A. van Moorselaar, MD P.F.A. Mulders, MD
Poland Curie Oncology Institute, Nowotworow Oncology Clinic, Warsaw Niepubliczny Zaklad Opieki Zdrowotnej Urology Center, Myslowice Regional Osrodek Oncology, Lodz EuroMediCare, Wroclaw Wojewodki Hospital Urology Clinic, Bialystok Wielkopolskie Oncology Center, Poznan LexMedica Rudolfa, Wroclaw	Tomasz Demkow, MD, PhD Adam Dobrowolski, MD Ewa Kalinka-Warzocho, MD Rafal Kmieciak, MD Robert Kozlowski, MD, PhD Piotr Milecki, MD, PhD Zenona Jablonska, MD
Romania Oncolab, Craiova Fundeni Clinical Institute, Bucharest Sf Ioan cel Nou Emergency County Hospital, Suceava The Oncology Institute, Cluj Napoca Opris Emergency County Hospital, Baia Mare Oncomed SRL, Timisoara Ianuli Medical Consult SRL, Bucharest Municipal Hospital Ploiesti, Ploiesti	Dan Lungulescu, MD Mihai Harza, MD Doina Ganea, MD, PhD Cristina Cebotaru, MD; Tudor Ciuleanu, MD, PhD Dumitru Filip, MD Cristina Oprean, MD Carmen Ianuli, MD Gabriel Doru Ghizdavescu, MD

(continued on following page)

Table A1. Phase III Study Sites (241) in 37 Countries (continued)

Site	Principal Investigator
Russia	
Omsk Healthcare, Oncology Center, Omsk	Evgeniy Kopyltsov, MD, PhD
Clinic Andros Lenina Street, St Petersburg	Alexey Plekhanov, MD, PhD
Leningrad Regional Oncology Center, St Petersburg	Denis Khvorostenko, MD
St Petersburg Healthcare City Hospital, St Petersburg	Vakhtang Shanava, MD, PhD
Vladimir Healthcare Oncology Center, Vladimir	Natalya Rodicheva, MD
Orkli, Sredniy Prospekt, St Petersburg	Vladimir Kheifets, MD, PhD
Federal State Institution Moscow Research Oncology Institute, Moscow	Boris Alekseev, MD, PhD
State Institution of Healthcare Sverdlovsk Regional Hospital, Ekaterinburg	Alexander Zyryanov, MD
Regional State Institution of Healthcare Novosibirsk Regional Oncology Centre, Novosibirsk	Marat Zaripov, MD
Slovak Republic	
Ambulatory Urology Clinic, Trecin	Roman Sokol, MD
Cuimed, Bratislava	Frederico Goncalves, MD, PhD
Spain	
Hospital Clinic 1 Provincial Oncology Servicio de Oncologia Medica, Barcelona	Begoña Mellado, MD
Corporacio Sanitaria Parc Tauli Hospital de Sabadell Servicio de Oncologia Medica, Barcelona	Enrique Gallardo, MD
Hospital Infanta Sofia, Madrid	Emilio Ríos, MD
Clinica Universidad de Navarra Servicio de Oncologia, Pamplona	Jose Luis Perez Gracia, MD
Hospital Universitario Virgen del Rocio Servicio de Oncologia Medica Ave Manuel Siurot, Sevilla	Begoña Pérez Valderrama, MD
Hospital Clinico Universitario de Valencia Servicio de Oncologia Medica, Valencia	Isabel Chirivella, MD
Hospital Universitario Marques de Valdecilla Servicio de Oncologia Medica, Santander	Marta Lopez-Brea Piqueras, MD
Hospital Universitario Vall D'Hebron Servicio de Oncologia-Unidad, Barcelona	Joan Carles Galceran, MD
Hospital de la Santa Creu, Barcelona	José Pablo Maroto, MD
Hospital Universitario Fundacion Alcorcon Servicio de Oncologia Medica, Alcorcon	Susana Hernando Polo Jesus, MD; Garcia-Donas Jimenez, MD
Hospital Clinico Universitario "Lozano Blesa" Servicio de Oncologia Medica, Zaragoza	Alberto Saenz Cusi, MD
Hospital Universitario Central de Asturias Servicio de Oncologia, Oviedo	Emilio Esteban Gonzalez, MD; Enrique Estrada, MD
Instituto Valenciano de Oncologia, Valencia	Eduardo Solsona Narbon, MD
Sweden	
Radiumhemmet Karolinska University Hospital, Stockholm	Sten Nilsson, MD
Sahlgrenska University Hospital, Gothenburg	Jan-Erik Damber, MD
Central Hospital Karlstad Oncology Clinic, Karlstad	Claes Ginman, MD
Turkey	
Istanbul University Cerrahpasa School of Medicine, Istanbul	Can Obek, MD
Ukraine	
Municipal Institution of Healthcare VI Shapoval Regional Clinical Centre of Urology and Nephrology Urology Department #4, Kharkiv	Igor Antonyan, MD, PhD
Ivano-Frankivsk Regional Oncology, Dispensary Clinical Mammology Centre, Department with Urology Beds Ivano-Frankivsk	Volodymyr Romanchuk, MD; Ipolit Kostinskyy, Professor, MD, PhD
Municipal Institution, Multifield City Clinical Hospital #4, Department of Chemotherapy, Dnipropetrovsk	Igor Bondarenko, Professor, MD, PhD
Municipal Institution, Zaporizhzhia Regional Clinical Hospital of Zaporizhzhia Regional Council, Urology Department; State Institution Zaporizhzhia Medical Academy of Postgraduate Education of Ministry of Health of Ukraine Chair of Urology, Zaporizhzhya	Olexiy Lyulko, Professor, MD
Kyiv City Clinical Hospital #3 Urology Department, Kyiv	Petro Ivashchenko, MD
Municipal Institution, Dnipropetrovsk Regional Clinical Hospital named after I.I. Mechnikov, Urology Department #2, Dnipropetrovsk State Medical Academy Chair of Urology, Operative Surgery and Topographic Anatomy, Dnipropetrovsk	Olexiy Lyulko, Professor, MD; Viktor Stus, Professor, MD
Regional Municipal Institution, Chernivtsi Regional Clinical Hospital, Chernivtsi	Valerii Zaitsev, Professor, MD
Kyiv Oleksandrivska Clinical Hospital, Urology Department #3, Kyiv	Sergii Pasichnikov, Professor, MD
Uzhgorod Central City Clinical Hospital, City Oncology Centre, Uzhgorod	Yevhen Hotko, MSD, MD, PhD
Municipal Clinical Medical and Preventive Treatment Institution, Donetsk Regional Antitumour Centre, Donetsk	Andriy Anishchenko, MD
Medical and Preventive Treatment Institution, Volyn Regional Oncology Dispensary, Lutsk	Orest Andrusenko, MD
Lviv State Oncology Regional Treatment and Diagnostic Center, Lviv	Yaroslav Shparyk, MD, PhD
Municipal Treatment-Prophylactic, Institution Central City Clinical Hospital, Donetsk	Yuri Semyak, MD

(continued on following page)

Tasquinimod in Metastatic Castration-Resistant Prostate Cancer

Table A1. Phase III Study Sites (241) in 37 Countries (continued)

Site	Principal Investigator
Municipal Institution of Kyiv Regional Council, Kyiv Regional Oncology Dispensary, Kyiv	Iurii Golovko, MD
Center of Reconstructive and Restorative Medicine (University Clinic) of Odesa National Medical University, Odesa	Natalia Tavartkiladze, MD
United Kingdom	
St James University Hospital, Leeds	William Cross, MD
Royal Marsden Hospital, Sutton	Robert Huddart, MD
Mount Vernon Hospital, Northwood	Peter Hoskin, MD
Oxford Cancer Centre, Headington	Andrew Protheroe, MD
St Richard's Hospital, Chichester	James Hicks, MD; Paul Carter, MD
Scunthorpe General Hospital, Scunthorpe	Sanjay Dixit, MD
Sarah Cannon Research, London	Simon Chowdhury, MA, MBBS, MRCP, PhD
University Hospitals Birmingham National Health Service Foundation Trust Queen Elizabeth Hospital, Birmingham	Nicholas James, MD
Nottingham University Hospitals National Health Service Trust, Nottingham	Santhanam Sundar, MD
Canada	
The Fe/Male Health Centre, Oakville, ON	Richard Casey, MD
Probit Medical Research, North York, ON	Stanley Flax, MB, BCH
Southern Interior Medical Research, Kelowna, BC	Thomas Kinahan, MD
Mor Urology, Newmarket, ON	Morrie Liquornik, MD
Pacific Urologic Research, Victoria, BC	Gary Steinhoff, MD
St Joseph's Lifecare Centre, Brantford, ON	Wilson Leung, MD
United States	
Duke University Medical Center, Durham, NC	Andrew Armstrong, MD
Peachtree Hematology-Oncology Consultants, Atlanta, GA	Vasileios John Assikis, MD
Urologic Consultants of Pennsylvania, Bala Cynwyd, PA	Laurence H. Belkoff, DO
Pacific Urology Institute, Santa Monica, CA	Stanley Brosman, MD
The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins University, Baltimore, MD	Michael Carducci, MD
Ventura County Hematology Oncology Specialists, Oxnard, CA	Kevin Q. Chang, MD
Vanderbilt University Medical Center, Nashville, TN	Sam Chang, MD
Clinical Trials Office, Dallas, TX	James Cochran, MD
University of Pittsburgh Physicians Department of Urology, Pittsburgh, PA	Jeffrey Gingrich, MD
Premier Medical Group, Poughkeepsie, NY	Evan R. Goldfischer, MD
Midwest Urology Associates, Melrose Park, IL	Richard G. Harris, MD
Lawrenceville Urology, Lawrenceville, NJ	Gary S. Karlin, MD
Capitol Comprehensive Cancer Care Clinic, Jefferson City, MO	Ali Khojasteh, MD
Carolina Urology Partners, Concord, NC	David U. Lipsitz, MD, FACS, CPI
Palm Beach Urology Associates, Wellington, FL	Georgis Patsias, MD
Roswell Park Cancer Center Institute, Buffalo, NY	Roberto Pili, MD
Grand Strand Urology, Myrtle Beach, SC	Neal Shore, MD, FACS
Lancaster Urology, Lancaster, PA	Paul R. Sieber, MD
Boise Urology, Meridian, ID	Joseph H. Williams, MD
Metropolitan Urology, Jeffersonville, IN	James L. Bailen, MD
Frankel, Reed & Evans, Burien, WA	Jeffrey M. Frankel, MD
Virginia Oncology Associates, Norfolk, VA	Mark T. Fleming, MD
Virginia Cancer Specialists, Fairfax, VA	Alexander I. Spira, MD
Texas Oncology-Baylor Charles A. Sammons Cancer Center, Dallas, TX	Thomas E. Hutson, DO
Blue Ridge Cancer Care, Roanoke, VA	Mark D. Kochenderfer, MD
Willamette Valley Cancer Institute and Research Center, Eugene, OR	Joseph A. Fiorillo, MD; John R. Caton Jr, MD
Comprehensive Cancer Centers of Nevada, Las Vegas, NV	Nicholas J. Vogelzang, MD
Raleigh Hematology Oncology Associates, DBA Cancer Centers of North Carolina, Raleigh, NC	William R. Berry, MD
University of Utah/Huntsman Cancer Center, Salt Lake City, UT	Neeraj Agarwal, MD
Associates in Oncology/Hematology, Rockville, MD	Manish Agrawal, MD
Oncology Specialists, Park Ridge, IL	Timothy Lestingi, MD; Chadi Nabhan, MD
John Theurer Cancer Center at Hackensack, Hackensack, NJ	Robert Alter, MD
Redwood Regional Medical Group, Santa Rosa, CA	Wes S. Lee, MD
Arizona Oncology Associates, Tucson, AZ	Christopher Di Simone, MD
Tufts Medical Center, Boston, MA	Paul Mathew, MD

Table A2. Other Secondary Efficacy End Points

Outcomes	Tasquinimod (n = 832)		Placebo (n = 413)		HR	95% CI	P
	Median (months)	95% CI	Median (months)	95% CI			
New bone lesion							
Local	8.3	6.0 to 9.5	4.5	3.1 to 5.6	0.723	0.616 to 0.848	< .001
Central	8.1	6.0 to 8.5	4.8	3.1 to 5.6	0.735	0.623 to 0.867	< .001
New soft tissue lesion							
Local	19.4	16.6 to 25.3	11.1	8.6 to 16.4	0.612	0.493 to 0.760	< .001
Central	20.5	19.3 to NR	19.1	11.5 to NR	0.678	0.531 to 0.866	.002
First radiologic or symptomatic progression							
Local	4.8	4.1 to 5.5	3.2	2.9 to 4.2	0.812	0.714 to 0.925	.002
Central	5.2	4.4 to 5.6	3.7	3.1 to 4.4	0.849	0.745 to 0.967	.013
First radiologic or symptomatic progression or death							
Local	4.8	4.0 to 5.5	3.2	2.9 to 4.1	0.812	.716 to .922	.001
Central	5.2	4.4 to 5.6	3.6	3.1 to 4.3	0.845	0.744 to 0.959	.009
Tumor-related pain progression*	5.6	4.9 to 6.0	8.3	6.7 to 10.8	1.259	1.097 to 1.445	< .001
KPS deterioration	11.7	10.3 to 13.6	17.4	14.5 to 19.1	1.292	1.110 to 1.505	< .001
PSA doubling time	5.2	4.5 to 5.6	3.3	2.9 to 4.0	0.734	0.631 to 0.853	< .001

Abbreviations: HR, hazard ratio; KPS, Karnofsky performance status; NR, not reached; PSA, prostate-specific antigen.

*Including palliative interventions.