




1980

## Immuno- and Bioassayable Extrapituitary Growth Hormone and Thyroid Stimulating Hormone from Discrete Areas of the Central Nervous System

Sally Ann Hojvat  
*Loyola University Chicago*

Follow this and additional works at: [https://ecommons.luc.edu/luc\\_diss](https://ecommons.luc.edu/luc_diss)

 Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#)

---

### Recommended Citation

Hojvat, Sally Ann, "Immuno- and Bioassayable Extrapituitary Growth Hormone and Thyroid Stimulating Hormone from Discrete Areas of the Central Nervous System" (1980). *Dissertations*. 1986.  
[https://ecommons.luc.edu/luc\\_diss/1986](https://ecommons.luc.edu/luc_diss/1986)

This Dissertation is brought to you for free and open access by the Theses and Dissertations at Loyola eCommons. It has been accepted for inclusion in Dissertations by an authorized administrator of Loyola eCommons. For more information, please contact [ecommons@luc.edu](mailto:ecommons@luc.edu).



This work is licensed under a [Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 License](#).  
Copyright © 1980 Sally Ann Hojvat

IMMUNO- AND BIOASSAYABLE EXTRAPITUITARY  
GROWTH HORMONE AND THYROID STIMULATING HORMONE  
FROM DISCRETE AREAS OF THE CENTRAL NERVOUS SYSTEM

SALLY A. HOJVAT

A Dissertation Submitted to the Faculty of the Graduate School  
of Loyola University of Chicago in Partial Fulfillment  
of the Requirements for the Degree of  
Doctor of Philosophy

## ACKNOWLEDGEMENTS

The author of this dissertation would like to gratefully acknowledge the supervision given throughout this thesis by Dr. Ann Lawrence and to the following colleagues for their interest and supportive help with the project:

G. Baker, L. Kirsteins, H. Patejak, B. Booker, H. Tonaki, K. Ernst, N. Emanuele, and J. Kislá, R. Oslapas.

Also to my husband Carlos, daughter Nara, and my mother, my thanks for their constant encouragement through both the high and low points encountered in this research project.

## LIFE.

Sally Ann Hojvat was born in Bangor, N. Wales, on August 19th, 1942. After graduating from Queen Elizabeth's Grammar School, Manchester, England, in 1960, she received her Bachelor of Science with Honors in Microbiology and Agriculture from the University of Wales, Aberystwyth, in 1964. The next two years were spent completing a Master of Science in Microbiology from the University of Alberta, Edmonton, Canada. For the following ten year period, Sally was employed as a research assistant in the Departments of Pharmacology at the University of British Columbia, Canada and the Université de Genève, Switzerland, and the Department of Immunology, McGill University, Montreal, Canada. In 1977, she began her doctoral thesis research at Loyola under the supervision of Dr. Ann Lawrence, studying the characterization of two pituitary peptides found in the rodent CNS.

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGEMENTS . . . . .	ii
LIFE . . . . .	iii
LIST OF TABLES . . . . .	iv
LIST OF FIGURES . . . . .	vii
INTRODUCTION . . . . .	1
1. STATEMENT OF PROBLEM . . . . .	1
2. RELEVANT LITERATURE REVIEW . . . . .	7
A. The Pituitary-Hypothalamic Link . . . . .	7
B. Pituitary Growth Hormone . . . . .	13
a. Molecular structure of growth hormone . . . . .	14
b. Growth hormone levels in plasma and pituitary . . . . .	16
c. Biological effects of growth hormone . . . . .	19
d. Regulation of growth hormone release . . . . .	25
C. Pituitary Thyroid Stimulating Hormone (TSH) . . . . .	30
a. Molecular structure of Pituitary TSH . . . . .	31
b. Thyroid stimulating hormone levels in plasma and pituitary . . . . .	32
c. Biological effects of TSH . . . . .	34
d. Regulation of TSH release from the pituitary . . . . .	38
D. The Amine Precursor Uptake and Decarboxylation Theory (APUD) . . . . .	41
E. Peptides Found in the Central Nervous System Hypothalamic Hormones: . . . . .	42
a. Gonadotropin-releasing hormone (GNRH) . . . . .	44
b. TRH (thyroid-releasing hormone) . . . . .	45
c. Somatostatin . . . . .	48
d. Bombesin . . . . .	50
e. Glucagon . . . . .	51
f. Insulin . . . . .	51
g. Gastrin and cholecystokinin (CCK) . . . . .	52
h. Vasoactive intestinal polypeptide (VIP) . . . . .	54
i. Substance P . . . . .	56
j. Neurotensin . . . . .	58
k. Renin-angiotensin . . . . .	60

(Table of Contents, continued)

	<u>Page</u>
l. Vasopressin . . . . .	63
m. A common precursor for some pituitary hormones	64
n. Brain ACTH . . . . .	68
o. $\alpha$ melanocyte-stimulating hormone ( $\alpha$ MSH) . . . . .	71
p. The endorphins . . . . .	74
q. Prolactin . . . . .	81
r. Thyroid stimulating hormone (TSH) . . . . .	81
s. Growth hormone (GH) . . . . .	82
3. PROPOSED INVESTIGATION . . . . .	83
Specific Objectives . . . . .	83
4. MATERIALS AND METHODS . . . . .	85
A. The Development of Radioimmunoassays (RIA) for Rat Growth Hormone and Thyroid Stimulating Hormone . . . . .	85
I. The Rat Growth Hormone Radioimmunoassay . . . . .	88
a. Production of specific antiserum to rat growth hormone . . . . .	88
b. Radioiodine labeling procedure for rat growth hormone . . . . .	94
(i) Labeling procedure . . . . .	95
c. Preparation of GH standards . . . . .	98
d. Separation of bound and free antigen in the GH RIA . . . . .	99
e. Assay protocol . . . . .	99
f. Treatment of data obtained from the GH RIA . . . . .	100
g. Validation of the GH radioimmunoassay . . . . .	101
II. The Rat Thyroid Stimulating Hormone Radioimmuno- assay . . . . .	109
a. Antibody to TSH . . . . .	109
b. Radioisotope labeling procedure for rat TSH . . . . .	109
c. Preparation of TSH standards . . . . .	112
d. Assay protocol . . . . .	112
e. Treatment of data obtained from the TSH radio- immunoassay . . . . .	113
f. Validation of the TSH radioimmunoassay . . . . .	113

(Table of Contents, continued)

	<u>Page</u>
III. Radioimmunoassays for Growth Hormone and TSH in Primate Brain Homogenates . . . . .	116
a. Growth hormone . . . . .	116
b. Thyroid stimulating hormone . . . . .	117
B. Description and Care of Animals Used Throughout This Study . . . . .	118
C. Brain Dissection Procedures . . . . .	118
D. Tissue Extraction Methods . . . . .	121
E. Tissue Culturing of Brain Cells . . . . .	123
F. Subcellular Fractionation of Brain Homogenate to Detect the Localization of Pituitary-Like Hormones in the Brain . . . . .	128
G. Immunohistochemistry of Pituitary and Brain Tissues . . . . .	131
H. Column Chromatography of Brain Extracts . . . . .	139
I. Studies to Determine the Immunological Identity of "Brain" Hormones . . . . .	141
J. Biological Activity Studies . . . . .	142
a. Growth hormone bioactivity in brain extracts . . . . .	142
b. Thyroid stimulating hormone bioactivity in brain extracts . . . . .	145
K. Examination of Rat Blood-Brain Barrier Intactness to Pituitary GH and TSH . . . . .	147
L. The Effects of Hypophysectomy on Brain-Based Pituitary-Like GH and TSH . . . . .	150
M. Thyroxine ( $T_4$ ) and $T_3$ Resin Uptake Assays . . . . .	151
a. Measurement of $T_4$ . . . . .	151
(i) Assay procedure for the detection of $T_4$ in serum . . . . .	152
b. Measurement of the radiolabeled $T_3$ resin uptake in serum samples . . . . .	152
(i) Assay procedure for the measurement of the $T_3$ resin uptake in serum . . . . .	153

(Table of Contents, continued)

	<u>Page</u>
N. Alteration of Feedback Control Mechanisms . . . . .	154
I. Target Organ Removal . . . . .	154
a. Oophorectomy . . . . .	155
b. Orchidectomy . . . . .	155
c. Adrenalectomy . . . . .	156
d. Thyroidectomy . . . . .	156
II. Thyroxine . . . . .	157
O. Ontogeny Studies . . . . .	157
P. Statistical Analysis of Data . . . . .	159
5. RESULTS . . . . .	160
A. Pituitary Growth Hormone and Thyroid Stimulating Hormone-Like Immunoreactive Peptide Distribution in Rat Brain . . . . .	160
a. Growth hormone-like immunoreactivity in the brain . . . . .	160
b. Thyroid stimulating hormone-like immunoreactivity in the rat brain . . . . .	165
B. Growth Hormone and Thyroid Stimulating Hormone-Like Immunoreactive Peptide Distribution in Primate Brain	170
a. Growth hormone-like immunoreactivity in the primate brain . . . . .	171
b. Thyroid stimulating hormone-like immunoreactivity in the primate brain . . . . .	171
C. Growth Hormone and Thyroid Stimulating Hormone-Like Immunoreactivity in Extrapituitary Regions Other Than the Rat CNS . . . . .	174
a. Growth hormone-like immunoreactivity in the pineal gland and GI tract . . . . .	174
b. Thyroid stimulating hormone immunoreactivity . . . . .	177



(Table of Contents, continued)

	<u>Page</u>
D. Growth Hormone and Thyroid Stimulating Hormone Immunoreactivity in Medium from Cultures of Monolayered Nervous Tissue . . . . .	177
a. Growth hormone immunoreactivity detected in tissue culture medium . . . . .	177
b. Thyroid stimulating hormone immunoreactivity detected in tissue culture medium . . . . .	183
E. The Effect of Known Hormonal Regulators on the Release of Immunoreactive GH and TSH from Tissue Cultured Cells of the Brain and Anterior Pituitary . . . . .	186
a. Somatostatin added to tissue cultured brain and pituitary cells . . . . .	186
b. Thyroxine (T <sub>4</sub> ) added to tissue cultured brain and pituitary cells . . . . .	186
F. Immunoreactive Thyroid Stimulating Hormone Levels Detected in a Cultured Neuroblastoma Cell Line N <sub>4</sub> TGI	189
G. Fractionation of Brain Homogenates by Ultracentrifugation . . . . .	189
a. Growth hormone: distribution in ultracentrifuged homogenate fractions . . . . .	191
b. Thyroid stimulating hormone: distribution in ultracentrifuged homogenate fractions . . . . .	193
H. Immunoperoxidase Staining of Brain Tissue for Pituitary-Like-Hormone Distribution . . . . .	193
a. Thyroid stimulating hormone . . . . .	193
I. Characterization of "Brain Hormones" by Gel Chromatography . . . . .	201
a. IRGH-like material in the rat brain . . . . .	201
b. IRTSH material in the rat brain . . . . .	201
c. IR pituitary-like hormones from a cloned mouse neuroblastoma line . . . . .	204
J. Immunological Characterization of Brain Pituitary-Like Hormones and Further Immunoassay Validation . . . . .	204

(Table of Contents, continued)

	<u>Page</u>
K. Biological Activity of Brain Pituitary-Like Hormones as Compared to Pituitary Hormones . . . . .	208
a. Growth hormone bioassay . . . . .	208
b. Gonadotrophic bioactivity in brain extracts . . . . .	210
c. Thyroid stimulating hormone bioactivity in brain extracts . . . . .	210
L. Testing the Intactness of the Blood-Brain Barrier in Both Adult and Young Animals . . . . .	215
M. The Effect of Hypophysectomy on Levels of Brain Pituitary-Like Hormones . . . . .	217
a. Growth hormone: effects of hypophysectomy . . . . .	220
b. Thyroid stimulating hormone: effect of hypophy- sectomy . . . . .	220
c. Validation of hypophysectomy procedure . . . . .	225
N. Target Organ Removal: Effect on Levels of Brain Hormones . . . . .	225
a. Growth hormone: effect of removal of target organs . . . . .	225
(i) Adrenalectomy . . . . .	229
(ii) Oophorectomy . . . . .	229
(iii) Orchidectomy . . . . .	234
(iv) Thyroidectomy . . . . .	234
b. Thyroid stimulating hormone: effect of removal of target organs . . . . .	234
(i) Adrenalectomy . . . . .	234
(ii) Oophorectomy . . . . .	241
(iii) Orchidectomy . . . . .	241
(iv) Thyroidectomy . . . . .	241
O. Thyroxine Levels: Effect of an Increase in Circulating T <sub>4</sub> on Serum and Tissue Levels of TSH and T <sub>4</sub> . . . . .	250
P. Ontogeny of Pituitary and Brain Pituitary-Like Hormones	252
a. Growth hormone . . . . .	252

(Table of Contents, continued)

	<u>Page</u>
(i) Growth hormone in brain and pituitary tissues .	252
(ii) Growth hormone in fetal and maternal sera and amniotic fluid . . . . .	256
b. Thyroid stimulating hormone . . . . .	256
(i) Thyroid stimulating hormone in brain and pituitary tissue . . . . .	256
(ii) Thyroid stimulating hormone in fetal and maternal sera and amniotic fluid . . . . .	258
c. Summary of fetal brain levels of immunoreactive GH and TSH data . . . . .	258
d. Immunological characterization of fetal brain hormones . . . . .	258
e. Molecular weight determinations of fetal brain hormones . . . . .	263
6. DISCUSSION OF DATA . . . . .	267
CONCLUSIONS . . . . .	291
REFERENCES . . . . .	293

LIST OF TABLES

	<u>Page</u>
<u>Table 1.</u> Recovery of added amounts of reference preparation GH in normal brain homogenate . . . . .	107
<u>Table 2.</u> Recovery of added amounts of reference preparation TSH in normal brain homogenate . . . . .	115
<u>Table 3.</u> Distribution of immunoreactive GH in adult rat brain (ng/mg protein) . . . . .	161
<u>Table 4.</u> Distribution of immunoreactive GH in adult rat brain (pg/mg wet weight tissue) . . . . .	162
<u>Table 5.</u> Distribution of immunoreactive GH in adult rat brain regions (ng/total brain region) . . . . .	163
<u>Table 6.</u> Distribution of immunoreactive TSH in adult rat brain (ng/mg wet weight tissue) . . . . .	166
<u>Table 7.</u> Distribution of immunoreactive TSH in adult rat brain (ng/mg protein) . . . . .	167
<u>Table 8.</u> Distribution of immunoreactive TSH in adult rat brain regions (ng/total brain region) . . . . .	168
<u>Table 9.</u> Distribution of immunoreactive GH in adult monkey brain (pg/mg wet weight tissue) . . . . .	172
<u>Table 10.</u> Distribution of immunoreactive TSH in adult monkey brain (pg/mg wet weight of tissue) . . . . .	173
<u>Table 11.</u> Distribution of immunoreactive GH in adult rat tissues (ng/mg protein) . . . . .	175
<u>Table 12.</u> Distribution of immunoreactive GH in adult rat tissues (pg/mg wet weight tissue) . . . . .	176
<u>Table 13.</u> Distribution of immunoreactive TSH in adult rat tissues (ng/mg protein) . . . . .	178
<u>Table 14.</u> Distribution of immunoreactive TSH in adult rat tissues (ng/mg wet weight) . . . . .	179
<u>Table 15.</u> Fractionation of brain homogenates by ultracentrifuga- tion of growth hormone: content of fractions (ng/ fraction and ng/mg protein) . . . . .	192

(List of Tables, continued)

	<u>Page</u>
<u>Table 16.</u> Fractionation of brain homogenates by ultracentrifugation of thyroid stimulating hormone: content of fractions (ng/fraction and ng/mg protein) . . .	194
<u>Table 17.</u> Body weight changes over 5-day biological experiment	209
<u>Table 18.</u> Bioassay of pituitary and amygdala extracts . . .	211
<u>Table 19.</u> Evaluation of TSH effectiveness to change the pattern of $^{131}\text{I}$ uptake into the thyroid . . . . .	214
<u>Table 20.</u> Tissue distribution of $^{125}\text{I}$ rat growth hormone injected IV into adult rats and IRGH content of tissues . . . . .	216
<u>Table 21.</u> Tissue distribution of $^{125}\text{I}$ rat thyroid stimulating hormone injected IV into adult rats and IRTSH content of tissue . . . . .	218
<u>Table 22.</u> Distribution of $^{125}\text{I}$ rat growth hormone in fetal and neonatal brain and other tissues after injection of radiolabeled hormone either intraperitoneally into pregnant rats (A) or into 21-day-old fetuses (B), or into neonatal rats (C) . . . . .	219
<u>Table 23.</u> Immunoreactive growth hormone in extracts from discrete brain parts after hypophysectomy . . . . .	221
<u>Table 24.</u> Immunoreactive thyroid stimulating hormone in extracts from discrete brain parts after hypophysectomy . . . . .	224
<u>Table 25.</u> Immunoreactive GH and TSH levels in serum of adult rats . . . . .	227
<u>Table 26.</u> Circulating thyroid hormone levels in serum of adult rats . . . . .	228
<u>Table 27.</u> Effect of adrenalectomy (4 weeks) on brain GH levels	231
<u>Table 28.</u> Effect of oophorectomy (4 weeks) on brain GH levels .	233
<u>Table 29.</u> Effect of orchidectomy (4 weeks) on brain GH levels .	236
<u>Table 30.</u> Effect of thyroidectomy (4 weeks) on brain GH levels	238

(List of Tables, continued)

	<u>Page</u>
<u>Table 31.</u> Effect of adrenalectomy (4 weeks) on brain TSH levels	240
<u>Table 32.</u> Effect of oophorectomy (4 weeks) on brain TSH levels	243
<u>Table 33.</u> Effect of orchidectomy (4 weeks) on brain TSH levels	245
<u>Table 34.</u> Effect of thyroidectomy (4 weeks) on brain TSH levels	247
<u>Table 35.</u> Significance of changes in brain GH after hypophy- sectomy and target organ removal (4 weeks) . . . .	248
<u>Table 36.</u> Significance of changes in brain TSH after hypophy- sectomy and target organ removal (4 weeks) . . . .	249
<u>Table 37.</u> GH- and TSH-like immunoreactive peptides in 21-day fetal CNS . . . . .	262
<u>Table 38.</u> Substances whose levels rise just prior to birth . . . .	282
<u>Table 39.</u> Substances whose levels rise after birth . . . . .	283

## LIST OF FIGURES

		<u>Page</u>
<u>Fig. 1.</u>	The major nuclear centers of the hypothalamus and its neural interconnections with the CNS . . . .	9
<u>Fig. 2.</u>	Frontal section showing relationships of nuclear centers of the hypothalamus . . . . .	11
<u>Fig. 3.</u>	Proteolytic processing of the 30K precursor in pituitary tissue . . . . .	65
<u>Fig. 4.</u>	Relationship between $\beta$ lipotropin amino acid sequence and other peptides of the brain and pituitary . . . . .	76
<u>Fig. 5.</u>	Schematic diagram illustrating the principle of the radioimmunoassay . . . . .	86
<u>Fig. 6.</u>	Standard curve for a radioimmunoassay . . . . .	87
<u>Fig. 7.</u>	Precipitin curve for a monospecific system: one antigen and the corresponding antibody . . . . .	89
<u>Fig. 8.</u>	Determination of the titer of GH antibody binding 35% of the total radiolabeled GH antigen from one test bleeding of 5 guinea pigs (12/18/78) . . . . .	93
<u>Fig. 9.</u>	Computerized output of GH standard curve . . . . .	102
<u>Fig. 10a.</u>	Best fit analysis for GH standard curve data . . . . .	103
<u>Fig. 10b.</u>	Weighted regression analysis of GH standard curve data . . . . .	103
<u>Fig. 11.</u>	GH standard curve plotted as B/BO vs. (ng/ml) . . . . .	104
<u>Fig. 12.</u>	GH standard curve plotted as logit vs. (ng/ml) . . . . .	104
<u>Fig. 13.</u>	Calculation of GH content of unknown samples . . . . .	105
<u>Fig. 14.</u>	TSH standard curve plotted as B/BO vs. (ng/ml) . . . . .	114
<u>Fig. 15.</u>	TSH standard curve plotted as logit vs. (ng/ml). . . . .	114
<u>Fig. 16.</u>	Isolated rat brain . . . . .	119
<u>Fig. 17.</u>	Coronal section of rat brain . . . . .	120

	<u>Page</u>
<u>Fig. 18.</u> Representative samples of 10-day old nervous tissue culture obtained from the amygdala of adult rats .	126
<u>Fig. 19.</u> Fractionation scheme utilized to determine the growth hormone and TSH content of subcellular fractions of rat brain . . . . .	129
<u>Fig. 20.</u> Components of the peroxidase-antiperoxidase complex unlabeled antibody stain used for the detection of rat TSH in brain and pituitary tissue . . . . .	133
<u>Fig. 21.</u> Distribution of immunoreactive growth hormone in the adult rodent brain . . . . .	164
<u>Fig. 22.</u> Distribution of immunoreactive TSH in the adult rodent brain . . . . .	169
<u>Fig. 23.</u> Cumulative immunoreactive growth hormone released from cells of the adult rat amygdaloid nucleus grown in tissue culture . . . . .	180
<u>Fig. 24.</u> Cumulative immunoreactive growth hormone released from cells of the 21-day post-hypophysectomized, adult rat brain regions grown in tissue culture .	182
<u>Fig. 25.</u> Cumulative immunoreactive thyroid stimulating hormone released from cells of the adult rat brain regions and pituitary grown in tissue culture . . .	184
<u>Fig. 26.</u> Cumulative immunoreactive thyroid stimulating hormone released from cells of the 21-day post-hypophysectomized, adult rat brain regions grown in tissue culture . . . . .	185
<u>Fig. 27.</u> Immunoreactive growth hormone released from cells of the adult rat brain and pituitary grown in tissue culture for 24 hours with or without somatostatin	187
<u>Fig. 28.</u> Immunoreactive TSH released from cells of the adult rat brain and pituitary grown in tissue culture for 24 hours with or without L-thyroxine . . . . .	188
<u>Fig. 29.</u> Cumulative immunoreactive thyroid stimulating hormone (TSH) released from the neuroblastoma cell line N <sub>4</sub> TGI . . . . .	190
<u>Fig. 30.</u> Immunohistochemical staining of pituitary thyrotrophs by the PAP technique (x 100) . . . . .	196
<u>Fig. 31.</u> Immunohistochemical staining of pituitary thyrotrophs by the PAP technique (x 400) . . . . .	197



(List of Figures, continued)

	<u>Page</u>
<u>Fig. 32.</u> Control pituitary stained by the PAP technique using absorbed TSH antiserum (x 400) . . . . .	197
<u>Fig. 33.</u> Immunohistochemical staining by the PAP technique of amygdaloid tissue for TSH ( x 1000) . . . . .	198
<u>Fig. 34.</u> Immunohistochemical staining by the PAP technique of amygdaloid tissue for TSH controls (x 1000) . . . . .	199
<u>Fig. 35.</u> Immunohistochemical staining of amygdaloid tissue for TSH (x 400) . . . . .	200
<u>Fig. 36.</u> Elution pattern of lyophilized and reconstituted amygdala extract IRGH and <sup>125</sup> I rat pituitary GH on a Sephadex G-100 column eluted with veronal buffer . . . . .	202
<u>Fig. 37.</u> Elution pattern of lyophilized and reconstituted amygdala extract IRTSH and <sup>125</sup> I rat pituitary TSH on a Sephadex G-100 column eluted with PO <sub>4</sub> buffer . . . . .	203
<u>Fig. 38.</u> Elution pattern of lyophilized and reconstituted N <sub>4</sub> TGI cell extract IRTSH and <sup>125</sup> I rat pituitary TSH on a Sephadex G-100 column eluted with PO <sub>4</sub> buffer . . . . .	205
<u>Fig. 39.</u> Logit log plot demonstrates parallel displacement for serial dilutions of pituitary and CNS growth hormone extracts . . . . .	206
<u>Fig. 40.</u> Logit log plot demonstrates parallel displacement for serial dilutions of pituitary and CNS TSH extracts . . . . .	207
<u>Fig. 41.</u> Histology of the thyroid gland of 14-day hypophysectomized animals treated with brain and pituitary extracts (x 400) . . . . .	213
<u>Fig. 42.</u> Immunoreactive growth hormone present in amygdaloid nucleus extracts of intact and hypophysectomized rats . . . . .	222
<u>Fig. 43.</u> Immunoreactive growth hormone present in caudate extracts of intact and hypophysectomized rats . . . . .	223
<u>Fig. 44.</u> Immunoreactive TSH present in extracts from various brain regions of intact and 28-day hypophysectomized rats . . . . .	226

(List of Figures, continued)

	<u>Page</u>
<u>Fig. 45.</u> Effect of adrenalectomy on GH levels in serum, pituitary, and brain . . . . .	230
<u>Fig. 46.</u> Effect of oophorectomy on GH levels in serum, pituitary, and brain . . . . .	232
<u>Fig. 47.</u> Effect of orchidectomy on GH levels in serum, pituitary, and brain . . . . .	235
<u>Fig. 48.</u> Effect of thyroidectomy on GH levels in serum pituitary and brain . . . . .	237
<u>Fig. 49.</u> Effect of adrenalectomy on TSH levels in serum, pituitary, and brain . . . . .	239
<u>Fig. 50.</u> Effect of oophorectomy on TSH levels in serum, pituitary, and brain . . . . .	242
<u>Fig. 51.</u> Effect of orchidectomy on TSH levels in serum, pituitary, and brain . . . . .	244
<u>Fig. 52.</u> Effect of thyroidectomy on TSH levels in serum, pituitary, and brain . . . . .	246
<u>Fig. 53.</u> The effect of injecting IM 0.4 $\mu$ g $T_4$ /gm body weight daily for 7 days into intact adult rats, on TSH levels in serum, pituitary, and amygdala . . . . .	251
<u>Fig. 54.</u> The ontogenic development of growth hormone levels in the CNS from fetal to adulthood in the rat . . . . .	253
<u>Fig. 55.</u> The ontogenic development of growth hormone levels in the pituitary from fetal to adulthood in the rat . . . . .	254
<u>Fig. 56.</u> The ontogenic development of growth hormone levels in the pituitary and CNS from fetal to adulthood in the rat . . . . .	255
<u>Fig. 57.</u> Levels of GH in amniotic fluid, fetal serum, and maternal serum from 17 days fetal to 2 days post-natal in the rat . . . . .	257
<u>Fig. 58.</u> The ontogenic development of TSH levels in the CNS from fetal to adulthood in the rat . . . . .	259
<u>Fig. 59.</u> The ontogenic development of TSH levels in the pituitary and CNS from fetal to adulthood in the rat . . . . .	260

(List of Figures, continued)

	<u>Page</u>
<u>Fig. 60.</u> Levels of TSH in amniotic fluid, fetal serum, and maternal serum from 17 days fetal to 2 days post-natal in the rat . . . . .	261
<u>Fig. 61.</u> Logit log plot demonstrates parallelism between pituitary and fetal CNS GH . . . . .	264
<u>Fig. 62.</u> Elution pattern of lyophilized 21-day fetal brain and <sup>125</sup> I adult rat pituitary on a Sephadex G-100 column eluted with Veronal buffer . . . . .	265
<u>Fig. 63.</u> Elution pattern of lyophilized 21-day fetal brain and <sup>125</sup> I adult rat pituitary on a Sephadex G-100 column eluted with phosphate buffer . . . . .	266

## INTRODUCTION

### 1. STATEMENT OF PROBLEM

Although the formal study of neuroendocrinology is a relatively new discipline, historical writings have long shown an awareness of interactions between the nervous and endocrine systems. The writings of Aristotle on pseudo-pregnancy described a relationship between hormones and emotional states, and the practice of primitive husbandmen to castrate bulls, thus altering aggressive behavior, are two such examples.

It was mainly the results of Geoffrey Harris's work in the late 1940's which ushered in the new era of neuroendocrinology. The anatomical studies of Green and Harris (1949) demonstrated that a direct connection between the central nervous system (CNS) and the endocrine pituitary gland existed in the form of the portal venous system. The portal veins were shown to drain capillaries that have traversed the median eminence region of the hypothalamus, an area of the central nervous system having complex neural connections with other brain areas. The veins themselves then break up into a capillary plexus in the pituitary which, in its turn, is drained by veins into the general venous circulation. Thus virtually all the afferent blood supply of the anterior pituitary was shown to have first been in contact with the hypothalamus.

Specific examples of the interaction between the central nervous and endocrine systems have been revealed by clinical observations

demonstrating that lesions of the hypothalamus can induce malfunction of the pituitary gland, such as suppression of the release of pituitary growth hormone, as observed by Frohman et al (1968). Other functions attributed to the pituitary gland which are modified by such lesions include the regulation of endocrine systems manufacturing steroids in the adrenals and thyroxine in the thyroid gland, as well as the regulation of other key metabolic processes more directly regulated through the pituitary gland's output of growth hormone and thyroid stimulating hormone.

The importance of the portal venous connection between hypothalamus and pituitary was shown by the subsequent work of Schally et al (1966), and Guillemin (1966) who demonstrated the existence of hypothalamic peptidergic hormones that act to control the release of pituitary hormones. These peptides are produced in the hypothalamus and are transported via the portal vascular system to the pituitary cells on which they exert their action. Gonadotropin releasing hormone (GnRH) appears to modulate the synthesis and release of two pituitary hormones, follicle stimulating hormone and luteinizing hormone; and thyroid releasing hormone (TRH), the secretion of two other pituitary hormones, prolactin and thyroid stimulating hormone (TSH). Both GnRH and TRH have been characterized chemically and synthesized by Enzmann et al (1971) and Matsuo et al (1971), respectively. The hypothalamus was also found to contain peptides that inhibited the release of pituitary hormones. The first of these to be isolated by Brazeau et al (1973), was the tetradecapeptide, somatostatin, which was shown to inhibit the secretion

of pituitary growth hormone (GH), TSH, and prolactin as well as the secretion of glucagon and insulin from the pancreas.

It appears from studies such as those of Krulich et al (1977), Schneider et al (1969), and Müller (1973) that the hypothalamic secretory cells producing such regulatory factors are themselves activated by neurotransmitters of CNS origin such as dopamine and norepinephrine, which are released at synaptic connections from the various afferent neurons known to converge on the hypothalamic region from brain areas such as the hippocampus, amygdala, cortex and thalamus (Szentágothai et al, 1968).

A somewhat simplified picture of the neuroendocrine system can therefore be summarized as involving the stimulation of hypothalamic neurosecretory cells by neurotransmitters, produced in other higher centers of the brain, to produce releasing or inhibitory factors, which regulate pituitary hormone secretion, which in turn largely controls growth and development and functions connected with the adrenals, thyroid, and reproductive glands. On the other hand, the hypothalamic inhibitory and releasing factors have been shown to have roles which are not restricted to the control of pituitary function. As reviewed by Prange et al (1975) in the article "Brain Behavioral Effects of Hypothalamic Releasing Hormones," peptides identical to hypophysiotropic hormones have been found in extrahypothalamic areas of the brain; TRH has been demonstrated in the cerebellum, thalamus, brain stem and cerebral cortex (Winokur et al, 1974, and Oliver et al, 1974); somatostatin has been found widespread throughout the CNS (Brown et al, 1975), and

GnRH appears to be present in the mammillary body, thalamus, amygdala, and hippocampus (Kubek et al, 1979). These findings have led to the suggestion by Krieger et al (1980) that these peptides may also act as synaptic transmitters, i.e., convey information between adjacent nerve cells or neuromodulators, which amplify or dampen neuronal activity. Such a neuromodulator role may include functions in relation to sensory conduction (Renaud et al, 1975), pain mechanisms (Segal et al, 1975), and behavioral responses (Moss, 1979) in various brain regions.

Schaeffer et al (1977) and Arimura et al (1975) have reported the presence of hypophysiotropic hormones outside the central nervous system, most notably in the gastrointestinal tract where, for example, the finding of TRH has led to the speculation that this peptide may modulate the responses of feeding and drinking behavior (Moss, 1979). There is growing evidence that the reverse situation is also true. In addition to the presence of brain peptides outside the CNS, other hormones, classically thought to have a site of origin in, for example, the kidney, such as renin (Haulica et al, 1975), and in the gastrointestinal tract, such as glucagon (Conlon et al, 1979), gastrin (Vanderhaeglin et al, 1975), or vasoactive intestinal protein (Said et al, 1976) and Substance P (Hökfelt et al, 1975), have now also been isolated from the CNS.

More recently there have been reports giving evidence for the presence of the classical anterior pituitary hormones in discrete regions of the CNS. Krieger et al (1977<sup>a</sup>) have reported the finding of adrenocorticotrophin (ACTH) in extrapituitary locations such as the arcuate

nucleus of the hypothalamus and the amygdala. Fuxe et al (1977) have shown prolactin containing cells in the hypothalamus, and Vaudry et al (1978) have demonstrated melanocyte stimulating hormone ( $\alpha$  MSH) in cells of the arcuate nucleus.

Once thought only to contain monoaminergic neurotransmitters or their precursors, the isolation of more than 20 different peptides in the brain in the last ten years has led to a new concept of the functioning of the CNS. Speculation, such as that of Krieger (1980), has led to the hypothesis that peptides, such as those of pituitary origin, exist in the brain as part of a peptidergic neural system, present in addition to the classically described motor, sensory, and autonomic systems.

During this time when ectopic sites of origin for many peptide hormones were being described, Lawrence et al (1974) reported the presence of a factor in human cerebrospinal fluid which actively promotes the in-vitro and in-vivo release of pituitary growth hormone.

In the course of seeking to identify the origin of this growth hormone releasing factor, Pacold et al (1976) found pituitary growth hormone-like immunoreactive material in dispersed rat brain cell cultures.

It is the further characterization of this anterior pituitary-like growth hormone in the CNS that will be the subject of this dissertation. Similar studies will also aim to characterize a pituitary TSH-like material which was also found during the course of this work in rat brain cell tissue cultures. The purpose of this dissertation will



therefore first be to review data drawn from classical studies of the two pituitary hormones, GH and TSH, with a view to characterizing their presence in the rat brain. An attempt will be made using biochemical, anatomical, and physiological methodology to determine whether these particular peptides correspond to their pituitary counterparts or not. The significance of these findings and the findings of other "brain peptides" will be reviewed and discussed.

## 2. RELEVANT LITERATURE REVIEW

### A. THE PITUITARY-HYPOTHALAMIC LINK

An understanding of the anatomy of the pituitary gland and its hypothalamic connections is crucial to the analysis of its physiological functions.

These functions include the secretion of at least 6 well-characterized hormones from the anterior lobe of the gland. Growth hormone (GH) and thyroid stimulating hormone (TSH) have important modulatory effects on carbohydrate, lipid, and protein metabolism, whereas ACTH, a linear polypeptide, once released from the pituitary, exerts its most potent effect on the adrenal cortex, where it stimulates the conversion of cholesterol to pregnenolone, a precursor of steroid hormones such as cortisol. Follicle stimulating hormone (FSH) and luteinizing hormone (LH) are glycoproteins produced in the pituitary and which act on the Leydig cells of the testis to stimulate androgen secretion and on the tubules to induce spermatogenesis in the male. In the female they regulate the ovulatory cycle. In addition, the anterior pituitary cells secrete prolactin, a polypeptide exhibiting a diversity of physiologic actions, including an effect on growth of mammary duct tissue and the maintenance of progesterone output by the corpus luteum of the ovary. Another group of pituitary peptides consists of  $\alpha$  and  $\beta$  melanocyte stimulating hormones ( $\alpha$  and  $\beta$  MSH),  $\beta$  and  $\gamma$  lipotropins ( $\beta$  and  $\gamma$  LPH), and the endorphins which, together with ACTH, originate from a common precursor molecule and exhibit a variety of physiological effects.

Two hormones, vasopressin and oxytocin, are released from the posterior lobe of the pituitary gland and are concerned with the maintenance of volume and tonicity of body fluids, and influence pregnant uterine contractions and post partum milk ejection, respectively.

The pituitary gland, itself long considered the "master gland" for reasons obvious from the preceding discussion, is located at the base of the skull, covered by a tough connective tissue diaphragm and lies in a bony cup, the sella turcica. It is connected by the pituitary stalk, containing vascular and neural elements, to the floor of the hypothalamic region of the diencephalic brain. Fig. 1.

This intimate relationship between the pituitary gland and the hypothalamus develops during embryonic life. The origins of the pituitary can be traced to Rathke's pouch, which begins as a cul-de-sac protruding from the roof of the stomoderm, the posterior nasopharynx. During development, Rathke's pouch loses its connection to the stomoderm and becomes a hollow spheroid collection of cells forming the primitive anterior pituitary. The posterior pituitary evaginates downward from the base of the primitive brain diencephalon as the anterior lobe grows upward from the buccal endoderm, forming the mature pituitary gland linked to the diencephalon by the pituitary stalk. The distinct origins of posterior and anterior pituitary lobes are reflected in their cellular anatomy, the posterior lobe consisting of up to 42% nerve axons and axon terminals whereas the anterior lobe contains a variety of hormone secretory cells and an apparent absence of neural input from the brain.




Figure 1. The major centers of the hypothalamus and its neural interconnections with the CNS.

The origin of the nerve terminals in the posterior pituitary are the supraoptic and paraventricular nuclei in the hypothalamus (Figs. 1 and 2). Nerve tracts descend from these hypothalamic nuclei via the pituitary stalk and the severance of this hypothalamic-pituitary connection causes atrophy and irreversible damage to posterior lobe function. This intimate anatomical link influencing the functioning of the posterior pituitary was defined by Scharrer in 1954, who showed that the two pituitary hormones, oxytocin and vasopressin, originate in the supraoptic- and paraventriculohypophyseal nuclei of the hypothalamus. Experiments in which the neurons of these nuclei were pulse-labeled with radioactive amino acids indicated that the two posterior pituitary hormones and their respective neurophysin-carrier proteins are synthesized as part of separate precursor proteins which are processed into the smaller biologically active molecules while being transported along the axons within the pituitary stalk (Sachs et al, 1969). A study of the development of this neural link by Diakoku in 1971 revealed that by the 18th day of fetal life, a rat already possesses some nerve fibers containing dense core, precursor-filled vesicles in the hypothalamus.

The link between the rat hypothalamus and the anterior lobe of the pituitary is vascular in nature. These connections via the pituitary stalk were first observed by Popa and Fielding in 1930. Although these early studies postulated that the flow of blood in the stalk vessel system was from pituitary to brain, Wislocki et al in 1936 demonstrated that the flow of blood was toward the pituitary. Green and Harris, in 1949,




Figure 2. Frontal section showing relationships of nuclear centers of the hypothalamus.

confirmed this observation and their classic anatomical studies identified the vascular connections as the long portal vessels, which were independent from the blood supply to the brain and upper hypothalamus and which divided into capillary tufts within the floor of the third ventricle of the diencephalon. The embryonic development of such a hypothalamic-pituitary vascular link was examined in the rat by Glydon in 1957, who showed that although the portal veins could be identified in the pituitary stalk at the time of birth, the primary capillary plexus in the hypothalamus did not actually attain full development until 5 days after birth.

In adult animals, loss of this vascular connection leads to a marked decrease in some of the anterior pituitary's functions such as the secretion of the hormone TSH. Transection of the rat pituitary stalk by Cahane in 1936 led to a degenerative change in the histological appearance of both the pituitary and thyroid, the latter being a target organ for pituitary TSH. As already reviewed, the finding of hypothalamic inhibitory and releasing factors controlling anterior pituitary hormone secretion by Schally (1966), Guillemin (1966), and Brazeau (1973) supports this idea that the brain, particularly the hypothalamus, is intimately involved in the functioning of the anterior pituitary.

Mechanisms controlling anterior pituitary secretion, other than those of hypothalamic origin, are those exerted by the target gland hormones such as thyroid, adrenal, and gonadal hormones. Changes in circulating levels of target organ hormones may either inhibit or encourage

the synthesis of pituitary hormones, and in this way an optimum level of hormone is maintained in the general circulation. These "feedback loops", as they are known, are also thought to affect the levels of hypothalamic-hypophysiotropic hormones on exposure of the hypothalamus to changing levels in circulating hormones such as  $T_4$  and the steroids.

This section has dealt with the hypothalamic-pituitary link and its role in the control of pituitary hormone release. The two anterior pituitary hormones, growth hormone, and thyroid stimulating hormone will be discussed in greater depth in the sections which follow, with respect to their molecular structure, normal levels found in the pituitary and serum, mechanism of action on target tissues or metabolic systems, and the regulation of their release from the normal pituitary gland.

#### B. PITUITARY GROWTH HORMONE

Synthesized and secreted by the eosinophylic somatotroph cells of the anterior pituitary, growth hormone cells account for 4% - 10% of the wet weight of the adult anterior pituitary. The hormone is produced in the Golgi area of the somatotroph, and the cell's microtubular system is involved in the movement of hormone-containing granules from this area to the plasma membrane, whose release into the circulation is associated with fusion of the secretory granules with the plasma membrane.

Circulating GH is needed for longitudinal growth of the skeleton, has both insulin-like and anti-insulin like effects on the metabolism of



glucose and enhances the incorporation of amino acid into protein. The hormone also induces lipolysis of stored fats as shown by Raben in 1959, who observed a rise in plasma-free fatty acids after administration of growth hormone intravenously to man.

a. Molecular Structure of Growth Hormone.

Rat growth hormone is a polypeptide with a molecular weight of 46,000 Daltons, an NH<sub>2</sub> terminal phenylalanine, and a sedimentation coefficient, S<sub>20</sub> of 3.21, as shown by Ellis et al (1968). Complete sequencing of rat growth hormone has not been reported; however, it is known from the work of Li et al (1971) that all mammalian growth hormones contain 2 disulfide bridges, one spanning 110 residues and joining cysteine residues 53-164; the other spans only 6 residues and joins residues 181 and 189. There is evidence that pituitary growth hormone contains 50% of  $\alpha$  helix, and all known species, such as human, porcine, and rodent, undergo similar structural transitions in acid or base. Of interest in this regard, two other peptide hormones, placental lactogen and prolactin, show remarkable structural similarities with growth hormone. This is reflected in their similar placement of disulfide bridges and residue homology, and has led to the suggestion of a common evolutionary origin for the 3 hormones by Niall et al (1971), which may have arisen from repeated tandem duplication of a gene coding for a relatively small peptide. These similarities in structure are not, however, reflected in the major biological activities of the 3 hormones. Growth hormone alone possesses the ability to increase the length of the skeleton, whereas

prolactin is found to be essential for such processes as mammary duct development, lactogenesis and, together with luteinizing hormone, for the maintenance of progesterone output of the corpus luteum. Metabolic actions resembling those of GH have been demonstrated for prolactin, but only after relatively large doses, and it is questionable whether prolactin makes any significant contribution to somatotrophic activity under normal circumstances. Placental lactogen, produced by the placenta, is highly lactogenic but cannot induce growth of the long bones. (Wallis et al, 1976 and Kelly et al, 1976) All three hormones have been found to exist in several forms with varying molecular weights in the serum. (Hambley et al, 1972, and Gordon et al, 1973)

The suggestion has been made by Sigel and Lewis (1979) that the predominant form of growth hormone in pituitary extracts is itself a prohormone, and that after limited proteolysis by plasmin (Ellis et al, 1968), smaller molecular weight sequences still retain biological activity. (Li et al, 1974) This heterogeneity with respect to size of circulating human growth hormone (hGH) has been studied by Gordon et al (1973) and Bala et al (1970), these groups showing forms with molecular weights both larger and smaller than monomeric hGH. Some of the larger species dissociate when treated with urea, whereas others require disruption of the interchain disulfide bonds for conversion to a molecular weight comparable to monomer hGH. (Beneviste et al, 1978) Bioassay discloses hGH-like activity in blood which does not correspond to immunoreactivity, indicating the presence of circulating forms with heterogeneous immunogenicity. Enzymatic modification of the major polypeptide

chain by chymotrypsin, papain, or subtilisin results in proteolytic cleavage occurring in the large disulfide loop to produce two acidic isohormones. These exhibit a several-fold higher biological activity than other cleavage products, as shown by Singh et al (1974); however, the mechanism of this enhanced activity is unknown. Frigen and Lewis (1980) have shown that a 20,000 Dalton variant hGH lacks various insulin-like and anti-insulin-like effects found with the principle form of hGH as reported by Goodman (1965); yet it is fully active in promoting growth by the GH bioassay measuring rat tibia.

Human, and perhaps rat growth hormone as usually isolated, can therefore be considered as a complex of proteins, perhaps serving different functions. The diversity of effects exhibited by growth hormone on metabolism and growth may perhaps be accounted for by this heterogeneity. However, as shown from Sigel's (1980) latest work, the specific physiological activities required to make a protein a growth hormone are essentially unknown.

Levels of circulating radioimmunoassayable growth hormone vary depending on such factors as age of the animal and the time of day the sample is taken. These factors are discussed in the next section on rat growth hormone.

b. Growth Hormone Levels in Plasma and Pituitary.

Birge et al (1967) found that GH was undetectable in the rat fetus in either plasma or pituitary before day 19 of gestation. Rieutort, in

1972, speculated that the origin of circulating GH after day 19 until birth at day 21-22 was the fetal pituitary, as growth hormone was absent in hypophysectomized animals, i.e., rats which had their pituitaries removed. Jost (1959), working with encephalotomized fetuses with intact pituitaries, showed a lowering of plasma GH in the fetus, suggesting that the CNS, and in particular the hypothalamus, have an early role on somatotroph development and growth hormone secretion. Hypothalamic control, in terms of that found in the adult, does not seem likely in the fetal rat, as somatostatin, the GH-inhibitory hypothalamic peptide, is undetectable in the CNS until the fifth day after birth. There is evidence that fetal and not maternal levels of circulating GH control the release of pituitary hormone, as Grumbach, in 1974, has shown little or no interchange of any pituitary hormone transplacentally. Gluckman et al (1979), Cornblath et al (1965), and Blasquez et al (1974) have also shown a rise in fetal plasma growth hormone in the last days of gestation of the sheep, rat, and human to levels exceeding maternal plasma hormone levels, indicating that the two circulatory systems are independent in terms of growth hormone levels. Postpartum levels of plasma and pituitary hormone reach adult levels between 15 and 20 days as shown by Ojeda et al (1977).

In the adult rat, a circadian variation in plasma hormone level is observed. Evidence indicates that all the pituitary hormones studied to date are secreted in such an episodic manner throughout the 24-hour day, with both plasma and urine concentrations manifesting a circadian rhythm (Saunders et al, 1976). It is now clear that, in the absence of stress,

whether physiological or psychological, endocrine activity, particularly the secretion of ACTH, GH, and prolactin, is extremely variable and under the control of some CNS mechanism other than a stimulus response and "feedback" system. Because of the absence of confirming evidence for a "steady state" system of neuroendocrine activity, a new hypothesis has developed that proposes that hormone secretion is temporally ordered and that the dominant determinant of this activity in man and other animals is the 24-hour sleep-wake rhythm (Weitzman, 1980). Studies by Weitzman's group have shown that, in man, the release of growth hormone coincides with the onset of definitive sleep, characterized by slow, synchronous electrocortical activity, REM sleep, suggesting that the same mechanism precipitates both events. Saunders et al (1976) showed a more regular pattern of episodic release of GH with intervals 3 hours apart in the rat. Peak levels occurred in late morning, indicating that species-to-species differences to occur.

In the female rat, the amplitude of pulses does not change during pregnancy (Willoughby et al, 1978) or with variation in temperature. This unresponsiveness to external stimuli is typical of circadian rhythms in both man and animals (Zucker, 1980). The significance of circadian rhythms to endocrine systems lies in the need for the synergistic effects of hormones on the precise timing of secretory activity.

Several CNS areas have been implicated in such rhythms, especially the suprachiasmatic and preoptic nuclei and the pineal gland. Destruction of the suprachiasmatic nucleus appears to render animal behavior

aperiodic, and adrenal corticosterone rhythms are also eliminated. Pineal dysfunction has been implicated in cases of infertility where abnormal variations in the ovulatory cycle were the underlying cause (Wurtman, 1980). While these are certainly not the only mammalian circadian clocks, the suprachiasmatic nucleus in particular probably functions to integrate the activity of other oscillating systems that in turn regulate the periodicity of numerous physiological and behavioral systems.

c. Biological Effects of Growth Hormone.

Initial recognition of the pituitary's relationship to growth came from the correlation of growth hormone producing pituitary tumors and gigantism with the pathology of the gland. Pituitary deficiency of growth hormone secretion was also found to be associated with dwarfism. The regulation of growth by growth hormone was first examined in the 1930's when pituitary extracts containing growth hormone became available (Silberberg, 1935). Growth of the mammalian skeleton was shown to occur at the epiphyseal plate where chondrocytes proliferate and synthesize a matrix composed of collagen and sulfated polysaccharides. Kibrick et al (1941) found that narrow epiphyseal plates in hypophysectomized rats became wider after growth hormone was given in vivo, a property which has since been used as a bioassay for GH. Other workers such as Murphy and Daughaday (1956) observed that cartilage metabolism could be measured by the

incorporation of radioactive sulfate into cartilage chondroitin sulfate. Cartilage from hypophysectomized rats had low sulfate uptake, which was increased by growth hormone administration given in vivo.

The classic experiments of Salmon and Daughaday in 1957 examined the mechanism by which growth hormone might stimulate growth. The in vitro addition of growth hormone to rodent excised rib cartilages in tissue culture did not accelerate radioactive sulfate incorporation; however, incorporation was stimulated by a component of normal serum. This sulfation factor was absent in the serum from hypophysectomized animals and was restored to serum by the administration of growth hormone to these hypophysectomized animals. From these observations, it was hypothesized that growth hormone stimulated skeletal growth indirectly through generation of a circulatory "sulfation factor" or somatomedin, as it was later termed, which acted directly to cause cartilage proliferation.

Van Wyk et al (1972) found initially that this active component in whole serum behaved as a large protein. After acid ethanol extraction, the active material had a molecular weight between 4,000 and 12,000 D., suggesting that somatomedin was aggregated or bound to a larger carrier protein. Two such carrier proteins were isolated by Kauffman et al (1977), and "somatomedin" was found to be not one but a family of small peptides. Three of these were named Somatomedins A → C, with molecular weights ranging from 5,000-7,000 Dalton by Van Wyk et al in 1974. Two other insulin-like growth factors have also been shown to stimulate cartilage proliferation by Ginsberg et al (1979). These growth factors resemble

proinsulin in amino acid sequence and tertiary structure but have limited cross-reactivity with insulin in binding to receptor sites on target tissues, as demonstrated by Blundell et al (1978).

Structurally similar to proinsulin, the somatomedins have also been shown to have anabolic, insulin-like actions on fat and muscle by Clemmons et al (1974) and Uthne et al (1974), as well as their property to enhance cartilage proliferation.

It is thought that growth hormone stimulates skeletal growth, in vivo, indirectly by regulating the level of the somatomedins. A contradiction to this idea is found in the observation that somatomedin levels are lower in the young animal than in the adult. This has led to speculation that growth may be less dependent on somatomedins or that somatomedins may be more available in the young due to reduced binding to a specific serum carrier protein or to an increase in somatomedin receptor sites as shown by Rosenfeld et al (1979). However, growth hormone excess in humans and rats is associated with an increase in somatomedins as shown by Firlanetto et al in 1977. Regulation of somatomedin production appears to occur mainly in the liver and reflects nutritional (Phillips et al, 1978) and hormonal status, possibly insulin and GH levels (Hintz et al, 1978), of the animal. Levels of somatomedins in perfusates of livers from hypophysectomized animals are below levels in normal animals and are restored to normal by treatment with growth hormone in vivo, as shown by Schalch et al (1977). Thus, the liver may respond to available hormone levels in altering the levels of circulating somatomedins.



In addition to somatomedin being a mediator of growth hormone's effects on collagen synthesis, it appears to mediate other actions of growth hormone, specifically, its anabolic effects on skeletal muscle. Salmon and DuVall (1970) have provided evidence that somatomedin is more potent than native growth hormone in increasing leucine incorporation into muscle protein. Earlier work by Kostyo et al (1966) had shown that leucine incorporation into protein, in excised diaphragm muscle from normal rats, to be twice that of tissue removed from hypophysectomized rats. Treatment of these animals without pituitaries with growth hormone restored amino acid incorporation into new protein. Korner et al (1965) showed increases in the total number of ribosomes in rat liver cells, an increased capacity of ribosomes for incorporating amino acids into protein, and an elevation in messenger RNA in vivo after growth hormone administration. Wikland et al (1980) have studied the response to growth hormone on amino acid transport and protein synthesis in vivo in the rat diaphragm and find that stimulation is transient and followed after about 3h by a period of refractoriness. In contrast, they found a much longer lasting stimulatory effect of growth hormone on protein synthesis in tissues of hypophysectomized rats. These "insulin-like" effects of growth hormone suggest that changes in responsiveness of tissues to growth hormone, or more likely to somatomedin, play a physiological role in the regulation of protein metabolism in muscle tissue.

As with effects on amino acid transportation, so also with respect to plasma free fatty acid levels, there is an early "insulin-like" action shortly after intravenous injection of growth hormone. Birnbaum and

Goodman (1979) showed that levels of free fatty acids fall within the first 1/2h of injection before showing the secondary rise typical of the lipolytic action of the hormone. Somatomedins have been shown to stimulate incorporation of free fatty acids into lipids. (Clemmons et al, 1974) This early antilipolytic effect of growth hormone could be mediated by somatomedin, or, Birnbaum and Goodman (1976) postulate, it may result from a decreased activity of protein kinase in the cytosol.

The delayed rise in plasma free fatty acids after growth hormone administration can be blocked by administration of glucose. Also giving GH simultaneously with insulin, a potent inhibitor of free fatty acid release from adipose tissue, obliterates the lipolytic action. (Merimee, 1979) It is possible that the inhibiting effect of glucose is mediated via stimulation of endogenous insulin release. Fain et al (1971), using isolated fat cells in the presence of dexamethasone, have demonstrated a requirement for RNA synthesis and for new protein synthesis in the lipolytic action of the hormone.

Growth hormone also has both an "insulin-like" and "anti-insulin" effect on the metabolism of glucose. An "insulin-like" effect such as a fall in blood glucose levels has been seen within 10 minutes of an I.V. injection of bovine growth hormone into a normal rat by Goodman (1967). An increase in glucose uptake and glucose oxidation due to an increase in cell permeability is postulated to account for this fall in serum glucose; however, Martin and Young (1965) have shown that very small concentrations of growth hormone depress glucose uptake when added to preparations of

excised rat diaphragm. The "insulin-like" effect of growth hormone on glucose metabolism may therefore be merely a pharmacological effect.

There is greater unanimity with regard to the less immediate effects of growth hormone administration on glucose metabolism, namely, its "contra-insulin" and its "insulinotropic" effect, i.e., enhancement of pancreatic release of insulin in response to a variety of stimuli as shown by Martin et al (1967). Daughaday and Kipnis (1966) compared the rate constant for glucose disappearance from human serum under control conditions with that observed when an I.V. glucose load was administered 2h after the administration of growth hormone (5 mg. I.V.). The clearance rate of glucose from the serum fell from control values after GH administration, substantiating a "contra-insulin" effect for the hormone. Precisely in what manner this "contra-insulin" effect is mediated is not clear.

A second aspect of the action of growth hormone on glucose metabolism is the "insulinotropic" action of the hormone. Pretreatment of normal subjects with human growth hormone results in an increase of insulin release in response to a large number of secretagogues such as glucose and arginine, as shown by Rimoin et al (1968). In vitro data from Martin et al (1967) also supports a direct effect of growth hormone on the  $\beta$  cell of the pancreas. Isolated islets from hypophysectomized rats have a significantly reduced insulin content, output, and synthesis, which can be corrected by growth hormone treatment.

The biological effects of growth hormone can therefore be viewed as the result of a dual action: those of growth hormone per se, and those actions of growth hormone dependent on somatomedins. Factors controlling the secretion of growth hormone by the pituitary are unusually complex; however, most stimuli appear to act via a neural mechanism through the hypothalamus to the pituitary. Such stimuli will be reviewed in the next section.

d. Regulation of Growth Hormone Release.

The direct release of growth hormone appears to be under the control of two substances: a growth-hormone-releasing factor, which has yet to be isolated, and an inhibitory agent, somatostatin.

While searching for the releasing factor, Brazeau (1973) observed that the addition of crude hypothalamic extract to cultures of rat pituitary cells decreased the secretion of growth hormone. Further purification resulted in the isolation of the linear tetradecapeptide, somatostatin. This peptide has been located by Sétáló et al (1978) in the external layer of the median eminence around the capillary loops of the portal vessels which eventually lead to the pituitary. Somatostatin has also been located in substantial amounts by similar immunohistochemical methods in the ventromedial nuclear area of the hypothalamus by Hökfelt et al (1977<sup>a</sup>), as well as other areas of the CNS.

Evidence for some form of "short-loop feedback" control mechanism for growth hormone involving somatostatin comes from experiments with

hypophysectomized rats performed by Baker et al (1976). A reduction in somatostatin levels in the median eminence was observed after pituitary removal, suggesting that this fall in inhibitory factor levels could be directly due to insufficient circulating growth hormone, and Hoffman et al (1977) claim to have been able to restore, partially to normal, levels of somatostatin by administration of exogenous growth hormone to hypophysectomized animals. Whether GH acts directly on somatostatinergetic neurons in the hypothalamus to restore protein synthesis (Karatsuka et al, 1979), or indirectly through somatomedin has not been determined. Baker et al (1976) also caution that the partial recovery of somatostatin levels could be due to the regeneration of neurons destroyed during hypophysectomy.

The mechanism by which somatostatin inhibits the release of growth hormone, induced by prostaglandins, dibutyryl cyclic AMP, and theophylline, is possibly due to inactivation of adenylyl cyclase. A rise in cyclic GMP due to somatostatin administration has also been observed in rat somatotropes. Takahara et al (1980) have recently proposed that the release of growth hormone induced by  $\gamma$ -aminobutyric acid (GABA), a neurotransmitter, may also be partly due to the inhibition of hypothalamic somatostatin release.

Evidence linking other neurotransmitters such as norepinephrine, dopamine, and serotonin with the release of GH has been reviewed by Martin (1973). These more indirect mechanisms of controlling growth hormone release are currently thought to be mediated via any of three

neural centers: the ventromedial nucleus (VMN) and arcuate nucleus (AC) of the hypothalamus, and the limbic system which includes the hippocampus and amygdaloid nuclei of the brain.

Although somewhat of an oversimplification, norepinephrine can be thought of as the dominant neurotransmitter for stimuli reaching the VMN, dopamine for the AC, and serotonin for stimuli mediated via the limbic system. One can thus divide stimuli for growth hormone release on the basis of which of these three systems they may be activating.

When lesions are placed in the VMN, suppression of growth hormone release has been observed to occur by Frohman et al (1968). If the neurotransmitter content in this area of the hypothalamus is norepinephrine, then excitation can be said to be via  $\alpha$  adrenergic mechanisms. Vasopressin (Heidingsfelder, 1968), exercise (Hansen, 1971), cholecystokinin, and renin (Vijayan, 1979) are thought to induce growth hormone release at the level of the VMN via such  $\alpha$  adrenergic mechanisms. Blackard et al (1968) also showed that GH release, mediated by insulin-produced hypoglycemia, could be prevented by blockage of  $\alpha$ -adrenergic receptors by phentolamine, an inhibitor of  $\alpha$  adrenergic transmission. The exact method by which norepinephrine and adrenergic blocking agents act to regulate GH secretion is unknown but could be due to an effect on the production of releasing factors in the hypothalamus. Ascending norepinephrine inputs to the hypothalamus have been shown by Jonsson and Hökfelt (1972) to terminate directly on the arcuate nucleus in the rat, and could therefore have some effect on somatostatinergic neurons.

This same group showed, by histochemical means, that dopaminergic nerve terminals surround the primary capillary plexus of the portal vascular system, and that the arcuate nucleus is rich in dopamine. Conflicting results are found in the literature concerning the effect of dopamine on GH release. Administration of L-dopa by Kansal et al (1972) to rhesus monkeys led to the release of GH from the pituitary; however, no such release was observed if L-dopa was administered to rats (Kato et al, 1973). Müller et al (1967) and Boyd et al (1970) thought that L-dopa is taken up by the monkey dopaminergic terminals in the arcuate nucleus from the circulation and converted to dopamine by dopa decarboxylase. Increased storage of the neurotransmitter could then be postulated to lead to the release of some growth-hormone-releasing hypothalamic factor into the portal system. However, more recent work by Steiner et al (1978) argued against the view that GH release by L-dopa was mediated by a dopamine receptor mechanism in the hypothalamus. Their work and the observation by Brown et al (1976) suggest that dopamine receptors are not found in the median eminence of the hypothalamus but only in the pituitary. This suggested to these authors and others such as Ben-Jonathan et al (1977) that dopamine in the median eminence may serve as a neurohormone carried by the portal vessels to the pituitary rather than as a neurotransmitter acting on the hypothalamic releasing and inhibitory factors.

In summary, the present data on the effects of the catecholamines, norepinephrine, and dopamine on GH release suggest that  $\alpha$ -adrenergic receptors in the hypothalamus stimulate the release of GH, whereas dopamine receptors may inhibit GH, this area being somewhat controversial, and

wide species differences may also exist.

The third area of the brain associated with growth hormone release is the limbic system. Having a role involving the maintenance of wake emotional and behavior states, the limbic system also appears to play an important role in slow wave sleep. As previously noted in humans, this sleep state is associated with a rise in GH levels. Since nerve terminals of the limbic system terminate in the hypothalamus and are, in general, serotonergic, the neurotransmitter serotonin may mediate the growth hormone release seen with slow wave sleep in man.

There are a number of stimuli to GH release from the pituitary, whose locus of action is unclear. These include stress, protein depletion, and the administration of morphine derivatives as shown by Greenwood et al (1966), Schalch (1969), and Pimstone et al (1966). As with stimulation, there are other conditions that inhibit growth hormone release such as obesity, elevation of free fatty acid levels, glucocorticoids, and hypothyroidism, whose mechanism of action is likewise unknown. (Pimstone, 1966; Unger, 1965; and Frantz, 1964) Fragments of ACTH (4-10) and ACTH (18-39),  $\alpha$  MSH and  $\beta$  MSH also can be shown to inhibit growth hormone secretion in the rat. Here, the mechanism of action studied by Bruni et al (1977) appears to be mediated via the hypothalamus, not the pituitary, as they reported a failure to elicit hormones in vitro from pituitaries.

Evidence also exists for a "short loop feedback" mechanism of growth hormone itself on both the hypothalamic and pituitary levels. For example, growth hormone pellets implanted into the hypothalamus can cause a



reduction in circulating pituitary GH levels (Voogt et al, 1970). Exogenous growth hormone can diminish the growth hormone response to insulin, exercise, or arginine in the rat (Müller et al, 1966). Similarly, Hagen et al (1972) observed a dampening of GH release to known stimuli such as arginine infusion in normal volunteers administered human growth hormone.

In summary, factors that affect the release of growth hormone from the pituitary are complex, some being mediated via hypothalamic inhibitory and possibly releasing factors, and others whose mechanisms of action remain unknown.

The next section will review the relevant literature relating to thyroid stimulating hormone (TSH), another hormone which, like growth hormone, is derived from the pituitary gland.

### C. PITUITARY THYROID STIMULATING HORMONE (TSH)

Thyroid stimulating hormone (TSH) is the principal hypophyseal hormonal regulator of thyroid growth and the biosynthesis and secretion of the thyroid hormones, thyroxine ( $T_4$ ) and triiodothyronine ( $T_3$ ).

Using immunohistochemical methodology, Moriarty (1973) showed that thyrotrophs constitute 3%-5% of the pituitary cell population. Polygonal cells, often appearing in clusters predominantly in the anteromedial portion of the gland, they contain small cytoplasmic granules. In the absence of the thyroid, these cells show a marked ballooning of the rough endoplasmic reticulum cisternae and a decrease in secretory granules,

implying a disruption of the transport processes of granules through the Golgi apparatus. (Farquhar et al, 1954) The concentration of TSH, which is synthesized on ribosomes and stored in the granules, may only be 5% of normal in a hypothyroid rat. The molecular structure of this pituitary hormone will be dealt with in the next section.

a. Molecular Structure of Pituitary TSH.

Thyroid stimulating hormone is a glycoprotein containing 15% carbohydrate and having a molecular weight of 23,300 Daltons. Treatment of the molecule with dilute acid by Cornell et al (1973) resulted in its dissociation into two dissimilar subunits or peptide chains, designated alpha ( $\alpha$ ) and beta ( $\beta$ ). The  $\alpha$  subunit consists of 96 residues, which are almost identical to  $\alpha$  chorionic gonadotrophin (CG),  $\alpha$  luteinizing hormone (LH), and  $\alpha$  follicle stimulating hormone (FSH) subunits. The  $\beta$  subunit is larger than the  $\alpha$ , being composed of 110 amino acid residues, and is structurally dissimilar from the  $\beta$  units of CG, LH, and FSH. Both subunits contain many half cystine residues, assumed by Pierce (1974) to be in a covalent-disulfide bond form.

Each TSH molecule contains three oligosaccharide moieties attached to the peptide via asparagine residues. The sugars and amino sugars include fructose, mannose, galactose, glucosamine, and galactosamine. Two of the three carbohydrate moieties are associated with the  $\alpha$  subunit, where sialic acid residues are similarly located. The significance of the carbohydrate constituents, in terms of biological expression on TSH

metabolism, has not been determined. The native structure of the molecule depends upon noncovalent interactions between the two subunits. A high glycine and proline content and the intrachain bonding preclude any long distances of  $\alpha$  helix. An isolated  $\alpha$  chain has no biological activity, and a separated  $\beta$  chain only a minimal ability to bind and activate TSH receptor sites. Recombination of  $\alpha$  and  $\beta$  chains fully reactivates the hormone. Studies by Vaitukaitis et al (1973), generating antisera to both intact molecules and to both subunits, revealed that most immunological determinants reside on the hormone-specific  $\beta$  chain. Although TSH $\beta$ , LH $\beta$ , and CG $\beta$  share extensive sequence homology, little immunological cross reactivity has been encountered. A common ancestral molecule for the four different hormones has been proposed to account for the similarities between individual  $\alpha$  and  $\beta$  subunits.

Levels of TSH present in the circulation vary with age and are dependent in the young rat on the development of the hypothalamic pituitary-thyroid axis.

b. Thyroid Stimulating Hormone Levels in Plasma and Pituitary

The development of the hypothalamic pituitary thyroid axis has been examined in the rat by several groups: Dussault et al (1973), Cons et al (1975), Kieffer et al (1976), and Friedman et al (1979). In all cases, pituitary TSH was found in low levels just before birth and postnatally reached a maximum at day 15 before falling slightly to adult levels.

At term, fetal rat serum TSH is found to be slightly higher than maternal serum levels. There is a falloff in hormonal levels of serum after birth followed by a surge to reach adult levels by day 12. A rise in human serum TSH is also observed at birth and is sustained for 2-3 weeks postpartum (Conklin et al, 1973). In this study and those of the rat, maternal TSH was found not to traverse the placenta.

Adult rats have been reported to show diurnal variation in their levels of serum TSH, a similar situation to that reviewed for GH except that Parket et al (1976) do not believe a link exists with sleep and TSH. Leppäluoto et al (1974) have reported diurnal variations in serum, peak levels being attained by late evening, the lowest levels in the afternoon, and a relatively high plateau between 8-10 a.m. The mechanisms responsible for this rhythm are not completely understood, although Niles et al (1979) have suggested that melatonin or the precursor of melatonin, n-acetyl serotonin, in the pineal, may have a role in the diurnal rhythm of TSH. Pinealectomy was found to increase circulating TSH in animals kept under one-hour-light/23-hour-dark lighting conditions, and a similar increase in TSH levels was observed after active immunization against melatonin and n-acetyl serotonin.

The overall concentration of TSH in the periphery reflects a dynamic equilibrium between TSH secretion and TSH distribution, degradation, and excretion. This regulated level of circulating TSH is dependent on many factors; however, the condition of the thyroid gland, which requires TSH for the maintenance of euthyroidism, is one of the most important factors.

c. Biological Effects of TSH

Effects of TSH on the thyroid gland include hyperplasia and hypertrophy, stimulation of  $I^-$  transport into thyroid cells and its incorporation into the thyroid hormones  $T_4$  and  $T_3$ , and the hormonogenesis and secretion of these two hormones. In order to comprehend the importance of pituitary TSH to the thyroid, a brief description of the thyroid and the production of  $T_4$  and  $T_3$  will be given.

The thyroid gland consists of two lobes connected by a narrow isthmus crossing the rat trachea. These lobes are composed of follicles, which are spheroid structures consisting of a single layer of epithelial cells enclosing a cavity or follicle lumen which is filled with a viscous, proteinaceous solution, the colloid. The shape of the follicle cells vary; however, in any one follicle, the cell height is almost uniform. In general, low cells are characteristic of hypoactive follicles, as seen in hypophysectomized animals, whereas high cells are hyperactive follicles. Administration of pituitary TSH to hypophysectomized animals can restore the height of thyroidal follicle cells to normal (Tice, 1974).

The specific physiological processes of the thyroid include trapping of iodide required for the thyroid hormones, the synthesis by the follicle cell of a specific thyroid glycoprotein, thyroglobulin; storage of thyroglobulin in the follicle lumen, reabsorption of thyroglobulin into the cell, degradation of this protein, and release of the thyroid hormones  $T_3$  and  $T_4$  into the circulation.

The kinetics of TSH action appear to be biphasic on iodide transport. An early depressive phase lasts 1-2h and is followed by overall enhancement, as shown by Williams et al (1975) using mouse thyroid tissue. The depressive phase is thought to be due to an acute increase in iodide efflux from the follicular cells, possibly reflecting a general increase in membrane permeability, and the enhancement phase by an increase in the unidirectional clearance of iodide by cells presumably reflecting the induction of iodide carriers.

The binding of iodide to protein and the formation of iodothyronine is a rapid process. Two enzymatic systems are involved in this step, the generation of  $H_2O_2$  and the oxidation of  $I^-$  by a peroxidase (Taurog, 1970). As no effect of TSH on peroxidase activity has been observed, it is probable that the  $H_2O_2$ -generating system is the hormone's site of activation as proposed by Ahn et al (1970).

Thyroglobulin is synthesized in the rough endoplasmic reticulum and passes via the Golgi apparatus to the cell surface in vesicles before reaching the follicular lumen by exocytosis. The influence of TSH in this process is observed in hypophysectomized animals where a decrease in the size and contents of the lumen is noted. Ekholm et al (1975) have demonstrated this phenomenon using radioautography to show that a rapid stimulation by TSH discharged radio-labeled thyroglobulin into the colloid space.

The first step in the release and secretion of thyroid hormones is marked by pseudopod formation by the apical plasma membrane protruding

into the follicle lumen and by endocytosis followed by portions of colloid reappearing in follicular cells. Pseudopods are formed within two minutes of TSH addition to incubated thyroid slices, and 10 minutes later  $T_4$  can be measured in the external medium (Ekholm, 1979).

Degradation of thyroglobulin in the follicle cell requires that hydrolytic enzymes gain access to the content of the colloid droplets; the source of these enzymes is the lysosomes, as shown by Wollman et al (1964). Lysosomes are found in all parts of the cell but accumulate towards the luminal cell surface after TSH stimulation (Seljelid, 1967). The consensus, therefore, is that proteolytic enzymes reach the thyroglobulin by colloid-droplet/lysosome fusion, whereby degradation of the protein and release of thyroid hormones are achieved.

The precise way in which TSH achieves this regulation of thyroid protein synthesis is not clear. Adiga et al (1971) showed that certain effects of TSH on the thyroid such as the stimulation of RNA synthesis and RNA polymerase in porcine thyroid could be inhibited by protein or RNA synthesis inhibitors, indicating the possibility that TSH acts at the level of transcription. Although a direct effect on the synthesis of thyroglobulin messenger RNA has not been observed by Larry et al (1971) in rats, the continuous presence of TSH seems to be required to sustain the expression of the thyroglobulin gene, as postulated by Pavlovic-Hournac et al (1971).

The stimulation of incorporation of amino acids into thyroid proteins in vivo (Pavlovic-Hournac et al, 1973), and in vitro (Lecocq et al,

1972) is thought to be an effect at the translational level by such authors. However, a direct stimulatory effect of cyclic adenosine monophosphate (cAMP), the postulated intracellular mediator of TSH actions (Rapaport, 1976), on protein synthesis in an acellular system, devoid of nuclei, has only been observed at unphysiological concentrations of the cyclic nucleotide by Wägar (1973).

With regard to thyroid metabolism, TSH enhances thyroid cell respiration, mainly mitochondrial, as shown by Dumond (1971), and enhances glucose uptake and its overall metabolism (Dumont et al, 1965). Dumont's group also showed an augmentation in the overall pentose phosphate cycle and the incorporation of glucose into protein after TSH administration. Glycogen and fatty acid levels in lipids decrease, reflecting an activated catabolism. TSH has been shown to enhance the incorporation of precursors into most phospholipids, as demonstrated by Scott et al (1970) using  $^{32}\text{P}$  incorporation into monophosphatidylinositol, phosphatidates, phosphatidylethanolamines, phosphatidylserine, and in some cases phosphatidylcholine. No effect of TSH was observed on  $^{32}\text{P}$  incorporation into sphingomyelin or phosphatidylglycerol in isolated cells or slices from pig thyroid.

Extrathyroidal actions of TSH, the significance of which are not well understood, include the ability to stimulate lipolysis (Hartree et al, 1972) and intracellular levels of prostaglandins (Vale et al, 1972) in human adipose and rat pituitary tissue.

The hormone can therefore be seen to have a wide range of effects both on the thyroid and other tissues. The secretion of TSH from the



pituitary gland can be regulated by several factors which will be discussed in the next section.

d. Regulation of TSH Release from the Pituitary.

The hypothalamic-pituitary-thyroid axis has been shown to be a dynamic "negative feedback" control loop, the activity of which is influenced by the peripheral consumption of thyroxine ( $T_4$ ) and modulated by hypothalamic control of pituitary responsiveness, by thyrotropin releasing factor (TRH).

Early experiments with pituitary stalk sectioning by Boler & Schally et al (1969) and Burgus & Guillemin et al (1969) led to the identification and synthesis of thyrotropin releasing hormone (TRH), a small peptide consisting of the 3 amino acids, Glu-His-Pro-NH<sub>2</sub>, which was inactivated within minutes by serum. Secreted into the portal blood vessels by hypothalamic cells, TRH travels to the pituitary and binds to the thyrotroph membrane on specific receptor sites with high affinity, stereospecificity, and reversibility, as shown by Wilber et al (1973<sup>a</sup>). The magnitude of TSH release from the pituitary cells is dependent on the increase in membrane binding of TRH (Wilber, 1973<sup>b</sup>). TRH stimulation also causes augmented TSH synthesis, as judged by the incorporation of <sup>14</sup>C alanine or <sup>14</sup>C glucosamine into pituitary slices (Wilber, 1971). An interesting immunohistochemical report of Moriarty et al (1978) suggests that TRH may also be located in the thyrotroph secretory granule within the pituitary gland. Whether this material is of hypothalamic origin is not known.

Control of TRH secretion is not well understood. Ablation of supra-optic and paraventricular nuclei results in suppression of thyroid function, and destruction of the median eminence and arcuate nucleus results in a fall of TSH and  $T_4$  levels (Gear, 1951; and Hefco et al, 1955<sup>a</sup>). Attempts to correlate the hypothalamic content of TRH with metabolic status indicate minimal or no variation (Montoya et al, 1975), although possibly the augmented TSH secretion induced in rats by cold exposure is via an elevation in TRH production (Hefko et al, 1975<sup>b</sup>). The synthesis of TRH is stimulated also by dopamine and norepinephrine and reduced by serotonin, as shown in vitro using mouse hypothalamic tissue by Grimm and Reichlin (1973). Further evidence of monoaminergic transmitter control of TRH comes from the work of Krulich et al (1977), who showed that systemic injection of  $\alpha$  methyl-p-tyrosine and inhibitors of dopamine  $\beta$  hydroxylase cause decreases in serum TSH, as does blockade of  $\alpha$  adrenergic receptors. Stimulation of dopamine receptors with apomorphine and L-dopa will similarly cause decreased release of TSH, leading to the conclusion that central noradrenergic systems have a stimulatory effect on serum TSH via TRH. The opiate peptide, leu-enkephalin, also appears to inhibit TSH via TRH as shown by May et al (1979).

Primary control of TSH secretion is, as described, from the positive input of TRH as well as the negative control exerted by the thyroid hormone ( $T_4$ ). The inability of pharmacological quantities of exogenous  $T_4$  to prevent an increase in TRH in response to cold exposure suggests that  $T_4$  does not alter TRH secretion (Montoya et al, 1974), but rather has an effect on the pituitary thyrotroph. Reichlin et al (1973), however,

showed that intrahypothalamic implants of  $T_4$  do depress thyroid function; the controversy of where  $T_4$  exerts its negative feedback control over TSH release still exists.

$T_4$  stimulates in the pituitary thyrotroph the synthesis of nucleic acids and protein which, in turn, affects the secretion of TSH. Observation by Samuels et al (1973) has shown that the effects of  $T_4$  on the pituitary are mediated by effects on transcription of genetic material to increase protein synthesis. Thus, a metabolic effect of thyroid hormone, rather than actual circulating level of hormone, modulates pituitary responsiveness.

Other regulators of TSH secretion than TRH and  $T_4$  are also known.

Knigge (1964) has shown that infusion of TSH itself into the cat hypothalamus will inhibit thyroid hormone release, and Motta et al (1969) have shown decreases in TRH after exogenous TSH was administered to thyroidectomized rats. Although TSH levels are not modified by activity, starvation, circulating free fatty acids, or amino acids, hypoxia can cause a lowering of TSH in rats, which is probably a stress phenomenon (Curbelo et al, 1979). This group also showed a decrease in circulating TSH after adrenalectomy.

The last two main sections of this literature review have dealt with the characterization of the two pituitary hormones, GH and TSH, whose presence in the brain is the subject of this dissertation. In order to introduce the next section, which reviews the present knowledge

on peptides present in the brain, it is necessary at this point to introduce the "amine precursor uptake and decarboxylation theory" of A. Pearse. Thyroid releasing hormone has been located in a number of other locations besides the hypothalamus, in particular, the pancreatic islet D cells. Known to contain primarily the hormone somatostatin, they are also known to contain TRH-like immunoreactive material (Morley et al, 1978). The finding of this and other peptides in both brain and gut has led to the "APUD" theory of Pearse (1966), Pearse (1977), and Pearse and Takor (1979).

#### D. THE AMINE PRECURSOR UPTAKE AND DECARBOXYLATION THEORY (APUD)

The idea that cells widely dispersed in the nervous and gastrointestinal system share the same ability to secrete similar peptides has been examined by Pearse. These cells, known as "APUD" cells, were postulated to be derived originally from the embryological neural crest. Such cells included thyroid C cells, ultimobranchial body C cells, suprarenal medullary cells, melanocytes, mast cells, pituitary corticotrophs, melanotropes, somatotropes, pinealocytes, and a wide range of endocrine cells in the gastrointestinal tract and pancreas. These regulatory molecules produced by such cells may enter the bloodstream to reach and alter the activity of distant target cells, an endocrine action; diffuse locally through interstitial spaces to reach and influence groups of neighboring cells, a paracrine action; or cross a synaptic junction to activate or inhibit the postsynaptic cell, a neurocrine action (Zimmerman, 1979).

Although the APUD-neural crest theory suggested the existence of a widely dispersed neuroendocrine system, offering a unifying concept for understanding many tumor-associated endocrine syndromes and the phenomenon of ectopic hormone production by a wide range of tumors; it seems evident from more recent studies that the possession of APUD characteristics by cells does not imply necessarily a neural crest origin. Cheng et al (1974) conclude that APUD endocrine cells of the pancreas and intestine arise from the endoderm of the developing gut wall, and that the development of "APUD" characteristics appears to be one of several pathways of differentiation available to primitive and uncommitted epithelial cells.

However, APUD characteristics do seem to be associated with tissues producing small polypeptide hormones, and it was therefore not surprising to find such hormones in both brain and gut. The next section of this review of literature discusses in more detail such findings, and demonstrates that many of the peptides found in brain and gut appear to have significant roles to play in the functioning of the brain.

#### E. PEPTIDES FOUND IN THE CENTRAL NERVOUS SYSTEM

The idea of chemicals being involved in neuronal communication is credited to Elliot (1904). His theories foresaw later results from micro-electrode recordings of single cells and the microiontophoretic application of peptides to neurons which showed that results could not be explained completely by the presence of purely electrical junctions.

The concept of the presence of chemical neurotransmitters, which convey information between adjacent nerve cells, and chemical neuromodulators, which amplify or dampen neural activity but which are not responsible for direct transfer of a nervous signal from the pre- to post-synaptic element, has developed over the last 60 years. Monoaminergic catecholamines such as dopamine and serotonin are known to be among the class of substances known as neurotransmitters. Whether peptides recently found in the brain such as renin and the hypothalamic peptides act as neurotransmitters or neuromodulators is not as yet known. It is clear that many such peptides affect the ionic potential of a receptive membrane as will be seen in the following review, and that these effects are not identical to those produced by conventional neurotransmitters.

Hökfelt et al in 1979 proposed that the peptide receptor is restricted to specialized areas of the post-synaptic membrane. Several theories have been discussed by this group, using the evidence available in 1979, for a neuromodulator role for neuropeptides. One such theory entertains that a peptidergic nerve terminal in the brain could release its product into the extracellular spaces adjacent to fenestrated capillaries. The peptide is then carried via the vascular system to a target, where it interacts with a receptor. Another possibility would be a synaptic arrangement, where the peptide and possibly a neurotransmitter are released into the synaptic cleft where interaction occurs with receptors on the post-synaptic membrane. A third possibility would be a parasynaptic arrangement, whereby the peptide, on release from the nerve terminal, would diffuse and interact with receptors at sites other than the post-

synaptic membrane, possibly nearby neural membranes, or on the peptidergic terminal itself. These are some of the ideas at present being discussed to explain the presence of the large number of peptide hormones being discovered in the CNS.

They could also be present in the brain due to incomplete suppression of the genome or as receptor-bound hormone originating from secretory tissue in areas other than the brain. In the following discussion of present-day knowledge of brain peptides, beginning with a review of the presence in extrahypothalamic areas of hypothalamic hormones, some of these points will be discussed.

#### HYPOTHALAMIC HORMONES

The localization of peptidergic neurons in the CNS had indicated an emerging pattern with respect to distribution. Thyroid-releasing hormone, somatostatin, and gonadotropin-releasing hormones have been found in cell bodies and nerve terminals, both intra- and extrahypothalamically. It appears that besides mediating anterior pituitary secretion, these peptides may also play a role in brain function by mediating extrapituitary events. They will be reviewed individually here.

##### a. Gonadotropin-Releasing Hormone (GnRH).

GnRH is a decapeptide with a blocked aminoterminal composed of pyroglutamic acid, which stimulates pituitary gonadotropins and was found in the hypothalamus by Matsuo and Schally (1971), and Burgus and

Guillemin (1971). Kubek et al (1979) have identified immunoreactive GnRH in extrahypothalamic areas such as the mammillary body, thalamus, amygdaloid nucleus, and hippocampus. Very small amounts or no hormone are found in the human cerebellum and cortex. Baker et al (1975) reported the hormone to be present in fibers leading to extrahypothalamic areas of the rat brain. Extraction and analysis of this material showed molecular similarities to the original hypothalamic hormone. Studies by Parker et al (1978) showed the presence of neurons, the terminals of which end in the median eminence, which contained GnRH-like material associated with dense cored granules. On fractionation, 52%-65% of the hormone was associated with the synaptosomes. In hypophysectomized, ovariectomized rats, the intraventricular application of synthetic GnRH produces a specific action on mating behavior in estrogen-primed animals (Foreman and Moss, 1977).

b. TRH (Thyroid-Releasing Hormone).

TRH-like material has been found in varying concentrations in the rat brain extrahypothalamically by several groups including Oliver et al (1974) and Winokur et al (1974). Although the greatest concentration for any one brain part is found in the hypothalamus, over two-thirds of the total brain content of TRH is found outside of the hypothalamus. Areas rich in TRH include the cerebellum, thalamus, brain stem and cerebral cortex. Little or no hormone is found in the pineal, amygdala, hippocampus, or caudate. These two groups differ as to whether TRH is present or absent in the pituitary; however, Moriarty et al (1978), as



previously reported, have shown histochemically, TRH associated with pituitary thyrotrophs.

The TRH-like material found extrahypothalamically possesses the same physiochemical and biological properties as the hypothalamic peptide (Nemeroff et al, 1979). The content of TRH in the CNS is not affected by hypophysectomy, thyroidectomy, or adrenalectomy (Nemeroff et al, 1979), whereas hypothalamic levels of TRH fall after such treatment. It was found by Nemeroff (1979) that hypothalamic TRH levels could be restored with a combination of  $T_4$  plus TSH treatments. In the rat brain, it appears that the synthesis of this tripeptide occurs in situ, independent of the possible transport of the hormone from the median basal hypothalamus (Jackson et al, 1977). Studies by Bennett et al (1975), Parker et al (1978), and Schaeffer et al (1977) indicate the presence of TRH in rat median eminence and hypothalamic synaptosomal fractions, and that depolarizing procedures can release the hormone from such preparations. Burt et al (1975) have shown the presence of binding receptors in rat brain membrane preparations. These receptors occur ubiquitously throughout the regions of the brain, except for their absence in the cerebellum, and give credence to a possible neurotransmitter, rather than neuromodulator, role for TRH.

A study by Renaud et al (1975) on single neuronal activity in various brain sites showed that microiontophoretic application of TRH, somatostatin, and GnRH reduced neuronal activity in the rat. Studies by Brown et al (1975) show that, although this effect may be similar for all

three hormones, the biological effects on the CNS suggest opposing mechanisms for the individual peptide. However, the independence of all these effects from those on the pituitary suggest a direct action on the CNS by the hypothalamic hormones.

Pentobarbital toxicity is reversed by TRH but enhanced by somatostatin (Prange et al, 1974), whereas a strychnine effect on convulsions is potentiated by TRH but antagonized by somatostatin (Moss, 1979). It has been shown that TRH not only increases norepinephrine turnover in the brain (Keller et al, 1974), but also potentiates the behavioral effect of L-dopa (Plotnikoff et al, 1972). Plotnikoff showed that this latter ability was reduced by castration or pinealectomy. It appears from the work of Metcalf (1974) that TRH markedly antagonizes the sedation and hypothermia induced by a wide variety of centrally acting depressants such as ethanol, and that adrenergic blocking agents such as propranolol do not alter this antagonistic effect. Cott et al (1976) have further shown that these effects of TRH are independent of TSH and  $T_4$  influences, since hypophysectomy and  $T_4$  administration failed to reverse these responses. Prompt emergence from ether anesthesia is also characteristic in rats following TRH treatment (Moss, 1979). Other behavioral effects seen after intraventricular injection of TRH by Schenkel-Hulliger et al (1974) include muscle tremor, excitation, tail lifting, and pilo erection. A noted resemblance of these effects to those induced by morphine withdrawal has been noted, and it is of interest that brain areas where naloxone, a competitor for morphine binding sites, precipitates withdrawal shaking, parallel the sites of TRH-induced shaking and endogenous

sites of TRH in the brain. Another behavioral effect by TRH administration was shown by Vijayan et al (1977) to be an apparent suppression of feeding and drinking behavior. Most catecholamine levels remain unchanged after TRH treatment; however, Nemeroff et al (1979) report an increase in norepinephrine release from presynaptic terminals.

The range of CNS effects which TRH exhibits is therefore large. Apart from the CNS, thyroid releasing hormone has been identified in the spinal cord (Hökfelt et al, 1975<sup>a</sup>), retina (Schaeffer et al, 1977), in the islet cells of the pancreas (Morley et al, 1978), and in the placenta (Shambaugh et al, 1978). Recent evidence from Engler et al (1980) has shown that circulating TRH in the neonatal rat is derived from the pancreas and GI tract.

The third hypothalamic peptide to be isolated extrahypothalamically is the GH-inhibiting factor or somatostatin.

c. Somatostatin.

This linear tetradecapeptide has been found widespread throughout the CNS using histochemical and radioimmunoassay techniques. Although the highest concentration is in the hypothalamus, other areas contain substantial amounts (Parsons et al, 1976). Subcellular studies by Epelbaum et al (1977) indicate that 70% of the somatostatin is localized in the synaptosome fraction. This peptide has also been isolated in both cells and fibers of the hypothalamus as well as from cell bodies throughout the rat brain, in particular the limbic system and cortical areas as

shown by the immunohistochemical studies of Hökfelt et al (1977<sup>a</sup>).

Somatostatin's presence in synaptosomes and its ability to act on single neuron units to inhibit spontaneous electrical activity imply a role in neural function (Brown et al, 1975). Tan et al (1977) have suggested that somatostatin inhibits the release of a pool of  $\text{Ca}^{++}$  taken up by synaptosomes. As  $\text{Ca}^{++}$  influx is closely related to nerve depolarization, neurosecretion in nerve cells, and secretory processes in endocrine cells, it is possible that somatostatin influences neuronal excitation through modification of  $\text{Ca}^{++}$  transport at the synaptic membrane level. Injection of somatostatin into the amygdaloid nucleus has been studied by Rezek et al (1977). They revealed a dose-dependent dissociation of effects on behavior and motor control. Lower doses induced a behavioral excitation accompanied by a variety of tremors, whereas higher doses prevented the expression of a normal behavioral pattern due to the severity of disturbances in motor control and the sleep-waking cycle. The effect on behavior is almost opposite to that of TRH. This includes a reduction in locomotion (Segal et al, 1974), prolongation of pentobarbital sedation (Prange et al, 1974), a decrease in the duration of strychnine-induced seizures, and sedation and hypothermia induction (Brown et al, 1975). In general, therefore, somatostatin exhibits a CNS-depressant form of action as shown by Segal et al (1975) and Prange (1974). Somatostatin also potentiates the behavioral effects of L-dopa and can act as a partial antagonist-agonist on opiate receptors.

From this review it is apparent that, as with the other hypothalamic peptides, a dual role exists for somatostatin. Extrahypothalamically, it

acts as a modulator of nervous function and hypothalamically as a peptide inhibiting the release of growth hormone, TSH, prolactin, glucagon, insulin, gastrin, and cholecystokinin.

Somatostatin-like immunoreactivity has also been located in thyroid C-cells (O'Brian et al, 1979), and in the stomach (Arimura et al, 1975) and pancreas of the rat by Luft et al (1974). In the pancreas, the anatomical positioning of islet D-cells between  $\alpha$  and  $\beta$  cells of the islets suggests that they may be involved in some form of control over insulin and glucagon secretion (Dubois, 1975) and (Goldsmith et al, 1975). Hökfelt et al (1977<sup>a</sup>) have suggested the possibility that somatostatin is present in a population of peripheral sympathetic noradrenergic neurons, a fact which would violate Dale's principle of "one neuron, one transmitter" to suggest concomitant storage of a small biologically active peptide and an amine in the same neuron.

Another group of hormones found in the CNS are those originally detected in the GI tract.

d. Bombesin.

Originally detected in frog skin and mammalian gut by Wharton et al (1978), bombesin has also been isolated from mammalian hypothalamus by Villareal et al (1978). The "brain" hormone is able to stimulate the secretion of growth hormone, glucagon, and prolactin, and is chemically and immunologically distinct from frog skin bombesin. If injected intraventricularly, a lowering of body temperature is noted and an

elevation in blood sugar levels also occurs (Brown et al, 1977).

A second GI tract hormone to be found in the CNS is glucagon.

e. Glucagon.

This pancreatic islet hormone has recently been detected in several areas of canine brain by Conlon et al (1979). Found in concentrations 100× the level in plasma, the glucagon-like peptide seems to be a variety of gut glucagon.

A second pancreatic hormone recently found in the CNS is insulin.

f. Insulin.

Insulin has been demonstrated in the brain in concentrations higher than plasma by Havrankova et al (1979). This "brain" insulin is indistinguishable chemically and immunologically from pancreatic insulin and does not vary with extreme changes in plasma insulin, indicating that brain insulin and its receptors are probably regulated independently of their counterparts elsewhere. Whether this hormone is synthesized locally is not known. Specific binding sites for insulin are localized on axons and axon terminals in the external median eminence and hypothalamic arcuate nucleus, suggesting that a discrete population of nerve terminals exist as potential targets for the direct effects of insulin (Van Houten et al, 1980). This group suggests that, after the initial binding, in the median eminence, insulin may be internalized and transported by retrograde flow to the arcuate nucleus.

Highest concentrations of insulin receptors are found extrahypothalamically in the cerebral cortex and olfactory bulb. Other areas such as the hippocampus and amygdala have fairly high numbers of receptors, whereas the retina, spinal cord, and cerebellum have the least. Speculation that insulin receptors are preferentially located within elements of the limbic system leads to an association between emotional behavior, feeding, and defense, characteristically limbic system functions, with insulin as a possible regulator of these actions.

The hormone has been shown to promote, in cells of the hypothalamus, local glucose uptake (Smith et al, 1972) and subsequent metabolism (Panksepp, 1973), to be capable of altering the electrical activity of hypothalamic neurons (Oomura, 1976), and to stimulate hypothalamic norepinephrine release (McCaleb, 1979).

A note of caution before assigning insulin as a peptide indigenous to the CNS is that apparently no close correlation between insulin levels and receptor content exists in the different regions so far studied.

Two other GI tract peptides, gastrin and cholecystokinin, are also found in the CNS.

g. Gastrin and Cholecystokinin (CCK).

Vanderhaeglin et al (1975) first reported the finding of a peptide in hog, dog, and human CNS which reacted with antigastrin antibodies. From chromatographic elution patterns, they concluded that the molecular weight of this peptide was less than gastrin (2-17) and more like that of

cholecystokinin. Since then, it is apparent that both gastrin- and cholecystokinin-like activity are present, but that they exhibit distinct distribution patterns in the CNS.

As reported by Rehfeld et al (1979), gastrin is found in the hypothalamus, pituitary, and peripheral nerves. Cholecystokinin distribution has been studied histochemically by Innis et al (1979), Larsson et al (1979<sup>a</sup>), and Strauss et al (1977). This peptide was found to be widely distributed in nerves throughout the body. In the CNS, neocortical as well as hippocampal, amygdaloid nucleus and hypothalamic regions are richly endowed with cholecystokinin-containing nerves and cell bodies.

Rehfeld et al (1979), using <sup>35</sup>S methionine incorporation, have shown in vivo biosynthesis of cholecystokinin in human and hog cortical cells. Brain CCK has been shown by Dockray (1980) to be mainly the COOH-terminal octapeptide (CCK 8) of the GI hormone. Cholecystokinin and CCK 8 are present in amounts comparable to those found in the GI tract. Malesci et al (1980) have shown that the brain contains at least two enzymes involved in the conversion of CCK to CCK 12 and CCK 8; the twelve residue peptide appears to be more biologically active than CCK 8.

In vivo and in vitro effects of CCK have been reported on gonadotropin, prolactin, growth hormone, and TSH release in the rat by Vijayan et al (1979). Intraventricular injections, followed by assay of plasma hormone levels, showed an increase in growth hormone and a decrease in TSH levels. The mechanism of release of pituitary hormone appears to be mediated via a hypothalamic intermediary. Gastrin exhibits similar



properties in the release of growth hormone and TSH (Vijayan et al, 1978). A possible role for CCK in appetite regulation, as proposed by Gibbs et al (1973), is more controversial, as results may be due to the peripheral effects of the peptide.

Evidence for serotonergic and opiate modulation of brain cholecystokinin has been presented by Morley et al (1980).

Vasoactive Intestinal Polypeptide (VIP), another GI tract peptide, has been isolated from CNS tissue.

#### h. Vasoactive Intestinal Polypeptide (VIP).

Vasoactive Intestinal Polypeptide (VIP) is a 28-amino-acid residue peptide originally found in porcine duodenum by Said et al (1972). It possesses similar amino acid sequences and pharmacology to secretin and glucagon.

Its actions peripherally, as shown by Said and his co-workers, are systemic vasodilation, stimulation of myocardial contractility, glycogenolysis, lipolysis, stimulation of insulin secretion, and inhibition of gastric acid production.

Said et al (1976) reported the presence of VIP in intra- and extra-hypothalamic sites in the CNS, in neural tumor cell lines, as well as in the GI tract. A recent paper by Sims et al (1980) has demonstrated histochemically the presence of four major VIP-containing systems in the brain. A major intracerebral cortical system, another innervating the

central amygdala and nucleus of the stria terminalis, a pathway originating in the suprachiasmatic nucleus of the hypothalamus, and another in the central gray of the midbrain. Specific cell body staining was seen in the limbic and neocortex, the suprachiasmatic nucleus of the hypothalamus, and the central gray of the midbrain. Giachetti et al (1977) have shown VIP's localization in synaptosomes, and a  $K^+$ -evoked,  $Ca^{++}$ -dependent release has been demonstrated by Emson et al (1978). Specific binding of VIP and stimulation of cAMP release has been demonstrated in brain synaptic membranes by Taylor et al (1979).

Intraventricular administration of VIP has been shown to stimulate the release of growth hormone, prolactin, and luteinizing hormone (Borghi et al, 1979) from the pituitary. Direct application of VIP to the pituitary has no effect, but Samson and McCann et al (1980) have shown that VIP receptors are present in the hypothalamus and that VIP can act to stimulate LH release from hypothalamic synaptosomes obtained from ovariectomized female rats. They propose a neuromodulator role on GnRH in the hypothalamus, as the mechanism for release of LH from the pituitary.

The heavy VIP innervation of the central nucleus of the amygdala and central gray of the midbrain is similar to the reported distribution of the opiate, enkephalin, and for neurotensin (Basbaum et al, 1978). As hypothesized for these two peptides, VIP may also have a role in pain sensation.

Evidence for synthesis within certain brain areas comes from the work of Besson et al (1979), who studied the effects of deafferentation

of the hypothalamus on levels of VIP. Two weeks after deafferentation, the VIP concentration of the caudal hypothalamus and stalk were lowered by 40%. In contrast, no change was shown in cortical VIP levels, suggesting that the peptide is derived from neural cell bodies mainly outside the median basal hypothalamus.

The next peptide to be discussed, Substance P, was known to be present for many years in other tissues before its recent isolation from the CNS.

i. Substance P.

Von Euler et al (1931) first isolated from equine tissues, other than brain, a factor which they designated Substance P and which caused intestinal contractions and lowered blood pressure by vasodilation. Substance P was not, however, more thoroughly characterized until 1971 with the establishment of the amino acid sequence of a sialogogic peptide isolated from bovine hypothalamus (Chang et al, 1971).

Using immunohistochemical techniques, Hökfelt et al (1977<sup>b</sup>) reported Substance P to be present in the central and peripheral nervous system, showing an uneven distribution in the CNS except in the cortex, cerebellum, and in the sympathetic ganglia of the guinea pig, cat, and rat. Emson et al (1978) have reported the presence of Substance P in the medial and central nuclei of the amygdala and the bed nucleus of the stria terminalis. Efferents lead from the amygdala to the hypothalamus via the stria terminalis. Highest concentrations are found in the substantia

nigra. The occurrence of Substance P in neurons of the medullary raphe nucleus projecting to the spinal cord is intriguing, since these same nuclei also have been shown to contain serotonin by Hökfelt's group, yet another possible violation of Dale's Law of "one neuron, one transmitter."

The levels of Substance P in the brain stem and spinal cord are higher in fetal than adult tissues, as shown by Gilbert et al (1979). Other areas of the brain such as the cortex, caudate, and hypothalamus show a rise in peptide level at 14-28 days postpartum and demonstrate a similar developmental pattern to that of VIP and noradrenaline.

Substance P has been shown to have some neuroendocrine functions, particularly those mediated by dopamine. It has been shown by Fisher et al (1974) that Substance P can release LH and FSH but not GH, PRL, or TSH, in vitro, from pituitary slices, whereas Euler et al (1979) showed stimulation of prolactin, growth hormone release, and inhibition of the release of insulin, in vivo.

This peptide fulfills several criteria to support its postulated role as a transmitter of pain impulses (Otsuka et al, 1975); however, it has yet to be shown to be synthesized in sensory nerves mediating pain transmission. A close analogy exists between Substance P and the opiate met-enkephalin. They both are present in the substantia gelatinosa of the spinal cord, and it may be that Substance P is a transmitter of brain signals carried by sensory nerves into the spinal cord and then relayed to the brain thalamus, where pain sensing terminals are located. Because

of their immediate proximity, it has been postulated that the endorphins may produce their analgesic effects by suppressing Substance P release in the spinal cord (Marx et al, 1979).

Evidence that Substance P modulates synaptic transmission of cat motoneurons comes from the work of Krivay et al (1979). This effect appears biphasic: low doses inhibiting and high doses facilitating synaptic transmission, suggesting a neurotransmitter role. However, the slow time-course of some Substance P responses (neurotransmitters usually induce rapid responses), and the finding that the peptide depresses responses to other putative neurotransmitters, suggests that the peptide may not be a true neurotransmitter (Otsuka et al, 1975). Krnjevic et al (1977) have proposed two operationally distinct actions of Substance P to explain these functional differences: a neurotransmitter-like action mediating excitatory synaptic events which are both rapid in onset and brief in duration, and actions of Substance P such as those involving the depression of glutamate-activated events, more of a neuro-modulator effect. The presence of Substance P, but not its exact role in the CNS, has clearly yet to be completely characterized.

During the course of the purification of Substance P from bovine hypothalamic extracts, a new tridecapeptide was detected, sequenced, and synthesized by Carraway and Leeman (1973) and is reviewed below.

j. Neurotensin.

Neurotensin has been found by Kobayashi et al (1977) to be in the

highest concentrations in the median eminence, but also to be present in the amygdala, nucleus accumbens, and caudate. Low levels are found in the cortex, hippocampus, and pineal, and trace amounts are present in the cerebellum. Carraway et al (1976) showed that receptor binding concentrations differ from this hormonal distribution pattern in that highest concentrations are in the thalamus and cortex as well as the hypothalamus. Adrenalectomy or hypophysectomy does not seem to alter overall levels of the hormone in the brain, and the intestine, stomach, and pancreas have higher levels than those found in brain tissue (Leeman et al, 1977).

Intraventricular doses of neurotensin were given by Carraway et al (1976) and found to lower body temperature, accentuate barbiturate sedation, cause hypotension, increase vascular permeability, cyanosis, and a marked hypoglycemia.

Disagreement exists as to its role in modulation of the secretion of the hormones of the pituitary. Rivier et al (1977<sup>a</sup>) have shown that neurotensin is able to stimulate the pituitary secretion of growth hormone, prolactin, luteinizing hormone, follicle stimulating hormone, and ACTH, postulating the effect is via a hypothalamic intermediate. However, in a recent abstract by Nemeroff et al (1980), intraventricular neurotensin modulated neurotransmitter activity in the mesolimbic but not the nigrostriatal dopamine system and reduced the secretion of TSH, GH, and prolactin.

Neurotensin can therefore be summarized as being present in both the CNS and GI tract and appears to have some biological effects on brain tissue.

The next peptides to be reviewed are those originally associated with the renin-angiotensin system of the kidney which have been isolated from CNS material over the last five years.

k. Renin-Angiotensin.

Angiotensin II is an octapeptide formed by the action of converting enzymes on the decapeptide Angiotensin I, which is normally the product of the enzymatic action of renin from the kidneys on angiotensinogen. Almost all parts of this renin-angiotensin system have now been isolated from brain homogenates.

A renin-like activity was found in dog, human, and rat brain by Haulica et al (1975), exhibiting molecular and immunological similarities to kidney renin. Renin levels were measured in hog brain by Hirose et al (1980), who showed that the pineal was the richest source of renin-like activity, followed by the hypothalamus, cerebellum, and amygdala.

Reid et al (1977) reported that the pH optimum of brain renin differed from that of kidney renin and that the renin-like brain peptide behaved more like Cathepsin D, a nonspecific acid protease. However, more recent studies have shown that "brain renin" is indistinguishable from renal renin both biochemically and serologically (Slater et al, 1980, and Inagami, 1978). Studies also demonstrated that the distribution of renin and Cathepsin D were distinct and showed no overlap, as shown by Govindrajan et al (1974) and Daul et al (1975). Renin has been identified by Slater and Zimmerman (1980) in many neurons throughout the human

brain by immunohistochemical methods.

Angiotensinogen, renin's natural substrate, was first located in brain tissue by Ganten et al (1976) and shown to have a molecular weight of 40,000-60,000 D, presumably too large to cross the blood brain barrier and, therefore, it was concluded by the authors to be endogenous to the brain. Wallis et al (1980) have shown a wide distribution for angiotensinogen in the rodent brain and that these levels can be modulated by glucocorticosteroids. The hypothalamus, preoptic area, and periaqueductal gray regions showed the most consistent reduction in brain angiotensinogen after 14 days' adrenalectomy, other brain regions showing little or no effect. Treatment with corticosterone reversed the effect of adrenalectomy in the four brain regions.

The "converting enzyme" for Angiotensin I to Angiotensin II is present in rat and human brain in large amounts and is widely distributed (Yang et al, 1972, and Poth et al, 1975).

Fuxe et al (1976) found positive fluorescence for angiotensin II-like material in the median eminence and amygdaloid nucleus. The octapeptide is localized in neuronal cell bodies, fibers, and terminals in the paraventricular and perifornical area of the hypothalamus, with highest concentrations in the terminals of the hypothalamic dorsomedial and basal nuclei.

Bennet et al (1976) and Siret et al (1977) have identified high affinity binding sites for <sup>125</sup>I angiotensin II on rat brain membranes which were saturable and reversible. Specific receptors exist only in



the midbrain, thalamus, and hypothalamus. Baxter et al (1980) have shown that these binding sites reach maximum numbers, ten times those found in adult animals, in the first week of post-natal life. Levels of renin and angiotensin II are also higher at birth than those found in adult brains.

Nephrectomy does not change the levels of angiotensin II in the brain, indicating an independence from the renal system. This peptide has been shown to exert several actions on the CNS after intracerebral injections. These include stimulation of drinking, a rise in blood pressure, and vasopressin and adrenocorticotropin (ACTH) release after intraventricular injection (Fitzsimon, 1972). The sites of action of intracerebral angiotensin II appear to be the same as for the circulating peptide, that is, close to the ventricles in the septal region, anterior hypothalamus, and preoptic areas (Johnson and Epstein, 1975). Destruction of tissue at these sites abolishes the rise in blood pressure and increase in water uptake, stimulated by angiotensin II (Anderson et al, 1975). Intracarotid injection of saralasin acetate, an inhibitor of angiotensin II activity, decreases blood pressure in genetically hypertensive rats (Phillips et al, 1977).

In this review, the site and mode of action of components of the brain iso-renin-angiotensin system have been discussed. Vasopressin, the posterior pituitary hormone, will now be reviewed.

## 1. Vasopressin.

Traditionally a posterior pituitary hormone, vasopressin is produced by neuronal cell bodies in the supraoptic and paraventricular nuclei of the hypothalamus and has recently been reported in extrahypothalamic areas of the brain. Kovács et al (1979<sup>a</sup>) have studied the stimulatory effect of minute quantities of vasopressin injected into certain limbic-midbrain structures such as the hippocampus, and showed facilitation of memory consolidation processes. Oxytocin, another posterior pituitary hormone, was shown to attenuate passive avoidance behavior in a time-dependent manner, suggesting that memory consolidation is influenced by this peptide (Kovács et al, 1979<sup>b</sup>).

Whether the presence of these peptide hormones extrahypothalamically represents in-situ synthesis or axonal transport from the hypothalamus is still unresolved, but results suggest that limbic-midbrain areas are involved in the mediation of action of both posterior pituitary neuropeptides on memory consolidation.

Not only posterior pituitary but anterior pituitary hormones are among those peptides being found localized in the CNS. The discovery of a family of related peptides in both brain and pituitary demands an introduction to their relationships in terms of molecular characteristics.

m. A common precursor for some pituitary hormones.

For many years it was known that the bioactive form of ACTH was only 39 amino acids long and yet antibodies to this peptide also reacted with a protein six times larger. The nature of this large precursor revealed new structural relationships among a group of peptide hormones.

Eipper et al (1975) and Mains et al (1977) found by studying the synthesis of ACTH, using pulse chase experiments with pituitary tumor cells incubated in  $^3\text{H}$  tyrosine-containing media, that the biologically active ACTH 4.5K fragment was cleaved from the middle of a 31K.ACTH precursor molecule (Fig. 3). At the same time, a close relationship had been observed between ACTH and the  $\beta$  lipotropin ( $\beta$  LPH) families of peptides. They had been shown to be released in roughly equivalent amounts by normal pituitary tissue, and also Moriarty (1973) and Pelletier et al (1977) had shown that the two peptides were localized in the same secretory granules in pituitary cells. It had also been noted that the 31K precursor ACTH molecule contained peptide segments similar to both ACTH and the opiate  $\beta$  endorphin. Further work by Mains et al (1977) showed that the peptides of an ACTH biosynthetic intermediate, plus the peptides from  $\beta$  LPH, summed to equal the sequences from pro-ACTH/endorphin, thus establishing a common precursor relationship.

The  $\beta$  LPH can further be broken down to yield  $\beta$  endorphin and also  $\beta$  melanocyte-stimulating hormone ( $\beta$  MSH). At the other end of the large precursor, a sequence of seven amino acids corresponded to the structure of  $\alpha$  MSH, being the breakdown product of the ACTH part of the molecule,




Figure 3. Proteolytic processing of the 30K precursor in pituitary tissue.

together with a 21-amino-acid peptide known as "CLIP." In summary, ACTH, the lipotropins, and the endorphins all originate as controlled cleavage products from a common 31,000 D glycopeptide.

The two lobes of the pituitary, anterior and intermediate, which contain the common precursor, exhibit differences in the levels of ACTH and  $\beta$ -lipotropin-related peptides, apparently due to different cleavages and other post-translational processing steps (Eipper and Mains, 1980). These peptides are not only found in the pituitary, but the precursor protein and many of its breakdown products have been found in the hypothalamus.

Highest concentrations of ACTH,  $\alpha$  MSH,  $\beta$  lipotropin, and the opiate peptide  $\beta$  endorphin have been found in the hypothalamus. However, significant amounts are also found in the limbic system and to a lesser extent in the midbrain, cortex, and cerebellum (Krieger et al, 1977<sup>b</sup>).

Cell bodies containing these peptides have been isolated, histochemically, lateral to the arcuate nucleus of the rat hypothalamus, and positively staining fibers have been shown projecting from this area to the amygdala, thalamus, and the stria terminalis by Zimmerman et al (1978) and Watson (1977 and 1978). Similar immunohistochemical studies on rat pituitaries, using antisera directed against  $\beta$ -lipotropin,  $\beta$ -lipotropin/ $\beta$ -endorphin, ACTH and  $\alpha$  MSH showed immunostaining of the same or similar cells in the anterior lobe of the gland (Jacobowitz et al, 1978, Martin et al, 1979).

Parallel synthesis of ACTH,  $\beta$  lipotropin, and  $\beta$  endorphin has been reported by Krieger and Liotta (1980) in the pituitary, and Gramsch et al (1980) have shown parallel patterns of distribution of  $\beta$  lipotropin,  $\beta$  endorphin, ACTH, and  $\alpha$  MSH in the pituitary and brain. Gramsch also showed that whereas  $\beta$  endorphin and  $\alpha$  MSH are predominant in the brain, in contrast, their respective precursors,  $\beta$  LPH and ACTH, were predominant in the pituitary of both rat and man. Unlike the pituitary, no cells or fibers in the CNS appear to contain both ACTH and  $\beta$  lipotropin together (Krieger et al, 1977<sup>b</sup>).

Local differences for concentrations of  $\beta$  LPH and ACTH have also been noted by Krieger et al (1979) in the regions of the CNS, such as lower  $\beta$  LPH than ACTH levels in hypothalamus, hippocampus, and amygdala, and higher  $\beta$  LPH than ACTH levels in the midbrain, pons, spinal cord, and medulla.

These data may be indicative of important differences in post-translational processing of the common precursor molecule between hypothalamus and pituitary tissues, and even between brain regions. Since the pituitary and hypothalamus are on opposite sides of the blood brain barrier, these two separate sources of peptides are therefore possibly regulated separately and exhibit differences in rates of synthesis, biological half-lives, and types of precursor present. Mains (1980) speculates that this compartmentalization may have evolved to allow the pituitary to stimulate glucocorticoid production by means of ACTH without over-producing opiate activity, and to allow the hypothalamus to achieve the maximum brain effect of its common precursor, derived opiate

peptides, without causing cortisol release.

Even though it seems from this discussion that the two pools of hormones are separate entities, the presence of pituitary-like hormones in the brain such as ACTH has been further investigated to prove this situation.

n. Brain ACTH.

The pattern of distribution of ACTH in the brain, as reported by Krieger (1977<sup>a</sup>), persists after hypophysectomy, adrenalectomy, and dexamethasone suppression. Immunohistochemical studies by Pelletier et al (1980) also confirm this observation. After long-term hypophysectomy, the level of hormone initially falls slightly in the median eminence but not in the median basal hypothalamus or other tissues (Liotta et al, 1978). An enhancement in ACTH concentration is, in fact, observed after removal of both pituitary and median eminence in the remaining hypothalamic tissue. This data by Liotta tends to eliminate a pituitary source for brain ACTH, whether by contact with hypophyseal tissue or retrograde transport from the pituitary. Other lines of evidence to demonstrate that ACTH present in the brain does not have a pituitary source comes from the work of Allen (1974), who showed that the hormone is incapable of penetrating the blood brain barrier. Other evidence that indirect transport via the systemic system can be eliminated as a contaminating source of ACTH is that, in cases where a high level of ACTH is present in the circulation, such as following adrenalectomy, no

corresponding rise in "brain ACTH" has been observed (Krieger et al, 1979).

There is a possibility, not proven, of retrograde transport of hormone up the portal vessels. Thus, brain ACTH could reflect both the contribution of a pituitary source, perhaps mediating a short-loop feedback suppression of pituitary ACTH and a brain source. Moldow et al (1978<sup>a</sup>) have suggested that all extrapituitary sites of ACTH are pituitary-derived via such retrograde flow. Evidence for this comes from their observations that the distance from the pituitary at which ACTH is found and the concentration gradient in brain tissues are roughly comparable. However, this would seem to be invalidated by the biosynthesis, in vitro, of an immunoreactive 3,000 D ACTH/endorphin-like material in bovine hypothalamic cells maintained in culture, which was measured by Liotta et al (1979<sup>a</sup>) by showing <sup>3</sup>H-labeled amino acid incorporation into a high molecular weight material. The suggestion was made in this paper that cells within the hypothalamus, but not cortex, synthesize material with the immunological characteristics of the 31D ACTH precursor. Characterization by Liotta et al (1979<sup>b</sup>) confirmed such similarities, which included apparent identical size, presence of oligosaccharide units, the basic nature of the protein, and the ability to be processed by trypsin-like enzymes to smaller molecular weight products. The same group has shown that rat arcuate nucleus cells, maintained in culture for 17 days, release twice the amount of ACTH and  $\beta$  endorphin than is present in the tissue on day one.



Other lines of evidence for a brain origin include studies by Krieger's group where destruction of the arcuate nucleus in newborn rats by monosodium glutamate was associated with a decrease in brain ACTH but no change in pituitary ACTH levels. Immunohistochemical studies by Watson et al (1978), using colchicine to block axonal transport along nerve fibers, have shown an increased density of staining of arcuate nucleus cell bodies after such treatment, indicating accumulation of the hormone in the cells and therefore probable in-situ synthesis. Either stress and chronic immobilization, which normally are associated with changes in ACTH pituitary levels, appears to have no effect on the levels of the brain hormone (Krieger, 1979).

These studies tend to weaken the hypothesis that the presence of ACTH extrapituitarily is due to diffusion from the pituitary alone, and strengthens the idea that the hormone may have specific functions on nervous tissue.

ACTH and ACTH fragments are reported to play a role in motivation, learning, and memory (Gispen and de Wied, 1977). Other fragments are reported to induce excessive grooming, stretching, and yawning when administered intraventricularly (Gispen et al, 1976<sup>a</sup>). These activities are not mediated by adrenal steroids. Sufficient evidence is available from Dunn et al (1977) to indicate that ACTH affects cerebral protein synthesis, RNA synthesis, protein phosphorylation, and cyclic nucleotide metabolism. It has been reported by Terenius et al (1975) that ACTH interacts with opiate receptors. This could possibly interfere with the analgesic action of morphine on the CNS. Naloxone can suppress the

excessive grooming seen after intraventricular administration of ACTH (Gispen et al, 1976<sup>b</sup>).

ACTH also affects the acetylcholine, norepinephrine, and serotonin content or turnover of various brain regions, selectively modifies unit activity in midbrain limbic structures, and enhances the dephosphorylation of membrane proteins leading to membrane permeability changes, properties reviewed by Gispen et al (1977).

These changes in behavior and the general metabolism of nervous tissue brought about by ACTH in the brain indicates its probable importance as a neuromodulator.

Another important product of the large precursor protein associated with ACTH synthesis is  $\alpha$  melanocyte-stimulating hormone or  $\alpha$  MSH. The primary function of pituitary  $\alpha$  MSH is known to be the regulation of pigmentation changes in many vertebrates.

o.  $\alpha$  Melanocyte-Stimulating Hormone ( $\alpha$  MSH).

Before the work concerning the existence of a large precursor protein for  $\alpha$  MSH in both brain and pituitary; the sole source of  $\alpha$  MSH, a basic tridecapeptide amide, was thought to be the pituitary gland.

Dube et al (1978) reported finding  $\alpha$  MSH-containing nerves in both rat and man. The source of  $\alpha$  MSH fibers appeared to be the  $\alpha$  MSH-containing perikarya of the arcuate nucleus in the hypothalamus (Vaudry et al, 1978), and the peptide has been localized within vesicles of nerve

endings by Barnea et al (1977). Warburg et al (1977) demonstrated the release of  $\alpha$  MSH from a synaptosome-enriched fraction prepared from hypothalamic tissue.

Over 40% of the immuno-assayable total brain content has been found in the hypothalamus, the limbic system containing substantial amounts also (Eskay et al, 1979<sup>a</sup>). Surgical isolation of the arcuate nucleus of the hypothalamus resulted in a significant fall in extrahypothalamic  $\alpha$  MSH levels, whereas no reduction in hypothalamic peptide content was noted. This data gives further evidence that  $\alpha$  MSH-containing cells in the arcuate region send projections to extrahypothalamic areas and are the source of  $\alpha$  MSH in the CNS. Treatment of rats with monosodium glutamate (MSG) results in destruction of the neurons in the arcuate nucleus (Eskay et al, 1979<sup>b</sup>). This study measured the levels of  $\alpha$  MSH in the hypothalamus and the pituitary after treatment with MSG and showed, again, that levels of peptide fell in brain tissue, whereas pituitary  $\alpha$  MSH levels remained constant. This same treatment by Hughes et al (1979) also resulted in a reduction of ACTH-like and endorphin-like concentrations in the hypothalamus, substantiating the theory of a common precursor origin of  $\alpha$  MSH, ACTH, and  $\beta$  endorphin.

Barnea et al (1977) examined the cosequestration of ACTH and  $\alpha$  MSH in both hypothalamic and hypophysial neuronal granules. The ratio of one peptide to the other found in the granules was dependent on location. In the hypothalamus, 5 times more  $\alpha$  MSH than ACTH was found, whereas 20 times more ACTH than  $\alpha$  MSH was seen in the pituitary. The peptide granules for each peptide were distinguishable, and the peptides

are likely to be released simultaneously. There must be differences in the intragranular conversion of ACTH and  $\alpha$  MSH, as has been alluded to previously, to account for the differences in hormone levels in pituitary and hypothalamic neurons. Similar data has recently been reported for human brain tissue (Kleber et al, 1980).

On the basis of chromatographic, electrophoretic, and biological characteristics,  $\alpha$  MSH present in the rodent brain has been shown by Oliver et al (1978) to be identical to synthetic  $\alpha$  MSH.

In support of the hypothesis that  $\alpha$  MSH is a neurotropic substance, its effect in vitro on neural tissue has been shown to alter electrical activity (Krivoy and Guillemin, 1961, and Denman et al, 1972); to alter biochemical events such as the metabolism of biogenic amines (Leonard et al, 1976); and to increase the levels of c-AMP in the occipital cortex (Christensen et al, 1976). Administration of  $\alpha$  MSH has also been shown to have an effect on various behavioral tasks such as avoidance behavior (Wimersma, 1977, and Greven et al, 1977).

Significant diurnal variation in  $\alpha$  MSH has been observed by O'Donohue et al (1979), and the hypothesis that the hormone may participate in the manifestation of the diurnal rhythms of such behavioral effects as arousal, attention, and vigilance has been proposed. Panksepp et al (1976) have also reported an  $\alpha$ -MSH effect on other obvious circadian rhythm patterns such as sleep-waking activity.

$\alpha$  MSH, therefore, appears to be a peptide found both in brain and pituitary, similar in structure, but each pool of hormone being independently regulated. Its importance as a neuromodulator has been discussed.

The next section deals with a group of endogenous substances which bind to opiate receptors and mimic the biological effects of morphine, the endorphins, which include  $\alpha$  and  $\beta$  endorphins, and the enkephalins.

P. The endorphins.

Endorphins have been isolated from various tissues including the brain by Hughes et al (1975), the pituitary by Teschenmacher and Goldstein (1975), and the GI tract (Elde et al, 1976).

Historically, opiate receptors were found using immunohistochemical techniques by Kuhar et al (1973) to be concentrated in the mesolimbic system of the brain. It seemed unlikely that such highly stereospecific receptors should be developed merely to interact with alkaloids from the opium poppy. Early experiments by Akil et al (1972) showed that analgesia produced by electrical stimulation of the periaqueductal gray region of the brain stem of rats could be reversed by the administration of naloxone, a pure opiate antagonist. Jacob et al (1974) found that mice could be made more sensitive to pain by injection of small amounts of naloxone. These observations indicated that there existed an endogenous analgesic opiate, the action of which could be blocked by naloxone.

By 1975, Hughes et al (1975<sup>b</sup>) isolated from pig brain a peptide with a molecular weight of 1000 D, which could combine as an agonist at opiate receptor sites and whose actions could be blocked by naloxone. Cox et al (1975) isolated a similar opiate peptide from human cerebral spinal fluid and pituitary extracts. Furthermore, in the brain, these opioid peptides

were localized in the same areas anatomically as Kuhar's opiate receptors, as shown by Pert and Sneider (1976).

Two related pentapeptides with opiate activity were isolated by Hughes (1975<sup>a</sup>). The most abundant, in pig brain, was named methionine-enkephalin and the other, leucine-enkephalin. As previously noted (Fig. 3), their sequences were contained within the structure of  $\beta$  lipotropin (Fig. 4). Low levels were also found in the pituitary. Evidence for separate populations in the brain of the met- and leu-enkephalin have been reported by Larsson et al (1979<sup>b</sup>) using histochemical techniques. The enkephalin-containing nerves also showed a distribution distinct from nerves containing  $\beta$  endorphin, a larger opiate peptide (Guillemin et al, 1976), ACTH, and  $\beta$  lipotropin (Bloom et al, 1978).

At the present time, there is no definitive evidence that the enkephalins in the brain are derived from  $\beta$  lipotropin (Guillemin, 1980).

The perinatal development of the endorphin- and enkephalin-containing systems has been studied in the rat brain by Bayon et al (1979). Endorphin levels were much higher than enkephalins in the 16-day fetus, dropping to lower levels by the 20th day of gestation. The enkephalin content rose at day 20 and continued to increase at a faster rate than the endorphins from the 6th to the 25th day postnatally to achieve adult levels. This would suggest an independent development of the endorphin and enkephalin systems due to the lack of correlation between rates of increase; however, the authors suggest that certain endorphin-like systems could develop differentially into enkephalin systems during gestation.




Figure 4. Relationship between  $\beta$  Lipotropin amino acid sequence and other peptides of the brain and pituitary.

This concept is in agreement with observations that small local interneurons, like the enkephalin-containing cells, frequently migrate and differentiate more than larger neuron-forming pathways.

Evidence for distinct pools of brain endorphins and pituitary endorphins comes also from the work of Kobayashi et al (1978) and Cheung et al (1976). These groups showed that hypophysectomy failed to alter the brain content of opioid peptides after as long as 34 days without a pituitary. Guillemin et al (1977) also demonstrated that  $\beta$  endorphin levels in the brain did not change in experiments causing marked variations in pituitary  $\beta$  endorphin levels.

Two functions have been suggested for opiate peptides in brain. They may act as hormones or neuromodulators transported to their target organs by CSF or the vascular system; or, they may act as neurotransmitters, liberated from nerve terminals and binding to postsynaptic membranes.

Several studies suggest a role as neurotransmitters. In support of this role, they have been found concentrated in the synaptosomal fraction of brain homogenates and are localized in nerve endings (Simantov et al, 1976<sup>b</sup>). They appear to be able to inhibit neurotransmitter release, a response apparently related to changes in the  $\text{Ca}^{++}$  flux. It has been shown that morphine can decrease acetylcholine (Yaksh et al, 1975), and norepinephrine release (Arbilla and Langer, 1978), an effect antagonized by  $\text{Ca}^{++}$ . Evidence by Guerrero-Munoz et al (1979) indicates that a single dose of  $\beta$  endorphin causes a decrease in  $\text{Ca}^{++}$  concentration in synaptosomal fractions, attributable to reduced  $\text{Ca}^{++}$  binding and uptake.



Sustained administration of  $\beta$  endorphin showed a reversal of this effect and the development of tolerance and physical dependence or addiction, supporting the concept that  $\beta$  endorphin may inhibit the release of transmitters by inhibition of  $\text{Ca}^{++}$  influx in a mode similar to that of morphine.

The rapid degradation of the pentapeptide endorphin in serum seems to exclude a hormonal function (Simantov and Snyder, 1976<sup>a</sup>), although Sarne et al (1978) have postulated that a larger enkephalin-like substance found in sera was resistant to degradation and could act as a neuropeptide stimulating pituitary GH and prolactin release, as shown by Rivier et al (1977<sup>b</sup>). Drouva et al (1980) have shown that this may be mediated by the inhibition of release of somatostatin in the hypothalamus by opiate peptides such as met-enkephalin. This inhibition of release of hypothalamic hormone was naloxone reversible and was consistent with the hypothesis that opiates interfere with  $\text{Ca}^{++}$  influx. Met-enkephalin and morphine were also shown to inhibit GnRH, and therefore LH release, from the pituitary, and opiate receptors were isolated on GnRH as well as somatostatin nerve endings in the hypothalamus.  $\beta$  endorphin has been reported to promote the secretion of prolactin, the exact mechanism of this being unclear (Rossier et al, 1980, and Spies et al, 1980).

Dawson et al (1979) have studied another aspect of the effect opiate peptides may have on nervous tissue. The presence of opiates in cultured mouse neuroblastoma cells modified the biosynthesis of both membrane glycosphingolipid and glycoprotein. These effects seem to be mediated by

opiate receptors, of which this cell line has a prolific quantity, as they were stereospecific and naloxone reversible. Neuroblastoma cells treated with c-AMP are known to undergo morphological differentiation and show enhanced glycosylation of a number of secretory proteins (Truding et al, 1975). As other studies have demonstrated that opiates and enkephalins inhibit PGE<sub>1</sub>-stimulated adenylate cyclase activity (Klee et al, 1976), it appears that the modifications of membrane glycosphingolipid and glycoprotein seen by Dawson et al (1979) may represent a c-AMP mediated event.

Interest in the endorphins stems from the possibility that they or their analogues may be non-addictive pain killers to replace opiates such as morphine. It has been demonstrated that they are distributed in the pain-mediated pathways of the CNS. However, a lack of effect in producing analgesia has been observed when enkephalin is injected intraventricularly into rat brain. In vitro, indications are that enkephalins have a more profound opiate effect and bind to receptors about half as avidly as morphine. All these effects and those such as the induction of a state resembling catatonic schizophrenia by  $\beta$  endorphin (Bloom et al, 1976) can be blocked by the specific antagonist of opiate action, naloxone.

Evidence is opposed to the fact that the opiates are non-addictive, as both enkephalins and  $\beta$  endorphins are found to produce tolerance and dependence effects (Pert et al, 1976, and Blasig et al, 1976).  $\beta$  endorphin appears to resemble morphine pharmacologically. As an analgesic, it is 20-30 times more potent when given intraventricularly (Li et al, 1977).

Recent work by Pasternak et al (1980) supports the concept of multiple opiate receptors and gives evidence for mediation of opiate analgesia by a subpopulation of high affinity opiate receptors.

A role for enkephalins in the mediation of behavioral reinforcement is supported by several lines of evidence. Central injection of enkephalin serves to reinforce self-administered behavior, and electrical stimulation of enkephalin-rich regions of the brain also serves to reinforce self-stimulated behavior. Both effects are blocked by naloxone. Gunne et al (1977) reported that chronic schizophrenics exhibit elevated levels of endorphins in their CSF during symptom-rich phases of their disease. When their condition improved, a return to low endorphin levels was observed, implying that some psychological symptoms might be due to excessive endorphin activity.

The endogenous opioids clearly have powerful effects, are involved in behavior, and interact with other neuronal systems. While enkephalins potentially appear to conform to definitions of a neurotransmitter,  $\beta$  endorphin may be a neuromodulator. The enkephalins seem to be situated in such a way to influence basic processes such as pain transmission, respiration, motor integration, endocrine responses, and limbic functions involved in the elaboration of emotions.  $\beta$  endorphin seems more likely to affect various structures by interacting with other local neuroregulators.

Apart from ACTH and  $\beta$  LPH, there have also been recent reports of other pituitary hormones, such as prolactin, in extrapituitary locations.

q. Prolactin.

Evidence from Fuxe et al (1977) showed immunohistochemically that prolactin appears to be present in the nerve terminals of several hypothalamic nuclei such as the arcuate and ventromedial areas. Furthermore, Toubeau et al (1979) have recently noted the similarity in distribution of prolactin and ACTH immunoreactivity throughout the rat brain. Adrenalectomy and hypophysectomy have been reported to produce no effect on levels of prolactin in the hypothalamus.

A direct effect of prolactin on the CNS is suggested by reports that impotence present in males with hyperprolactinemia and normal testosterone concentrations may be corrected by lowering the prolactin concentrations by bromocryptene administration (Carter et al, 1977). Little else is known of the effect of prolactin on the CNS, as there is of the finding of TSH in extrapituitary tissues.

r. Thyroid stimulating hormone (TSH).

As long ago as 1967, using a bioassay, Bakke et al reported the presence of TSH in the rat median eminence, which was apparently independent of circulating TSH levels. In more recent findings, Moldow et al (1978<sup>b</sup>) attributed the presence of TSH outside the pituitary to retrograde flow from the pituitary. Further investigation into these observations will be particularly considered in this dissertation as will the locating and biochemical characterization of radioimmunoassayable growth hormone-like material in the CNS.

s. Growth Hormone (GH)

Pacold et al (1976) were the first to report the finding that dispersed rat brain cells grown under standard tissue culture conditions produced an immunoassayable growth-hormone-like material, indicating the possibility that GH could be added to the growing list of brain peptides.

The continuance of these observations, together with the finding of an immunoassayable TSH-like peptide in the rat brain, represents the thrust of the independent investigations that form the basis of this thesis.

To reiterate, my research has been oriented to the further confirmation of the presence of pituitary-like GH and TSH within discrete areas of the rat brain. Having developed a substantial background literature review of recent developments dealing with the unusual, if not unexpected, finding of diverse peptides in the brain, subsequent consideration will deal with my independent investigations dealing with the distribution of GH- and TSH-like peptides in the rat brain, and with studies designed to ascertain the physiological meaning of these novel observations.

### 3. PROPOSED INVESTIGATION

#### Specific Objectives

The distribution in the rat and primate brain of the anterior pituitary hormones, growth hormone, and thyroid-stimulating hormone will be studied. Experimental work will involve the development of working, sensitive, and specific radioimmunoassays for two rat hormones, TSH and GH. Another line of approach will be the development of primary cell cultures of neuronal tissue from various brain regions and the use of immunohistochemical staining techniques to further characterize the distribution of pituitary-hormone-like material in the adult rat brain. Ultracentrifugation of brain homogenates will determine the association or lack of association of the hormones with a particular brain fraction, either nuclear, mitochondrial-synaptosomal, or microsomal. Also to be examined will be the ontogenetic developmental pattern of these hormones in fetal, neonatal, and adult brain regions.

In an effort to determine the similarities or dissimilarities between pituitary and brain hormones, a part of this study will attempt to characterize biochemically and biologically the brain hormones. Using gel chromatography, the molecular weights will be determined, parallelism studies with the radioimmunoassay will examine immunological similarities, and bioassays will show the extent of biological activity of brain hormones as compared to hormones of pituitary origin.

Although it appears unlikely that these brain-based hormones could be of pituitary origin, one aspect of this investigation will be to test that concern. This will be accomplished by examining the intactness of the blood-brain barrier to circulating radiolabelled pituitary hormones. Furthermore, the use of hypophysectomized animals will provide the opportunity to compare levels of brain hormones in animals with and without an intact pituitary gland. In other studies, neural cells from hypophysectomized rats will be cultured in monolayers to test their ability to release GH and TSH hormone in vivo.

If two distinct pools of hormone, brain and pituitary, exist, these may turn out to be regulated in different ways. In an attempt to discern these interrelationships, experiments such as endocrine target organ ablation studies will be undertaken in order to understand this problem and its ramification.

From these different study areas it is hoped that the objective of this work will gain some insight into the physiological implications involved in the finding of pituitary-like hormones in discrete areas of the rat brain.

#### 4. MATERIALS AND METHODS


##### A. THE DEVELOPMENT OF RADIOIMMUNOASSAYS (RIA) FOR RAT GROWTH HORMONE AND THYROID STIMULATING HORMONE

The radioimmunoassay, as introduced by Yalow and Berson (1959) for the measurement of serum insulin concentrations, is a method of detecting trace concentrations of any substance that can serve as an antigen.

Based on competition for antibody; directed against the antigen; between a standard amount of radioactive antigen,  $C^*$ , and its unlabeled counterpart  $C$ , at unknown concentration in the test sample (Fig. 5). The higher the concentration of unlabeled competitor, the lower the ratio of bound  $C^*$  to unbound  $C^*$  which can be measured. Separation of bound from unbound  $C^*$  can be achieved by the precipitation of bound labeled and unlabeled antigen by a second antibody directed against the antibody moiety of the immune complex, and measurement of the radioactivity representing bound  $C^*$  in this precipitate. The higher the concentration of unlabeled competition, the lower the ratio of bound  $C^*$ , ( $B$ ) to unbound  $C^*$  ( $B_0$ ).

In order to determine the concentration of unknown  $C$  in a sample of blood, urine, or tissue extract, the amount of competitive inhibition observed in the unknown is compared with that obtained in separate solutions to which known amounts of purified  $C$  is added. A standard curve plotting the known concentrations of  $C$  against the ratio of bound to unbound  $C^*$  can thus be obtained (Fig. 6), and the level of unknown  $C$  in a sample determined by its ratio of bound  $C^*$  to unbound  $C^*$ .





$C =$  COMPOUND TO BE MEASURED (ANTIGEN)

$C^* =$  SAME COMPOUND TAGGED WITH RADIOACTIVE TRACER ( $^{125}\text{I}$ )

$AB =$  SPECIFIC ANTIBODY TO  $C$

Figure 5. Schematic diagram illustrating the principle of Radioimmunoassay.




Figure 6. Standard calibration curve for radioimmunoassay.

Basic requirements for a radioimmunoassay therefore include a specific antigen-directed antiserum; a radioactive-labeled, purified antigen; appropriate standard reference preparations of the antigen, and a method for the clear, clean and consistent separation of antigen-antibody complexes from the unbound fraction.

In the double antibody method of separation, which was selected for the RIA's for rat GH and TSH in this study, the first antibody, which is specific to the antigen to be determined, is added at a level that enhances competition between labeled and unlabeled antigens or ligands. That is, the antibody/antigen ratio corresponds to the antigen-excess side of the equivalence zone in a liquid precipitin reaction (Fig. 7). The second antibody (anti-antibody) is introduced in excess to ensure complete precipitation of the first antibody and its complexes with antigen, that is, the antibody-excess side of the equivalence zone of the curve. The method of RIA is exceedingly discriminating because of the possibility to choose antisera with great specificity and sensitivity for the test activity.

## I. THE RAT GROWTH HORMONE RADIOIMMUNOASSAY

### a. Production of specific antiserum to rat growth hormone.

The larger molecular weight protein hormones, such as growth hormone (GH) and thyroid-stimulating hormone (TSH), generally elicit a good antigenic response, and it was therefore decided to produce our own rat growth hormone antibody from guinea pigs. Since a very pure antigen is




Figure 7. Precipitin curve for a monospecific system: one antigen and the corresponding antibody.

radioactively labeled and used as the radioactive tracer in the RIA, the presence in the antibody preparation of other antibodies does not interfere with the reaction between labeled antigen and its antibody (Straus et al, 1977). It is therefore possible to use a relatively impure antigen, in this case growth hormone, for immunization. Extraction and isolation of crude rat pituitary GH was accomplished using a modification of the method of Stroud et al (1973). Approximately 50 rat pituitaries were removed from decapitated animals, weighed, and homogenized in a 3:2 mixture of 10% ammonium acetate and 96% ethanol. The mixture was stirred for two hours at 4° C, centrifuged for 20 minutes at 2000 rpm at 4° C, and the supernatant decanted off. The residue was stirred overnight in 30 mls of 0.1N NaOH, pH 10.0. After adjustment to pH 8.4 with 4N HCl, the slurry was stirred for 30 minutes and then centrifuged for 15 minutes at 3000 rpm at 4° C. The supernatant was reserved, and the residue re-extracted overnight with 0.1N NaOH, pH 10.0 and centrifuged at 4° C, after which the supernatants were pooled.

Using 4N HCl, the pH of the pooled supernatants was adjusted to 4.8 and stirred for 30 minutes. The heavy precipitate that formed was removed by centrifugation for 15 minutes at 3000 rpm at 4° C and the supernatant reserved. The precipitate was then dissolved in 0.1N NaOH and the solution adjusted to pH 4.8 by addition of 4N HCl and stirred for 30 minutes. After centrifugation, the precipitate was discarded and the supernatant combined with the previous pooled supernatant. The volume of the resultant solution was measured and 96% v/v alcohol, cooled to -30° C, was added slowly to give a final alcohol concentration of 25% v/v. After

stirring for a further 30 minutes, the precipitated crude growth hormone was collected by centrifugation for 15 minutes at 3000 rpm at 4° C. A total of twelve guinea pigs each received 0.5 ccs. of crude Bordatella pertussis vaccine (E. Lilly Corp., Indianapolis, Indiana) intradermally into 30-50 sites over a small area of their backs.

Twenty-four hours later, each received, subcutaneously, approximately 1 mg of the crude rat growth hormone, prepared as described above, dissolved in 0.5 ccs of saline together with an equal volume of complete Freund's adjuvant (Difco). Five subsequent similar booster immunizations were given with this crude extract over a five-month period.

Within 10 days of each injection, the guinea pigs were bled from a leg vein and serum collected for evaluation of antibody "titre", which reflects the binding affinity of the antibody to labeled antigen. The "sensitivity" of the antiserum was judged as the ability of small amounts of unlabeled hormone to displace the radioactive labeled hormone, and the "specificity" or degree of cross-reactivity of an antiserum to one hormone with other closely related hormones was examined.

Effective competition upon addition of unlabeled hormone occurs only under conditions of limited antibody, and in the RIA, that amount of antibody which shows a binding of 35%-50% of labeled hormone is usually chosen, depending on the levels of unknown hormone to be measured. Where levels of unknown hormone to be measured are expected in the lower range, 35% binding conditions are used, as they are known to give a more sensitive standard curve for the lower concentrations of standard hormone.

In order to determine the best titre of antibody for a sensitive assay, a level of 35% binding of labeled hormone was therefore set for testing the individual test bleedings for antibody titre.

Dilutions of antiserum were prepared from 1:1000 to 1:1,000,000, and the binding and dissociation characteristics of radiolabeled purified growth hormone were examined, as shown for 5 different animals at one test bleeding (Fig. 8). Antibody binding characteristics not only varied between animals but between bleedings from the same animal. Serum from guinea pig 11, at a dilution of 1:65,000, was eventually selected for the growth hormone RIA on the basis also of the "sensitivity" of this particular antiserum.

The "sensitivity" of a particular antiserum was assessed by adding graded amounts of purified standard GH and testing their ability to displace the radiolabeled hormone under the conditions of our radioimmunoassay. If a standard curve was too flat, the antiserum was rejected as being "non-sensitive". A "sensitive" curve was one with a steep fall in the range of hormone concentration expected from our samples.

The "specificity" of the antibody from guinea pig 11 was evaluated by examining the degree of cross-reactivity with other molecules similar in their primary structure to GH, such as rat prolactin. A standard curve using growth hormone antiserum #11, radiolabeled GH, and known quantities of purified prolactin showed that less than 1% of the purified prolactin was being measured. The antiserum to GH from #11 guinea pig was therefore sensitive and specific and could be used in a radioimmunoassay. The second essential ingredient of an RIA is the radiolabeled hormone.




Figure 8. Determination of titer of G.H. antibody binding 35% of the total radiolabelled G.H. antigen from one test bleeding of five guinea pigs (12/18/78).



b. Radioiodine labeling procedure for rat growth hormone.

<sup>125</sup>I has proved to be the radioisotope most suitable for labeling peptides. The labeling reaction consists of substitution of radioactive iodine, produced by iodide oxidation in the presence of protein, for the tyrosine in peptide linkages.

The chloramine-T technique of isotope labeling developed by Hunter et al (1962) was used. This method produces radiolabeled GH with a high specific activity, that is, with a high amount of radioactivity per amount of hormone, while using smaller amounts of isotope than other techniques. This is important, as to achieve maximum sensitivity the concentration of labeled antigen must be kept below the minimum concentration of antigen to be measured. Rat GH was obtained from the NIAMDD Pituitary Hormone Distribution Program through Dr. A. F. Parlow.

In this method, radioiodide is oxidized at pH 7.6, with chloramine-T to iodine, which in turn iodinate the protein's tyrosyl and histidine residues, specifically.

The unreacted radioiodine is reduced to iodide by the addition of sodium metabisulfite to the reaction mixture, and albumin is added to protect the hormone from further radiation damage by a mechanism not entirely clear. The labeled protein is immediately separated from unlabeled iodide and damaged protein fractions to protect the labeled hormone from further radiation damage. This separation is achieved using gel filtration on coarse Sephadex G-50.

(i) Labeling Procedure.

Lyophilized, purified growth hormone was obtained from the NIAMDD, Bethesda, and 100  $\mu\text{g}$  was weighed out and dissolved in 10  $\mu\text{l}$  0.01N NaOH and 90  $\mu\text{l}$  distilled water. This was aliquoted in 20  $\mu\text{l}$  amounts and stored at  $-70^{\circ}\text{C}$ .

At all times, adequate radiation safety procedures were observed and the iodination procedure carried out under a hood to avoid contamination of the surrounding area by volatilized material. Iodination was performed in a glass vial 5 cm x 0.5 cm by addition of reagents in the following order:

1. 20  $\mu\text{l}$  GH (purified from NIAMDD).
2. 50  $\mu\text{l}$  Phosphate buffer. 0.5 M pH 7-6.
3. 10  $\mu\text{l}$   $^{125}\text{I}$ , approximately 1 m curie (New England Nuclear, Boston, Mass.).
4. Mixed well.
5. 10  $\mu\text{l}$  Chloramine-T solution (35 mg Chloramine-T, Eastman Kodak, Rochester, N. Y.) in 10 ml freshly prepared 0.05 M  $\text{PO}_4$  buffer, pH 7.6).
6. 45 seconds contact by swirling.
7. 15  $\mu\text{l}$  Sodium metabisulfite solution (75 mg sodium metabisulfite, Fisher Scientific, New Jersey) in 10 ml freshly prepared 0.05M  $\text{PO}_4$  buffer, pH 7.6.
8. 100  $\mu\text{l}$  2-5% bovine serum albumin (Miles Laboratories, Inc., Elkhart, Ind.) in 0.05 M  $\text{PO}_4$  buffer, pH 7.6.

This mixture was then transferred to a freshly prepared Sephadex G-50 coarse 0.9 x 20 cm plastic column (Pharmacia Fine Chemicals, Uppsala, Sweden), which had previously been coated with a few drops of 22% bovine serum albumin (BSA). The column was eluted with 0.075 M veronal buffer, pH 8.6, and 15 drop fractions were collected in albumin-coated glass tubes. The two tubes showing the highest radioactivity in the first peak were saved, diluted with 10 ml 1% BSA, 0.01 M phosphosaline buffer, pH 7.4, and stored in 1 ml amounts at  $-20^{\circ}$  C. The stability of this iodinated growth hormone could be maintained for approximately 4 weeks. After this, there was an increase in free radioactive iodide indicating a breakdown of the labeled hormone, perhaps due to radiation damage. An approximate specific activity between 90-110 mC/mg was obtained by this procedure.

Prior to setting up each radioimmunoassay, the labeled rat growth hormone was repurified using a Sephadex G-100 column 0.9 x 50 cm. The Sephadex was first equilibrated with 0.075 M veronal buffer and approximately 1 ml containing  $1 \times 10^6$  cpm labeled hormone applied. One ml fractions were collected and the top two tubes of the second peak used for the assay. The labeled hormone was immediately diluted in 0.01 M "working" phosphosaline buffer, pH 7.4, to approximately 12,000 cpm/100  $\mu$ l.

Solutions Used in Labeling Procedure for Rat GHPhosphate BufferStock solution, 0.1 M24.2 g  $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ 2.62 g  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ Distilled  $\text{H}_2\text{O}$  to 1 litre, pH 7.6Working solution: 1% BSA, phosphosaline buffer(0.15 M NaCl, 0.01 M  $\text{PO}_4$ ), pH 7.65.14 gm  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ 

26.6 gm NaCl

22.7 mls of 22% BSA/litre

0.1% Thimerosal (Sigma)

Final volume, 3.0 litres, pH 7.6.

Veronal Buffer 0.075 M , pH 8.6

292 g Sodium diethyl barbiturate (Fisher)

52.44 g Barbituric acid (Fisher)

4 liters distilled  $\text{H}_2\text{O}$ 

Heat to dissolve

Make up to 19 litres

Stored at 4° C

c. Preparation of GH standards.

For validation of the RIA procedure, the unknown GH substance to be assayed must be immunochemically identical with the substance used as an antigen standard. For this purpose, lyophilized reference preparations of rat growth hormone were obtained from the NIAMDD, the stock standard solutions were prepared as follows:

- 100  $\mu$ g was weighed using a Mettler balance, dissolved in 100  $\mu$ l 0.01 N NaOH, and made up to 1 ml with 0.01 M phosphate buffer, pH 7.6.
- Aliquots of 50  $\mu$ l containing 5  $\mu$ g were then stored at  $-20^{\circ}$  C for up to 6 months without loss of activity.

The standard curve for a RIA was made up using the following dilutions of stock hormone as growth hormone standards in working phosphosaline buffer, pH 7.6:

50 $\mu$ l stock + 2.5 cc buffer	=	2000 ng/ml
Standard 9	=	100 ng/ml
Standard 8	=	50 ng/ml
Standard 7	=	40 ng/ml
Standard 6	=	20 ng/ml
Standard 5	=	10 ng/ml
Standard 4	=	8 ng/ml
Standard 3	=	5 ng/ml
Standard 2	=	4 ng/ml
Standard 1	=	2 ng/ml

d. Separation of bound and free antigen in the GH RIA.

In a radioimmunoassay, the labeled peptide-antibody complexes are generally soluble, and some technique must be employed to distinguish between the bound and the unbound labeled peptide. One of the most accepted and least troublesome methods is that depending on precipitation of the complex with an antibody directed against the  $\gamma$  globulin of the first antibody in the complex, the "double antibody" method. The precipitate formed is isolated by centrifugation. After several trial runs, a 1:10 dilution of goat anti-guinea pig IgG (Pel-Freeze) was found to give the optimum degree of precipitation in the RIA for GH.

e. Assay protocol.

The final protocol for the rat growth hormone double antibody RIA was as follows, and the assay was incubated throughout at 4° C in 12 x 75 mm glass tubes. All samples were run in triplicate with individual non-specific binding controls (NSB).

DAY 1 100  $\mu$ l GH antibody 1:65,000, produced in guinea pig, made as follows:

- (1) 10  $\mu$ l stock GH antibody in 6.5 ml working  $\text{PO}_4$  buffer +  
5  $\mu$ l/ml normal guinea pig serum (Miles Laboratories,  
Elkhart, Indiana)

or

100  $\mu$ l non-specific binding buffer, made as follows:

5  $\mu$ l normal guinea pig serum/1 ml working  $\text{PO}_4$  buffer.

- (2) 100  $\mu$ l standard or unknown.

DAY 5

- (3) 100  $\mu$ l  $^{125}\text{I}$  GH, 12,000 cpm.

DAY 7

- (4) 100  $\mu$ l goat anti-guinea pig serum, 1:10.
- (5) 200  $\mu$ l working  $\text{PO}_4$  buffer.

Incubation for  $4\frac{1}{2}$  hours at  $4^\circ$  C.

- (6) 1 ml ice cold saline.

Centrifugation for 30 minutes at 2,200 rpm using an IEC 418 head in a refrigerated IEC Model PR-6000 centrifuge.

The supernatant was decanted and the precipitate counted in a Micromedic System 2/200 automatic gamma counter.

f. Treatment of data obtained from the GH RIA.

Calculation of data was handled using a Hewlett-Packard Clinical Laboratory Radioimmunoassay computerized program cassette with a Hewlett-Packard Model 7830A calculator and printout. This program performed a weighted and unweighted least squares regression analysis using the log of dose vs. % bound, logit of % bound, probit of % bound, and arcsin of % bound. The "best fit" of these linearizing transforms was automatically

selected and all useful standard curve statistics and plots were outputted to verify acceptability of each dose response variable.

Typical data processing of standards and unknowns are seen in Figs. 9 - 13. In Fig. 9, the standard curve data is shown as the total amount of cpm of tracer present per tube, and nonspecific binding (NSB) as cpm/tube, representing the bound antigen. The concentration in GH ng/ml of each set of three tubes per standard is given plus the cpm in the bound fraction for each standard assayed. Fig. 10a illustrates the computer's assessment of the correlation, or how this data best fits a straight line, using different ways of handling the data such as "Logit," "Probit," and "Arcsin." In this case, Logit was chosen as the best fit, and a weighted regression was performed on the data of the standard curve as shown in Fig. 10b. The coefficient of variation for each point in this assay was acceptable, i.e. under 10%, and Logit transformation was used to plot the curve, shown in Fig. 12, from which the unknown samples were calculated, as in Fig. 12. Fig. 13 shows the standard curve plotted using the data  $B/B_0$  vs. ng GH/ml. The "sensitivity" of this assay appeared good over the range of 2-100 ng of growth hormone.

g. Validation of the GH radioimmunoassay.

After the reagents were obtained and the method established, it was necessary to validate the RIA of GH in brain extracts. Primarily, assay validation consists of a demonstration that the assay is measuring what it is designed to be measuring, in this case, rat growth hormone. Multiple dilutions of brain extract and pituitary extract should give lines



RGH 68D  
3/6/79 TECHNOLOGIST S. FOJVAT

## TOTALS (CPM)

11586  
11599  
11725

TUBE NO.	CONCENTRATION IN NG/ML	CPM
1	NSB	410
2	NSB	333
3	NSB	309
4	0	4034
5	0	4116 B.
6	0	4194
7	2	3987
8	2	3951 B
9	2	3790
10	4	3522
11	4	3574
12	4	3730
13	5	3542
14	5	3589
15	5	3612
16	8	3203
17	8	3326
18	8	3351
19	10	3263
20	10	3167
21	10	3141
22	20	2273
23	20	2416
24	20	2408
25	40	1398
26	40	1403
27	40	1425
28	50	1174
29	50	1073
30	50	1080
31	100	764
32	100	743
33	100	705

Figure 9. Computerized output of G.H. standard curve.

Figure 10a.

ASSAY: RGH 68D  
3/6/79 TECHNOLOGIST S. FOJVAT

MODEL	INTERCEPT	SLOPE	CORRELATION
RAW DATA	1.257956973	-0.599259588	0.989075873
LOGIT	4.084257617	-3.134433938	0.997675286
PROBIT	7.430118155	-1.876054213	0.996794585
ARCSIN	1.431518075	-0.699741408	0.972278768

BEST FIT FOR RGH 68D IS LOGIT

WEIGHTED REGRESSION

Figure 10b.

ASSAY RGH 68D  
3/6/79 TECHNOLOGIST S. FOJVAT

DOSE	B/B0	STD. DEV.	VARIANCE	STD ERROR	Coeff OF VAR
0.00	1.0000000	0.0213562	0.0004561	0.0123300	2.1356199
2.00	0.9454481	0.0279642	0.0007820	0.0161452	2.9577765
4.00	0.8655685	0.0288450	0.0008320	0.0166536	3.3324872
5.00	0.8582182	0.0095649	0.0000915	0.0055223	1.1145030
8.00	0.7817924	0.0211258	0.0004463	0.0121970	2.7022286
10.00	0.7544279	0.0171477	0.0002940	0.0099002	2.2729378
20.00	0.5353348	0.0214009	0.0004580	0.0123558	3.9976582
40.00	0.2810840	0.0038443	0.0000148	0.0022195	1.3676850
50.00	0.2014701	0.0150043	0.0002251	0.0086627	7.4474094
100.00	0.1028162	0.0078120	0.0000610	0.0045102	7.5979878

## X LOGIT TRANSFORM

2	2.933497406
4	1.878566729
5	1.80200039
8	1.278911035
10	1.123850082
20	0.141728422
40	-0.939142958
50	-1.378810207
100	-2.168243684

Figure 10a. Best fit analysis for growth hormone standard curve data.

Figure 10b. Weighted regression analysis of G.H. standard curve.

ASSAY RGH 68D  
 3/6/79 TECHNOLOGIST S. HOJVAT





Figure 11. Growth hormone standard curve plotted as B/B<sub>0</sub> vs (ng/ml).

Figure 12. Growth hormone standard curve plotted as logit vs (ng/ml).

TUBE#	CPM	N <sub>50</sub> .	CONCENTRATION IN NG/ML	(+ OR -) 95% CONFIDENCE
34		396		
35		395		
36	2804			
37	2850			
38	2931			
			12.5451	1.3877
			-----	
39		448		
40		380		
41	2132			
42	2301			
43	1964			
			22.9888	6.0996
			-----	
44		455		
45		440		
46	1423			
47	1436			
48	1414			
			43.4068	0.9736
			-----	

Figure 13. Calculation of growth hormone content of unknown samples.

parallel to the standard curve line of purified pituitary rat growth hormone, which would indicate immunological similarities between the brain and pituitary hormones. Physiological changes in the rat's circulating GH levels are brought about by hypophysectomy. These changes should be reflected in pre- and post-surgical serum samples, and were found to be distinguishable by this rat GH RIA. Another method of controlling and detecting interassay variation is the use of internal standards of pooled brain extract in each assay performed. A possible error source frequently encountered in RIA's are "false positive" measurements due to damage of the labeled antigen during the iodination process, or by enzymatic breakdown by tissue homogenates, which leads to a reduction in bound labeled hormone and a misinterpretation of these low counts, so obtained, as indicative of large amounts of growth hormone in the sample.

Recovery studies were therefore performed in order to test the assay for this type of error. Purified rat growth hormone was added in graded, known amounts to brain extracts in triplicate, each sample with its own nonspecific binding control (NSB). These NSB tubes, generally run in duplicate, contained no first antibody in order to detect the binding of labeled antigen nonspecifically to the walls of the assay test tube or to some protein, for which it has a nonimmunological affinity, and which is present in the sample.

As shown in Table 1, the detection of added known amounts of purified GH was approximately 100% from this growth hormone RIA, indicating no change in the hormone molecule or to the destruction of the labeled

TABLE 1

RECOVERY OF ADDED AMOUNTS OF REFERENCE PREPARATION GH  
IN NORMAL BRAIN HOMOGENATE

<u>ng added. GH</u>	<u>Recovery</u>		<u>O/E x 100</u>
	<u>obtained (O)</u>	<u>expected (E)</u>	
0	8.0 = amount of hormone indigenous to brain homogenate		
2	11.9	10.0	119.0%
4	16.6	12.0	138.3%
5	11.9	13.0	91.5%
8	16.2	16.0	101.3%
10	13.9	18.0	77.2%
20	24.3	28.0	86.8%
40	60.7	48.0	126.5%
50	63.2	58.0	109.0%
100	127.5	108.0	118.1%
		Average	107.5%

growth hormone by brain homogenate. From previous "specificity" studies and the "recovery" studies of purified GH, it seemed that this RIA, as developed for rat GH, was a valid assay and the results were not subjected to any artifactual problems.

The radioimmunoassay to detect thyroid-stimulating hormone in brain homogenates was developed in a similar way to that of GH and will be described in detail in the section to follow.

## II. THE RAT THYROID STIMULATING HORMONE RADIOIMMUNOASSAY

### a. Antibody to TSH.

For this assay, antiserum to rat TSH, prepared in rabbits, was obtained from Dr. Parlow, NIAMDD. This was stored as a 1:10 dilution at  $-70^{\circ}$  C in a phosphosaline EDTA buffer (0.01 M  $\text{PO}_4$ , 0.15 M NaCl) with 0.05 M EDTA and 3% normal rabbit serum, pH 7.6. For the TSH RIA, a dilution of antibody 1:15,000 in the same buffer was found to give the most sensitive assay.

### b. Radioisotope Labeling Procedure for Rat TSH.

A purified rat TSH was obtained from NIAMDD and stored in 20  $\mu\text{l}$  aliquots containing approximately 5.0  $\mu\text{g}$  of hormone in 0.01 M phosphosaline buffer, pH 7.6 at  $-70^{\circ}\text{C}$ . The Chloramine-T oxidation method was again successfully employed for labeling of TSH with radioactive iodine (Hunter et al, 1962). The procedure was carried out using similar radiation safety measures as that of GH and was as follows:

1. 20  $\mu\text{l}$  stock rat TSH (5.0  $\mu\text{g}$ , NIAMDD).
2. 10  $\mu\text{l}$   $^{125}\text{I}$  - approximately 1 m curie (New England Nuc.)
3. 25  $\mu\text{l}$   $\text{PO}_4$  buffer, 0.05 M, pH 7.6.
4. 10  $\mu\text{l}$  Chloramine-T (25 mg Chloramine-T in 10 ml freshly prepared 0.05 M  $\text{PO}_4$  buffer, pH 7.6).

Mixed for 40 seconds.



5. 25  $\mu$ l Sodium metabisulfite (25 mg sodium metabisulfite in 10 ml freshly prepared 0.05 M  $\text{PO}_4$  buffer, pH 7.6).
6. 50  $\mu$ l 0.4% KI (Fisher).

This mixture was then transferred to a Sephadex G-50 coarse, 0.9 x 20 cm plastic column, coated with bovine serum albumin, and eluted with 0.01 M phosphosaline buffer, pH 7.6. Fifteen-drop fractions were collected in 16 x 100 cm glass tubes, and those tubes showing the highest radioactivity in the first peak were diluted in 0.01 M phosphosaline, 1% albumin buffer, pH 7.6, pooled and stored in 1ml aliquots at  $-20^\circ$  C. The stability of this iodinated TSH was for approximately two weeks at this temperature.

Repurification of labeled hormone was performed the day of an assay using a Concanavalin-A, Sepharose column (Pharmacia Fine Chem.). The principles behind this method of purification of TSH are described briefly here.

"Affinity chromatography" using Con-A Sepharose is a well established preparative technique which can be used for isolating glycoproteins which contain  $\alpha$ D-glucose or  $\alpha$ D-mannose residues. In such a procedure, glycoproteins in a sample are washed into the column and are retained due to their affinity for the lectin, Concanavalin-A, which has been previously immobilized on the Sepharose beads. These glycoproteins can now be separated from other proteins in the sample which are not retained on the beads. After a thorough washing of the column with a phosphosaline buffer to remove any traces of non-glycoprotein, the bound glycoprotein may

be eluted off the column by 1-0 methyl  $\alpha$ -D-glucopyranoside, pH 9.0 (Shore et al, 1973). The glucopyranoside competes with bound glycoprotein for Con-A binding sites and at sufficient concentration can remove all glycoprotein from the ligand.

The following procedure, adapted from the method of Patritti-Laborde et al (1979), was employed to recover repurified  $^{125}\text{I}$  TSH using a small Con-A Sepharose column, 0.9 x 19 cm. Glass wool was placed on the scintered glass disc at the base of the column and the gel poured to a height of 9 cm. The column was then washed continuously with 20 mls of 0.5 M phosphosaline, 1% BSA buffer, pH 7.6, and then approximately 1 ml of freshly  $^{125}\text{I}$  labeled purified TSH applied. The column was then washed with 60 ml of 0.5 M phosphate buffer and the radiolabeled bound glycoprotein TSH, subsequently eluted with 0.2 M 1-0-methyl- $\alpha$ -D-glvcopyranoside (Sigma Chemical Company, St. Louis, Missouri) in 0.5 M phosphosaline buffer, pH 7.6.

Three ml aliquots were then collected in 16 x 100 cm glass tubes and the two peak radioactive tubes pooled and diluted to yield approximately 9,000 cpm/100  $\mu\text{l}$  in "TSH buffer." The specific activity of the  $^{125}\text{I}$  labeled TSH ranged from 80-95 mC/mg.

#### Solutions:

##### Phosphate buffer - 0.5 M

6.9 gm  $\text{NaH}_2\text{PO}_4$   $\text{H}_2\text{O}$  in 100 ml

7.1 gm  $\text{NaHPO}_4$  in 100 ml

1:9 monobasic to dibasic, pH adjusted to 7.6.

0.05 M  $\text{PO}_4 = 0.5$  M diluted 1:10 with distilled  $\text{H}_2\text{O}$ .

"TSH buffer" = 1% BSA, 0.01 M phosphate, 0.15 M saline.

2 mls 0.5 M phosphate buffer  
 100 mls 0.155 M saline (9g NaCl/litre)  
 4.5 mls 22% BSA  
 pH 7.6

c. Preparation of TSH Standards.

Rat TSH reference preparations from NIAMDD were stored as 10 µg TSH/100 µl of "TSH" buffer in aliquots at -20° C. Standards were prepared by double dilution of this stock solution in the same buffer, and ranged from Standard 1 = 39 ng/ml to Standard 8 = 5000 ng/ml.

d. Assay Protocol.

The rat TSH double antibody assay was run at room temperature and required 3 days of incubation in 12 x 75 mm glass tubes.

- DAY 1
1. 200 µl TSH antibody, 1:15,000 rabbit anti-rat TSH (NIAMDD), made in 3% normal rabbit serum phosphosaline EDTA buffer;  
 or  
 200 µl non-specific binding buffer (3% normal rabbit serum in phosphosaline EDTA buffer).
  2. 100 µl standard or unknown.
  3. 100 µl <sup>125</sup>I TSH, 9,000 cpm.
  4. 200 µl "TSH" buffer.

Mixed well.

DAY 2            5. 100  $\mu$ l anti-rabbit IgG produced in goat (PelFreeze, Roger, Arkansas) 1:10 dilution in "TSH" buffer.

DAY 3            6. 1 ml saline.

The samples were spun for 30 minutes at 2,200 rpm, decanted, and the antigen-antibody precipitate counted.

e. Treatment of Data Obtained from the TSH Radioimmunoassay.

Data was treated in a similar way to that described previously for GH. Typical standard, and logit vs. conc. curves, for the TSH radioimmunoassay are given in Figs. 14 and 15. The sensitivity of this assay was in the range of 39-5,000 ng/ml.

f. Validation of the TSH Radioimmunoassay.

Attempts to validate the TSH assay were similar to those described for GH. Cross-reactivity to FSH and LH was less than 0.14%, and recovery studies using <sup>125</sup>I TSH added in varying amounts to brain homogenates showed an average 97.3% recovery (Table 2).

ASSAY TSH 12E  
 10/29/79 TECHNOLOGIST S. HOJVAT





Figure 14, TSH standard curve plotted as B/B<sub>0</sub> vs (ng/ml).

Figure 15, TSH standard curve plotted as logit vs (ng/ml).

TABLE 2

RECOVERY OF ADDED AMOUNTS OF REFERENCE PREPARATION TSH  
IN NORMAL BRAIN HOMOGENATE

<u>ng added TSH</u>	<u>Recovery</u>		<u>O/E x 100</u>
	<u>obtained(O)</u>	<u>expected(E)</u>	
0	266.8 = amount of hormone indigenous to brain homogenate.		
39	265.0	305.8	86.7%
78	348.2	344.8	101.0
156	389.4	422.8	92.1
312	548.1	578.8	94.7
625	850.6	891.8	95.4
1250	1577.3	1516.8	104.0
2500	2902.7	2766.8	104.9
5000	5248.4	5266.8	99.7
Average			97.3%

### III. RADIOIMMUNOASSAYS FOR GROWTH HORMONE AND TSH IN PRIMATE BRAIN HOMOGENATES

#### a. Growth Hormone

A solid phase radioimmunoassay in antibody-coated tubes was set up after the method of Catt et al (1967). The principle of this solid phase radioimmunoassay is based on the ability of antibody-coated polymers, such as plastic test tubes, to bind radioactive tracer antigen. The use of antibody in this form is a technique, other than the "double antibody" method, for the rapid removal of unbound from bound radioactive tracer antigen by washing of the solid phase material with water on completion of the immune reaction. The solid phase consists at this point of bound radioactive tracer antigen-antibody complexes and competing bound unlabeled sample antigen-antibody complexes. This solid phase is then counted, after removal of unbound antigen by water, for the quantitation of bound tracer which varies inversely with the total quantity of antigen present in the original mixture.

In our assay, one ml of first antibody, anti-human growth hormone produced in rabbits and obtained from L. Kirsteins, was used at a 1:1500 dilution in 0.05 M bicarbonate buffer, pH 9.6, to coat the sides of 12 x 75 mm plastic tubes overnight at 37° C. An irreversible antibody-polymer complex was thus formed. After washing, to remove non-complexed antibody, radioiodinated growth hormone (Abbot) was added at a concentration of 10,000 cpm to each coated tube containing either standard growth hormone,

internal controls, or unknown tissue homogenate, plus bovine serum. This mixture was incubated at 37° C for 36 hours and, after rinsing, the tubes were counted for bound radioactivity. Levels of primate growth hormone were calculated for each tissue homogenate as pg/mg wet weight of monkey brain and pituitary tissue. The range of this assay was 1 to 100 ng of growth hormone.

b. Thyroid Stimulating Hormone

Samples of monkey brain were homogenized, and a double antibody radioimmunoassay for human TSH was carried out by the late Dr. A. Guan-  
sing, Wood Veterans Administration Center, Milwaukee, Wisconsin. The data obtained were calculated as pg/mg wet weight of tissue.



## B. DESCRIPTION AND CARE OF ANIMALS USED THROUGHOUT THIS STUDY.

Sprague Dawley male and female rats were used for these studies. All animals were kept at day/night ambient light and temperature regimes and were fed Purina rat chow with water ad libitum. Pregnant female Sprague Dawley rats were housed in separate cages throughout pregnancy, and no more than three animals per cage were allowed for other studies.

Hypophysectomized and thyroidectomized rats were purchased from Hormone Assay Laboratories, Chicago, Illinois, and were fed a liquified rat chow and orange slice diet for the first week post-operatively.


Animal sacrifice was routinely performed between 8 a.m.-10 a.m. using a guillotine.

## C. BRAIN DISSECTION PROCEDURES.

After allowing at least a one-hour period of acclimatization to a new locale, animals were killed by instantaneous decapitation using a guillotine, and trunk blood was collected by gravity flow from the severed neck. Handling prior to decapitation was kept to a minimum to prevent possible anomalous, stress-mediated rises in pituitary hormones. To minimize the possibility of retrograde portal blood flow after death, the occipital-parietal calvarium was cut away within two seconds of decapitation and the frontal lobes of the brain gently lifted in order to separate the brain from the pituitary stalk without disruption of the diaphragm sellae (Figs. 16 and 17).

CORTEX

HYPOTHALAMUS



CEREBELLUM

Figure 16. Isolated rat brain.




Figure 17. Coronal section of rat brain.

For tissue extraction studies, the brain was placed rapidly onto ice and, using separate instruments, individual brain parts were dissected free and immediately frozen on dry ice to minimize proteolytic enzyme activity. Brain parts intended for tissue culture experiments were placed directly from the animal into Basal Media Eagles (Gibco, Grand Island, New York) at room temperature before further processing.

Brains from the primate, Macaca mullata, were obtained from the Department of Dentistry, Loyola University of Chicago, after a lethal pentobarbital dose, dissected, and placed in acetone before extraction and hormone analysis.

For the harvesting of pituitary glands from intact rats and primates, the diaphragm sella was gently excised and the anterior hypophysis lifted from the sella turcica and placed on dry ice.

All separately dissected specimens of rat brain and pituitary were stored in aluminum foil at  $-70^{\circ}$  C and serum samples at  $-20^{\circ}$  C until radioimmunoassayed or bioassayed for the detection of hormone content. The method of extracting hormones from brain and pituitary tissues is outlined in the next section.

#### D. TISSUE EXTRACTION METHODS.

Individual rat brain parts and pituitaries were thawed at room temperature. The anterior pituitary, hypothalamus, and serum samples were assayed individually. However, GH and TSH concentrations in other brain regions required pooling tissue from at least five animals in order

to extract sufficient material to be measured within the sensitivity limits of the radioimmunoassay.

Fetal animals of appropriate gestational ages were obtained by maternal decapitation and hysterectomy. The brains of younger fetuses were frozen within six seconds of maternal death. The older fetuses were carefully removed from the uterine horns and kept warm and under as little stress as possible before decapitation.

Individual brain parts and pooled material were weighed and homogenized in 1 ml of ice cold buffer (PBS 0.01M, pH 7.6) for twenty seconds using a Tekmar homogenizer. The samples were spun at 3,000 rpm in a refrigerated centrifuge for 30 minutes, supernatants removed, and the pellets resuspended in 1.0 ml of cold buffer and stored overnight at 4° C. Tubes were recentrifuged in the cold and the two supernatants pooled. These were either assayed immediately or stored at -70° C until the time of assay.

In addition to being radioimmunoassayed for hormone content, materials for assay were thawed and a determination of protein content of tissue made on each sample using the method of Wadell (1956). Hormone levels for each brain area were determined using the radioimmunoassays of rat GH and TSH, and the results were expressed as either ng of hormone/mg wet weight tissue, or ng of hormone/mg protein or ng hormone/total brain part. In this manner, the distribution of pituitary-like hormones was mapped for the rat brain. A rough mapping of the primate Macaca mu-  
lata brain was also determined.

## E. TISSUE CULTURING OF BRAIN CELLS

Culture of tissue falls into two broad areas: the use of explant cultures of small fragments, or single cell preparations from tissue dissociated by mechanical or enzymatic procedures. The preparation and culture of single cell suspensions was the method of choice for this study in which distinct brain areas and pituitary tissue were further tested for their ability to release hormone into the growth media. The principle behind single cell suspensions, that of removing tissue from the influence of the whole organism and placing it in a more defined environment, has been validated by the demonstration that neurons in culture can develop physiological properties similar to those seen in comparable cells in situ, including the formation of new synaptic connections with one another and small aggregates (Fischbach, 1972, and Dichter et al, 1975). Some latent difficulties are, however, involved in the tissue culturing of nervous tissue. Cell division is a limited phenomenon resulting in the inability of primary cell cultures to remain viable indefinitely. Furthermore, long term subculturing of nervous tissue generally yields a nonfunctioning population of cells, as shown by Yasamura (1968). One alternative to these problems has been to culture functioning differentiated neoplasms which can be serially propagated in tissue culture indefinitely without losing functionality, for reasons not fully understood.

To obtain successful cell separation from primary tissue for tissue culture, it is necessary to begin with a suspension of single viable cells, relatively free of damaged cells, cellular debris, and red and white blood cells. Brain parts and pituitaries are dissected, using sterile instruments,

from 30-day-old male rats or 30-60-day-old males who have been hypophysectomized for various time periods beforehand. Tissue from approximately 10 animals is placed in a petri dish containing sterile Basal Medium Eagle plus 10% fetal calf serum (Gibco), and cleaned of any adhering meninges or connective tissue. To remove debris and red blood cells, the tissue is washed four times in sterile media before being cut into small 1 mm cubed pieces using two sharp scalpels to avoid tearing. Dissociation is carried out using 10 mls of 0.25% Pronase (Calbiochem-Behring Corp., San Diego, California) in an Erlenmeyer flask shaken gently in a 37° C water bath. Freshly released cells are harvested at 10-minute intervals and stored at 4° C with an equal volume of growth medium. The separated cells are triturated with a Pasteur pipette approximately five times and clumps of cells removed by passing the suspension through a sterile nylon sieve of 35 micron pore size (Tetko, Inc., Elmsford, N.Y.). The cells are spun gently for 10 minutes at 1,000 rpm in a Sorvall GLC-2 centrifuge at room temperature. After four washes with growth medium to remove Pronase, the cells are counted in a Petroff Hauser counting chamber. Viability is simultaneously checked by the trypan blue exclusion method, where only dead cells take up the blue dye and therefore can be distinguished from live cells. From 90%-95% viability is generally recorded with this methodology. After gentle mixing, the cells are diluted up to  $10^6$ /ml and 1 ml aliquots plated with 2 mls of growth medium in plastic tissue culture flasks, 25 cm<sup>2</sup> (Falcon, Oxnard, California). These flasks and controls containing no cells are then placed in a horizontal position and incubated at 95% humidity, 37° C, and 94% air/6% CO<sub>2</sub>.

The medium is changed every 3-5 days and the spent medium removed and frozen to await radioimmunoassay for both GH and TSH. Cell viability is checked at this time in each culture. It was determined that results obtained from such cultures were markedly altered and variable with cultures over 30 days old, and therefore only media harvested by 30 days is generally used. Cellular morphology is examined throughout the entire 30 days. Initially, the nervous tissue is plated as a population of 5-10  $\mu$ m spherical cells and after 3-4 days the cells are observed to adhere to the walls of the flask. By 8 days, a differentiated population of neurons and glia are present which resemble cells from the original tissue (see Fig. 18 ).

Pituitary cells remained as spherical cells throughout culture and became attached to the culture flask after 3-4 days.

A mouse neuroblastoma cell line, N4TGI, was obtained from Dr. R. Miller, University of Chicago, and cultured in a similar fashion to that described for primary tissue. The medium from such cultures was tested for immunoassayable GH and TSH. Cells from these cultures were also homogenized and extracted, as previously described for other tissues, and the supernatants analyzed for their hormone content by RIA.

Pooled samples of media from actively growing cultures of this cell line were lyophilized and reconstituted in 1 cc of 0.01 M phosphate buffer, pH 7.6, before elution on a Sephadex G-100 column to test the molecular identity of the immunoassayable TSH-like material.







Figure 18. Representative samples of 10 day old nervous tissue culture obtained from the amygdala of adult rats.

Somatostatin, 1  $\mu\text{g/ml}$  (Bachem Inc., Marina del Rey, California, 90201), and L-thyroxine, 1  $\mu\text{g/ml}$  (Sigma) can be added to selected primary, neural, and pituitary cultures to observe any possible regulatory mechanisms governing GH and TSH release from monolayered cells in tissue culture.

Data from all tissue culture experiments are expressed as the average amount of hormone present/ml of medium taken from five individual Falcon flasks. The amount of hormone found intrinsically in the medium, due to the serum content, can be determined from the controls containing no cells and subtracted from all results obtained from flasks containing cells.

Solutions used for tissue culture experiments involving brain and pituitary tissue:

Phosphosaline buffer 7.4 (PBS)

8.0 g NaCl

0.2 g KCL

2.0 g  $\text{Na}_2\text{HPO}_4$

0.4 g  $\text{K H}_2\text{PO}_4$

pH adjusted with 7.5% sodium bicarbonate (Gibco)

Pronase solution

1.25 g Pronase (Calbiochem)

500 ml PBS

Mix and allow to stand 30' RT<sup>o</sup>.

Centrifuged for 10 minutes and sterilized by filtration

(Millipore, Filter Corp., Bedford, Mass., 0.22 pore size).

Store at -20<sup>o</sup> C in aliquots.

Trypan Blue

1 ml Trypan Blue (Gibco)

3 ml Growth Medium

Filter before use.

Growth Medium

100 ml Basal Medium Eagle + 10% Fetal Calf Serum (Gibco)

1 ml L-glutamine, 200 mM (Gibco)

1 ml MEM non-essential amino acids, 10 mM (Gibco)

0.1 ml Gentamicin Sulfate, 5 mg/ml (Sigma)

pH adjusted with 7.5% sodium bicarbonate to 7.4.

F. SUBCELLULAR FRACTIONATION OF BRAIN HOMOGENATE TO DETECT THE LOCALIZATION OF PITUITARY-LIKE HORMONES IN THE BRAIN

Subcellular fractions of brain homogenates were prepared by differential centrifugation, as outlined in Fig. 19 .

In a typical preparation, 10 adult male rats are sacrificed by decapitation, their brains rapidly removed and the hypothalamus, amygdaloid nucleus, and thalamus immediately dissected and separated out at 4° C. Homogenization is accomplished for each pooled brain part in 1 ml ice-cold 0.32 M sucrose, 10  $\mu$ M CaCl<sub>2</sub>, pH 7.4, by 12-15 up and down strokes using a glass teflon homogenizer, 0.22  $\mu$ m clearance, driven at 800 rpm. The homogenates are sedimented at 1,000 g for 15 minutes in a Sorvall RC 5B refrigerated centrifuge. Supernatant, S<sub>1</sub>, is removed and the pellets, P<sub>1</sub>, resuspended in 0.01 M phosphosaline buffer, pH 7.6, for




Fig. 19

Fractionation Scheme utilized to determine the growth hormone content of subcellular fractions of rat brain.

radioimmunoassay. Pellet, P<sub>1</sub>, contains cellular debris, nuclei, and some myelin. Supernatants, S<sub>1</sub>, are sedimented at 20,000 g for 20 minutes in a FA 65 rotor using a Beckman LS-65 preparative ultracentrifuge. The second pellet, P<sub>2</sub>, primarily contains synaptosomes, mitochondria, and myelin and is resuspended for further analysis. Supernatant, S<sub>2</sub>, is sedimented at 150,000 g for 60 minutes. The resulting pellet, P<sub>3</sub>, consists of microsomal and other membrane fragments.

Due to the possibility of contamination of low-speed pellets with high speed components, the pellets are washed 3-4 times before each step of the centrifugation scheme. Samples of all three pellets and supernatants are then analyzed for both protein, TSH, and growth hormone content.

Furthermore, addition of <sup>125</sup>I GH and <sup>125</sup>I TSH during one subcellular fractionation can be used to check the possibility of hormone binding preferentially to a particular fraction and to determine whether peptidase activity is present in any fraction. No enhanced degradation of <sup>125</sup>I hormones and no selective binding to any particular fraction was observed using this fractionation scheme in this study, thus validating the data obtained.

The detection of antigenic material in cells of various tissues has been aided by the development of various immunohistochemical techniques.

### G. IMMUNOHISTOCHEMISTRY OF PITUITARY AND BRAIN TISSUES

Microscopic sections of tissues whose cells contain various antigenic substances, such as hormone can, in the simplest case, be incubated with specific antibodies to such antigens, and antigen-antibody complexes will form within the sectioned tissue's cells. If this second antibody is conjugated to a fluorescent or colored probe, these antigen-antibody complexes will be easily detectable in antigen or hormone-containing cells when examined under the microscope.

In order to study the tissues of various rat brain and pituitary parts for the localization of the peptide hormones GH and TSH, such immunohistochemical techniques were employed. One of these, the unlabeled antibody-enzyme-peroxidase method of Sternberger et al (1970), has provided a highly sensitive histochemical technique for the localization of antibody-antigen complexes. Its permanent reaction products provide an advantage over immunofluorescent histochemical techniques, in terms of long-term study of preparations, as no rapid fading of the labeled complex occurs with immunoperoxidase techniques as it does in immunofluorescence-treated material.

The antibody-enzyme peroxidase method of Sternberger allows the possibility of attaching an enzyme such as horseradish peroxidase to tissue antigens via a specific antibody without the use of a covalent labeling reaction. In this method, an antibody bridge system involving a soluble peroxidase-antiperoxidase complex (PAP) is bound to the primary, specific, tissue-bound antibody through a second antibody bridge. Cells

containing the specific bound-antibody are stained dark brown by the precipitated peroxidase. Schematically, the components of the peroxidase-antiperoxidase complex, unlabeled antibody stain for microscopic sections of tissue are given in Fig. 20 .

As a guide to the concentrations of antibody and times of incubation to be used in such a technique to detect TSH in tissue sections, the optimum conditions for the TSH radioimmunoassay were used. Sprague Dawley male rats, weighing approximately 200 g, were used for this study of the immunohistochemical localization of TSH in the CNS and pituitary. These animals were anesthetized using 0.1 ml/100 gm body weight, sodium pentobarbital (Holmes Serum Co., Springfield, IL). The thorax was quickly opened and the left ventricle of the exposed heart cannulated. The right atrium was cut and, using a large glass syringe, 50-75 ml of ice-cold calcium-free Tyrodes solution was very slowly perfused through the aorta. This was followed by a 30-minute perfusion of 300-500 ml ice-cold paraformaldehyde in Sorenson's buffer into the rat. The animal was kept on ice throughout this procedure. Following perfusion, the entire brain and pituitary were removed and immersed in the same fixative for 90 minutes, after which they were transferred to 5% sucrose in 0.1 M Sorenson's phosphate buffer and stored at 4<sup>o</sup> C. Before sectioning, the tissue was routinely dehydrated through a series of graded alcohols and xylene before embedding in paraffin.

Sections 6 microns in thickness were cut through the whole brain and pituitary and placed on gelatin-coated glass slides. After drying, these sections were gently heated to 40-45<sup>o</sup> C to melt the paraffin and the slide-




Figure 20. Components of the peroxidase-antiperoxidase complex unlabeled antibody stain used for the detection of rat TSH in brain and pituitary tissue.



bound tissue rehydrated through a series of xylene, alcohols (absolute - 70%), and finally in distilled water before commencing immunohistochemical staining. Staining procedures for TSH were modifications of histochemical protocols given by Moriarty (1973) and Moi Yoi et al (1979).

Deparaffinized sections were incubated in methanolic peroxide (1 vol. 3% H<sub>2</sub>O<sub>2</sub>/5 vol. methanol) for 30 minutes at room temperature to eliminate endogenous peroxidase activity. After washing in phosphate buffer (0.1 M), these sections were incubated for a further 30 minutes at room temperature with 5% aqueous egg albumin (Schwarz-Mann, Orangeburg, N.Y.) to minimize background nonspecific binding of antibody. Without washing, the sections were treated as in step 1 of the immunocytochemical staining procedure, by covering the tissue mount with the first antibody, rabbit anti-rat TSH antibody at a 1:15,000 dilution. Incubation of the slides was then carried out at room temperature for 48 hours in a moist chamber consisting of a sealed box containing dampened foam rubber so that the sections were never allowed to dry out throughout the entire procedure. After this two-day incubation, slides were thoroughly washed with 0.05 M phosphate buffer and then overlaid with the second antibody, a 1:10 dilution of goat anti-rabbit IgG. Incubation with this antibody was for 18 hours at room temperature, and after a wash step, sections were incubated for a further 30 minutes with a 1:50 dilution of rabbit peroxidase anti-peroxidase (PAP) complex in phosphate-albumin buffer. After rinsing in phosphate buffer (0.05 M), the peroxidase was localized by incubating the tissue in a solution containing hydrogen peroxide and 3,3' diaminobenzidine tetrahydrochloride for 30 minutes. Oxidation of the reagent formed a brown

precipitate at the site of the peroxidase - anti-peroxidase complexes, which was intensified by incubation in 1% OsO<sub>4</sub> for 30 seconds. The slides were then washed in phosphate buffer and mounted in Permount (Fisher) before examination microscopically using a Leitz Diavert inverted microscope. Sections were photographed using Kodak Ektachrome fast color 400 daylight film with a 35 mm microscope-mounted Leitz camera. Validation of the specificity of the reaction was tested via the following controls, which were stained for the same length of time for each tissue examined.

Negative controls were those in which the first antibody was replaced by buffer or by normal rabbit serum, diluted to 1:15,000. In some sections, the second antibody step was also replaced by addition of buffer. Immun-absorption of the anti-rat TSH serum before use in the procedure with purified TSH, overnight at room temperature at a concentration of 100 µg/TSH/ml of diluted primary antisera, was also used to test the specificity of the reaction.

The methodology described so far has applied to the histochemical detection of TSH in slices of brain and pituitary. Histochemical detection of GH in brain slices was also attempted with less success, using a less sensitive method of immunoperoxidase staining than the method of Sternberger, the "indirect method of staining."

Although the method of section preparation was similar for GH as that for TSH detection, no PAP-complex was used in the immunohistochemical staining procedure. Instead, the immune reaction was terminated at the

point of formation of an antigen-antibody complex between the second antibody, labeled with peroxidase, and the initial complex formed between first antibody and tissue-bound GH. Although cells containing a dark-colored complex could be observed, this work will have to be repeated using Sternberger's technique in an effort to obtain better histochemical resolution.

Solutions used for the histochemistry study to detect TSH and GH in brain tissue.

Sorenson's phosphate buffer 0.2 M

8.06 g  $\text{KH}_2\text{PO}_4$

37.75 g  $\text{Na}_2\text{HPO}_4 \cdot 7 \text{H}_2\text{O}$

Made up to 1 litre with distilled water, pH = 7.2.

Paraformaldehyde in Sorenson's buffer

Dissolve 40 g paraformaldehyde in 450 ml distilled water.

Heat to 55-60° C.

Add 2 N NaOH dropwise until solution clears.

To the above paraformaldehyde, add 500 ml 0.2 M Sorenson's phosphate buffer, pH 7.2.

Bring final volume to 1 litre.

Yields 1 litre of 4% paraformaldehyde in 0.1 M buffer.

Filter and store at 4° C.

Calcium (Ca<sup>++</sup>) free Tyrodes

Add 800 mls distilled water to a 1-litre flask.

NaCl	6.80 g	Glucose	1.00 g
K Cl	0.40 g	NaHCO <sub>3</sub>	2.20 g
MgCl <sub>2</sub>	0.15 g	Swirl after each component.	
MgSO <sub>4</sub> 7 H <sub>2</sub> O	0.10 g	Dilute to 1 litre. Gas with	
NaH <sub>2</sub> PO <sub>4</sub> 2 H <sub>2</sub> O	0.19 g	95% O <sub>2</sub> /5% CO <sub>2</sub> on ice.	

Slide gelatinization technique

For 100 ml:

0.5 g swine skin gelatin (Sigma)

0.05 g chromium potassium sulphate 12 H<sub>2</sub>O (Fisher)

Heat 100 ml water, dissolve gelatin and chromium potassium sulphate.

Dip slides quickly, dry in open air.

Store at 4° C wrapped in Parafilm.

Phosphate buffer 0.1M

Used for wash after fixation.

Solution A:

27.80 g NaH<sub>2</sub>PO<sub>4</sub> · H<sub>2</sub>O

1000 mls distilled H<sub>2</sub>O

Solution B:

53.65 Na<sub>2</sub>HPO<sub>4</sub> 7 H<sub>2</sub>O

1000 mls distilled H<sub>2</sub>O

(Solution B, continued)

19 mls of A + 81 mls B diluted to 200 mls, pH 7.4.

For wash between stains, diluted with water to 0.05 M.

For dilution of antisera and PAP: 0.05 M. phosphate buffer +  
2.5 mg/ml bovine serum albumin.

#### The "PAP" complex

Made in rabbit from the method of Sternberger by Miles Laboratories,  
Elkhart, Indiana.

An antigen-antibody complex dissociated and solubilized in excess  
peroxidase antigen forming a stable soluble complex.

Maintained at  $-70^{\circ}$  C in 0.1 ml aliquots.

#### Tris buffer 0.5 M

30.285 g THAM (Tris /Hydroxy-methyl/ amino methane)

250 ml H<sub>2</sub>O

pH with 1 N HCl to 7.6

Add 250 ml H<sub>2</sub>O

#### Diaminobenzidine - H<sub>2</sub>O<sub>2</sub> solution

Always freshly made; good for 60-90 minutes.

6 mg 3 - 3' diaminobenzidine hydrochloride (Sigma)

10 ml Tris buffer

1-2 drops 3% H<sub>2</sub>O<sub>2</sub> (not more than 2 months old)

Filtered and kept stirred throughout staining reaction.

Dehydration protocol for all tissues

2 x 80% alcohol	30 minutes
2 x 95% alcohol	30 minutes
2 x absolute alcohol	30 minutes
2 x chloroform	30 minutes
Paraffin	

Hydration protocol for all tissues

2 x xylene	3 minutes
2 x absolute alcohol	3 minutes
2 x 95% alcohol	3 minutes
2 x 80% alcohol	3 minutes
2 x distilled H <sub>2</sub> O	3 minutes

The next studies to be described were designed to determine the approximate molecular weight of the brain hormone in comparison to that of the hormone of pituitary origin.

H. COLUMN CHROMATOGRAPHY OF BRAIN EXTRACTS

A calibrated column of Sephadex gel provides a simple way of determining the molecular weights of proteins such as GH and TSH. Sephadex is a bead-formed gel prepared by cross linking dextran with epichlorohydrin. As a solute passes down a vertical column packed with such chromatographic beads, its movement depends on the bulk flow of the mobile phase and upon

the Brownian movement of the solute molecules. These movements cause the diffusion of molecules in the solute both in and out of the stationary phase of the beads. The separation of solutes in gel filtration depends on the different abilities of the various sample molecules to enter pores which contain the stationary phase. Very large molecules which never enter the stationary phase therefore move through the chromatographic gel bed faster than smaller molecules which can enter the gel pores, and thereby move more slowly through the column since they spend a part of their time in the stationary phase. Molecules are therefore eluted in order of decreasing molecular size. Results are usually expressed in the form of an elution diagram showing the variation of solute concentration in the eluent with the volume of eluent passed through the column. Sephadex G-100 was the gel of choice for these experiments, being able to separate clearly molecules with a molecular weight range between 4,000 - 150,000 Daltons. This range was chosen in order to detect the possibility of any larger prohormone-like molecules that may have been present in brain homogenates.

From a total of 40 male, 200 gm rats, brains were dissected into the amygdaloid nucleus, hypothalamus, and thalamus. Each brain region was pooled and kept on dry ice before processing. Extraction was as described in section D, except that the final supernatant was spun at 3,000 g for 10 minutes in a FA-65 rotor of a Beckman ultracentrifuge to remove nuclei and other debris. This supernatant was then lyophilized and reconstituted either in 1 ml veronal buffer, pH 8.6, for GH studies, or in 1 ml 0.01 M phosphosaline buffer, pH 7.6, for TSH column work.

Sephadex G-100, 0.9 x 50 cm columns were prepared, coated with albumin, and calibrated using blue dextran 2000 (Pharmacia) bovine serum albumin, mol. wt. 68,000 D, and <sup>125</sup>I insulin, mol. wt. 5,700 D, to give reliable calibration points.

A trace amount, approximately 7,000 cpm in 100  $\mu$ l buffer of either <sup>125</sup>I rat pituitary GH or TSH, was applied to the column together with 1 ml of reconstituted brain homogenate. One-ml fractions were collected in a Gilford automatic fraction collector. These were analyzed for both radioactivity, using a Micromedic gamma counter, or for hormone content, using the radioimmunoassays for TSH or GH. In this way the elution pattern of the radioactive-labeled pituitary growth hormone could be compared to that of the radioimmunoassayable brain hormone. Similar studies were performed using material from fetal brain and pituitaries.

Another parameter of the brain "pituitary-like" hormones examined was their immunological similarity to their respective pituitary counterparts.

#### I. STUDIES TO DETERMINE THE IMMUNOLOGICAL IDENTITY OF "BRAIN" HORMONES

Immunological similarities between brain and pituitary hormones were determined by preparing serial dilutions of both tissue extracts and subjecting them to radioimmunoassay. Parallelism, or lack of parallelism, between logit log transformed displacement curves produced for each tissue was used as an indicator of probable similar or different interaction with antibody sites to a particular hormone.



The biological activity of brain hormones was also compared with that of pituitary GH and TSH.

## J. BIOLOGICAL ACTIVITY STUDIES

### a. Growth Hormone Bioactivity in Brain Extracts.

The traditional bioassay for the measurement of pituitary growth hormone has been that perfected by Greenspan et al in 1949. It had been shown earlier by Li et al (1945) that pituitary growth hormone extract stimulated growth of the bony epiphyseal plate of hypophysectomized rats. This observation formed the basis for the Greenspan assay which is based on the ability of growth-hormone-like material in tissue extracts to increase the width of the epiphyseal cartilage in hypophysectomized animals, a parameter measured quantitatively with a microscope micrometer. Li et al (1945) showed that a total dose of 100  $\mu$ gms of pure growth hormone administered to 30-day post-hypophysectomized adult rats was necessary to produce a significant response in an increase in body weight bioassay for GH, whereas in the tibia epiphyseal cartilage test, as noted by Greenspan, a total dose of only 5  $\mu$ gms would produce a significant response. Therefore without necessarily obtaining a growth hormone response in terms of weight gain, the use of the tibia bioassay provides a characteristic of growth hormone action which is able to be measured.

This experiment was repeated four times in hypophysectomized rats, twice using female and twice using male recipients. For each experiment, 250 donor rats were sacrificed by decapitation and the amygdaloid nuclei

and pituitaries harvested individually and placed immediately on dry ice. Each tissue was then weighed, pooled, and homogenized separately, taking care to avoid any possible contamination of brain homogenate with pituitary tissue. The supernatants obtained from extraction of the tissues with 0.01 phosphate buffer, pH 7.6, were spun at 5000 g to eliminate any debris and nuclear material, dialyzed against 0.15 M saline, lyophilized, and reconstituted in 20 ml of physiological saline (0.15 M, pH 7.4). The final concentration of GH in both pituitary and brain extracts was measured by the RIA described earlier and appropriate dilutions to give approximately 20-30  $\mu$ g of immunoreactive GH/ml obtained. A more concentrated pituitary extract was found to give no advantage over the lower pituitary dosage.

For the actual bioassay, forty-two-day-old male and female rats, which had been hypophysectomized at age 28 days, were kept in groups of five in separate cages and fed a diet of oranges and rat chow mash. Groups of five animals received twice a day for four days either: 0.5 ml saline, 0.5 ml rat pituitary extract, or 0.5 ml of amygdaloid nucleus extract intraperitoneally. Animals were weighed daily at the time of injection to measure any weight gain. Twenty-four hours after the last injection, all animals were weighed and sacrificed using sodium pentobarbital injection. One or both tibias were dissected free of soft tissue and split with a sharp razor in the mid-sagittal plane. The two-bone halves were either stained immediately or fixed in 10% neutral formalin. Thyroids, ovaries, and testes were similarly dissected from all control and assay animals and fixed in 10% neutral formalin.

The staining procedure for the tibia epiphyseal cartilage halves is as described by Evans et al (1943) and is as follows:

1. Bone washed in H<sub>2</sub>O - 30 minutes
2. Acetone - 60 minutes
3. Wash in H<sub>2</sub>O - 30 minutes
4. Freshly prepared 2% silver nitrate - 1-3 minutes
5. Wash in H<sub>2</sub>O and while under water, expose to a strong light until calcified portions appear dark brown.
6. 10% sodium thiosulfate - 25-30 seconds
7. Wash in running H<sub>2</sub>O - 30 minutes
8. Store in 80% alcohol in dark at room temperature.

The width of the uncalcified epiphyseal cartilage, which remained unstained and easily observable, is measured under the low power of an American Optical microscope using a micrometer eye piece calibrated with a stage micrometer so that results can be expressed as micra units. A minimum of 10 measurements should be made along each epiphyseal cartilage in areas selected at random. This study was carried out "blind", and two independent observers used. The data between saline-, pituitary-, and amygdaloid-injected groups was statistically analyzed using the Student T test. The cartilage plate increases in width slightly from anterior to posterior, and there is therefore an inherent error in the measurement of mean width. For this reason, all readings should always be from narrow to wider areas and the variation accounted for by the multiple readings. There was usually no significant difference between left and right tibias, but only one tibia, usually the left, was used for the

measurements. A comparison in the total body weights and wet weights of testes and ovaries from both control and hypophysectomized rats was also statistically analyzed.

b. Thyroid stimulating hormone bioactivity in brain extracts.

The effects of TSH on the histology of the thyroid follicle cell were previously discussed in the Review of Literature section of this dissertation. Briefly, the presence of TSH influences the size and shape of the follicle epithelial cells and the distribution of colloid in the follicle lumen. Follicles from hypophysectomized rat thyroids contain cells which are flattened, and little colloid is found in the central lumen of the follicle, as compared to normal, euthyroid animal thyroid follicles.

A minimum dose of 20 ng immunoreactive pituitary TSH/hypophysectomized animal, given over a four-day period, is necessary to restore the histology of the thyroid to that of an intact rat. To demonstrate another biological effect of pituitary TSH on the thyroid, that of an increase in thyroidal radioiodide uptake, at least 10  $\mu$ gm of TSH/rat are required.

For our experiments, aimed at assaying TSH-like activity of CNS extracts, thyroids from the three groups of animals used in the GH bioactivity experiments were used, i.e., hypophysectomized animals receiving saline, or CNS extract, or pituitary extract. A fourth group of hypophysectomized animals were also used in this TSH bioassay which received a higher dose of pituitary extract than the animals used in the GH bioassay.

In terms of levels of radioimmunoassayable TSH, these four groups of rats received 0  $\mu$ g of TSH in the saline-injected group; 4.5  $\mu$ g TSH, from amygdaloid tissue, in the CNS-injected group; 5  $\mu$ g TSH, obtained from pituitary extracts, in the third group; and 100  $\mu$ g TSH, from pituitary extracts, in the fourth group of rats.

Twenty-four hours after the last injection, the thyroids were removed, en bloc, with the trachea, fixed in neutral formalin, and subsequently paraffin-sectioned at a width of 6 microns and stained with hematoxylin eosin. Sections were photographed in order to examine the follicle cells for any histological changes which could be related to the type of material injected. This method of bioassaying TSH represents a modification of the work of Tice (1974).

Another test of TSH's biological effectiveness was to examine this hormone's property to facilitate the uptake of circulating radioiodine by the thyroid. Hypophysectomized animals' ability to take up radio-labeled iodine into the thyroid with, and without, injections of immunoassayable TSH in extracts from brain and pituitary were compared. Similarly, the effectiveness of brain and pituitary extracts to increase the circulating levels of  $T_4$  in hypophysectomized animals was examined as a measure of the bioactivity of brain TSH-like material.

In these two groups of experiments, i.e., measurement of iodine uptake and measurement of circulating  $T_4$  levels, four groups of 5 adult, 14-day, post-hypophysectomized animals received one of the following treatments over a 4-day period: either an IP injection of 0.5 ml of

saline daily, or a total dose of 100  $\mu\text{g}$  immunoassayable TSH from pituitary extracts in 0.5 ml saline, or a total dose of 2  $\mu\text{g}$  TSH from pituitary extracts, diluted saline, or 4.3  $\mu\text{g}$  TSH obtained from amygdaloid tissue homogenates diluted in saline. Eighteen hours before sacrifice, approximately 1.5  $\mu$  curies <sup>131</sup>I was injected intraperitoneally into all groups, and after sacrifice, thyroid and muscle levels of radioactivity were measured. Thyroxine ( $T_4$ ) levels of samples of blood, taken from the trunk by gravity after decapitation, were measured by means of a commercial RIA kit (Tetra-Tab), which is described in detail in section K of this Materials and Methods chapter.

Although the blood-brain barrier prevents the passage of a majority of blood-borne materials into the brain, its effectiveness in preventing the contamination of CNS tissue by pituitary-derived GH and TSH was tested by methods described in the following section.

#### K. EXAMINATION OF RAT BLOOD-BRAIN BARRIER INTACTNESS TO PITUITARY GROWTH HORMONE AND THYROID STIMULATING HORMONE

Numerous chemical substances do pass from the bloodstream into the brain at rates which are slower than for entry into all other organs in the body. There are similarly slow rates of transport between the CSF and the brain. The permeability barrier appears to be the end result of numerous contributing factors that present diffusional obstacles to chemicals on the basis of molecular size, charge, solubility, and specific carrier systems. The difficulty has not been in establishing the existence of these barriers, but rather in determining their mechanisms. When the relatively small protein, 43,000 D, horseradish peroxidase is injected

intravenously into rats, its intracellular location within most tissues, such as muscle, can be demonstrated histochemically with the electron microscope. As opposed to the easy transvascular movement of horseradish peroxidase across muscle capillaries into muscle tissue, in brain the peroxidase molecule is unable to penetrate through the continuous layer of vascular endothelial cells constituting the blood brain barrier. This single layer of brain capillary endothelium differs from those of other tissue capillaries such as muscle and heart, in that the intercellular zones of cell membrane apposition are much more highly developed in the brain and are virtually continuous along the entire surface of these cells. They also show a lack of pinocytotic vesicles. These vesicles, characteristic of capillaries found elsewhere throughout the body, have been related to transvascular carrier systems of both large and small molecules. Since the horseradish peroxidase enzyme marker can neither go through nor between the endothelial cells comprising the vascular system of the CNS, an operationally defined barrier is said to exist, i.e., the blood-brain barrier.

For a solute to enter the brain easily, it appears to have to be either highly lipid-soluble and be able to diffuse through cell membranes or have an affinity for certain specific and selective carrier-mediated transport systems present in the endothelial cell plasma membrane, as shown by Oldendorf (1977).

In order to test the possibility that circulating GH or TSH of pituitary origin could be reaching the CNS tissue via possible weak areas in the blood-brain barrier, systemically administered radiolabeled

hormones were used as markers of distribution, as described by Stern et al (1975). Using freshly iodinated GH or TSH, five adult animals per hormone were injected intravenously via the tail vein with approximately  $1 \times 10^6$  cpm hormone in 200  $\mu$ l phosphosaline buffer, pH 7.6. Thirty minutes later the animals were decapitated and the brain areas and various peripheral tissues dissected out. All tissue samples were weighed and counted in a gamma counter to detect the presence of radioactivity arising from the labeled hormone injection. Results were expressed as cpm/mg wet weight of tissue. Control rats injected with 200  $\mu$ l of buffer alone were similarly killed and the tissues dissected out for detection of immunoassayable hormone, expressed as ng/mg wet weight of tissue.

For the studies involving fetuses, to test the possibility that the blood-brain barrier develops with age, the following protocol was used.

Pregnant females in the 20-21st day of pregnancy were decapitated and the fetuses in the uterine horns gently lifted out of the body cavity and kept warm. Once released from the uterus, these full-term fetuses, released from their attached placentas, were able to survive for several hours. To test the intactness of the blood brain barrier at this early age,  $1 \times 10^5$  cpm of either labeled GH or TSH were injected intraperitoneally in 10  $\mu$ l PBS into the fetuses. Thirty minutes later, the fetuses were sacrificed by decapitation with scissors and various brain parts and organs dissected out to determine the location of the radiolabeled tag. If there was the possibility of some breakdown in the blood brain barrier, and if the blood content of the brain is approximately 50  $\mu$  litres/gm of



brain tissue, by assaying the hormonal content of sera, it was possible to calculate the possible level of contamination from circulating hormone of brain tissue.

#### L. THE EFFECTS OF HYPOPHYSECTOMY ON BRAIN-BASED PITUITARY-LIKE GH AND TSH.

In this series of three experiments, the effect of removing the pituitary on levels of immunoreactive hormone in various areas of the brain and in serum were compared to levels obtained from intact, sham-operated controls of equivalent age. For each experiment, approximately 80 hypophysectomized animals and 80 controls were obtained from Hormone Assay Lab., Inc., Chicago. The hypophysectomized animals were operated on at 28 days of age, and batches of 20 animals were kept for 7, 14, 28, and 48 days post-surgery on a diet of oranges and mashed moist rat chow before sacrifice. Salinated water was found to be unnecessary for these animals.

Equivalent groups of intact controls, in terms of age, were sacrificed within one hour of the sacrifice of the hypophysectomized rats. Trunk blood was collected for hormone analysis and the brain parts dissected out on dry ice as previously described in section C. Extraction and homogenization were also performed as previously noted in section D of this dissertation, and levels of hormone were calculated, after radioimmunoassay, as ng hormone/mg protein or ng hormone/mg wet weight of tissue.

The effectiveness of hypophysectomy was determined by gross examination of the sella turcica with a dissecting microscope and also by measuring the levels of circulating hormones GH, TSH, and the thyroid hormone T<sub>4</sub>,

as well as the " $T_3$  resin uptake," another indicator of thyroid and indirectly pituitary function, which will be described in the next section.

M. THYROXINE ( $T_4$ ) AND  $T_3$  RESIN UPTAKE ASSAYS

The levels of circulating total thyroxine or  $T_4$  were obtained using a commercial  $T_4$  diagnostic kit, "Tetra-Tab RIA" (Nuclear Medical Lab., Inc., Dallas, Texas), and the " $T_3$  uptake" estimation of unsaturated thyroid binding globulin capacity was also run using a kit, "Tri-Tab," from the same company.

a. Measurement of  $T_4$

Thyroxine circulates in the bloodstream as free (1%) and thyrobinding globulin-bound hormone. In the  $T_4$  radioimmunoassay, total thyroxine,  $T_4$ , is released from endogenous thyroxine-binding proteins present in the serum by the chemical inactivation of the binding protein by 0.025 N HCl. The  $T_4$  so released is then available for assay of total  $T_4$ . The principle of the RIA is as described for GH and TSH. Specifically for  $T_4$  detection in the assay, a serum sample is mixed with 200  $\mu$ l of 0.025 N HCl and a constant amount of iodinated  $T_4$  is added and the mixture incubated at room temperature for 30 minutes with a highly specific antibody to  $T_4$  prepared in rabbits. Free and antibody-bound fractions are then separated by ammonium sulfate precipitation and centrifugation of the antibody-bound fraction. Following centrifugation, the supernatant was discarded and the radioactivity in the precipitate measured directly in a gamma counter. A standard curve was prepared calculating the percent of the

total radioactivity remaining in the bound fraction for the five serum standards ranging from 1 - 28  $\mu\text{g T}_4/\text{dl}$  and plotted against  $\text{T}_4$  concentration. Unknown  $\text{T}_4$  values were calculated for the experimental serum samples directly from the standard curve as  $\mu\text{gm}/\text{dl}$ .

(i) Assay procedure for the detection of  $\text{T}_4$  in serum

Assay carried out in triplicate in 12 x 75 mm plastic tubes, and reagents added in the following order:

1. 200  $\mu\text{l}$  - 0.025 N HCl.
2. 10  $\mu\text{l}$  - standard or serum sample.

Standards = 1 - 28  $\mu\text{g T}_4/\text{dl}$ .

All tubes shaken 10-15 seconds and incubated at RT for 60 minutes.

3. 400  $\mu\text{l}$  <sup>125</sup>I  $\text{T}_4$ .

All tubes shaken and incubated for 60 minutes.

4. 2.0 ml - 35.5% ammonium sulfate containing 0.4% calf serum.

All tubes mixed gently and incubated 20 minutes before centrifugation for 10 minutes at 1000 g.

The supernatant was decanted and the precipitate counted, accumulating a minimum of 20,000 counts/tube. Reproducibility was assessed using frozen serum pools as internal standards for each assay.

b. Measurement of the radiolabeled  $\text{T}_3$  resin uptake in serum samples.

The radiolabeled  $\text{T}_3$  resin uptake test which is not a measurement of circulating  $\text{T}_3$  or  $\text{T}_4$  hormones is designed to assess the unsaturated

binding capacity of certain serum proteins, primarily thyroxine-binding globulin (TBC). This assessment is accomplished in the assay by the addition of an excess of  $^{125}\text{I}$ -labeled L-triiodothyronine ( $\text{T}_3$ ) to serum in order to saturate all available TBG binding sites in the serum sample. If the endogenous  $\text{T}_4$  level is increased, as in hyperthyroidism, the TBG will be relatively saturated. More  $^{125}\text{I}$ -labeled  $\text{T}_3$  remains in a free state to be adsorbed by a secondary silicate binder. Conversely, when  $\text{T}_4$  output is low, as in hypothyroidism, more of the added labeled  $\text{T}_3$  will bind to the serum TBG, yielding a comparatively low percentage of the remaining radioactivity to be adsorbed secondarily. For a secondary binding agent, the commercial kit uses an inorganic adsorbent, silicate, as binding of  $^{125}\text{I}$ -labeled  $\text{T}_3$  reaches over 99% of the maximum uptake within five minutes and remains constant for up to three hours. Basically, this test involves mixing a serum sample with excess  $^{125}\text{I}$ -labeled  $\text{T}_3$  which occupies all unsaturated binding sites on the TBG in the serum.  $^{125}\text{I}$   $\text{T}_3$  not bound to TBG was removed by addition of the secondary silicate binder. Following centrifugation, the supernatant was discarded, and the radioactivity in the sample compared to the amount of radioactivity in a normal control serum, with an established percent uptake. Interassay reproducibility was assessed using a pooled, frozen internal serum standard.

(i) Assay procedure for the measurement of the  $\text{T}_3$  resin uptake in serum

Assay carried out in triplicate in 12 x 75 mm plastic tubes at room temperature.

1. 100  $\mu$ l standards or serum sample.
2. 2 ml -  $^{125}$ I  $T_3$  in 0.07 M barbital buffer containing 0.125% BSA.  
Mixed thoroughly.
3.  $T_3$  adsorbent tablet containing 10%-40% silicate by weight added.  
Mixed for 5 seconds and incubated 10 minutes.

Centrifuged 5 minutes at 1000 g, supernatant discarded and the pellet counted, in a gamma counter, each tube accumulating at least 20,000 counts. The percent  $T_3$  uptake is determined by dividing the test counts/the total counts x 100. An accurate and practical assessment of thyroid status, and therefore indirectly of pituitary TSH status, can be made by calculating the product of the radiolabeled  $T_3$  resin uptake and the total  $T_4$  value, the "Free Thyroxine Index."

#### N. ALTERATION OF FEEDBACK CONTROL MECHANISMS

##### I. TARGET ORGAN REMOVAL

The effect on pituitary hormones of removing a target organ has previously been alluded to in the discussion on feedback control mechanisms. For example, removal of the thyroid gland serves to remove the "negative feedback" control which thyroxine,  $T_4$ , the major product of the thyroid gland, has upon TSH production from the pituitary. Adrenalectomy and gonadal removal also affect the levels of GH and TSH in the pituitary, indicating that circulating steroid hormones have a certain amount of influence in the release of hormones from the pituitary. The question posed by this series of experiments was whether target gland removal would

affect the levels of pituitary-like peptide hormone found in the central nervous system in a similar way to effects of target gland removal on pituitary hormone content and release. If the answer was "no", it would be another indication supporting a separate and distinct pool of pituitary-like peptides in the brain.

a. Oophorectomy.

Groups of 20, sixty-day-old female rats were oophorectomized under light sodium pentobarbital anesthesia through bilateral paraspinal 1-cm incisions. A single ligature was made behind the ovary around the fallopian tube and the ovary with its surrounding capsule and fat cut away. Sham-operated animals were similarly handled without the tying of the fallopian tubes or removal of the ovaries. Both sham and operated animals were sacrificed either one or four weeks post-operatively.

b. Orchidectomy.

Groups of 20, 60-day-old male rats were lightly anesthetized and a longitudinal incision made in each scrotal sac. The testicle, along with the epididymus, were exposed and vascular structures were ligatured. The testicle was removed and the scrotal sac reanastomosed with 2 0 silk. Sham operated animals were similarly handled without ligaturing. As in the oophorectomized animals, both sham and castrated rats were sacrificed either one or four weeks post-operatively.

c. Adrenalectomy.

Under light anesthesia, groups of 20, sixty-day-old male rats were adrenalectomized or sham adrenalectomized. Bilateral paraspinal 1-cm incisions were made to expose the kidney and adjacent adrenal gland embedded in perirenal fat. Care was taken to remove the adrenal gland as gently as possible with curved forceps, in order to avoid disruption and dispersal of adrenal cells. Adrenalectomized animals were kept on 0.9% saline and normal rat chow. Both sham and adrenalectomized rats were sacrificed after one or four weeks post-operatively.

d. Thyroidectomy.

Thyroidectomized animals were obtained from Hormone Assay Labs, Inc., and kept for one or four weeks before sacrifice. Sixty-day-old male rats, equivalent to the thyroidectomized animals, were sham operated by making an incision over the trachea and gently parting the muscles overlaying the trachea to reveal the thyroid gland. Sacrifice was as for the thyroidectomized animals. Efficiency of thyroidectomy was confirmed by analysis of the levels of TSH,  $T_4$ , and  $T_3$  uptake obtained from serum samples.

In all target organ experiments, the animals were sacrificed by decapitation, trunk blood collected, and brain parts and pituitary removed for tissue homogenization and radioimmunoassay. The levels of pituitary hormone and similar CNS peptides in intact and operated animals were compared after one week and four weeks of target organ absence.

## II. THYROXINE

The effect of injecting intramuscularly 0.4  $\mu\text{g}$   $\text{T}_4$ /gm body weight daily for seven days into groups of 20 hypophysectomized or intact animals on levels of brain and pituitary hormones was also examined using saline-injected animals as controls.

Trunk blood was collected at sacrifice and TSH,  $\text{T}_3$  uptake, and  $\text{T}_4$  levels analyzed for all groups.

Brain parts were homogenized and treated as before with levels of immunoassayable hormone being detected by radioimmunoassay.

### Preparation of thyroxine:

9 mg L-thyroxine/sodium salt-pentahydrate (Sigma)

90 mls phospho-buffered saline 0.01 M, pH 7.6

The thyroxine required dissolving in a few drops of 1 N NaOH before addition of phosphosaline buffer to make a 0.1 mg in 1 ml solution. Appropriate dilutions with buffer were made to give the correct dosage for individual rats according to weight. Control rats were injected with buffer only.

## O. ONTOGENEY STUDIES

Accumulating knowledge of the importance of the fetal endocrine system on morphogenesis, differentiation, and parturition in both man and other mammals prompted this study of the ontogenesis of brain pituitary-like hormones. It was also hoped that the presence of CNS peptides could



be detected earlier than pituitary peptides which would give further credence for their separate identities.

Eight pregnant rats were housed in separate cages from the day of insemination, which was taken as day one of gestation. Staging of estrous cycle periods and confirmation of fertilization was determined from daily microscopic examination of vaginal scrapings suspended in saline and viewed with a 10 x ocular. Fetal and neonate animals were decapitated at various stages of development, pre- and post-partum. Trunk blood, amniotic fluid, and samples of brain and pituitary tissue were frozen prior to determination of GH and TSH levels by radioimmunoassay. Levels of hormone in maternal blood and brain were also determined. Levels of hormone were expressed as ng/mg protein. Immunologic and physiochemical characteristics of the two fetal brain hormones, GH and TSH, were compared with those of pituitary origin in a similar manner to adult hormones, i.e., by column chromatography and radioimmunoassay parallelism studies. Attempts to trace the origin of fetal brain hormones centered on the measurement of placental, amniotic, fetal, and maternal sera levels of hormone, and the intactness of the fetal blood brain barrier was examined as described in section I. In order to confirm the presence of a placental barrier to material pituitary hormones,  $1 \times 10^6$  cpm of  $^{125}\text{I}$  GH and TSH were injected into the pregnant mothers at 21 days of gestation in 200- $\mu\text{l}$  amounts via the tail vein. Thirty minutes later, fetuses were removed and the kidney, pituitary, and brain checked for radioactivity. The presence of radioactivity would indicate the possibility of pituitary hormones crossing the placental barrier; its absence would indicate barrier intactness.

P. STATISTICAL ANALYSIS OF DATA

Except for the target organ experiments, all data was analyzed by the two-tailed Students T Test which compared the mean and standard error of the mean to give the probability or "p" values of any difference between two groups being significant. This "p" value can be expressed, as for example  $p = < 0.05$ , which would indicate that the two sets of data compared have less than a 5% chance of being the same, or if  $p = < 0.01$ , in which case the two sets of data would have a less than 1% chance of being the same (Steel and Torrie, 1960).

As there was unexplainably wide variation in the data from the target organ experiments, it was decided to handle the data from these experiments using the Mann-Whitney  $\mu$ -Two-Tailed Statistical Test in which the non-parametric confidence interval is based on the rank sum test (Mann and Whitney, 1947).

## 5. RESULTS

### A. PITUITARY GROWTH HORMONE AND THYROID STIMULATING HORMONE-LIKE IMMUNOREACTIVE PEPTIDE DISTRIBUTION IN RAT BRAIN

Brain extracts from specific parts of the CNS of adult rats were examined for their hormonal content using the radioimmunoassays for rat growth hormone (GH) and rat thyroid stimulating hormone (TSH). The results are depicted in Tables 3 - 8 and Figures 21 and 22.

#### a. Growth Hormone-Like Immunoreactivity in the Rat Brain

The data in Tables 3 and 4 demonstrate the levels of immunoreactive growth hormone present in various brain regions and in the pituitary of the adult male rat expressed in terms of ng/mg protein or pg/mg wet weight of tissue, respectively. The amygdaloid nucleus was found to contain the largest amounts of this particular immunoreactive material per mg of protein,  $5.28 \pm 0.21$  ng, and  $284 \pm 10$  pg/mg wet weight of tissue. The hypothalamus,  $3.16 \pm 0.33$  ng/mg protein,  $119 \pm 6$  pg/mg wet tissue, and the caudate,  $2.09$  ng/mg protein,  $117 \pm 5$  pg/mg wet tissue, had lower levels. Even smaller quantities were detected in the hippocampus:  $1.35 \pm 0.04$  ng/mg protein,  $95 \pm 3$  pg/mg wet tissue; cortex,  $1.23 \pm 0.13$  ng/mg protein,  $48 \pm 7$  pg/mg wet tissue; and thalamus,  $1.59 \pm 0.15$  ng/mg protein,  $99 \pm 8$  pg/mg wet tissue, with barely detectable amounts being found in the cerebellum-- $0.03 \pm 0.01$  ng/mg protein, and  $5 \pm 2$  pg/mg wet tissue. These values are to be contrasted to the amount of GH found in the anterior pituitary gland, i.e.,  $1.3 \times 10^6 \pm 0.45$  ng/mg protein and  $2.8 \times 10^5 \pm 0.26$  pg/mg wet tissue.

TABLE 3

## DISTRIBUTION OF IMMUNOREACTIVE GH IN ADULT RAT BRAIN

(ng/mg protein)\*\*

	<u>n</u>	<u>GH (ng/mg protein)**</u>
Amygdaloid nucleus	13*	5.28 ± 0.21
Hypothalamus	50*	3.16 ± 0.33
Caudate	10*	2.09 ± 0.24
Thalamus	13*	1.59 ± 0.15
Hippocampus	11*	1.35 ± 0.04
Cortex	12*	1.23 ± 0.13
Cerebellum	13*	0.03 ± 0.01
Anterior pituitary	30	1.3 x 10 <sup>6</sup> ± 0.45 x 10 <sup>6</sup>

n = number of experiments.

\* represents a pool of 10 animals.

\*\* Mean ± S. E. M.

TABLE 4

## DISTRIBUTION OF IMMUNOREACTIVE GH IN ADULT RAT BRAIN

(pg/mg wet weight tissue)\*\*

	<u>n</u>	<u>GH (pg/mg wet wt.)**</u>
Amygdaloid nucleus	13*	284 ± 10
Hypothalamus	50*	119 ± 6
Caudate	10*	117 ± 5
Thalamus	13*	99 ± 8
Hippocampus	11*	95 ± 3
Cortex	12*	48 ± 7
Cerebellum	13*	5 ± 2
Anterior pituitary	30	$2.8 \times 10^5 \pm 0.26 \times 10^5$

n = number of experiments.

\* represents a pool of 10 animals.

\*\* Mean ± S. E. M.

TABLE 5

## DISTRIBUTION OF IMMUNOREACTIVE GH IN ADULT RAT BRAIN REGIONS


(ng/total brain region)\*\*

n = 12	<u>ng GH/brain region</u>
* Cortex	38.40 ± 0.74
* Amygdala	14.20 ± 1.20
* Hippocampus	6.11 ± 0.56
* Caudate	5.40 ± 0.11
Hypothalamus	4.76 ± 0.27
* Thalamus	3.96 ± 0.63
* Cerebellum	1.00 ± 0.05
Anterior pituitary	3.6 x 10 <sup>6</sup> ± 0.39 x 10 <sup>6</sup>


n = number of experiments.

\* represents a pool of 10 animals.

\*\* Mean ± S. E. M.



CORONAL SECTION



GH ng/mg PROTEIN  
■ > 5  
▨ > 2  
▤ > 1  
□ NOT DETECTABLE

Figure 21. Distribution of immunoreactive growth hormone in the adult rodent brain.

When expressed as the total amount of hormone present in an entire brain region, as shown in Table 5, the cortex was shown to contain the largest overall amount of immunoreactive growth hormone-like immunoreactive material in the brain. Nevertheless, the total amount of growth hormone-like material present in an adult rat brain is still  $1.5 \times 10^4$  less than that found in the adult rat pituitary.

The amount of immunoreactive growth hormone found in the systemic circulation was shown to be approximately  $249.80 \pm 35.32$  ng/ml, a value similar to what has been previously reported (Ojeda *et al.*, 1977). The amount of hormone per mg of brain tissue which could be attributed to circulating hormone contamination was determined to be approximately 5-12 pgs/mg wet weight of tissue, assuming the blood content of brain tissue to be 50  $\mu$ l/gm. Taking into account this level of possible contamination and ignoring the blood-brain barrier effectiveness factor, the amount of hormone detected in cerebellar tissue could therefore be considered artifactual or merely the amount of immunoreactive GH present in the vasculature of the cerebellum. However, all other brain areas have immunoreactive material in amounts considerably greater than that possibly due to contamination of the tissue by blood-borne products.

b. Thyroid Stimulating Hormone-Like Immunoreactivity in the Rat Brain.

When the levels of immunoreactive TSH were similarly mapped for the brain areas of the adult rat, the results obtained are shown in Tables 7 and 6 for ng/mg protein or ng/mg wet weight of tissue, respectively. The distribution of TSH-like material/brain region was calculated as shown in



TABLE 6

## DISTRIBUTION OF IMMUNOREACTIVE TSH IN ADULT RAT BRAIN

(ng/mg wet wt. tissue)\*\*

	<u>n</u>	<u>TSH (ng/mg wet wt. tissue)**</u>
Hypothalamus	50	5.443 ± 0.56
Amygdala	17*	1.100 ± 0.42
Thalamus	10*	0.967 ± 0.10
Caudate	11*	0.810 ± 0.12
Cerebellum	12*	0.799 ± 0.05
Hippocampus	9*	0.798 ± 0.04
Cortex	13*	0.726 ± 0.09
Anterior pituitary	50	1.9 × 10 <sup>5</sup> ± 0.50 × 10 <sup>5</sup>

n = number of experiments.

\* Represents a pool of 10 animals.

\*\* Mean ± S. E. M.

TABLE 7

## DISTRIBUTION OF IMMUNOREACTIVE TSH IN ADULT RAT BRAIN

(ng/mg protein)\*\*

	<u>n</u>	<u>TSH (ng/mg protein)**</u>
Hypothalamus	50	204.20 ± 3.6
Amygdala	17*	29.42 ± 6.3
Cerebellum	12*	16.98 ± 2.5
Thalamus	10*	16.50 ± 1.4
Caudate	11*	15.21 ± 2.6
Cortex	13*	13.89 ± 0.5
Hippocampus	9*	11.88 ± 2.4
Anterior pituitary	50	1.2 × 10 <sup>6</sup> ± 0.75 × 10 <sup>6</sup>

n = number of experiments.

\* Represents a pool of 10 animals.

\*\* Mean ± S. E. M.

TABLE 8


DISTRIBUTION OF IMMUNOREACTIVE TSH IN ADULT RAT BRAIN REGIONS  
(ng/total brain region)\*\*

n = 15	<u>ng TSH/brain region**</u>
* Cortex	580.21 ± 5.6
Hypothalamus	217.72 ± 7.3
* Cerebellum	158.03 ± 2.5
* Hippocampus	49.47 ± 3.0
* Amygdala	44.88 ± 3.6
* Thalamus	39.06 ± 4.1
* Caudate	37.42 ± 3.9
Anterior pituitary	2.7 x 10 <sup>6</sup> ± 0.65 x 10 <sup>6</sup>


n = number of experiments.

\* represents a pool of 10 animals.

\*\* Mean ± S. E. M.



CORONAL SECTION



TSH ng/mg PROTEIN

- > 200
- ▣ > 20
- ▤ > 15
- ▥ > 10
- < 10

Figure 22. Distribution of immunoreactive TSH in the adult rodent brain.

Table 8. If measured in terms of total amount of hormone/brain region, the cortex contains the highest level of any other brain region,  $580.21 \pm 5.6$  ng/cortex, although the total amount of brain TSH is  $10^3$  times less than TSH found in the anterior pituitary,  $2.7 \times 10^6 \pm 0.65$  ng/pituitary. When calculated as per mg of protein or as wet weight of tissue, the hypothalamus shows by far the highest concentration of immunoreactive TSH,  $204.20 \pm 3.6$  ng/mg protein and  $5.443 \pm 0.56$  ng/mg tissue, followed by the amygdaloid nucleus,  $29.42 \pm 6.3$  ng/mg protein and  $1.10 \pm 0.42$  ng/mg wet tissue. Lesser amounts were detected in other regions of the brain and varied from the cerebellum,  $16.50 \pm 1.4$  ng/mg protein,  $0.799$  ng/mg wet tissue, to the hippocampus,  $11.88 \pm 2.4$  ng/mg protein and  $0.798 \pm 0.04$  ng/mg wet tissue.

Assuming the level of TSH in serum to be approximately  $489.67 \pm 52.59$  ng/ml, as measured by the RIA, and using the same factor as for growth hormone, the amount of hormone which could be contributed by possible blood borne contamination of brain tissue would be  $24.5$  pg/mg wet weight of tissue. Thus, significant amounts of a TSH-like material were found to be present in all brain regions examined.

#### B. GROWTH HORMONE AND THYROID STIMULATING HORMONE-LIKE IMMUNOREACTIVE PEPTIDE DISTRIBUTION IN PRIMATE BRAIN

These data are considered preliminary as only 3-8 brains were examined; however, because they show findings comparable to those observed in the rodent brain and extend relevancy to these observations, they will be reported here. Variables beyond the control of the experimenter such

as prior treatment of the animals and the use of fixed material may have been present to alter these results.

a. Growth Hormone-Like Immunoreactivity in the Primate Brain

From a total of 8 adult monkeys, the amount of immunoreactive growth hormone in various brain regions was calculated and expressed in Table 9 as pg/mg wet tissue. In the primate, the hypothalamus appeared to contain the largest concentration of growth hormone-like material,  $164.01 \pm 9.5$  pg/mg tissue, although fairly substantial amounts were also shown to be present in the amygdala,  $56.20 \pm 7.6$  pg/mg tissue and many other brain regions. As with the rat, GH-like material in the monkey cerebellum showed the lowest amounts of GH/mg of tissue, only 19.66 pg/mg of brain tissue.

b. Thyroid Stimulating Hormone-Like Immunoreactivity in the Primate Brain

Measured as pg/mg wet tissue in Table 10, the hypothalamus and cortex were shown to contain the highest amounts of thyroid stimulating hormone-like activity,  $133.92 \pm 9.20$ , and  $54.10 \pm 2.36$  pg/mg wet weight tissue, respectively. Several other areas of the primate brain showed little or no immunoreactive TSH activity such as the amygdala, cerebellum, thalamus, and putamen.

TABLE 9

## DISTRIBUTION OF IMMUNOREACTIVE GH IN ADULT MONKEY BRAIN

(pg/mg wet wt. tissue)\*\*

n = 8	GH (pg/mg wet wt. tissue)**
Hypothalamus	164.01 ± 9.5
Amygdala	56.20 ± 7.6
Pallidus	39.20 ± 2.5
Thalamus	36.80 ± 2.1
Cortex	35.50 ± 1.8
Hippocampus	30.88 ± 3.3
Caudate	25.80 ± 1.2
Putamen	23.80 ± 3.6
Cerebellum	19.66 ± 2.4
Pituitary	$1.78 \times 10^4 \pm 0.51 \times 10^4$

n = number of experiments.

\*\* Mean ± S. E. M.

TABLE 10

## DISTRIBUTION OF IMMUNOREACTIVE TSH IN ADULT MONKEY BRAIN

(pg/mg wet weight of tissue)\*\*

n = 3	TSH (pg/mg wet wt. tissue)**
Hypothalamus	133.92 ± 9.20
Cortex	54.10 ± 2.36
Mid Brain	13.28 ± 4.63
Hippocampus	5.92 ± 2.79
Amygdala	NM
Caudate	NM
Cerebellum	NM
Pallidus	NM
Putamen	NM
Thalamus	NM
Pituitary stalk	$1.5 \times 10^4 \pm 0.23 \times 10^4$

n = number of experiments.

NM = not measurable.

\*\* Mean ± S. E. M.



C. GROWTH HORMONE AND THYROID STIMULATING HORMONE-LIKE IMMUNOREACTIVITY  
IN EXTRAPITUITARY REGIONS OTHER THAN THE RAT CNS

a. Growth Hormone-Like Immunoreactivity in the Pineal Gland and GI  
Tract

In Tables 11 and 12, the levels of immunoreactive GH, expressed as ng/mg protein and as pg/mg wet weight tissue, respectively, for extrapituitary regions other than the CNS, show that substantial amounts of material exist in such areas. Certain areas of the gastrointestinal tract (GI) such as the small intestine,  $1.23 \pm 0.3$  ng/mg protein, 42.6 pg/mg tissue, and the pancreas,  $1.12 \pm 0.05$  ng/mg protein,  $30.2 \pm 7.6$  pg/mg tissue, contain fairly high levels of a growth hormone-like material.

In an effort to determine whether these results were artifactual, due to enzymatic destruction of the labeled growth hormone in the RIA, a study of the recovery of added amounts of purified pituitary growth hormone to tissue extracts was carried out. Near to 100 % recovery of added hormone revealed no breakdown of the radioactive tracer, strongly indicating that these levels of immunoreactive growth hormone in the GI tract are probably not due to an artifact in the system. One month after hypophysectomy, levels of GH found in these areas did not alter or change.

The pineal gland was also shown to contain substantial amounts of immunoreactive growth hormone,  $77.45 \pm 8.3$  ng/mg protein and  $41.63 \pm 2.7$  pg/mg tissue. Of interest are preliminary studies which appear to show that these levels are subject to diurnal variation and that they drop significantly after removal of the pituitary.

TABLE 11

## DISTRIBUTION OF IMMUNOREACTIVE GH

## IN ADULT RAT TISSUES

(tissues perfused for 10 min. with phosphosaline buffer, pH 7.6)

(ng/mg protein)\*\*

	<u>n</u>	<u>GH (ng/mg protein)**</u>
Muscle	3	0.02 ± .005
Kidney	3	0.40 ± .008
Liver	3	0.93 ± .02
Pancreas	16	1.12 ± .05
Small intestine	16	1.23 ± .03
Large intestine	16	0.05 ± .02
<hr/>		
Pineal	30	77.45 ± 8.3
<hr/>		

n = number of experiments.

\*\* Mean ± S. E. M.

TABLE 12

## DISTRIBUTION OF IMMUNOREACTIVE GH

## IN ADULT RAT TISSUES

(tissues perfused with phosphosaline buffer, pH 7.6, for 10 mins.)

(pg/mg wet weight tissue)\*\*

	<u>n</u>	<u>GH (pg/mg wet wt.)**</u>
Amygdaloid nucleus	13	284 ± 10.0
Muscle	3	3.0 ± 0.5
Kidney	3	4.5 ± 2.1
Liver	3	21.0 ± 5.2
Pancreas	16	30.2 ± 7.6
Small intestine	16	42.6 ± 3.5
Large intestine	16	3.2 ± 0.35
<hr/>		
Pineal	30	41.63 ± 2.7
<hr/>		

n = number of experiments.

\*\* Mean ± S. E. M.

b. Thyroid Stimulating Hormone Immunoreactivity

Tables 13 and 14 demonstrate the amounts of immunoreactive TSH found in extrapituitary, non-brain tissues of the rat. As with immunoreactive GH, substantial amounts of TSH-like material were detected in GI tract tissues, especially in the pancreas,  $77.0 \pm 5.21$  ng/mg protein,  $12.42 \pm$  ng/mg tissue, and in the small intestine,  $23.8 \pm 7.62$  ng/mg protein and  $12.42 \pm 2.54$  ng/mg tissue. Similar to the results obtained for GH, the enzymes of the GI tract did not appear to interfere with the RIA for TSH, as good recoveries of added pure TSH to intestinal extracts were obtained in recovery studies. Hypophysectomy did not cause any appreciable fall in levels of immunoreactive TSH in these tissues 30 days after pituitary removal.

The pineal gland also demonstrated high levels of immunoreactive TSH, 388 ng/mg protein,  $1.49 \pm 0.43$  ng/mg tissue, which appears to be subject to diurnal variation being significantly higher in the afternoon than in the morning.

D. GROWTH HORMONE AND THYROID STIMULATING HORMONE IMMUNOREACTIVITY IN MEDIUM FROM CULTURES OF MONOLAYERED NERVOUS TISSUE

a. Growth Hormone Immunoreactivity Detected in Tissue Culture Medium

When the media from actively growing tissue cultures of amygdaloid tissue was examined for immunoreactive growth hormone-like material, the results, expressed as cumulative hormone present over a 30-day growth period, were determined and are shown in Fig. 23 .

TABLE 13

## DISTRIBUTION OF IMMUNOREACTIVE TSH

## IN ADULT RAT TISSUES

(tissues perfused for 10 min. with phosphosaline buffer, pH 7.6)

(ng/mg protein)\*\*

	<u>n</u>	<u>ng TSH/mg protein**</u>
Muscle	3	5.2 ± 0.30
Kidney	3	4.3 ± 0.25
Liver	3	4.8 ± 0.32
Pancreas	16	77.0 ± 5.21
Small intestine	16	23.8 ± 7.62
<hr/>		
Pineal	30	388 ± 25.1
<hr/>		

n = number of experiments.

\*\* Mean ± S. E. M.

TABLE 14

## DISTRIBUTION OF IMMUNOREACTIVE TSH

## IN ADULT RAT TISSUES

(ng/mg wet weight tissue)\*\*

	<u>n</u>	<u>ng TSH/mg wet wt.**</u>
Hypothalamus	50	5.433 ± 0.560
Muscle	3	0.010 ± 0.002
Kidney	3	0.040 ± 0.01
Liver	3	0.063 ± 0.03
Pancreas	16	12.42 ± 2.54
Small intestine	16	10.78 ± 3.26
Large intestine	16	0.18 ± 0.02
Stomach	16	0.18 ± 0.02
<hr/>		
Pineal	12	1.49 ± 0.43
<hr/>		

n = number of experiments.

\*\* Mean ± S. E. M.




Figure 23. Cumulative immunoreactive growth hormone released from cells of the adult rat amygdaloid nucleus grown in tissue culture. In these five different experiments, medium was changed at 5, 10, 20, 25 and 30 days. Each symbol represents the average growth hormone released from approximately  $2 \times 10^6$  viable cells plated in each of 5 Falcon culture flasks containing 3ml. media.

From a total of five experiments, in which the medium was changed every five days, approximately  $2 \times 10^6$  viable amygdaloid cells released 47-65 ng of a growth hormone-like material into the tissue culture medium over the 30-day period of the study. Each symbol in Fig. 23 represents one experiment, and individual points, the mean levels of immunoreactive growth hormone present in the medium of five culture flasks obtained when a complete exchange of growth medium was carried out approximately every five days.

When brain tissue was obtained from 21-day post-hypophysectomized rats and plated out in monolayer cultures, the cumulative amount of immunoreactive growth hormone found in the media, over a 30-day period, was determined by RIA and is shown graphically in Fig. 24. Four different brain areas are represented and each symbol shows the mean and standard error of the mean level of immunoreactive growth hormone released from a total of 15 plates from three different experiments. Medium was changed every five days and, in these studies using brain tissue from hypophysectomized rats, it was shown that substantial amounts of immunoreactive growth hormone were released from all brain tissues into the medium over 30 days of study. Little difference was observed between the amounts obtained from hypophysectomized amygdaloid tissue (Fig. 24) and amygdaloid tissue from normal intact animals (Fig. 23). The cumulative amounts of immunoreactive growth hormone released into the tissue culture medium from hypophysectomized brain tissue ranged from a mean of 20 ng in the cortex to 55 ng in amygdaloid cell cultures over three different experiments. Viability checks throughout all tissue culture experiments showed






Figure 24. Cumulative immunoreactive growth hormone released from cells of the 21 day post-hypophysectomised, adult, rat brain regions grown in tissue culture. Each symbol represents the mean and S.E.M. of hormone released in 3 experiments from approximately  $2 \times 10^6$  cells plated in 5 culture flasks containing 3 ml. media. In these experiments, medium was changed at 5, 10, 15, 20, 25 and 30 days.

between 95%-98% viable cells present at each sampling date.

b. Thyroid Stimulating Hormone Immunoreactivity Detected in Tissue Culture Medium

Studies of brain tissue monolayers, in which the measurement of TSH-like activity present in the growth medium was determined after a certain time period, are plotted in Fig. 25. Here cumulative TC medium-immunoreactive TSH is plotted against days grown in tissue culture of dispersed cells obtained from specific brain parts from normal adult rat brain and from anterior pituitaries. All tissues tested were shown to release a TSH-like material into the medium over the 30-day period of study, ranging from a total of 2,200 ng for the pituitary to 350 ng for thalamus cells. Each symbol in Fig. 25 represents tissue from a different brain area or anterior pituitary and the individual points, the mean and standard error of the mean, cumulative value for immunoreactive TSH released from such monolayered cells. When cells were taken from 21-day post-hypophysectomized adult rats and plated in tissue culture medium, the amount of cumulative immunoreactive TSH released over a 30-day period was also determined. The results are shown in Fig. 26 and ranged from a total of 750 ng of TSH from hippocampal tissue to 550 ng of TSH from amygdaloid tissue after 30 days in culture. This demonstrates that even after hypophysectomy, brain tissue possessed the ability to release immunoreactive TSH into the medium, as long as the viability of the cultured cells was maintained. Viability throughout these experiments was maintained between 93%-98% of the original cell number.




Figure 25. Cumulative immunoreactive thyroid stimulating hormone released from cells of the adult rat brain regions and pituitary grown in tissue culture. Each symbol represents the mean and S.E.M. of hormone released in 5 experiments, from approximately  $2.5 \times 10^6$  cells plated in 5 culture flasks containing 3 ml media. In these experiments, medium was changed at 5, 10, 20 and 28 days.




Figure 26. Cumulative, immunoreactive thyroid stimulating hormone released from cells of the 21 day post-hypophysectomised, adult, rat, brain regions in tissue culture. Each symbol represents the mean and S.E.M. of hormone released in 3 experiments from approximately  $2 \times 10^7$  cells plated in 5 culture flasks containing 3 ml media. In these experiments, medium was changed at 5, 10, 15, 22 and 30 days.

E. THE EFFECT OF KNOWN HORMONAL REGULATORS ON THE RELEASE OF IMMUNOREACTIVE GH AND TSH FROM TISSUE CULTURED CELLS OF THE BRAIN AND ANTERIOR PITUITARY

a. Somatostatin Added to Tissue Cultured Brain and Pituitary Cells

Somatostatin, a known inhibitor of growth hormone release from intact pituitaries (Brazeau, 1973) was shown to be effective at a concentration of 1  $\mu\text{g/ml}$  medium in significantly suppressing the release of immunoreactive GH from both pituitary and various brain tissues grown in culture. In Fig. 27, the levels of growth hormone present in the medium of control and somatostatin-spiked tissue cultures are shown after a 24-hour period of incubation. The results delineated in Fig. 27 represent the mean, and standard error of the mean, of immunoreactive growth hormone levels from three different experiments each containing five separate tissue culture flasks, each containing approximately  $6 \times 10^6$  viable cells.

b. Thyroxine ( $T_4$ ) Added to Tissue Cultured Brain and Pituitary Cells

Levels of thyroid stimulating hormone-like activity in actively growing cultures of brain tissue ( $10^7$  cells), incubated for 24 hours with and without  $10^{-9}$  M L-thyroxine are shown in Fig. 28. Although this concentration of thyroxine ( $T_4$ ) was able to significantly suppress the release of TSH from cultured pituitary cells from a mean of 1,700 ng/24 hours to a mean of 1,400 ng/24 hours, it was not, however, able to prevent the release of immunoreactive TSH from any part of the brain grown in monolayers under similar conditions. Results in all cases were expressed as the mean,




Figure 27. Immunoreactive growth hormone released from  $6 \times 10^6$  cells of the adult rat brain and pituitary grown in tissue culture for 24 hrs with or without somatostatin (1  $\mu\text{g}/\text{ml}$ ). Results represent the mean and S.E.M. from 3 experiments.




Figure 28. Immunoreactive thyroid stimulating hormone released from  $10^7$  cells of the adult rat brain and pituitary grown in tissue culture for 24 hrs with or without L-thyroxine, ( $T_4$   $10^{-9}$  M).

and standard error of the mean, levels of immunoreactive TSH detectable after 24 hours from at least five plates in three different experiments as shown in Fig. 28 .

F. IMMUNOREACTIVE THYROID STIMULATING HORMONE LEVELS DETECTED IN A CULTURED NEUROBLASTOMA CELL LINE N<sub>4</sub>TGI

Over a ten-day growth period, in which media was changed at days 1, 4, and 9, the levels of immunoreactive TSH released into the growth medium from actively growing cultures of the neuroblastoma cell line N<sub>4</sub>TGI were determined by RIA and are shown in Fig. 29 . The mean and standard error of the mean values for cumulative hormonal levels from 5 - 8 individual flasks in three different experimental runs are represented by each point. A total mean of 760 ng of immunoreactive TSH was found in the medium after 10 days growth of this cell line. Throughout the experiment, viability was checked by the trypan blue exclusion method and found to remain between 95% - 98% of the original number of viable cells plated at day one.

It was apparent that cells of this neuroblastoma cell line are capable of releasing a TSH-like immunoreactive material indefinitely, if one extrapolates, into the growth medium.

G. FRACTIONATION OF BRAIN HOMOGENATES BY ULTRACENTRIFUGATION

Assay of three pellets and three supernatants obtained by differential ultracentrifugation of brain extracts (as described in Materials and






Figure 29. Cumulative immunoreactive thyroid stimulating hormone TSH released from  $5 \times 10^6$  cells of the neuroblastoma cell line  $N_4TG1$ . Each point represents the mean and S.E.M. of hormone released from cells in 3 experiments. The media was changed at 1, 4 and 9 days.

Methods) for their content of either immunoreactive growth hormone or thyroid stimulating hormone, gave the data shown in Tables 15 and 16, respectively.

a. Growth Hormone: Distribution in Ultracentrifuged Homogenate Fractions

In Table 15, the quantity of GH is expressed either as ng/fraction, taking into account the volume of the fraction or as ng/mg protein, accounting for the amount of protein present. For amygdala and thalamic homogenates, the greatest concentration of immunoassayable growth hormone was clearly detectable in the second pellet  $P_2$ , i.e.,  $P_1 = 6.119$ ,  $P_2 = 17.879$ ,  $P_3 = 0.906$ ,  $S_1 = 18.346$ ,  $S_2 = 8.242$ , and  $S_3 = 5.477$  ng GH/fraction. This second pellet is known to represent the synaptosomal-mitochondrial fraction of the brain homogenate after ultracentrifugation. For the hypothalamic homogenate, however, there appeared to be an equal distribution of immunoreactive GH between the  $P_2$  pellet and the final supernatant. This would suggest, that in the hypothalamus there is a fraction of immunoreactive growth hormone that is not associated with a specific structural entity, such as the synaptosomes or mitochondria but present instead in some free state, whereas a second fraction which sediments out is presumably associated with a specific entity present in the  $P_2$  pellet preparation.

TABLE 15

## FRACTIONATION OF BRAIN HOMOGENATES BY ULTRACENTRIFUGATION:

## GROWTH HORMONE

(Content of fractions ng/fraction and ng/mg protein)

ng GH/fraction				
AMYGDALA	P <sub>1</sub>	6.119	S <sub>1</sub>	18.346
	P <sub>2</sub>	17.879	S <sub>2</sub>	8.242
	P <sub>3</sub>	0.906	S <sub>3</sub>	5.477
THALAMUS	P <sub>1</sub>	13.100	S <sub>1</sub>	30.296
	P <sub>2</sub>	21.602	S <sub>2</sub>	6.663
	P <sub>3</sub>	1.626	S <sub>3</sub>	7.166
HYPOTHALAMUS	P <sub>1</sub>	10.168	S <sub>1</sub>	70.762
	P <sub>2</sub>	36.661	S <sub>2</sub>	41.237
	P <sub>3</sub>	2.775	S <sub>3</sub>	36.718

ng/mg protein/fraction				
AMYGDALA	P <sub>1</sub>	2.769	S <sub>1</sub>	4.504
	P <sub>2</sub>	4.659	S <sub>2</sub>	2.601
	P <sub>3</sub>	1.839	S <sub>3</sub>	3.649
THALAMUS	P <sub>1</sub>	1.160	S <sub>1</sub>	4.483
	P <sub>2</sub>	3.673	S <sub>2</sub>	2.134
	P <sub>3</sub>	0.797	S <sub>3</sub>	2.020
HYPOTHALAMUS	P <sub>1</sub>	0.664	S <sub>1</sub>	24.824
	P <sub>2</sub>	6.886	S <sub>2</sub>	11.034
	P <sub>3</sub>	4.462	S <sub>3</sub>	5.669

P = pellet; S = supernatant

b. Thyroid Stimulating Hormone: Distribution in Ultracentrifuged Homogenate Fractions

Similar ultracentrifugal fractionation studies of brain homogenate for the detection of TSH-like material, by radioimmunoassay, gave the data shown in Table 16.

Here again, as shown for the amygdaloid homogenate, the greatest quantity of immunoreactive TSH was associated with the P<sub>2</sub> pellet whereas the hypothalamic fractionation pattern seemed to show two, rather than one, distinct pool of immunoreactive material--one associated with the P<sub>2</sub> pellet and the other with the final supernatant preparation.

Because of this heterogenous distribution of both immunoreactive GH and TSH in the hypothalamic homogenate, further fractionation of the P<sub>2</sub> pellet by sucrose density gradients might allow better discrimination of whether the two hormones are actually associated with a particular cellular entity such as the synaptosomes.

H. IMMUNOPEROXIDASE STAINING OF BRAIN TISSUE FOR PITUITARY-LIKE HORMONE DISTRIBUTION

a. Thyroid Stimulating Hormone

To further pinpoint the location of thyroid stimulating-like hormone in rat brain tissue, immunohistochemical staining of paraformaldehyde-fixed tissue was attempted. The conditions noted in the "Materials and Methods" section revealed a dense uptake of peroxidase-stained

TABLE 16

## FRACTIONATION OF BRAIN HOMOGENATES BY ULTRACENTRIFUGATION:

## THYROID STIMULATING HORMONE

(Content of fractions ng/fraction and ng/mg protein)

ng TSH/fraction				
AMYGDALA	P <sub>1</sub>	79.360	S <sub>1</sub>	251.071
	P <sub>2</sub>	225.632	S <sub>2</sub>	52.768
	P <sub>3</sub>	49.354	S <sub>3</sub>	6.811
HYPOTHALAMUS	P <sub>1</sub>	97.385	S <sub>1</sub>	347.631
	P <sub>2</sub>	126.783	S <sub>2</sub>	210.295
	P <sub>3</sub>	108.127	S <sub>3</sub>	116.534
ng TSH/mg protein				
AMYGDALA	P <sub>1</sub>	7.638	S <sub>1</sub>	13.364
	P <sub>2</sub>	9.999	S <sub>2</sub>	4.713
	P <sub>3</sub>	3.172	S <sub>3</sub>	2.008
HYPOTHALAMUS	P <sub>1</sub>	5.705	S <sub>1</sub>	96.730
	P <sub>2</sub>	48.630	S <sub>2</sub>	52.119
	P <sub>3</sub>	10.714	S <sub>3</sub>	42.588

P = pellet; S - supernatant

immunocomplex in cells of the pituitary, indicating the presence of TSH. Shown under low power (X 100) in Fig. 30 , and at higher magnification (X 400) in Fig. 31 , are representative sections of pituitary with the thyrotrophs staining a dark brown against a nonhormonal containing, unstained, cell background. All controls revealed a similar appearance to Fig. 32 , that is, no uptake of PAP complex by any cell due to the conditions set by the experiment. For the pituitary, therefore, it was demonstrated that the incubation times and concentrations of antibodies and other substituents in this immunohistochemical staining experiment to detect TSH activity were sufficient and satisfactory.

When amygdaloid tissue was treated in a similar manner and examined under a high magnification (X 1000), Figs. 33 and 34 , or at a lower magnification (X 400), Fig. 35 , certain cell bodies were seen to stain dark brown, indicating the presence of TSH-like immunoreactive material. Controls again provide evidence that this finding was not an artifactual observation. Pre-treatment with colchicine would perhaps have shown a more detailed picture of the involvement of the type of cell, neurone, or glia which was being stained. Cell bodies staining for TSH were also detected in the hypothalamus and hippocampal regions of the brain, with a few isolated bodies being also distinguished in the cerebral cortex and thalamic areas. In the amygdaloid area, all of the individual nuclei appeared to contain TSH-like hormone. Those which were more densely stained included the nucleus amygdaloides corticalis, amygdaloides basalis, pars mediales, and pars lateralis. In the hypothalamus, areas in which stained cell bodies were prominent were the dorsomedial and ventromedial nuclei.




Figure 30. Immunohistochemical staining of pituitary thyrotrophs by the PAP technique. (Blue filter, x 100).




Figure 31. Immunohistochemical staining of pituitary thyrotrophs by the PAP technique (x400).




Figure 32. Control pituitary stained by the PAP technique using absorbed TSH antiserum ( x 400).



Cell bodies containing  
TSH

Control immunoabsorbed  
1st. Ab.






Figure 33. Immunohistochemical staining by the PAP technique of amygdaloid tissue for TSH ( x 1000)




A




B

Figure 34. Immunohistochemical staining of amygdaloid tissue for TSH by the PAP technique. Controls (x 1000). A. Immunoabsorbed first antibody. B. First antibody replaced by normal rabbit serum.



A



B

Figure 35. A. Immunohistochemical staining for TSH in amygdaloid tissue (x400).  
B. Control. First antibody replaced by normal rabbit serum.

Similar studies with GH, not detailed in this thesis, have also detected GH-like material in cells of the amygdaloid nucleus by an indirect immunoperoxidase technique. Thus, confirming the association of hormone-like material with a particulate fraction of nervous tissue, these histochemical studies provide further evidence of a brain source for pituitary-like hormones.

## I. CHARACTERIZATION OF "BRAIN HORMONES" BY GEL CHROMATOGRAPHY

### a. IRGH-like Material in the Rat Brain

The elution pattern obtained when a sample of amygdaloid extract, containing immunoreactive GH-like material was placed on a Sephadex G-100 column together with a trace amount of  $^{125}\text{I}$  labeled rat pituitary GH, is shown in Fig. 36. The ordinates are given as the concentration of immunoreactive GH and cpm of radiolabeled pituitary growth hormone. On the abscissa, the tube number of the fractions collected is noted.

The peak of GH immunoreactivity coeluted with the trace amount of radiolabeled pituitary growth hormone, indicating molecular identity between the hormonal material found in the brain homogenate with that isolated from the pituitary. A molecular weight of approximately 46,000 D can therefore be assigned to the brain GH-like material.

### b. IRTSH Material in the Rat Brain

Fig. 37 illustrates a similar experiment in which the amygdaloid extract was fractionated on a Sephadex G-100 column, together with  $^{125}\text{I}$





Figure 36. Elution pattern of lyophilised and reconstituted amygdala extract IRGH and  $^{125}\text{I}$  rat pituitary GH on a Sephadex G-100 column eluted with veronal buffer. pH 8.6.



**Figure 37.** Elution pattern of lyophilized and reconstituted amygdala extract IRTSH and  $^{125}\text{I}$  rat pituitary TSH on a Sephadex G-100 column eluted with 0.01 M  $\text{PO}_4$  buffer pH 7.6.

rat pituitary TSH. Again, coelution of immunoreactive and labeled pituitary TSH material was observed, indicating molecular similarity between the hormones isolated from these two distinct sources. A probable molecular weight of 28,000 D can therefore be assigned to the brain TSH-like hormone. Similar results for both GH and TSH hormones were demonstrated for all brain regions examined. The existence of a prohormone or breakdown product of pituitary hormones in the brain thus seems unlikely.

c. IR Pituitary-Like Hormones from a Cloned Mouse Neuroblastoma Line

When a concentrated sample of neuroblastoma N4TGI cell extract was similarly fractionated with a trace amount of radiolabeled pituitary TSH, coelution of immunoreactive and tracer hormone was shown, as in Fig. 38 .

Similar results were obtained when non-cell line tissue extracts from amygdaloid tissue cell cultures and pituitary cell cultures were fractionated and compared. This would indicate that the immunoreactive hormonal activity detected in the in vitro experiments were also molecularly similar, in terms of molecular weight, to their pituitary counterparts in vivo.

J. IMMUNOLOGICAL CHARACTERIZATION OF BRAIN PITUITARY-LIKE HORMONES AND FURTHER IMMUNOASSAY VALIDATION

Logit log plots demonstrating parallel displacement for serial dilutions of pituitary and CNS extracts are shown in Figs. 39 and 40 for GH and TSH, respectively. Although values actually fell along the same line, the hippocampal data is displaced for visual emphasis of the

IMMUNOREACTIVE TSH ●—●  
ng/ml CELL EXTRACT

\*--\*  $^{125}\text{I}$  PITUITARY TSH  
C. P.M.




Figure 38. Elution pattern of lyophilised, reconstituted  $\text{N}_4\text{TG1}$  cell extract IRTSH and  $^{125}\text{I}$  rat pituitary TSH on a Sephadex G-100 column eluted with 0.01 M  $\text{PO}_4$  buffer pH 7.6






Figure 39. Logit log plot demonstrates parallel displacement for serial dilutions of pituitary and CNS growth hormone extracts.




Figure 40. Logit log plot demonstrates parallel displacement for serial dilutions of pituitary and CNS thyroid stimulating hormone extracts.

data's linearity. Such linearity between pituitary and brain hormones demonstrated probable immunological identity.

K. BIOLOGICAL ACTIVITY OF BRAIN PITUITARY-LIKE HORMONES AS COMPARED TO PITUITARY HORMONES

a. Growth Hormone Bioassay

When adult, 14-day post-hypophysectomized male and female rats received intraperitoneally injected doses of pituitary, amygdaloid extract or saline over a 5-day period, their body weights and width of the tibia epiphyseal cartilage were examined as indicators of biological activities usually attributable to growth hormone.

Table 17 shows the changes in body weights recorded over the 5-day injection period before sacrifice from two individual experiments. It is apparent that, whereas the extract from rat pituitary was able to produce an increase in body weight response, neither the amygdaloid extract nor saline-treated animals showed a rise in total body weight over the same time period. It was, however, apparent that the weight loss was substantially less in amygdaloid-treated animals when compared to that experienced by saline-treated, hypophysectomized controls. These data were not altogether surprising because, as noted before, a 100  $\mu$ g growth hormone dose is usually required before a weight gain is observable. In neither experiment was sufficient immunoassayable GH given in the brain extract to the test animals.

TABLE 17

BODY WEIGHT CHANGES OVER 5-DAY  
 BIOLOGICAL EXPERIMENT PERIOD (gms)

		Total Quantity of IR GH ( $\mu$ g) Injected Over 5 Days		
Treatment		Saline (0)	RAM (73)	RAP (113)
Exp 1	♂	- 5.26 $\pm$ 0.10	- 2.3 $\pm$ 0.21	+ 3.02 $\pm$ 0.25
Treatment		Saline (0)	RAM (90)	RAP (95)
Exp 2	♀	- 6.17 $\pm$ 1.35	- 2.3 $\pm$ 0.95	+ 3.17 $\pm$ 0.79

The mean and SEM of the difference in body weight between Day 1 of the biological experiment and Day 5. Each point represents 5 adult animals, 14 days after hypophysectomy.

RAP = rat anterior pituitary extract

RAM = rat amygdaloid nucleus extract

However, a much lower amount of growth hormone is required to elicit a response in the tibia epiphyseal cartilage bioassay, and this was reflected in the results obtained in Table 18a. In both male and female groups of hypophysectomized animals from all four experiments, a statistically significant increase in the epiphyseal cartilage width of amygdaloid extract-treated animals over those of saline-treated controls was observable. This ability to induce cartilage growth was equivalent to the effect produced by the pituitary extract on hypophysectomized animals under similar conditions. Thus, this experiment revealed that the GH-like material present in the brain extracts was capable of eliciting a response normally attributable to, and characteristic of, pituitary growth hormone.

b. Gonadotrophic Bioactivity in Brain Extracts

From the data examining the overall weights of testes and ovaries in all three groups of treated hypophysectomized animals, Tables 18 b and c, it was apparent that the brain extracts also contained gonadotrophic bioactivity. This observation has subsequently been substantiated by our laboratory (Emanuele et al, 1980) where significant amounts of immunoreactive luteinizing hormone have been found in certain areas of the rat brain.

c. Thyroid Stimulating Hormone Bioactivity in Brain Extracts

The thyroids of hypophysectomized animals treated with either saline, a high or low dose of rat pituitary extract, or amygdaloid tissue extract were examined biologically for structural changes.

TABLE 18

## BIOASSAY OF PITUITARY AND AMYGDALA EXTRACTS

a.	<u>DOSE/RAT</u> <u>µg RGH</u>	<u>SEX</u>	<u>TIBIA EPIPH.</u> <u>Width m.u.</u>
	Saline-0	M	8.3 ± 0.5
	RAP-113	M	10.7 ± 0.4**
	RAM-73	M	10.3 ± 0.2**
	Saline-0	F	11.1 ± 2.0
	RAP-95	F	13.6 ± 3.2
	RAM-90	F	15.1 ± 3.2**

b.	<u>DOSE/RAT</u> <u>µg RGH</u>	<u>SEX</u>	<u>TESTIS</u> <u>mg</u>
	Saline-0	M	154 ± 0.7
	RAP-113	M	193 ± 4.0**
	RAM-73	M	204 ± 6.2**

c.	<u>DOSE/RAT</u> <u>µg RGH</u>	<u>SEX</u>	<u>OVARIES</u> <u>mg</u>
	Saline-0	F	15.2 ± 3.2
	RAP-95	F	19.1 ± 4.9*
	RAM-90	F	29.7 ± 4.5**

m.u. = 0.005 mm

\*\* p = &lt; 0.01

\* p = &lt; 0.05

n = 4-5

RAP = rat anterior pituitary extract

RAM = rat amygdaloid nucleus extract

Mean and standard deviations derived from bioassay of anterior pituitary and amygdaloid nucleus extracts in adult, 14-day, post-hypophysectomized male and female rats. Four to five animals were included in each group, and experimental animals received comparable amounts of immunoreactive growth hormone in four daily divided doses administered intraperitoneally.

Details of the results observed are shown in the composite photograph, Fig. 41 .


Sections of thyroid from saline-treated animals revealed the typical follicular characteristics of hypothyroidism, that is, low cuboidal epithelial cells with flattened nuclei and a shrunken colloidal mass in the follicular lumen. One of the characteristics of TSH activity, as already discussed, is to maintain thyroid function. In both groups receiving pituitary extracts, thyroids revealed this response as demonstrated by the appearance of tall, cuboidal epithelial follicle cells; large, round nuclei filling the cells, and well-maintained follicular collagen and vacuolization of colloid.

The thyroid tissue dissected from amygdaloid extract-treated, hypophysectomized animals also showed these characteristics, indicating, on the basis of biological activity, the presence of a TSH-like material.

Evaluation of this TSH-like effectiveness was also attempted using the parameter of  $^{131}\text{I}$  uptake into the thyroid as an activity indicator. Table 19 shows the data obtained in terms of  $^{131}\text{I}$  uptake into various tissues as well as the  $\text{T}_4$  levels of circulating serum after five different treatments. It was apparent that only the high dose of pituitary extract, 100  $\mu\text{g}$ /hypophysectomized animal, was sufficient to produce an effect comparable to that demonstrated in the intact saline-treated control. In this short, four-day study, neither 2  $\mu\text{g}$  of pituitary TSH nor 4.3  $\mu\text{g}$  of amygdaloid extract TSH, were sufficient to increase the labeled iodide uptake into the thyroid or to raise circulating  $\text{T}_4$  levels, an indirect

SALINE (0)\*

LOW DOSE RAP (5)



HIGH DOSE RAP (100)

RAM (4.5)

Figure 41. Histology of the thyroid gland of 14 day hypophysectomised animals treated with brain and pituitary extracts (x 400).

\* ( ) =  $\mu\text{g}$  immunoreactive TSH administered over a 4 day period to adult rats, 14 days after hypophysectomy. Note the low cuboidal epithelium lining of the thyroid follicles of saline treated animals, and the increased height of epithelium of pituitary and amygdala treated animals.



TABLE 19

EVALUATION OF TSH EFFECTIVENESS TO CHANGE THE  
PATTERN OF  $^{131}\text{I}$  UPTAKE INTO THE THYROID

	Total Dose TSH $\mu\text{g}$	n	(cpm $^{131}\text{I}/\text{gm}$ ) Tissue		Serum
			Thyroid	Muscle	$\text{T}_4$ $\mu\text{gm}/\text{dl}$
Hypophysectomized +	Saline	5	4,447 $\pm$ 206	273 $\pm$ 20	0.8 $\pm$ 0.1
Hypophysectomized +	100 $\mu\text{g}$ RAP TSH	5	57,453 $\pm$ 129	237 $\pm$ 30	2.2 $\pm$ 0.3
Hypophysectomized +	2 $\mu\text{g}$ RAP TSH	5	3,000 $\pm$ 198	254 $\pm$ 16	0.9 $\pm$ 0.1
Hypophysectomized +	4.3 $\mu\text{g}$ RAM TSH	5	2,795 $\pm$ 216	212 $\pm$ 22	0.9 $\pm$ 0.2
Intact +	Saline	5	61,145 $\pm$ 598	189 $\pm$ 27	4.7 $\pm$ 0.3

Four groups of adult, 14-day, post-hypophysectomized rats received either saline, 100  $\mu\text{g}$  pituitary extract TSH, 2  $\mu\text{g}$  pituitary extract TSH, or 4.3  $\mu\text{g}$  amygdaloid nucleus extract TSH over a four-day period, intraperitoneally. Eighteen hours before sacrifice, each animal received 1.5  $\mu\text{c}$ uries of radioactive iodine IP.

indicator of a well-maintained thyroid. On reviewing the literature, this result was not surprising, as it was found that at least 100  $\mu\text{g}$  of pituitary TSH is necessary to show an increased  $^{131}\text{I}$  uptake into the thyroid of hypophysectomized rats (Cons et al, 1975).

#### L. TESTING THE INTACTNESS OF THE BLOOD-BRAIN BARRIER IN BOTH ADULT AND YOUNG ANIMALS

When  $^{125}\text{I}$  labeled pituitary hormones were injected IV into the tail veins of adult rats and 30 minutes allowed to pass before sacrifice, levels of labeled hormone were measured in various body tissues and brain in order to observe if radiolabeled growth and thyroid stimulating hormone were capable of passing through the blood brain barrier. Results of these experiments are given in Tables 20 and 21 for GH and TSH, respectively, as cpm/mg of wet tissue. The immunoreactive hormone content of saline-injected control rat tissue was also determined, in order to compare the sites of receptor-bound circulating hormone with sites of immunologically active, extractable growth and thyroid stimulating hormones.

As shown in Table 20, the large organs of the body readily bound circulating, labeled pituitary growth hormone. The pituitary itself had a fairly high level of  $^{125}\text{I}$  hormone/mg tissue,  $11.64 \pm 2.5$  cpm/mg, whereas little or no labeled hormone was detected in areas of the CNS,  $0.00 \rightarrow 0.90$  cpm/mg. By comparison, levels of immunoreactive growth hormone, as measured in tissue extracts by RIA, was highest in the CNS tissues such as amygdala,  $212 \pm 5.4$  pg/mg tissue, and lowest in the large organs such as the kidney,  $2.9 \pm 0.2$  pg/mg tissue.

TABLE 20.

TISSUE DISTRIBUTION OF  $^{125}\text{I}$ -RAT GROWTH HORMONE  
 INJECTED IV INTO ADULT RATS AND IR GH CONTENT  
 OF TISSUES (pg/mg wet weight)

	<u>n</u>	<u>cpm/mg*</u>	<u>GH pg/mg wet wt.*</u>
Kidney	5	433.84 ± 104.6	2.9 ± 0.2
Liver	5	55.56 ± 10.9	17.3 ± 2.6
Muscle	5	1.45 ± 1.0	2.2 ± 0.5
Pituitary	5	11.64 ± 2.5	1.9 x 10 <sup>5</sup> ± 0.7
Amygdala	5	0.90 ± 0.1	212 ± 5.4
Thalamus	5	0.89 ± 0.1	89 ± 3.6
Cortex	5	0.00	52 ± 5.2

n = number of experiments.

\* Mean ± S. E. M.

A similar picture was shown for the effectiveness of the blood brain barrier in preventing  $^{125}\text{I}$  labeled TSH from entering the brain via the peripheral circulation, as demonstrated in Table 21, where no radio-labeled hormone was shown to be detected in the CNS structures.

In studies with fetal material, there has been some discussion of possible leakages in the immature brain blood barrier. In order to determine whether or not this was the case, 21-day-old fetuses were given radiolabeled GH or TSH by an intraperitoneal route and the distribution of label was detected in various body tissues and brain 30 minutes after administration. Although results are only given for GH in Table 22b, similar data were obtained for TSH, revealing the total effectiveness of the 21-day-old fetal blood brain barrier to prevent the entry of circulating pituitary hormones into the brain.

Shown in Table 22a are the results of a similar experiment involving another such barrier, the fetal-placental barrier. This was also examined by injecting females in their 21st day of pregnancy with radio-labeled hormone. Dissection of the fetuses after a 30-minute delay revealed no label in any part of the fetus, demonstrating an inability of GH and TSH to cross the placental barrier.

#### M. THE EFFECT OF HYPOPHYSECTOMY ON LEVELS OF BRAIN PITUITARY-LIKE HORMONES

Animals which had been hypophysectomized for varying lengths of time were sacrificed and the levels of immunoreactive hormone present in

TABLE 21.

TISSUE DISTRIBUTION OF  $^{125}\text{I}$  RAT THYROID STIMULATING HORMONE  
 INJECTED IV INTO ADULT RATS AND IR TSH CONTENT  
 OF TISSUE (ng/mg wet weight)

	<u>n</u>	<u>cpm/mg*</u>	<u>IR TSH ng/mg wet wt.*</u>
Kidney	5	678.71 ± 10.5	0.040 ± 0.01
Liver	5	72.65 ± 12.3	0.075 ± 0.02
Muscle	5	18.64 ± 5.6	0.002 ± 0.01
Pituitary	5	17.33 ± 2.7	1.8 × 10 <sup>5</sup> ± 0.42
Amygdala	5	0.00	1.521 ± 0.36
Thalamus	5	0.00	0.795 ± 0.27
Cortex	5	0.00	0.651 ± 0.43

\* Mean ± S. E. M.

TABLE 22

DISTRIBUTION OF  $^{125}\text{I}$  RAT GROWTH HORMONE IN FETAL AND NEONATAL BRAIN AND OTHER TISSUES  
AFTER INJECTION OF RADIOLABELED HORMONE EITHER INTRAPERITONEALLY INTO PREGNANT RATS (A)

INTO 21-DAY-OLD FETUSES (B)

OR INTO NEONATAL RATS (C)

$^{125}\text{I}$  GH, cpm/gm Tissue

		<u>Kidney</u>	<u>Pituitary</u>	<u>Hypothalamus</u>	<u>Amygdala</u>
Rat Fetal Age (days):					
A.	21 (IP mother)	0	0	0	0
B.	21 (IP fetus)	3000	2	0	0
Rat Neonatal Age (days):					
C.	1	6676	318	0	0
	4	8759	378	0	0
	10	3764	183	0	0

the different brain regions compared to intact controls.

a. Growth Hormone: Effects of Hypophysectomy

Table 23 depicts the combined results from three separate experiments, in terms of mean hormone levels, ng/mg protein, per group of 20 animals, for the various brain regions extracted. Animals, hypophysectomized for periods of 7, 14, 28, and 48 days, were compared to intact, similar aged controls. An analysis of the data revealed that in some brain regions, namely the amygdala, hypothalamus, and thalamus, an initial fall was observed in hormonal tissue levels after one week of hypophysectomy, e.g., amygdala,  $6.25 \pm 0.46 \rightarrow 1.25 \pm 0.50$  ng/mg protein. However, after 28 days of hypophysectomy, these regions were found to contain more hormone than was found in comparable intact controls, e.g. amygdala,  $5.75 \pm 0.50 \rightarrow 15.00 \pm 3.10$  ng/mg protein. This is illustrated in Fig. 42 for mean and standard errors of the mean levels of growth hormone, ng/mg protein, found in amygdaloid tissue. Other areas of the brain, such as the caudate, hippocampus, and cortex, showed no fall in growth hormone levels throughout the entire 48 days of hypophysectomy examined. This kind of pattern is illustrated in Fig. 43, showing mean and standard error of the mean levels of hormone, ng/mg protein, assayed in caudate tissue from hypophysectomized and from caudate tissue of intact rats.

b. Thyroid Stimulating Hormone: Effect of Hypophysectomy

Similar experiments were developed to determine changes in levels of immunoassayable TSH in brain tissue. Combined results are given in Table 24 for hormonal levels in hypophysectomized and intact animals,

TABLE 23

IMMUNOREACTIVE GROWTH HORMONE (ng/mg protein)\*  
IN EXTRACTS FROM DISCRETE BRAIN PARTS

Days, Post-Hypophys.	HYPOPHYSECTOMIZED ANIMALS				INTACT CONTROLS			
	7	14	28	48				
	A	B	C	D	A	B	C	D
AMYGDALA	1.25 ± 0.50	3.25 ± 0.50	15.0 ± 3.10	13.25 ± 1.56	6.25 ± 0.46	5.75 ± 0.88	5.75 ± 0.50	6.25 ± 0.75
HYPOTHAL-AMUS	0.57 ± 0.20	1.73 ± 0.20	7.24 ± 0.50	6.53 ± 0.90	1.99 ± 0.30	2.89 ± 0.15	3.13 ± 0.40	2.96 ± 0.25
THALAMUS	0.95 ± 0.10	1.84 ± 0.02	2.07 ± 0.60	2.01 ± 0.50	1.94 ± 0.50	1.58 ± 0.10	1.86 ± 0.10	1.75 ± 0.04
CAUDATE	1.45 ± 0.12	1.81 ± 0.15	1.75 ± 0.09	1.91 ± 0.12	1.50 ± 0.15	1.75 ± 0.24	1.66 ± 0.27	2.00 ± 0.12
HIPPO--CAMPUS	1.36 ± 0.04	1.42 ± 0.05	1.29 ± 0.20	1.28 ± 0.40	1.40 ± 0.20	1.48 ± 0.10	1.39 ± 0.94	1.25 ± 0.01
CORTEX	0.95 ± 0.03	0.72 ± 0.04	0.99 ± 0.10	1.24 ± 0.06	1.32 ± 0.40	1.26 ± 0.05	0.90 ± 0.08	1.18 ± 0.02
CEREBELLUM	0.05 ± 0.02	0.20 ± 0.04	0.03 ± 0.10	0.00	0.05 ± 0.01	0.00	0.04 ± 0.01	0.03 ± 0.01

\*Mean ± S.E.M. of 3 experiments, 20 animals/group of hypophysectomized or intact controls.



## AMYGDALA




Figure 42. Immunoreactive growth hormone present in amygdaloid nucleus extracts of intact and hypophysectomised rats.

## CAUDATE




Figure 43. Immunoreactive growth hormone present in caudate extracts of intact and hypophysectomised rats.  
NS = not significant.

TABLE 24

IMMUNOREACTIVE THYROID STIMULATING HORMONE (ng/mg protein)\*  
IN EXTRACTS FROM DISCRETE BRAIN PARTS

Days Post-Hypophys.	HYPOPHYSECTOMIZED ANIMALS				INTACT CONTROLS			
	7	14	28	48				
	A	B	C	D	A	B	C	D
AMYGDALA	18.95 ± 2.10	27.58 ± 0.30	26.79 ± 2.70	24.64 ± 1.50	21.50 ± 2.10	25.95 ± 0.50	25.79 ± 3.31	26.32 ± 2.90
HYPOTHALAMUS	38.52 ± 0.20	31.43 ± 0.70	23.33 ± 3.40	25.95 ± 0.60	125.60 ± 1.50	143.50 ± 2.70	193.33 ± 1.66	139.60 ± 5.90
THALAMUS	16.55 ± 0.50	19.24 ± 1.50	17.32 ± 2.10	15.95 ± 1.00	17.21 ± 0.40	19.36 ± 0.80	18.30 ± 2.11	16.20 ± 1.10
CAUDATE	14.71 ± 0.80	12.92 ± 0.90	15.63 ± 0.90	14.99 ± 0.80	12.95 ± 0.20	14.83 ± 0.40	14.79 ± 0.60	15.60 ± 0.32
HIPPOCAMPUS	12.69 ± 0.80	9.76 ± 0.20	10.30 ± 1.20	11.51 ± 0.70	12.58 ± 1.10	11.63 ± 0.70	9.62 ± 0.91	10.60 ± 1.30
CORTEX	9.51 ± 0.40	11.18 ± 0.50	10.62 ± 3.20	10.98 ± 1.20	8.79 ± 0.90	10.51 ± 0.50	11.66 ± 1.60	12.68 ± 0.63
CEREBELLUM	13.95 ± 0.30	12.69 ± 2.20	14.95 ± 1.20	15.63 ± 1.10	14.23 ± 1.50	13.65 ± 0.80	16.35 ± 0.90	17.26 ± 0.50

\* Mean ± S.E.M. of 3 experiments, 20 animals/group of hypophysectomized or intact controls.

over different time periods, post-operatively.

All tissue extracts analyzed for TSH, except hypothalamic tissue, were found to show no change in immunoreactive hormone levels over the entire experimental period. However, a significant,  $p = < 0.02$ , fall in hypothalamic levels of TSH was noted from 7 - 28 days post-hypophysectomy. These results are illustrated for 28-day post-hypophysectomized and intact rats in Fig. 44 . The data are expressed as the mean and standard error of the mean IR hormone level per group of 20 animals for three different long-term hypophysectomy experiments.

c. Validation of Hypophysectomy Procedure

The efficiency of hypophysectomy was verified not only by microscopic examination of the sella turcica, but by determination of levels of circulating TSH,  $T_4$ , GH, and %  $T_3$  resin uptake, in intact and hypophysectomized rats. Results of these assays are given in Tables 25 and 26, and demonstrate that hypophysectomy could be considered complete for the animals used in these experiments, as levels of circulating pituitary hormones and thyroid hormones all fell to levels indicating complete loss of the pituitary.

N. TARGET ORGAN REMOVAL: EFFECT ON LEVELS OF BRAIN HORMONES

a. Growth Hormone: Effect of Removal of Target Organs

Removal of a target organ may or may not have an effect on levels of pituitary growth hormone. Lacking any one specific target organ, it




Figure 44. Immunoreactive TSH present in extracts from various brain regions of intact and 28 day hypophysectomised rats. NS = not significant.

TABLE 25

## IMMUNOREACTIVE GH AND TSH LEVELS IN SERUM OF ADULT RATS

(ng/ml)\*

GROWTH HORMONE		THYROID STIMULATING HORMONE	
<u>Intact</u> (n = 200)	<u>30 days Hypophysectomized</u> (n = 125)	<u>Intact</u> (n = 390)	<u>30 days Hypophysectomized</u> (n = 120)
249.86 ± 35.32	20.28 ± 5.80	489.67 ± 52.59	59.79 ± 3.13

TABLE 26.

## CIRCULATING THYROID HORMONE LEVELS IN SERUM OF ADULT RATS

<u>T<sub>4</sub> μg/dl</u>		<u>% RESIN UPTAKE T<sub>3</sub></u>	
<u>Intact</u> (n = 40)	<u>30-day Hypophysectomized</u> (n = 40)	<u>Intact</u> (n = 40)	<u>30-day Hypophysectomized</u> (n = 40)
64.02 ± 0.4	47.600 ± 2.1	4.5 ± 0.3	0.63 ± 0.01

becomes more difficult to assess the possible effect of organ removal on GH levels. However, the main objective in this series of experiments was to reveal possible differences between brain and pituitary hormonal levels after one or four weeks of target organ removal with respect to any effects in sham-operated controls. The four-week data proved the most consistent throughout all three separate experiments, and results for both TSH and GH are given as the mean obtained from three experiments of 20 animals/group/experiment.

(i) Adrenalectomy

As shown in Fig. 45 , the effect of removal of the adrenal gland on levels of immunoreactive GH in serum, pituitary, hypothalamus, and amygdala was demonstrated to have no significant effect when compared to sham-operated controls. Levels of GH in other brain areas showed a similar lack of change, as given in Table 27, in terms of ng/mg protein.

(ii) Oophorectomy

When the ovaries of female adult rats were removed for four weeks, a significant drop ( $P = < 0.01$ ) was observed in circulating GH, whereas pituitary levels of hormone remained unchanged (Table 28 and Fig. 46 ). Contrary to this observation was that shown for the amygdala and hypothalamus, which in all three experiments demonstrated a significant rise ( $P = < 0.01$ ) in GH levels present in brain tissue as compared to sham-operated controls (Fig. 46 ). Other brain areas did not show this rise in GH following the removal of ovaries but remained unchanged four weeks postsurgery (Table 28).



## GH LEVELS 4 WEEKS AFTER ADRENALECTOMY




Figure 45. Effects of adrenalectomy on growth hormone levels in serum, pituitary, and brain. NS = not significant.

TABLE 27

## EFFECT OF ADRENALECTOMY (4 WEEKS) ON BRAIN GH LEVELS

(ng/mg protein)\*\*

	<u>n</u>	<u>Sham</u>	<u>Adrenalectomized</u>	<u>Statistical Significance</u>
AMYGDALA	60	5.1 ± 0.26	5.4 ± 1.23	NS
HYPOTHALAMUS	60	3.2 ± 1.50	4.0 ± 0.50	NS
CAUDATE	60	0.95 ± 0.26	1.10 ± 0.51	NS
THALAMUS	60	1.32 ± 0.37	1.51 ± 0.76	NS
HIPPOCAMPUS	60	1.29 ± 0.08	1.24 ± 0.10	NS
CORTEX	60	1.110 ± 0.24	0.94 ± 0.08	NS
CEREBELLUM	60	0.04 ± 0.01	0.01 ± 0.006	NS
ANTERIOR PITUITARY	60	1.0 × 10 <sup>6</sup> ± 0.51	0.98 × 10 <sup>6</sup> ± 0.32	NS
<u>GH ng/ml **</u>				
SERUM	60	183 ± 33.0	179 ± 27.0	NS

n = total animals

\*\* = mean ± S. E. M. of 3 experiments

GH LEVELS 4 WEEKS AFTER OOPHORECTOMY




Figure 46. Effects of oophorectomy on growth hormone levels in serum, pituitary and brain. NS = not significant.

TABLE 28.

## EFFECT OF OOPHORECTOMY (4 WEEKS) ON BRAIN GH LEVELS

(ng/mg protein)\*\*

	<u>n</u>	<u>Sham</u>	<u>Adrenalectomized</u>	<u>Statistical Significance</u>
AMYGDALA	60	4.32 ± 1.21	9.21 ± 0.64	p = < 0.01
HYPOTHALAMUS	60	2.81 ± 0.63	5.42 ± 1.21	p = < 0.01
CAUDATE	60	2.09 ± 0.31	1.49 ± 0.52	NS
THALAMUS	60	1.31 ± 0.26	0.94 ± 0.20	NS
HIPPOCAMPUS	60	1.86 ± 0.30	1.44 ± 0.15	NS
CORTEX	60	0.81 ± 0.03	0.95 ± 0.11	NS
CEREBELLUM	60	0.11 ± 0.06	0.3 ± 0.01	NS
ANTERIOR PITUITARY	60	8.1 x 10 <sup>5</sup> ± 0.3	8.6 x 10 <sup>5</sup> ± 0.25	NS
GH ng/ml**				
SERUM	60	267 ± 22.6	68.90 ± 10.5	p = < 0.01

n = total animals

\*\* = mean ± S. E. M. of 3 experiments

(iii) Orchidectomy

Unlike oophorectomy, removal of the testes did not produce any significant change in immunoreactive GH levels in serum, pituitary, or in other brain tissues (Fig. 47 , table 29).

(iv) Thyroidectomy

Four weeks after removal of the thyroid, a significant drop ( $P = < 0.01$ ) was observed in both serum and pituitary immunoreactive GH levels. Levels of immunoreactive hormone in the hypothalamic and amygdaloid regions showed an opposite effect, that of a significant ( $P = < 0.01$ ) increase in GH levels in operated animals, as compared to sham controls. This data is illustrated in Fig. 48 . All other brain areas showed no significant change in immunoreactive GH content when compared to sham controls four weeks after thyroidectomy, as shown in table 30.

b. Thyroid Stimulating Hormone: Effect of Removal of Target Organs

(i) Adrenalectomy

As shown in Fig. 49 , four weeks after adrenalectomy, levels of circulating immunoreactive TSH remained constant, whereas a significant fall in pituitary TSH levels was observed ( $P = < 0.05$ ). Assay of hormone levels in various brain regions showed that only in the hypothalamus was any significant change brought about by removal of the adrenal glands (Table 31).

## GH LEVELS 4 WEEKS AFTER ORCHIDECTOMY




Figure 47. Effects of orchidectomy on growth hormone levels in serum, pituitary, and brain. NS = not significant.

TABLE 29

## EFFECT OF ORCHIDECTOMY (4 WEEKS) ON BRAIN GH LEVELS

(ng/mg protein)\*\*

	<u>n</u>	<u>Sham</u>	<u>Castrated</u>	<u>Statistical Significance</u>
AMYGDALA	60	6.12 ± 0.35	5.80 ± 0.10	NS
HYPOTHALAMUS	60	3.60 ± 1.20	3.95 ± 0.37	NS
CAUDATE	60	1.29 ± 0.51	1.02 ± 0.81	NS
THALAMUS	60	2.10 ± 0.39	1.95 ± 0.26	NS
HIPPOCAMPUS	60	1.13 ± 0.04	1.42 ± 0.31	NS
CORTEX	60	1.72 ± 0.33	1.75 ± 0.62	NS
CEREBELLUM	60	0.04 ± 0.02	0.06 ± 0.01	NS
ANTERIOR PITUITARY	60	1.86 × 10 <sup>5</sup> ± 0.40	1.72 ± 0.31	NS
<u>GH ng/ml**</u>				
SERUM	60	185.0 ± 22.0	177 ± 33.0	NS

n = total animals

\*\* = mean ± S. E. M. of 3 experiments.

## GH LEVELS 4 WEEKS AFTER THYROIDECTOMY




Figure 48. Effect of thyroidectomy on growth hormone levels in serum, pituitary, and brain.



TABLE 30

## EFFECTS OF THYROIDECTOMY (4 WEEKS) ON BRAIN GH LEVELS

(ng/mg protein)\*\*

	<u>n</u>	<u>Sham</u>	<u>Thyroidectomized</u>	<u>Statistical Significance</u>
AMYGDALA	60	4.52 ± 0.63	8.21 ± 0.96	p = < 0.01
HYPOTHALAMUS	60	2.75 ± 0.25	4.82 ± 0.65	p = < 0.01
CAUDATE	60	1.35 ± 0.05	1.39 ± 0.10	NS
THALAMUS	60	1.45 ± 0.12	0.95 ± 0.31	NS
HIPPOCAMPUS	60	1.87 ± 0.31	2.10 ± 0.26	NS
CORTEX	60	1.63 ± 0.20	1.24 ± 0.41	NS
CEREBELLUM	60	0.16 ± 0.05	0.05 ± 0.01	NS
ANTERIOR PITUITARY	60	9.5 × 10 <sup>5</sup> ± 0.7	1.1 × 10 <sup>5</sup> ± 0.4	p = < 0.01
GH ng/ml**				
SERUM	60	212 ± 25.0	93.0 ± 10.6	p = < 0.01

n = total animals

\*\* = mean ± S. E. M. of 3 experiments

## TSH LEVELS 4 WEEKS AFTER ADRENALECTOMY




Figure 49. Effect of adrenalectomy on TSH levels in serum, pituitary and brain.  
NS = not significant.

TABLE 31

## EFFECT OF ADRENALECTOMY (4 WEEKS) ON BRAIN TSH LEVELS

(ng/mg protein)\*\*

	<u>n</u>	<u>Sham</u>	<u>Adrenalectomized</u>	<u>Statistical Significance</u>
HYPOTHALAMUS	60	176.25 ± 5.31	225.65 ± 6.73	p = < .02
AMYGDALA	60	27.52 ± 2.63	32.5 ± 3.79	NS
CEREBELLUM	60	22.26 ± 6.33	20.09 ± 2.10	NS
THALAMUS	60	16.53 ± 0.59	16.75 ± 0.26	NS
CAUDATE	60	11.96 ± 0.73	12.13 ± 0.98	NS
CORTEX	60	14.25 ± 1.76	12.75 ± 0.52	NS
HIPPOCAMPUS	60	10.76 ± 0.92	9.83 ± 1.79	NS
ANTERIOR PITUITARY	60	3.6 x 10 <sup>5</sup> ± 0.26	2.5 x 10 <sup>5</sup> ± 0.50	p = < .05
<u>TSH ng/ml**</u>				
SERUM	60	826.91 ± 50.75	750.26 ± 75.63	NS

n = total animals

\*\* = mean ± S. E. M. of 3 experiments

As shown in Fig. 49 , a significant ( $P = < 0.02$ ) rise in immunoreactive TSH levels occurred in adrenalectomized animals' hypothalamic tissue, as compared to sham-operated controls.

(ii) Oophorectomy

Removal of the ovaries did not produce any significant change in TSH immunoreactive hormone levels in serum, pituitary, or brain tissue, as illustrated by the data obtained in Fig. 50 and Table 32.

(iii) Orchidectomy

Whereas removal of the testes caused a significant fall ( $P = < 0.05$ ) in immunoreactive TSH levels in serum and pituitary, a significant rise ( $P = < 0.02$ ) was shown for hypothalamic tissue four weeks postoperatively (Fig. 51 ). Other brain regions showed no significant change in TSH levels (Table 33).

(iv) Thyroidectomy

When immunoreactive TSH was measured four weeks after thyroidectomy, it was shown that serum levels of the immunoreactive hormone rose dramatically ( $P = < 0.001$ ) as compared to controls, whereas a significant fall in pituitary levels ( $P = < 0.02$ ) was observed (Fig. 52 ). This fall in pituitary levels was shown in all three experiments and is contrary to what would be expected after loss of  $T_4$  production by thyroidectomy. However, as noted from the literature previously, it is believed that the turnover of TSH is so great in the pituitary, under these conditions, as to result in measured low basal levels. Although the hypothalamus showed

## TSH LEVELS 4 WEEKS AFTER OOPHORECTOMY




Figure 50. Effect of oophorectomy on TSH levels in serum, pituitary and brain.  
 NS = not significant.

TABLE 32

## EFFECTS OF OOPHORECTOMY (4 WEEKS) ON BRAIN TSH LEVELS

(ng/mg protein)\*\*

	<u>n</u>	<u>Sham</u>	<u>Oophorectomized</u>	<u>Statistical Significance</u>
HYPOTHALAMUS	60	209.63 ± 5.11	212.90 ± 6.73	NS
AMYGDALA	60	33.96 ± 2.59	35.05 ± 3.72	NS
CEREBELLUM	60	16.22 ± 0.79	17.69 ± 0.95	NS
THALAMUS	60	18.13 ± 2.11	17.98 ± 0.63	NS
CAUDATE	60	12.93 ± 0.89	14.60 ± 0.84	NS
CORTEX	60	12.75 ± 2.30	11.98 ± 0.68	NS
HIPPOCAMPUS	60	11.38 ± 1.62	12.95 ± 0.98	NS
ANTERIOR PITUITARY	60	2.4 × 10 <sup>5</sup> ± 0.9	2.2 × 10 <sup>5</sup> ± 0.5	NS
TSH ng/ml **				
SERUM	60	640 ± 120.0	602 ± 35.0	NS

n = total animals

\*\* = mean ± S. E. M. of 3 experiments

## TSH LEVELS 4 WEEKS AFTER CASTRATION




Figure 51. Effect of orchidectomy on TSH levels in serum, pituitary and brain.

TABLE 33

## EFFECTS OF ORCHIDECTOMY (4 WEEKS) ON BRAIN TSH LEVELS

(ng/mg protein)\*\*

	<u>n</u>	<u>Sham</u>	<u>Castration</u>	<u>Statistical Significance</u>
HYPOTHALAMUS	60	205.61 ± 7.39	280.00 ± 11.65	p = < 0.02
AMYGDALA	60	26.32 ± 4.61	35.92 ± 1.95	NS
CEREBELLUM	60	13.40 ± 0.16	14.24 ± 1.36	NS
THALAMUS	60	12.46 ± 2.51	15.36 ± 0.95	NS
CAUDATE	60	16.39 ± 1.55	13.97 ± 2.60	NS
CORTEX	60	14.61 ± 0.78	12.65 ± 1.20	NS
HIPPOCAMPUS	60	12.59 ± 0.90	12.78 ± 1.10	NS
ANTERIOR PITUITARY	60	2.4 x 10 <sup>5</sup> ± 0.51	1.2 x 10 <sup>5</sup> ± 0.42	p = < 0.05
TSH ng/ml**				
SERUM	60	640.00 ± 79.6	405.00 ± 59.70	p = < 0.05

n = total animals

\*\* = mean ± S. E. M. of 3 experiments



## TSH LEVELS 4 WEEKS AFTER THYROIDECTOMY




Figure 52. Effect of thyroidectomy on TSH levels in serum, pituitary and brain.  
 NS = not significant.

TABLE 34

## EFFECTS OF THYROIDECTOMY (4 WEEKS) ON BRAIN TSH LEVELS

(ng/mg protein)\*\*

	<u>n</u>	<u>Sham</u>	<u>Thyroidectomized</u>	<u>Statistical Significance</u>
HYPOTHALAMUS	60	158.36 ± 10.63	207.00 ± 21.35	NS
AMYGDALA	60	33.39 ± 1.65	36.89 ± 2.71	NS
CEREBELLUM	60	15.96 ± 1.30	14.63 ± 2.50	NS
THALAMUS	60	18.15 ± 0.45	17.10 ± 2.73	NS
CAUDATE	60	16.72 ± 0.87	15.96 ± 0.55	NS
CORTEX	60	14.10 ± 2.61	16.59 ± 0.71	NS
HIPPOCAMPUS	60	13.78 ± 0.60	12.11 ± 1.24	NS
ANTERIOR PITUITARY	60	3.5 × 10 <sup>5</sup> ± 0.80	1.9 × 10 <sup>5</sup> ± 0.52	p = < 0.02
TSH ng/ml**				
SERUM	60	462.00 ± 59.60	2226.00 ± 102.3	p = 0.001

n = total animals

\*\* = mean ± S. E. M. of 3 experiments

TABLE 35

SIGNIFICANCE OF CHANGES IN BRAIN GH AFTER HYPOPHYSECTOMY AND TARGET ORGAN REMOVAL (4 WEEKS)

<u>PROCEDURE</u>	<u>Pituitary</u>	<u>Serum</u>	<u>Hypothalamus</u>	<u>Amygdala</u>	<u>Other Brain Regions</u>
HYPOPHYSECTOMY	--	p = < 0.001	p = < 0.01	p = < 0.01	Thalamus: p = < 0.05 Others: NS
ADRENALECTOMY	NS	NS	NS	NS	NS
OOPHORECTOMY	NS	p = < 0.01	p = < 0.01	p = < 0.01	NS
ORCHIDECTOMY	NS	NS	NS	NS	NS
THYROIDECTOMY	p = < 0.01	p = < 0.01	p = < 0.01	p = < 0.01	NS

TABLE 36

SIGNIFICANCE OF CHANGES IN BRAIN TSH AFTER HYPOPHYSECTOMY AND TARGET ORGAN REMOVAL (4 WEEKS)

<u>PROCEDURE</u>	<u>Pituitary</u>	<u>Serum</u>	<u>Hypothalamus</u>	<u>Amygdala</u>	<u>Other Brain Regions</u>
HYPOPHYSECTOMY	--	p = < 0.01	p = < 0.02	NS	NS
ADRENALECTOMY	p = < 0.05	NS	p = < 0.02	NS	NS
OOPHORECTOMY	NS	NS	NS	NS	NS
ORCHIDECTOMY	p = < 0.05	p = < 0.05	p = < 0.02	NS	NS
THYROIDECTOMY	p = < 0.02	p = < 0.001	NS	NS	NS
T <sub>4</sub> ADMINIS- TRATION	p = < 0.002	p = < 0.002	NS	NS	NS

a slight rise in TSH levels, it was not statistically significant, as was the case for all other brain areas, as shown in Table 34. Levels of circulating  $T_4$  fell from 3.42 to 0.02  $\mu\text{g}/\text{dl}$ , while the %  $T_3$  resin uptake remained constant in thyroidectomized animals.

The significance of the changes discussed in GH and TSH after hypophysectomy and target organ removal are shown in Tables 35 and 36, respectively.

0. THYROXINE LEVELS: EFFECT OF AN INCREASE IN CIRCULATING  $T_4$  ON SERUM AND TISSUE LEVELS OF TSH AND  $T_4$

The effect of injecting IM, 0.4  $\mu\text{g}$   $T_4$ /gm body weight, daily for 7 days into intact adult rats on TSH levels in serum, pituitary, and amygdaloid tissue, is illustrated in Fig. 53. Whereas approximately 200  $\mu\text{g}$   $T_4$  caused a dramatic, significant ( $P = < 0.002$ ) fall in circulating TSH (400  $\rightarrow$  58 ng/ml) and pituitary TSH (300  $\rightarrow$  50  $\mu\text{g}/\text{mg}$  protein), presumably by a "negative feedback" effect, no significant change in TSH levels was seen in amygdaloid or other brain tissues, including the hypothalamus. Circulating  $T_4$ 's were increased from 4.70  $\rightarrow$  17.42  $\mu\text{g}/\text{dl}$  in treated animals, whereas the %  $T_3$  resin uptake remained constant. In hypophysectomized animals, however, the effect of a similar dose of  $T_4$  on brain levels of TSH was to cause a statistically significant ( $P = < 0.05$ ) rise in the levels of IRTSH in the hypothalamus as compared to saline-injected hypophysectomized animals.  $T_4$  levels in such animals rose from 0.8  $\rightarrow$  24  $\mu\text{g}/\text{dl}$ , and similarly %  $T_3$  uptake also rose from 48.5  $\rightarrow$  70.229. Other brain regions showed no change in immunoreactive TSH levels after thyroxine treatment.




Figure 53. The effect of injecting IM. 0.4 µg T<sub>4</sub>/gm body weight daily for 7 days into intact adult rats, on TSH levels in serum, pituitary and amygdala. NS = not significant.

P. ONTOGENY OF PITUITARY AND BRAIN PITUITARY-LIKE HORMONES

In order to examine the levels of hormone at various stages of development, rat fetuses from ten days to birth and pups from birth to 30-day-old adults were sacrificed and brain homogenates analyzed for pituitary hormonal content by RIA.

a. Growth Hormone

(i) Growth hormone in brain and pituitary tissues

In Fig. 54 , levels of immunoreactive growth hormone, expressed as ng hormone/mg protein, in terms of fetal and neonatal age of the animal, are shown. This data demonstrated the appearance of immunoreactive hormone at fetal day 10 and a remarkable rise in the level of detectable hormone extracted from the hypothalamus and amygdala just prior to birth, or day 21 of gestation in the rat. Postpartum, hormone concentration in these brain regions fell abruptly and then rose, gradually attaining adult levels at around 30 days of age, essentially at the time of puberty. All other brain parts--thalamus, caudate, and cortex--showed a similar, although not so pronounced, preparturitional rise over adult levels of growth hormone. The hippocampus showed the most dramatic preparturitional increase of all brain regions compared to adult levels.

Levels of pituitary growth hormone were also measured over the same age span, as shown in Fig. 55 . Contrasting to the finding of pituitary-like hormones in the rodent brain as early as fetal day 10, growth hormone was not detectable until day 12 of fetal life in the pituitary anlage

# GROWTH HORMONE ng/mg PROTEIN




Figure 54. The ontogenetic development of growth hormone levels in the CNS from fetal to adulthood in the rat.






Figure 55. The ontogenetic development of growth hormone in the pituitary from fetal to adulthood in the rat.




Figure 56. The ontogenetic development of growth hormone levels in the pituitary and CNS from fetal to adulthood in the rat.

and showed no preparturitional surge, but rather a gradual rise toward adult levels between 10 and 20 days after birth. These two sets of data can be superimposed, as shown in Fig. 56 , to show the relationship which exists between the hormonal levels in brain versus pituitary tissue.

(ii) Growth hormone in fetal and maternal sera and amniotic fluid

Levels of GH, in terms of ng of hormone/ml, of amniotic fluid, fetal sera, and maternal sera were also measured at day 17, 19 and 21 days gestation, and for several days postpartum in the case of sera. As shown in Fig. 57 , amniotic fluid levels continued to rise steadily until parturition, and levels of fetal serum growth hormone also rose slowly until the first day after delivery and then fell to adult levels. At day 21, immunoreactive growth hormone levels were substantially higher in fetal serum than in maternal serum, which essentially showed no change throughout pregnancy and after delivery. Levels of extracted placental immunoreactive growth hormone were also measured and remained constant and at levels approximating those of maternal serum.

b. Thyroid Stimulating Hormone

(i) Thyroid stimulating hormone in brain and pituitary tissue

A similar pattern, in terms of development, was observed when brain TSH was measured in 10-day fetal and up to 30-day old adult rat brain tissue. All brain regions containing this material showed a dramatic preparturitional rise and abrupt fall just after birth. Fig. 58 shows this phenomenon for the hypothalamus and amygdala, in terms of ng immunoreactive TSH/mg protein. When pituitary, immunoreactive TSH levels,




Figure 57. Levels of growth hormone (ng/ml) in amniotic fluid, fetal serum, and maternal serum from 17 days fetal to 2 days postnatal in the rat.

determined over the same time period, are superimposed on this data, as in Fig. 59, the relationship between the development of hormonal levels in these two distinct tissues can be seen. Here again, TSH could be measured by day 10 of fetal life in the brain tissue extracts, whereas immunoassayable TSH was detectable by day 15-17 in the fetal pituitary.

(ii) Thyroid stimulating hormone in fetal and maternal sera and amniotic fluid

Unlike growth hormone, TSH amniotic fluid levels showed a fall before birth, whereas fetal serum levels remained constant and then dropped quickly after delivery. No change in maternal TSH serum levels was noted during the entire study period as shown in Fig. 60. Placental TSH levels remained constant throughout gestation, levels of hormone corresponding to those found in maternal sera.

c. Summary of Fetal Brain Levels of Immunoreactive GH and TSH Data

In Table 37, GH and TSH levels in various regions of the 21-day fetal CNS were compared in terms of the percent rise above levels normally found in the adult rat brain. The hippocampus, hypothalamus, and amygdaloid regions showed the greatest preparturitional surge in pituitary-like hormones.

d. Immunological Characterization of Fetal Brain Hormones

When the immunological characteristics of fetal brain growth hormone were compared with those from fetal pituitary homogenates, parallel




Figure 58. The ontogenetic development of TSH levels in the CNS from fetal to adulthood in the rat.




Figure 59. The ontogenetic development of TSH levels in the pituitary and CNS from fetal to adulthood in the rat




Figure 60. Levels of TSH in amniotic fluid, fetal serum, and maternal serum from 17 days fetal to 2 days postnatal in the rat.



TABLE 37

GH- AND TSH-LIKE IMMUNOREACTIVE PEPTIDES  
IN THE 21-DAY FETAL CNS

	<u>% ABOVE ADULT LEVELS</u>	
	<u>GH</u>	<u>TSH</u>
Hippocampus	2000	1000
Hypothalamus	300	1000
Amygdala	600	600
Cerebellum	10	300
Cortex	800	100
Caudate	15	10

dose-response displacement curves were obtained as shown in Fig. 61 .

Fetal TSH from brain tissue showed similar parallelism with pituitary TSH. This indicated immunological similarity between the hormones from the two different sources. Although the brain curve was in fact superimposable on the pituitary curve, it had been displaced downward for visual effect.

e. Molecular Weight Determinations of Fetal Brain Hormones

The possibility of heterogenous forms of these brain peptides occurring in brain and pituitary tissue of fetuses and adult rats was also examined. As shown in Fig. 62 , using a G-100 Sephadex column, 22-day-old fetal brain extracts were co-chromatographed with  $^{125}\text{I}$  labeled adult rat pituitary GH. It can be seen that the peak of immunoreactive GH-like material co-eluted with the radioactive marker.

In Fig. 63 , a similar column chromatography experiment demonstrated the co-elution of immunoreactive TSH-like material from fetal CNS extracts with  $^{125}\text{I}$  TSH from adult pituitaries, suggesting molecular similarity between fetal and adult brain pituitary-like hormones.




Figure 61. Logit log plot demonstrates parallelism between pituitary and fetal CNS growth hormone.




Figure 62. Elution pattern of lyophilized 21 day old fetal brain and  $^{125}\text{I}$  adult rat pituitary growth hormone on a Sephadex G-100 column eluted with veronal buffer pH 8.6.




Figure 63. Elution pattern of lyophilised 21 day old fetal brain and  $^{125}\text{I}$  adult rat pituitary TSH on a Sephadex G-100 column eluted with 0.01M  $\text{PO}_4$  buffer pH 7.6.

## 6. DISCUSSION OF DATA

Initially prompted by the observation of the presence of immunoreactive growth hormone-like material in the media of cultured rat nervous tissue reported by Pacold et al (1976), this study has attempted to characterize, in various ways, this growth hormone-like material. In addition, we have sought to characterize a TSH-like material, also found in the tissue culture medium of cultured rat nervous tissue, and discovered in the course of these original studies. The relevance of the data obtained from these investigations will be discussed in the light of other recent observations also dealing with the discovery of diverse peptides in the central nervous system, a phenomenon only recently appreciated. Previously, monoaminergic substances were considered the only chemical neurotransmitters, whereas it now seems abundantly likely that peptides of neuronal origin from cells within the CNS also have neurotransmitter roles. In addition, new and unusual roles may be assigned which could include effects on brain development and growth, perception of pain, normal and disturbed behavior, and even effects on intelligence to mention but a few possible roles for these recently discovered brain peptides.

The finding of such peptides and the speculation that they may play a physiological role in neural function is a relatively recent suggestion by de Wied (1980) and is now supported by several lines of evidence, such as the induction of lordosis by intracerebral injection of luteinizing-releasing hormone in female rats (Foreman, 1977). Observations such as these have led to the concept that many peptides, whatever

their traditional tissue of origin, exert effects upon the brain which lead to behavioral consequences apart from their classical endocrine effects.

The main objectives of this particular study concerning the presence of two such peptides, GH and TSH, traditionally of pituitary origin, in rodent and also in primate brain tissue, is to show that these hormones are probably of CNS origin rather than diffusion contaminants from the pituitary, that their presence is largely independent of pituitary function, and finally to speculate on the possible reasons for their presence in the brain.

Initial experiments were concerned with the systematic extraction and mapping of different areas of the rodent and primate brain. Both a growth hormone-like and thyroid stimulating-like material were found, by radioimmunoassay, to be unevenly distributed throughout the central nervous system of both rat and primate. These data immediately suggested a CNS origin rather than a pituitary origin for such extractable hormone-like materials, as an overall uniform distribution of hormone concentration throughout the CNS would have been indicative of contamination from circulating pituitary hormones.

Immunoreactive growth hormone-like material was found in greatest concentrations, in terms of ng hormone/mg protein in the amygdaloid nuclear region of the rat CNS by tissue extraction experiments. The rat hypothalamus was also found to contain substantial amounts of such immunoreactive material.

Extractable immunoreactive TSH-like hormone was also detectable in both rodent and primate brain tissue. The rodent hypothalamic region was found to contain the largest amount of detectable material, and a TSH-like substance was also present in the amygdaloid nucleus and other brain regions to a lesser extent. Large differences between pituitary and brain levels of both hormones were observed, even though these two tissues underwent similar extraction procedures. Several explanations for such wide variations can be envisaged, such as the possibility of differences in the rates of peptide synthesis in each tissue, and the biological half-lives of hormones of pituitary and brain origin. Another possibility for widely variant concentrations could be that differences in tissue extractable levels of hormone are due to the types of precursors present and the subsequent post-translational processing of such precursors. De Wied (1980), for example, when discussing the two apparent pools of hypothalamic and pituitary ACTH, suggests that such processing of brain peptides may give rise to fragments of hormone which, although having biologically significant effects on the surrounding CNS milieu, are not recognizable by either immunological or biological assays. Similar arguments could be made to speculate on the apparent differences in the magnitude found between growth hormone and thyroid stimulating hormone-like concentrations in brain and pituitary pools.

Preliminary studies showed that, in the case of both immunoreactive GH- and TSH-like material in primate brains, the hypothalamus exhibited the highest concentrations of these particular CNS peptides. Although growth hormone was found in most other areas of the primate brain, the



distribution of TSH was such that large areas of the primate CNS appeared to contain non-measurable quantities.

Further speculation on this data from primate CNS will have to await further in-depth investigation. Its measured presence, however, does indicate that certain pituitary-like peptides apparently are not restricted to the rodent brain.

Mapping locations of GH- and TSH-like peptides in the rodent central nervous system, using immunohistochemical techniques, demonstrated the presence of TSH in various areas of the central nervous system. In the hypothalamus, TSH was distinguishable in cell bodies of the dorsomedial and ventromedial nuclei, and most of the nuclei of the amygdala showed positive staining for the presence of TSH. Other areas of the CNS shown to have extractable hormone, but which did not show positive staining by the immunohistochemical technique, are known to be innervated by fibres originating from cell bodies found in the hypothalamus and the amygdaloid nuclei. Axon collaterals of hypothalamic tuberinfundibular neurons extend to brain sites quite distant from the hypothalamus, such as the thalamus, the limbic system, other than the amygdala, and the cortex (Martin et al, 1975), and therefore represent the means whereby hormone, possibly produced in the cell bodies of the hypothalamus, might be axonally transported to other brain regions. In preliminary studies, the amygdala and hypothalamus also showed the largest number of positively staining cell bodies for immunoreactive GH-like material.

At this level of investigation, it is difficult to associate any

particular cell type, either glial or neuronal, with a higher than average hormonal content, although from the immunohistochemistry data, neurons were more likely involved. However, in tissue culture studies with the neuroblastoma cell line N<sub>4</sub>TGI, TSH-like material was found in neuronal-type cells. Despite this, the tumor origin of such cell lines cautions its use as a positive indication that normal, non-tumor neuronal cells, rather than glial or interstitial cells, are involved in hormone synthesis or storage in the CNS. It can only be concluded from our tissue culture studies, therefore, that primary mixed cell cultures of rat nervous tissue appear to release both growth hormone and thyroid stimulating hormone-like peptides into the growth media over a 30-day time period. Further studies utilizing techniques to propagate pure neuronal or glial primary cell cultures are needed to further this study.

In order to pinpoint the association of these peptides with a particular cell fraction, our ultra-centrifugation studies showed that both TSH- and GH-like immunoreactive material were isolated together with the synaptosomal-rich fraction of CNS tissue homogenates, suggesting the possibility that, like other peptides such as somatostatin, TRH and VIP, these pituitary-like hormones may be associated with pre- or post-synaptic membranes in vivo.

There exists the possibility, as shown by these same ultra-centrifugation studies, that part of the extractable pool of GH- and TSH-like material found in the hypothalamus, may be present unassociated with any specific cellular fraction. Whether the origin of this free pool of hormone is the brain or pituitary is open to speculation.

The studies which were undertaken to characterize these extractable immunoassayable peptides from rat brain tissue in a chemical, immunological, and biological manner revealed that hormones of brain origin were indistinguishable from those found in the pituitary, in terms of molecular weight, as shown by column chromatography, and in terms of immunological identity, as shown by parallelism studies. Their biological activity also demonstrated a similarity to their respective pituitary counterparts in terms of possessing the capability of eliciting a simple TSH-like response on the histology of the thyroid or, for growth hormone-like material, to increase the width of epiphyseal cartilage in the tibias of hypophysectomized animals.

Such immuno- and biological similarities between pituitary and brain hormones would suggest, without further evidence, the possibility that the presence of pituitary-like hormones in the brain is merely due to pituitary hormone contamination of these CNS tissues by circulating pituitary hormone.

Experiments designed to test this concept were carried out in this study using radioactively-labeled pituitary hormone in order to test the effectiveness of the blood-brain barrier. They show that it seems unlikely that significant quantities of pituitary TSH or GH are transported to the brain either by the general circulation or portal vessel routes. Similarly, Mezey et al (1979), showed that whole brain uptake of  $^3\text{H}$  ACTH, given intravenously to adult rats, was less than 0.07% of the initial radioactive hormone dose given. In the experiments reported here, no radioactivity was found in brain tissue of either young or adult rats

after systemic injection of  $^{125}\text{I}$ -labeled pituitary GH or TSH. This indicates that, in the particular rodent model system which was used throughout this study, the blood-brain barrier was apparently intact throughout life, eliminating any possibility of cross-contamination of brain tissue by pituitary hormones. Other indications of the bidirectional intactness of this barrier were that no increase or decrease in circulating hormone levels after experimental manipulations, such as hypophysectomy and target organ removal, produced any subsequent rise in the hormonal levels found in brain tissue. Similarly, no increase or decrease in brain tissue levels produced a concomitant increase or decrease in the concentration of circulating pituitary hormone. Experiments by Weiss et al (1975) substantiate this data in that pituitary cells implanted in the rat brain ventricles of hypophysectomized rats did not produce normalization of circulating pituitary hormone levels, even though it could be demonstrated that such implants continued to produce growth hormone and thyroid stimulating hormone. Whatever the role of the pool of brain hormones, it is unlikely to be as a reserve pool which can be drawn on in times of pituitary deficiency.

Besides the demonstration of blood-brain barrier intactness to circulating pituitary hormone, evidence of a CNS origin of brain peptides, other than GH and TSH, comes from similar experiments comparing brain tissue levels of hormone-like material before and after removal of the pituitary gland. That the concentration of ACTH,  $\alpha$  MSH,  $\beta$  endorphin, and prolactin, as reported previously by Liotta et al (1978), Kobayashi et al (1978), and Toubeau et al (1979), are essentially unchanged in most

of the brain areas examined after such treatment, is consistent with the possibility of synthesis of these peptides by the brain tissue itself..

Such variations in hormonal level that do occur after hypophysectomy in the data obtained in this study for TSH could be the result of several factors, such as changes brought about in the levels of a known regulatory factor to pituitary hormone, e.g., dopamine, by hypophysectomy. The slight decrease in TSH levels seen 28 days after hypophysectomy are most probably too late an event to be the result of tissue damage incurred by surgical intervention but could be a reflection of falling dopamine levels, which have been observed in the hypothalamus after long-term pituitary removal (Nagatsu et al, 1964). A fall in dopamine levels would have an adverse effect on the content of TSH in a normal pituitary; however, whether similar effects can be induced on all or part of the hypothalamic pool of TSH-like peptide can only be speculated upon at this time.

Another controlling factor, in terms of hypothalamic levels of TSH after hypophysectomy, could be the effect of falling TRH levels, reported by Nemeroff et al (1979) to occur in the hypothalamus, but not in other brain regions, after pituitary removal. These levels were shown by Nemeroff to be restorable by  $T_4$  therapy. In a study for this dissertation using long-term hypophysectomized animals which received  $T_4$  substitution treatment, levels of extractable immunoreactive TSH in the hypothalamic region were shown to be restored by such therapy. The possibility, therefore, exists of a TRH-influenced control over part of the hypothalamic TSH pool and would provide an explanation for the fall seen

in hypothalamic TSH levels and not other CNS pools of TSH after long-term hypophysectomy.

Examination of the data concerning immunoreactive GH levels in brain tissue after hypophysectomy show no change in hormonal levels for most of the brain areas examined. Exceptions were the levels of growth hormone in hypothalamic and amygdaloid tissues, which demonstrated an initial fall in hormone levels after 7 days of hypophysectomy, followed by a remarkable rise in hormone content after 28 or more days without a pituitary.

The lack of such an early fall in GH measured from tissue-cultured amygdaloid and hypothalamic tissue derived from long-term hypophysectomized animals indicates the possibility that after a certain time period of hypophysectomy, such as 7 days, a growth hormone inhibitory factor, e.g., somatostatin, may be present in sufficiently lowered amounts as to be unable to cause a normal depression of growth hormone levels in vivo. This theory is substantiated by experiments, previously quoted by Baker et al (1976), showing that somatostatin levels do fall in the hypothalamus about 10 days after hypophysectomy. The ability of somatostatin to suppress "brain" growth hormone has been shown in our study using tissue cultured normal CNS tissue. Perhaps this initial decrease in amygdaloid and hypothalamic tissue levels of growth hormone is due to somatostatin, and the increased levels after 10-14 days hypophysectomy are due to the disappearance of this inhibitory factor from the CNS milieu because of pituitary ablation. Certainly the implication that animals in such experiments were incompletely hypophysectomized (Moldow et al, 1978<sup>a,b</sup>) has

not been substantiated either by microscopic inspection of the empty sella turcica, by levels of circulating  $T_4$ , TSH, and GH, which were measured in the serum of operated animals, or by gross examination of atrophied target glands such as the thyroid, adrenals, and gonads in hypophysectomized rats.

Enhancement of levels of ACTH-like material in rat hypothalamus, but not in other brain areas, 12 days after hypophysectomy has been demonstrated by Krieger and Liotta (1979), which tends to support our data for GH-like levels in the CNS after pituitary removal. These experiments add further evidence to the possibility that these pituitary-like peptides in the CNS are not of pituitary origin.

Other data, also corroborated by Krieger and Liotta's group, are those obtained from the tissue culture of cells from rat central nervous system tissues, demonstrating the presence of immunoreactive GH and TSH in the growth medium for up to 30 days of cell culture. The ability of cells derived from the arcuate nucleus of the hypothalamus to release ACTH into tissue culture media for up to 17 days, has similarly been reported (Liotta et al, 1979<sup>a</sup>). Using this model system of dispersed hypothalamic cells, it has been possible to demonstrate that <sup>3</sup>H-labeled amino acids can be incorporated into material with the characteristics of the pituitary precursor of ACTH, giving credence to a CNS origin for this traditionally thought of pituitary hormone (Liotta et al, 1979<sup>b</sup>). Evidence also exists that an ACTH-like pituitary peptide other than GH and TSH persists after long-term hypophysectomy in CNS tissues and that it is most likely synthesized in situ within the brain.

Other experiments have sought to investigate the mechanisms of regulation of these pituitary-like peptides, compared to those regulating pituitary hormones. Such experiments have shown, for example, that concentrations of hypothalamic immunoreactive ACTH-like material are not significantly altered 24 hours or 2 weeks after adrenalectomy, after dexamethasone or corticosteroid administration, or by the stress of ether administration (Liotta et al, 1978). All of these procedures are normally associated with alteration in the levels of ACTH in the pituitary gland and serum. Alterations of levels of ACTH in the pituitary and serum due to adrenalectomy are the result of feedback mechanisms.

The secretory functions of the anterior pituitary appear to be modulated by hormonal signals originating in such peripheral target glands as the adrenals, thyroid, and gonads, and in the pituitary itself, through complicated networks of feedback mechanisms called, respectively, "long" and "short." Within each category of feedback systems, "negative" as well as "positive" mechanisms have been described. A feedback system in which the input, represented by a hormone originating in a peripheral target gland, reduces the output of pituitary hormone is called "negative" and where a feedback system, in which increased secretion of a peripheral hormone stimulates anterior pituitary function, it is generally called "positive." The unaltered hypothalamic levels of ACTH after adrenalectomy, in contrast to increased levels found in the pituitary due to a lack of negative feedback of circulating adrenal cortisone, reflect a difference in the control mechanisms of brain-situated hormone compared to pituitary hormones. Similarly, Rossier et al (1977<sup>a</sup>) have observed no change in



the concentrations of immunoreactive  $\beta$ -endorphin-like material in the brains of adrenalectomized animals. Animals subjected to the stress of foot shock showed some decrease in  $\beta$ -endorphin-like immunoreactivity in the hypothalamus, but not in other brain areas (Rossier et al, 1979<sup>b</sup>). Other studies by Havrankova et al (1979) have demonstrated that "brain insulin" concentrations are also unaffected by extreme changes in circulating hormone concentrations.

Removal of such target organs as the adrenals and testes, in our experiments, appeared to have no effect on the levels of growth hormone in either serum, pituitary, or other brain areas analyzed. Obviously, no feedback mechanisms occur in vivo involving these peripheral organs and pituitary or brain growth hormone. Oophorectomy induced a significant fall in the levels of circulating GH, but no significant difference in pituitary levels of controls or in operated animals. A significant rise in the levels of hypothalamic and amygdaloid immunoassayable GH concentrations was seen, however, after the removal of the ovaries. Other brain areas appeared unaffected by such ovarian ablation.

Thyroidectomy produced a similar picture of a fall in serum and pituitary levels of immunoassayable growth hormone and a rise in the hypothalamic and amygdaloid levels of this immunoreactive peptide. As with the gonadectomized GH content, brain regions other than hypothalamus and amygdala appeared unaffected by removal of the thyroid. What seems most striking from data derived from these experiments is the demonstration of an apparent independence between the levels of, e.g., growth hormone found in the pituitary and in serum, compared to levels found

in specific areas of the brain.

The possibility of the existence of a short loop feedback regulatory system of some kind between the hypothalamus and the anterior pituitary would be a possible explanation of the data we have described from the present studies. It is disturbing, however, that GH and TSH levels, in areas other than the hypothalamus and amygdala, are unaffected by specific target organ removal and/or trophic hormone treatments, especially if we hypothesize from the immunohistochemical data, that extractable hormone levels in one region may be derived from cell bodies in the hypothalamus and amygdaloid nuclei. The possibility, therefore, exists that in situ synthesis of peptides in these brain areas exists at a level undetectable by the immunohistochemical technique performed in these studies. Further investigations should therefore aim at improving the sensitivity of this particular technique for detecting GH and TSH in brain areas other than the hypothalamus and amygdala.

Adrenalectomy and orchidectomy in adult male rats produced similar divergent results between pituitary and serum TSH levels and those measured in the hypothalamus. Thyroidectomy resulted in a significant rise in circulating TSH and a lowering of pituitary TSH levels. No effect on brain levels of TSH-like peptides was observed after removal of the thyroid.

Immunoreactive GH and TSH of the brain, like ACTH and  $\beta$  endorphin, appear, therefore, to be unaffected in any particular way by the positive and negative hormonal feedback signals normally affecting the concentration of these peptides in the anterior pituitary gland and in the

circulation. This apparent dissociation was also reflected in the tissue culture studies described in this study, where  $T_4$  added to the growth media of growing anterior pituitary or brain tissue monolayers had little or no effect on levels of immunoreactive TSH found in the culture medium of normal brain cells. TSH levels from medium of anterior pituitary cells, so treated, reflected the anticipated decline in TSH production due to a negative long feedback loop operating on pituitary TSH secretion in the presence of high levels of the thyroid hormone,  $T_4$ .

Speculation on the presence of a short loop feedback between pituitary and hypothalamic pools, similar to that postulated between pituitary and pituitary stalk TSH pools by Baake et al (1967), was not substantiated by data from these studies. Our data confirms, rather, the findings of others who have attempted direct implantations of TSH into the hypothalamus in order to demonstrate a feedback control mechanism on the pituitary. Levels of TSH, so implanted, apparently fail to affect the concentration of pituitary TSH by a negative short feedback loop, making TSH unique in this regard among the anterior pituitary hormones.

Further evidence for the independence of pools or in situ synthesis of brain and pituitary growth and thyroid stimulating hormones comes from the results of the ontogenetic studies reported in this dissertation, where brain, serum, and pituitary levels of GH and TSH were examined during gestation, in the neonatal period, and into adulthood.

In such ontogenetic studies, it was apparent from the data that a striking preparturitional surge of immunoreactive GH and TSH occurs in

the 24 hours immediately before delivery, and is followed by a fall in the first day of extrauterine life. These peptides were found to possess similar molecular and immunological characteristics to their pituitary counterparts. The abrupt rise of a substance within the developing fetal CNS and serum just before birth is not an unusual finding. Several reports from other investigators have noted comparable patterns to those described here for GH and TSH. Some of these are shown in Table 38, and include such identifiable substances as glycogen (Kohle et al, 1977), histamine (Pearce et al, 1969), estrogen (Dohler et al, 1975), prostaglandins (Vale et al, 1972), and receptors for serotonin and estrogen (McClusky et al, 1979), as well as angiotensin II (Baxter et al, 1980). Other data show that the levels of other substances such as somatostatin (Blasquez et al, 1974),  $T_4$  and  $T_3$  (Rabie et al, 1979), and TRH (Schaeffer et al, 1980) rise in the CNS in the first few days after birth, as shown in Table 39.

Changes in hormone concentrations of TSH or GH in the fetal brain did not appear to reflect pituitary secretions as the patterns of acquisition were quite different. Growth hormone and TSH were first detected in the fetal brain at 10 days of fetal age and could not be detected in the fetal pituitary until 12-15 days of age. Furthermore, pituitary hormone concentrations did not show the dramatic preparturitional surge noted in the brain.

Radiolabeled GH and TSH given intraperitoneally to either the fetus, or pregnant rat, did not reach the fetal rat brain tissue in either case, indicating the intactness of both the fetal blood-brain barrier as well as

TABLE 38

## SUBSTANCES WHOSE LEVELS RISE JUST PRIOR TO BIRTH

IN THE CNS		IN THE FETAL SERUM	
<u>RECEPTORS FOR:</u>			
Angiotensin II	Baxter <u>et al</u> , 1980	Testosterone	Weisz <u>et al</u> , 1980
Renin	Baxter <u>et al</u> , 1980	ACTH	Rose <u>et al</u> , 1978
Estrogen	McClusky <u>et al</u> , 1979	Corticoids	Rose <u>et al</u> , 1978
		Growth Hormone	Bassett <u>et al</u> , 1970
Glycogen	Kohle <u>et al</u> , 1977		
Spermidine	Pearce <u>et al</u> , 1969	LH	) Courbier <u>et al</u> , 1978
Histamine	Pearce <u>et al</u> , 1969	FSH	
Substance P	Gilbert <u>et al</u> , 1979	TSH	
TRH in telencephalon	Schaeffer <u>et al</u> , 1980		
Estrogen	Dohler <u>et al</u> , 1975		
Vasotocin	Pavel <u>et al</u> , 1976		
PGF <sub>2</sub> $\alpha$ , PGE, PGE <sub>2</sub>	Vale <u>et al</u> , 1972		
	Liggins <u>et al</u> , 1973		
Ketone Bodies	Haymond <u>et al</u> , 1979		
Membrane Phospholipids/ Cholesterol	Kapitulnik <u>et al</u> , 1979	Prolactin	Riddick <u>et al</u> , 1979
		Estrogen	Dohler <u>et al</u> , 1975
<u>ENZYMES:</u>			
Glucose-6-PO <sub>4</sub> tase	Oliver <u>et al</u> , 1963		
Dehydrogenase	Andres <u>et al</u> , 1980		
Monoamine Oxidase	Snyder <u>et al</u> , 1979		
Fructose 1-6 Diphosphatase	Haymond <u>et al</u> , 1979		
Pyruvate Carboxylase	Haymond <u>et al</u> , 1979		
UDP-glucose-glycogen Transglucosylase	) Kohle <u>et al</u> , 1977		
Phosphoglucomutase			

TABLE 39

## SUBSTANCES WHOSE LEVELS RISE JUST AFTER BIRTH

<u>C N S</u>	<u>DAY FIRST DETECTABLE</u>	<u>SOURCE</u>
Monoamines	1 - 2	Coyle <u>et al</u> , 1973
Somatostatin	5	Blasquez <u>et al</u> , 1974
TRH in hypothalamus	6	Schaeffer <u>et al</u> , 1980
Androgen receptors	5	Vito <u>et al</u> , 1979
T <sub>4</sub> and T <sub>3</sub>	2 & 12	Rabié <u>et al</u> , 1979
Glycogen phosphorylase	10 min.	Jacquot <u>et al</u> , 1964

the fetal-maternal-placental barrier, to circulating pituitary hormones at the time of the preparturitional brain surge. It is also known that the pituitary portal vessel-hypothalamic-link is not fully developed in the rat until Day 5 of life (Glydon, 1957), another indication that fetal brain hormone is not of fetal pituitary origin.

The gradual rise of levels of growth hormone and TSH in the fetal serum and their fall after birth is thought by some authors to be due to this immaturity of the pituitary-hypothalamic regulatory system. Somatostatin, for example, has failed to be detected before the fifth day of post-uterine life. The significant fall in serum levels of pituitary hormones after birth has also been postulated to be mediated by adrenergic mechanisms (Basset et al, 1970), as infusion of catecholamine and isoprenaline into fetal lambs was shown to decrease serum GH concentrations.

The dramatic rise in fetal brain levels of TSH and GH does not, however, reflect the same pattern displaced by serum and amniotic fluid and, as previously concluded with reference to pituitary levels of these hormones, the CNS concentrations are probably independent of such extra CNS pools of hormone. Any speculation on the role of this precipitous rise in fetal brain levels of immunoassayable GH and TSH should be discussed in the light of fetal morphology and physiology.

During the transition from intrauterine to extrauterine life, a number of metabolic changes and adaptations occur in fuel homeostasis. These adaptations allow the fetus to convert from a state in which it is totally dependent on maternal fuel sources, to a state in which it relies

on endogenous substrate for the maintenance of normal cellular growth. In order to maintain normal cellular metabolism and anabolic growth, the newborn must have at least three requirements: an adequate store of hepatic and brain glycogen, muscle protein and fat, the enzymatic capacity to release and utilize these substrates, and thirdly, to be able to regulate such enzyme induction, substrate mobilization, and peripheral utilization of metabolic fuels.

Unlike glycogen in the liver, the functional significance of glycogen in the brain is poorly understood, although its potential as an energy source may explain the greater resistance of young animals to hypoxia. Glycogen levels in the brain are shown by Kohle et al (1977) to rise abruptly at day 20-21 of gestation and then to decrease rapidly after birth until, 7 days postnatally, very low levels are observed. An increase in glycogen-synthesizing enzymes has also been observed to follow a similar developmental trend. Although glycogen breakdown appears to commence the day of birth, 26% of the total fetal brain glycogen store is utilized within 10 minutes of birth. This glycogenolysis is in response to accelerated cerebral glycolysis resulting from the cerebral hypoxia of vaginal delivery.

A role for fetal brain growth hormone in such adaptations to neonatal life could possibly be one in which the fetal hormone exhibits "insulin-like" properties by increasing the permeability of cells in order to feed the rapid rise in glycogenolysis. Growth-promoting hormones, in particular GH and  $T_4$ , have been shown by Levi-Montalcini et al (1968) to have complementary and synergistic effects on cell growth and the regulation of



somatic development. In the developing CNS, many more neurons are produced in fetal life than survive; in fact, cell death of up to 50% of original neurons before birth is a common feature. McClusky et al (1979) have speculated that the rapid increase in estrogen receptors and estrogen levels, reported two days prior to parturition, and their rapid disappearance after birth, could be linked to a role for estrogen in preventing cell death and stabilizing surviving neurons at this critical period of development. Perhaps a similar argument could be made for the preparturitional rise in brain growth hormone. Furthermore, it is thought that nerve cells die when they fail to contact other cells during brain development, and it is possible that growth hormone and perhaps TSH may increase nerve cell contacts by causing enlargement of cells and cell-to-cell surface contacts.

If indeed these brain hormones possess the same effects on cell metabolic activities as do their pituitary counterparts, other hormone-promoted activities, such as increased protein synthesis and m.RNA content for GH, increased mitochondrial respiration, and the increased uptake of precursors of phospholipids for possible cell wall synthesis by TSH, could perhaps contribute to the general maintenance of neuronal tissues in fetal and adult tissues.

However, in any effort to assess the functional significance of pituitary-like hormones such as GH and TSH in the brain, the possibility that the role of the hormones within the CNS differs from their currently assigned physiological role within the pituitary, should not be overlooked. For example, it is known that TRH-like activity is present in the neural

tissue of certain invertebrates, even though evidence of a role for it in the regulation of pituitary TSH has only been found in animals higher in the evolutionary scale (Scharrer, 1978).

Another line of approach to the question of function of neuronal peptides has been the concept that "peptidergic neurons" refer not only to the neurosecretory cells of the hypothalamus producing vasopressin and hypophysiotropic hormones, but also to those neurons containing additional peptides such as Substance P, neurotensin, angiotensinogen, gastrointestinal peptides, and now more recently the hormones of the anterior pituitary such as ACTH, GH, TSH, LH, and  $\beta$ -lipotrophin. There is the possibility that, in the case of hypothalamic ACTH, GH, and TSH, extractable levels of hormone could be derived from two different sources. Those of CNS origin could be part of a peptidergic neural system present in addition to the classically described motor, sensory, and autonomic nervous systems, but which interrelates with such systems. Such peptides are speculated to be secreted at peptidergic terminals, and may act on a variety of effector cells, including neurons, functioning either as neurotransmitters or neuromodulators, rather than within the normal definition of a "hormone." It has been hypothesized by Barker (1977) that whereas neurotransmitters are involved in the momentary mediation of single cell-to-cell interaction, the neurohormones may be important in the sustained modulation of specific sets of target neurons. Neuromodulation implies an action of the secreted peptide in altering the effect of the classical neurotransmitter on its receptors. Possible examples of such interactions have been shown by the reported presence by Hökfelt et al (1979) of

neurons containing both peptides and classic monoaminergic neurotransmitters, and by the effects which other peptides, such as ACTH and vasopressin, have on acetylcholine, norepinephrine, and serotonin content or turnover in various brain regions (Gispen and de Wied et al, 1977). De Wied has also shown that these peptides, vasopressin and ACTH, can enhance the dephosphorylation of membrane proteins leading to a change in membrane permeability.

The reports of opioid-peptide effects on pain perception, addictive states, and psychiatric disorders have previously been reviewed, as have those reports concerning the effects which ACTH and MSH, given intravenicularly, appear to have on rat behavior. Behavioral studies suggesting a role for vasopressin in learning and memory (De Wied and Versteeg, 1979) showed that impaired learning behavior in Brattleboro rats, which lack vasopressin-synthesizing mechanisms, can be reversed by intraventricular injections of vasopressin. In normal rats, peripheral vasopressin injections appear to influence learning and memory, and in clinical trials, vasopressin may improve memory in brain-damaged human subjects. Studies with TRH demonstrate that this hypothalamic hormone can elicit behavioral excitation and anorexia in animals and may cause mood enhancement in humans (A. J. Prange et al, 1975).

In addition to such postulated behavioral effects assigned to peptide hormones in the brain, such hormones may also have an effect on endocrine regulatory processes. Experiments with intrahypothalamically injected ACTH have been reported to demonstrate inhibitory effects on pituitary ACTH release, although unphysiological doses and long periods

of treatment tend to dispute this data of Yates et al (1974). However, the possibility of such short-loop feedback mechanisms has also been alluded to in this discussion as a possible role for hypothalamic GH. It remains to be determined if such observations reflect a dual effect of these compounds on releasing factors or on the neurotransmitters that regulate such releasing factors.

The finding of GH- and TSH-like hormones in both the CNS and in the gastrointestinal tract in this study is not terribly surprising, considering the number of other peptides recently reported in such dual locations. Such peptides as somatostatin (Arimura, 1975), glucagon (Conlon et al, 1979), and VIP (Said et al, 1972) are found in the GI tract and, from this study, immunoreactive GH- and TSH-like material in extrapituitary sites such as the pancreas and small intestine. These data give further credence to one of the objectives of this dissertation, i.e., that of determining whether "brain" hormones are of CNS origin rather than the result of retrograde transport via the portal vessels from the pituitary to hypothalamus.

The finding of immunologically active, pituitary-like hormones in such extrapituitary, extra-CNS sites as the GI tract and pineal gland, leads to speculation that these peptide-containing cells can be classified as belonging to the amine-precursor uptake and decarboxylation series of cells, a concept originally developed and outlined by Anthony Pearse in 1977.

Other lines of approach to explain this apparent multisite distribution of pituitary-like peptides would be that the presence of such peptides is due primarily to incomplete gene suppression in those particular tissues, or that the detectable levels merely represent bound, circulating, pituitary peptide to tissues with appropriate hormone receptors. Removal of the pituitary as a source of circulating hormone for up to 28 days did not appreciably lower the levels of extractable hormone in those extrapituitary sites studied. The possibility that the observed immunoreactive material was merely receptor-bound is therefore unlikely in the light of the in vivo hypophysectomy experiments reported here.

In conclusion, results obtained in this study appear to add yet two more peptides, formerly thought to have been of non-neural origin, to the ever-growing list of brain peptides. Many of these, such as Substance P, have been speculated to be potential neurotransmitter or neuromodulator candidates, and represent an active field of research at the present time seeking to unravel the parameters involved in the development and functioning of the CNS.

Brain growth hormone and TSH await further investigation before specific functions can be assigned to their presence in rodent and primate brain. However, their apparent independence from patterns of pituitary hormone secretion, and their probable synthesis within the CNS tissues are tantalizing observations that await further conceptual organization.

## CONCLUSIONS

The following conclusions can be drawn from the data obtained in this dissertation.

1. Immunoreactive growth hormone (GH)- and thyroid stimulating hormone (TSH)-like peptides were found to be unevenly distributed throughout the rodent and primate central nervous system (CNS) and gastrointestinal tract.
2. Highest levels of extractable GH-like material were obtained from rodent amygdaloid tissue, whereas the hypothalamus contained the largest concentration of a TSH-like peptide.
3. Both brain peptides showed a similarity to their pituitary counterparts in terms of molecular weight, immunological affinity, and biological activity.
4. Histochemical mapping of the CNS for the TSH-like peptide using a peroxidase anti-peroxidase technique showed hormonal material concentrated in neuronal cell bodies of the rodent amygdaloid nucleus and hypothalamus.
5. Ultra-centrifuged homogenates of amygdaloid and thalamic tissues showed an association between GH- and TSH-like immunoreactivity and the synaptosomal-mitochondrial pellet (SMP). In ultra-centrifuged hypothalamic tissue, the immunoreactivity was distributed evenly between the SMP and the final supernatant.

5. Tissue cultured CNS cells from intact and hypophysectomized animals released both GH- and TSH-like material into the growth medium over a 30-day period. This would indicate the CNS, rather than the pituitary, as the source of both brain hormones.
6. Contamination of CNS tissues by pituitary GH and TSH was shown to be unlikely by a demonstration of blood-brain barrier intactness throughout life, and a lack of change in brain hormone levels after hypophysectomy.
7. Target organ ablation studies showed an apparent independence between regulatory mechanisms controlling pituitary and brain hormone levels.
8. The appearance of detectable hormone in the fetus and the subsequent pattern of development differed between brain and pituitary hormone pools. A preparturitional rise in brain hormone levels was detected, followed by a fall to normal levels in the CNS after birth.
9. In conclusion, this study verified the presence in the rodent and primate brain of GH- and TSH-like materials which appear to be of CNS rather than of pituitary origin.

## REFERENCES

- Adiga, P.R., Murthy, P.V.N. and McKenzie, S.M. (1971). Stimulation by thyrotropin, LATS and dibutyryl 3',5' AMP of protein and RNA synthesis and RNA polymerase activity in porcine thyroid in vitro. Biochem. 10: 702-710.
- Ahn, C.S. and Rosenberg, I.N. (1970). Iodine metabolism in thyroid slices: Effects of TSH, dibutyryl cyclic 3',5'-AMP, NaF and prostaglandin E<sub>1</sub>. Endocrinol. 86: 396-405.
- Allen, J.P., Kendall, J.W., McGilvra, R. and Vancera, C. (1974). Immunoreactive ACTH in cerebrospinal fluid. J. Clin. Endoc. Metab. 38: 586-593.
- Akil, H., Mayer, D.J. and Liebskind, J.C. (1972). Comparison chez le rat entre l'analgésie induite par stimulation de la substance grise péri-aqueducal et l'analgésie morphinique. C.R. Acad. Sci. Ser. D. 274: 3603-3613.
- Andrés, A., Satrústegui, J. and Machado, A. (1980). Development of NADPH producing pathways in rat heart. Biochem. J. 186: 799-603.
- Arbilla, S. and Langer, S. (1978). Morphine and  $\beta$ -endorphin inhibit release of noradrenaline from cerebral cortex but not of dopamine from rat striatum. Nature (Lond.) 271: 559-560.
- Arimura, A., Sato, H. and Dupont, A. (1975). Somatostatin: Abundance of immunoreactive hormone in rat stomach and pancreas. Science 189: 1007-1009.
- Baker, B.L., Dermody, W.C. and Rieb, J.R. (1975). Distribution of gonadotropin releasing hormone in rat brain as observed with immunocytochemistry. Endocrinol. 97: 125-135.
- Baker, B.L. and Yen, Y.Y. (1976). The influence of hypophysectomy on stores of somatostatin in the rat hypothalamus and pituitary. Proc. Soc. Exp. Biol. and Med. 151: 599-602.
- Bakke, J.L. and Lawrence, N. (1967). Thyrotropin (TSH) in the rat median eminence. Neuroendocrinol. 2: 315-325.
- Bala, R.M., Ferguson, K.A. and Beck, J.C. (1970). Plasma biological and immunoreactive human growth hormone-like activity. Endocrinol. 87: 506-516.



- Barker, J.L. (1977). Physiological roles of peptides in the nervous system. In: Peptides in Neurobiology. H. Gainer (Ed.), pp. 295-322, Plenum Press, New York.
- Barnea, A., Oliver, G. and Porter, J.C. (1977). Subcellular localization of  $\alpha$ -melanocyte stimulating hormone in the rat hypothalamus. J. Neurochem. 29: 619-624.
- Barnea, A. and Cho, G. (1979). Co-sequestration of immunoreactive ACTH and  $\alpha$ MSH in hypothalamic and hypophysial granules. Endocrin. Soc. Ann. Meeting Abstract. 343.
- Basbaum, A.I. and Fields, H.L. (1978). Endogenous pain control mechanism: Review and hypothesis. Ann. Neurol. 4: 451-462.
- Basset, J.M., Thornburn, G.D. and Wallace, A.L. (1970). Plasma growth hormone concentrations in the foetal lamb. J. Endoc. 48: 251-263.
- Bayon, A., Shoemaker, W.S., Bloom, F.E., Mauss, A. and Guillemin, R. (1979). Perinatal development of the endorphin and enkephalin-containing systems in the rat brain. Brain Res. 179: 93-101.
- Baxter, C.R., Horvath, J.S., Duggen, G.C. and Tiller, D.J. (1980). Effect of age on specific angiotensin II binding sites in rat brain. Endocrinol. 106: 995-999.
- Beneviste, R. and Frohman, L.A. (1978). An immunological approach to the study of protein conformational heterogeneity: Its application to growth hormone. Endocrinol. 102: 198-209.
- Ben-Jonathan, N.C., Oliver, C., Weiner, H.J., Mical, J.S. and Porter, J.C. (1977). Dopamine secretion into hypophysial portal plasma of the rat during the estrous cycle and throughout pregnancy. Endocrinol. 100: 452-458.
- Bennet, J.B. and Snyder, S.H. (1976). Angiotensin II binding to mammalian brain membranes. J. Biol. Chem. 251: 7423-7436.
- Besson, J., Rotszteym, W., Laburthe, M., Epelbaum, J., Beaudet, A., Kordon, C. and Rosselini, G. (1979). VIP. Brain distribution, subcellular localization and effect of deafferentation of the hypothalamus in male rats. Brain Res. 165: 79-85.
- Bennett, G.W., Edwardson, J.A., Holland, D., Jeffcoate, S.L. and White, N. (1975). Release of immunoreactive luteinizing hormone, releasing hormone and thyroid-releasing hormone from hypothalamic synaptosomes. Nature (Lond.) 257: 323-325.

- Birge, C.A., Peake, G.T., Mariz, J.K. and Daughaday, W.H. (1967). Radioimmunoassayable growth hormone in the rat pituitary gland. Effects of age, sex and hormonal state. Endocrinol. 81: 195-204.
- Birnbaum, R.S. and Goodman, H.M. (1979). Comparison of several insulin-like effects of growth hormone. Horm. Metab. Res. 11: 136-142.
- Blackard, W.G. and Heidingsfelder, S.A. (1968). Adrenergic receptor control mechanism for growth hormone secretion. J. Clin. Invest. 47: 1402-1414.
- Blasig, J. and Herz, T. (1976). Tolerance and dependence induced by morphine-like pituitary peptides in rats. N.S. Arch. Physiol. 294: 297-300.
- Blasquez, E., Simon, F.A., Blasquez, M. and Foa, P.P. (1974). Changes in serum growth hormone levels from fetal to adult age in the rat. Proc. Soc. Exp. Biol. and Med. 147: 780-783.
- Bloom, F., Segal, D., Ling, N. and Guillemin, R. (1976). Endorphins: Profound behavioural effects in rats suggest new etiological factors in mental illness. Science 194: 630-632.
- Bloom, F., Battenberg, E., Rossier, J., Ling, N. and Guillemin, R. (1978). Neurons containing  $\beta$  endorphin in rat brain exist separately from those containing enkephalin. Immunocytochemical studies. Proc. Natl. Acad. Sci. U.S.A. 75: 1591-1595.
- Blundell, T.L., Bedarker, S., Rinderknecht, E. and Humbel, R.E. (1978). Insulin-like growth factor: A model for tertiary structure accounting for immunoreactivity and receptor binding. Proc. Natl. Acad. Sci. U.S.A. 75: 180-184.
- Boler, J., Enzmann, F., Folkers, K., Bowers, C.Y. and Schally, A.V. (1969). The identity of chemical and hormonal properties of the thyrotropin releasing hormone and pyroglutamyl-histidyl-proline amide. Biochem. Biophys. Res. Comm. 37: 705-717.
- Borghi, C., Nocasias, S., Giachetti, A. and Said, S.J. (1979). Adenylate cyclase of rat pituitary: Stimulation by V.I.P. F.E.B.S. Letters 108: 403-407.
- Boyd, A.E., Lebowitz, H.E. and Pfeiffer, J.G. (1970). Stimulation of human growth hormone secretion by L-DOPA. N. Engl. J. Med. 283: 1425-1426.

- Brazeau, P., Vale, W., Burgus, R., Ling, N., Butcher, M., Rivier, J. and Guillemin, R. (1973). Hypothalamic polypeptide that inhibits secretion of immunoreactive growth hormone. Science 179: 77-79.
- Brown, M. and Vale, W. (1975). Central nervous system effects of hypothalamic peptides. Endocrinol. 96: 1333-1336.
- Brown, W.A. and Williams, B.W. (1976). Methylphenidate increases serum growth hormone concentrations. J. Clin. Endocrinol. Metabol. 43: 937-939.
- Brown, M., Rivier, J. and Vale, W. (1977). Bombesin; potent effects on thermoregulation in the rat. Science 196: 998-1000.
- Bruni, J.F., Van Gught, D., Marshall, S. and Meits, J. (1977). Effects of naloxone, morphine and metenkephalin on serum prolactin, luteinizing hormone, FSH, TSH and growth hormone. Life Sci. 21: 461-466.
- Burgus, R., Dunn, F.T., Desiderio, D. and Guillemin, R. (1969). Structure moléculaire du facteur hypothalamique hypophysiotrope TRF d'origine ovine: mise en évidence par spectrométrie de masse. C.R. Acad. Sci. Paris 269: 1870-1877.
- Burgus, R., Dunn, T.F., Desiderio, D., Ward, D.N., Vale, W. and Guillemin, R. (1970). Characterization of ovine hypothalamic hypophysiotropic (TSH) releasing factor. Nature 226: 321-325.
- Burgus, R., Butcher, M., Ling, N., Monatas, M., Rivier, J., Tellows, R., Amoss, M., Blackwell, R., Vale, W. and Guillemin, R. (1971). Structure moléculaire du factor hypothalamique (LRF) d'origine ovine controlant la secretion de l'hormone gonadotrophe, hypophysaire de luteinisation (LH). C.R. Acad. Sci. Paris 273: 1611-1613.
- Burt, D.R. and Snyder, S.H. (1975). Thyrotropin-releasing hormone (TRH): Apparent receptor binding in rat brain membranes. Brain Res. 93: 309-328.
- Cahane, M. and Cahane, T. (1936). Sur certain modification des glands endocrine après une lésion diencephalique. Rev. Fr. Endocrinol. 14: 472-487.
- Carraway, R.E. and Leeman, S.E. (1973). The isolation of a new hypotensive peptide, neurotensin, from bovine hypothalami. J. Biol. Chem. 248: 6854-6861.

- Carraway, R.E., Demans, L.M. and Leeman, S.E. (1976). Hyperglycemic effect of neurotensin, a hypothalamic peptide. Endocrinol. 99: 1452-1462.
- Carter, J.N., Tyson, J.E., Tolis, G., Van Vliet, S., Faiman, C. and Friesen, H.G. (1977). Prolactin-secreting tumours and hypogonadism in 22 men. N. Engl. J. of Med. 299: 847-852.
- Catt, K., Niall, H.D. and Tregear, G.W. (1967). A solid disc radio-immunoassay for human growth hormone. J. Lab & Clin. Med. 70: 820-830.
- Chang, M.M., Leeman, S.E. and Niall, H.D. (1971). Amino acid sequence of substance P. Nature New Biol. 232: 86-87.
- Cheng, H. and Leblond, C.P. (1974). Origin, differentiation and renewal of the four main epithelial cell types in the mouse small intestine. V. Unitarian theory of the origin of the four epithelial cell types. Am. J. Anat. 141: 537-561.
- Cheung, A.L. and Goldstein, A. (1976). Failure of hypophysectomy to alter brain content of opioid peptides (endorphins). Life Sci. 19: 1005-1008.
- Christensen, C.W., Harston, C.T., Kastin, A.J., Kostrzewa, R.M. and Spirtes, M.A. (1976). Investigations on  $\alpha$ MSH and MIF. I. Effects on cyclic AMP levels in rat brains. Pharmacol. Biochem. Behav. 5: 117-120.
- Clemmons, D.R., Hintz, R.L., Underwood, L.E. and Van Wyk, J.J. (1974). Common mechanism of action of somatomedin and insulin on fat cells: Further evidence. Isr. J. Med. Sci. 10: 1254-1262.
- Conklin, P.M., Schundle, W.J. and Hull, S.F. (1973). Hypothalamic thyrotropin releasing factor. Neuroendocrinol. 11: 197-211.
- Conlon, J.M., Samson, W.K., Dobbs, R.E., Orci, L and Unger, R.H. (1979) Glucagon-like polypeptides in canine brain. Diabetes 28: 700-702.
- Cons, J.M., Umezu, M. and Timiras, P.S. (1975). Developmental patterns of pituitary and plasma TSH in the normal and hypothyroid female rat. Endocrinol. 97: 237-240.
- Cornblath, M., Parker, M.L., Reisner, S.H., Forbes, A.E. and Daughaday, W.H. (1965). Secretion and metabolism of growth hormone in premature and full term infants. J. Clin. Endoc. Metab. 25: 209-218.

- Cornell, J.S. and Pierce, J.G. (1973). The subunits of human pituitary thyroid stimulating hormone: Isolation, properties and composition. J. Biol. Chem. 248: 4327-4333.
- Cott, J.M., Breese, G.R., Cooper, B.R., Barlow, T.S. and Prange, A.J. (1976). Investigation into the mechanism of reduction of ethanol sleep by thyrotropin releasing hormone. J. Pharm. Exp. Ther. 196: 594-604.
- Courbier, P., Kerdethune, B., Picon, R. and Roffii, J. (1978). Changes in testicular weight and serum gonadotropin and testosterone levels before, during and after birth in the prenatal rat. Endocrinol. 103: 1985-1991.
- Cox, B.M., Opheim, K.E., Teschemacher, H. and Golstein, A. (1975). A peptide-like substance from pituitary that acts like morphine. 2. Purification and properties. Life Sci. 16: 1777-1782.
- Coyle, J.T. and Henry, D. (1973). Catecholamines in fetal and newborn rat brain. J. of Neurochem. 21: 61-67.
- Curbelo, H.M., Karliner, E.C. and Houssay, A.B. (1979). Effect of acute hypoxia on blood TSH levels. Horm. Metab. Res. 11: 155-157.
- Daughaday, W.H. and Kipnis, D.M. (1966). The growth-promoting and anti-insulin actions of somatotropin. Rec. Prog. Horm. Res. 22: 49-99.
- Daul, C.B., Heath, R.G. and Gary, R.E. (1975). Angiotensin forming enzyme in human brain. Neuropharm. 14: 75-80.
- Dawson, G., McLawhon, R. and Miller, R.J. (1979). Opiates and enkephalins inhibit synthesis of gangliosides and membrane glycoproteins in mouse neuroblastoma cell line N<sub>4</sub>TGI. Proc. Natl. Acad. Sci. 76: 605-609.
- Denman, P., Miller, L.H., Sandman, C., Schally, A.V. and Kastin, A.B. (1972). Electrophysiological correlates of MSH activity in the frog. J. Comp. Physiol. Psychol. 80: 59-65.
- DeWied, D. and Versteeg, D.H.G. (1979). Neurohypophyseal principles and memory. Fed. Proc. 38: 2348-2354.
- DeWied, D. (1980). Hormonal influences on motivation, learning, memory and psychosis. In: Neuroendocrinology. D. Krieger and J. C. Hughes (Eds.), pp. 194-204, Sinauer Assoc, Sunderland, Massachusetts.

- Dichter, M. (1975). Physiological properties of vertebrate nerve cells in tissue culture. In: Brain Mechanisms in Mental Retardation. N. Buchwald and M. Brazier (Eds.), pp. 59-67, Academic Press, New York & London.
- Dockray, G.J. (1980). Cholecystokinins in rat cerebral cortex: Identification, purification and characterization by immunochemical methods. Brain Res. 188: 155-165.
- Dohler, K.D. and Wuttke, W. (1975). Changes with age levels of serum gonadotropins, prolactin and gonadal steroids in prepubertal male and female rats. Endocrinol. 97: 898-907.
- Drouva, S.V., Epelbaum, J., Héry, M., Laplante, E. and Kordon, C. (1980). Ionic channels, opiates and the release of gonadotropin releasing hormone and somatostatin from perfused hypothalamus. Endoc. Soc. 62nd. Ann.Meeting, Abstract, #138, Washington.
- Dube, D., Lissitzky, J.C., Leclerc, R. and Pelletier, G. (1978). Localization of  $\alpha$  melanocyte stimulating hormone in the rat brain and pituitary. Endocrinol. 102: 1283-1291.
- Dubois, M. (1975). Immunoreactive somatostatin: Presence in discrete cells of the endocrine pancreas. Proc. Natl. Acad. Sci. U.S.A. 72: 1340-1343.
- Dumont, J.E. (1971 ). The action of thyrotropin on thyroid metabolism. Vitamins-Hormones 29: 287-412.
- Dumont, J.E. and Tondeur-Montenez, T. (1965). Action de l'hormone thyrotrope sur le métabolisme énergétique du tissu thyroïdien. III. Evaluation au moyen du <sup>14</sup>C glucose des voies du métabolisme du glucose, dans le tissu thyroïdien de chien. Biochem. Biophys. Acta 3: 258-272.
- Dunn, A.J. and Gispen, W.H. (1977). How ACTH acts on the brain. Biochem. Rev. 1: 15-23.
- Dussault, J.H. and Labrie, F. (1973). Development of the hypothalamic-pituitary thyroid axis in the neonatal rat. Endocrinol. 97: 1321-1324.
- Eipper, B.A. and Main, R.E. (1975). High molecular weight forms of adrenocorticotrophic hormones in a mouse pituitary cell line. Biochem. 14: 3836-3844.

- Eipper, B.A. and Mains, R.E. (1980). Structure and biosynthesis of pro adrenocorticotropin, endorphin and related peptides. Endocrin. Rev. 1: 1-27.
- Ekholm, R., Engström, G., Ericson, L.E. and Melander, A. (1975). Exocytosis of protein into the thyroid follicle lumen. An early effect of TSH. Endocrinol. 97: 337-346.
- Ekholm, R. (1979). Thyroid physiology: Anatomy and development. In: Endocrinology. L.J. Degroot (Ed.), pp. 307-311, Grune and Stratton,
- Elde, R., Hökfelt, T., Johansson, O. and Terenius, L. (1976). Immunohistochemical studies using antibodies to leucine-enkephalin: Initial observations on the nervous system of the rat. Neurosci. 1: 349-351.
- Elliot, T.R. (1904). Ileo-colic sphincter. J. Physiol. (Lond.) 31: 157-168.
- Ellis, S., Nuenke, J.M. and Grindeland, R.E. (1968). Identity between growth hormone degrading activity of pituitary gland and plasmin. Endocrinol. 83: 1029-1042.
- Emanuele, N., Connick, E., Hojvat, S., Baker, G., Kirsteins, L. and Lawrence, A. (1980). Brain LH: Distribution, characteristics and presence after hypophysectomy. Biol. of Reprod. (submitted for publication).
- Emson, P.E., Fahrenkrig, D., Schaffalitzley de Michadell, O.B., Jessell, T.M. and Iversen, L.L. (1978). Vasoactive intestinal peptide, vesicular localization and  $K^+$  evoked release from rat hypothalamus. Brain Res. 143: 174-178.
- Engler, D. and Jackson, I.M.D. (1980). Evidence that circulatory TRH in the neonatal rat is derived from the pancreas and gastrointestinal tract. Endoc. Soc. 62nd. Ann Meeting, Abstract, #126, Washington.
- Enzmann, F., Böler, J. and Folkers, K. (1971). Structure and synthesis of the thyrotropin releasing hormone. J. Med. Chem. 14: 409-472.
- Epelbaum, J., Brazeau, P., Tsang, D., Braver, J. and Martin, J.B. (1977). Subcellular distribution of radioimmunoassayable somatostatin in rat brain. Brain Res. 126: 309-323.

- Eskay, R.L., Giraud, P., Oliver, C. and Brownstein, M.J. (1979a). Distribution of  $\alpha$  melanocyte stimulating hormone in the rat brain. Evidence that  $\alpha$ MSH-containing cells in the arcuate nucleus region send projections to extrahypothalamic areas. Brain Res. 178: 55-67.
- Eskay, R.L., Brownstein, M.J. and Long, R.T. (1979b).  $\alpha$  Melanocyte stimulating hormone: Reduction in adult rat brain after monosodium glutamate treatment of neonates. Science 205: 827-828.
- Euler, V.S. and Pernow, B. (1979). Functions of peptides in the brain: Substance P. Fed. Proc. 38: 2304-2305.
- Evans, T.M., Simpson, M.E., Marx, W. and Kibrick, P.A. (1943). Bioassay of the pituitary growth hormone. Width of the proximal epiphyseal cartilage of the tibia in hypophysectomised rats. Endocrinol. 32: 13-16.
- Fain, I.N. and Saperstein, R. (1971). The involvement of RNA synthesis and cyclic AMP in the activation of fat cell lipolysis by growth hormone and glucocorticoids. In: Adipose Tissue Regulation and Metabolic Findings. pp. 20-22, Academic Press, New York.
- Farquhar, M.G. and Rinehart, J.F. (1954). Electron microscopic studies of the anterior pituitary gland of castrate rats. Endocrinol. 54: 516-541.
- Fischbach, G. (1972). Synapse formation between dissociated nerve and muscle cells in low density cultures. Devel. Biol. 28: 407-429.
- Fisher, G., Humphries, J. and Folkes, K. (1974). Synthesis and some biological activities of substance P. J. Med. Chem. 17: 843-846.
- Fitzsimons, J.T. (1972). Thirst. Physiol. Rev. 52: 468-561.
- Foreman, M.M. and Moss, R.L. (1977). Effects of subcutaneous injection and intrahypothalamic infusion of releasing hormones upon lordotic response to repetitive coital stimulation. Horm. Behav. 8: 219-234.
- Frantz, A.G. and Rabkin, M.T. (1964). Human growth hormone: Clinical measurement, response to hypoglycemia and suppression by corticosteroids. N. Engl. J. of Med. 271: 1375-1381.



- Friedman, D.J. and Zeman, F.J. (1979). Thyroid, pituitary and hypothalamic hormone levels in neonatal rat pups following maternal protein deficiency. Proc. of Soc. Exp. Biol. and Med. 161: 275-279.
- Frohman, L.A. and Bernadis, L.L. (1968). Growth hormone and insulin levels in weanling rats with ventromedial hypothalamic lesions. Endocrinol. 82: 1125-1130.
- Furnaletto, R.W., Underwood, L.E., Van Wyk, J.J. and D'erirole, A.J. (1977). Estimation of somatomedin-C levels in normals and patients with pituitary disease by radioimmunoassay. J. Clin. Invest. 60: 648-657.
- Fuxe, K., Ganther, D., Hökfelt, T. and Bolne, P. (1976). Immunohistochemical evidence for the existence of angiotensin II containing nerve terminals in brain and spinal cord of the rat. Neurosci. Letters 2: 229-236.
- Fuxe, K., Hökfelt, T., Eneroth, P., Gustafson, J.A. and Skett, P. (1977). Prolactin-like immunoreactivity: Localization in nerve terminals of rat hypothalamus. Science 196: 899-900.
- Ganten, D., Schally, P., Vecaci, P. and Ganten, U. (1976). Isorenin of extrarenal origin. The tissue angiotensinogenase system. Am. J. Med. 60: 760-772.
- Giachetti, A., Said, S.I., Reynolds, R.C. and Koniges, F.C. (1977). Vasoactive intestinal polypeptide in brain: Localization and release from isolated nerve terminals. Proc. Natl. Acad. Sci. U.S.A. 74: 3424-3428.
- Gibbs, J., Young, R.C. and Smith G.P. (1973). Cholecystokinin decreases food intake in rats. J. Comp. Physiol. Psychol. 84: 488-495.
- Gilbert, R.F.T. and Emson, P.C. (1979). Substance P in rat CNS and duodenum during development. Brain Res. 171: 166-170.
- Ginsberg, B.H., Kahn, C.R., Roth, J., Megyesi, K. and Baumm, G. (1979). Identification and high yield purification of insulin-like growth factors (nonsuppressible insulin-like activities and somatomedins) from human plasma by use of endogenous binding proteins. J. Clin. Endocrinol. Metab. 48: 43-49.

- Gispen, W.H., Weigant, V.M., Greven, H.M. and Deweid, D. (1976a). The induction of excessive grooming in the rat by intraventricular application of peptides derived from ACTH: Structure activity studies. Life Sci. 17: 645-652.
- Gispen, W.H. and Wiegant, V.M. (1976b). Opiate antagonists suppress ACTH 1-24 induced excessive grooming in the rat. Neurosci. Letters 2: 159-164.
- Gispen, W.H., Reith, M.E., Schotman, P., Wiegant, V.M., Zweiss, H. and Dewied, D. (1977). CNS and ACTH-like peptides: Neurochemical response and interaction with opiates. In: Neuropeptide Influence on the Brain and Behaviour. L.H. Miller, C.A. Sandman, and A.J. Kastin (Eds.), pp. 61-68, Raven Press, New York.
- Gluckman, P.D., Mueller, P.L., Kaplan, S.L., Rudolph, A.M. and Grumbach, M.M. (1979). Hormone ontogeny in the ovine fetus. I. Circulating growth hormone in mid and late gestation. Endocrinol. 104: 162-168.
- Glydon, R. St. J. (1957). The development of the blood supply of the pituitary in the albino rat, with special reference to the portal vessels. J. Anat. (Lond.) 91: 237-244.
- Goldsmith, P.C., Rose, J.C., Arimura, A. and Ganong, W.F. (1975). Ultrastructural localization of somatostatin in pancreatic islets of the rat. Endocrinol. 97: 1061-1064.
- Goodman, H.M. (1967). A comparative study of the effects of insulin and growth hormone on hexose metabolism in adipose tissue. Endocrinol. 80: 45-51.
- Goodman, H.M. (1968). Growth hormone and the metabolism of carbohydrate and lipid in adipose tissue. Ann. N.Y. Acad. Sci. 148: 419-440.
- Gorden, P., Hendricks, C.M. and Roth, J. (1973). Evidence for "big" and "little" components of human plasma and pituitary growth hormone. J. Clin. Endocrinol. Metabol. 36: 178-184.
- Govindrajan, D.R. and Clausen, J. (1974). Regional distribution of cathepsin B, and D in human brain. Brain Res. 67: 141-146.
- Gramsch, C., Kleber, G., Hout, V., Pasi, A., Mehracin, P. and Herz, A. (1980). Pro-opiocortin fragments in human and rat brain. Brain Res. 192: 109-119.

- Green, J.D. and Harris, G.W. (1949). Observation of the hypophysoportal vessels of the living rat. J. Physiol. 108: 359-361.
- Greenspan, F.S., Li, H.C., Simpson, M.E. and Evans, H.M. (1949). Bioassay of hypophysial growth hormone: The tibia test. Endocrinol. 45: 455-463.
- Greenwood, F.C. and Landon, J. (1966). Growth hormone secretion in response to stress in man. Nature 210: 540-541.
- Greer, M.A. (1951). Evidence of hypothalamic control of the pituitary release of thyrotropin. Proc. Soc. Exp. Biol. and Med. 77: 603-608.
- Greven, H.M. and Dewied, D. (1977). Influences of peptides structurally related to ACTH, and MSH on active avoidance behaviour in rats. In: Melanocyte Stimulating Hormone: Control, Chemistry and Effects. Frontiers in Horm. Res. F.J. Tilden, D.F. Swaab and Tj.B. Wimersma (Eds.), 4: 140-152, Karger, Basel.
- Grimm, Y. and Reichlin, S. (1973). Thyrotropin releasing hormone TRH: Neurotransmitter regulation of secretion by mouse hypothalamic tissue in vitro. J. Clin. Endocrinol. Metabol. 39: 406-409.
- Grumbach, M.M., Roth, J.C., Kaplan, S.L. and Kelch, R.P. (1974). Growth hormone; maternal-fetal relationship. In: Control of Onset of Puberty. G.D. Grave and F.E. Mayer (Eds.), pp. 115-120, Wiley, New York.
- Guerrero-Munoz, F., Guerrero, M. and Way Kong, E. (1979). Effect of  $\beta$ -endorphin on Ca uptake in brain. Science 206: 89-91.
- Guillemin, R., Lenz, N. and Burgus, R. (1976). Endorphins, peptides d'origine hypothalamique et neurohypophysaire á activit  morphinomimetique. Isolement et structure molculaire d'  $\alpha$  endorphine. C.R. Acad. Sci. Ser. D. 282: 783-785.
- Guillemin, R., Vargo, T., Rossier, J., Menick, S., Lung, N., Rivier, C., Vale, W. and Bloom, F. (1977).  $\beta$ -Endorphin and adrenocorticotropin are secreted concomitantly by the pituitary gland. Science 197: 1367-1369.
- Guillemin, R. (1980). Beta lipotropin and endorphins: Implications of current knowledge. In: Neuroendocrinology. D.T. Krieger and J. C. Hughes (Eds.), pp. 96-104, Sinauer Assoc. Inc., Massachusetts.

- Gunne, L.H., Lindstrom, L. and Terenius, L. (1977). Naloxone-induced reversal of schizophrenic hallucinations. J. Neur. Trans. 40: 13-22.
- Hagen, T.C., Lawrence, A.M. and Kirsteins, L. (1972). Autoregulation of growth hormone. Secretion in normal subjects. Metabol. 21: 603-610.
- Hambley, J. and Grant, D.B. (1972). Apparent heterogeneity of serum chorionic somatomammotrophin on gel filtration. Acta Endoc. 70: 43-47.
- Hansen, A.P. (1971). The effect of adrenergic receptor blockade on the exercise induced serum growth hormone rise in normals and juvenile diabetics. J. Clin. Endocrinol. Metabol. 33: 807-812.
- Haulica, I., Branisteanu, D.D., Rosca, V., Stratone, A., Bebelev, V., Balan, G. and Ionescu, L. (1975). A renin-like activity in pineal gland and hypophysis. Endocrinol. 96: 508-510.
- Havrankova, J., Brownstein, M.J. and Roth, J. (1979). Concentrations of insulin and of insulin receptors in the brain are independent of peripheral insulin levels. J. Clin. Invest. 64: 636-642.
- Haymond, M.W. and Pagliara, A.S. (1979). Endocrine and metabolic aspects of fuel homeostasis in the fetus and neonate. In: Endocrinology. D. Degroot (Ed.), pp. 1779-1804, Grune and Stratton, New York.
- Hefco, E., Krulich, L. and Aschenbrenner, J.E. (1975a). Effect of hypothalamic deafferentation on the secretion of thyrotropin in resting conditions in the rat. Endocrinol. 97: 1226-1233.
- Hefco, E., Krulich, L. and Aschenbrenner, J.E. (1975b). Effect of hypothalamic deafferentation on the secretion of thyrotropin during thyroid blockade and exposure to cold in the rat. Endocrinol. 97: 1234-1240.
- Heidingsfelder, S.A. and Blackard, W.G. (1968). Adrenergic control mechanism for vasopressin-induced plasma growth hormone response. Metabol. 17: 1019-1024.
- Hintz, R.L., Suskind, R., Amtayakul, K., Thanangkul, O. and Olson, R. (1978). Plasma somatomedin and growth hormone values in children with protein-calorie malnutrition. J. Pediatr. 92: 153-156.

- Hirose, S., Yokosawa, H., Inagami, T. and Workman, R. (1980). Renin and prorenin in hog brain: Ubiquitous distribution and high concentration in the pituitary and pineal. Brain Res. 191: 489-499.
- Hoffman, D.L. and Baker, B.L. (1977). Effect of treatment with growth hormone on somatostatin in the median eminence of the rat. Proc. Soc. Exp. Biol. and Med. 156: 265-271.
- Hartree, A.S., Thomas, M., Furnival, B., Burns, T.W. and Langley, P. (1972). Thyroid stimulating and lipolytic activities of purified preparations of human T.S.H. J. Endoc. 53: 95-100.
- Hökfelt, T., Fuxe, K., Johansson, O., Jeffcoate, S. and White, N. (1975-a). Thyrotropin releasing hormone (TRH)-containing nerve terminal in certain brain stem nuclei and in the spinal cord. Neurosci. Letters 1: 133-139.
- Hökfelt, T., Kellerth, J.O., Nilsson, G. and Pernow, B. (1975b). Experimental immunohistochemical studies on the localization and distribution of substance P in cat primary sensory neurons. Brain Res. 100: 235-252.
- Hökfelt, T., Elfvin, L.G., Elde, R., Schultzberg, M., Goldstein, M. and Luft, R. (1977a). Occurrence of somatostatin immunoreactivity in some peripheral sympathetic noradrenergic neurons. Proc. Natl. Acad. Sci. U.S.A. 74: 3587-3591.
- Hökfelt, T., Elfvin, L.G., Schultzberg, M., Goldstein, M. and Nilsson, G. (1977b). On the occurrence of substance P containing fibres in sympathetic ganglia: Immunohistochemical evidence. Brain Res. 132: 29-41.
- Hökfelt, T. and Elde, R. (1979). Localization of hypophysiotropic peptides and other biologically active peptides within the brain. Ann.Rev. Physiol. 41: 587-602.
- Hughes, J., Smith, T.W., Kosterlitz, H.W., Fothergill, M.A., Morgan, B. A. and Morris, H.P. (1975a). Identification of 2 related pentapeptides from the brain with potent opiate agonist activity. Nature 258: 577-579.
- Hughes, J., Smith, T., Morgan, B. and Fothergill, L. (1975b). Purification and properties of enkephalin. The possible endogenous ligand for the morphine receptor. Life Sci. 16: 1753-1758.
- Hughes, J. (1979). Opioid peptides and their relatives. Nature (Lond.) 278: 394-395.

- Hunter, W.M. and Greenwood, F.C. (1962). Preparation of iodine-131, labeled human growth hormone of high specific activity. Nature (Lond.) 194: 495-496.
- Inagami, T., Yokosawa, H. and Hirose, S. (1978). Definitive evidence for renin in rat brain by affinity chromatographic separation from protease. Clin. Sci. Molec. Med. 55: 121S-123S.
- Innis, R.B., Correa, F.M.A., Uhl, G.R., Schneider, B. and Snyder, S.H. (1979). Cholecystokinin octapeptide-like immunoreactivity. Histochemical localization in rat brain. Proc. Natl. Acad. Sci. U.S.A. 76: 521-525.
- Jackson, I.M.D. and Reichlin, S. (1977). Brain thyrotropin releasing hormone is independent of the hypothalamus. Nature (Lond.) 267: 853-854.
- Jacob, J.J., Tremblay, E.C. and Colombel, M.C. (1974). Facilitation de reactions nociceptives par la naloxone chez le souris et chez le rat. Psychopharmacologie 37: 217-223.
- Jacobowitz, D.M. and O'Donahue, T.L. (1978).  $\alpha$  Melanocyte stimulating hormone: Immunohistochemical identification and mapping in neurons of rat brain. Proc. Natl. Acad. Sci. (Wash.) 75: 6300-6304.
- Jacquot, R. and Kretchmer, N. (1964). Effect of fetal decapitation on enzymes of glycogen metabolism. J. Biol. Chem. 239: 1301-1305.
- Johnson, A.K. and Epstein, L. (1975). The role of cerebral ventricular system in angiotensin induced thirst. In: Control Mechanisms of Drinking. Peters, Fitzimons and Peters-Haejeli (Eds.), pp. 117-122, Springer-Verlag, Heidelberg.
- Jost, A. (1959). The role of fetal hormones in prenatal development. The Harvey Lectures, series 55, Academic Press.
- Jonsson, G., Fuxe, K. and Hökfelt, T. (1972). On the catecholamine innervation of the hypothalamus, with special reference to the median eminence. Brain Res. 40: 271-281.
- Kansal, P.C., Buse, J. and Talbert, O.R. (1972). The effect of L-dopa on plasma growth hormone, insulin and thyroxine. J. Clin. Endocrinol. Metabol. 34: 99-105.
- Kapitulnik, J. and Tshershedsky, M. (1979). Fluidity of rat liver microsomal membrane: Increase at birth. Science 206: 843-844.

- Karnatsuka, A., Makimo, H., Matsushima, Y., Osegawa, M., Yamamoto, M. and Kunagai, A. (1979). Effect of hypophysectomy and growth hormone administration on somatostatin content in rat hypothalamus. Neuroendocrinol. 29: 186-190.
- Kato, Y., Dupré, J. and Beck, J.C. (1973). Plasma growth hormone in the anesthetized rat: Effects of dibutyryl cAMP, prostaglandins E, adrenergic agents, vasopressin, chlorpromazine, amphetamine and L-Dopa. Endocrinol. 93: 135-146.
- Kaufmann, U., Zapf, J., Torretti, B. and Froesch, E.R. (1977). Demonstration of a specific serum carrier protein of non-suppressible insulin-like activity in vivo. J. Clin. Endocrinol. Metabol. 44: 160-166.
- Kelly, P.A., Tsushima, T., Shuiu, R.P.C. and Frisson, H.G. (1976). Lactogenic and growth hormone-like activities in pregnancy determined by radioreceptor assay. Endocrinol. 99: 765-774.
- Kibrick, E.A., Becks, H., Marx, W. and Evans, H.M. (1941). Effect of different dose levels of growth hormone on the tibia of young hypophysectomised female rats. Growth 5: 437-447.
- Kieffer, J.D., Mover, H., Panfilo, F. and Maloof, F. (1976). Pituitary-thyroid axis in neonatal and adult rats. Comparison of the sexes. Endocrinol. 98: 295-304.
- Kleber, G., Gramsch, C., Höelt, V., Mehraein, P., Pasi, A. and Herz, A. (1980). Extrahypothalamic corticotropin and  $\alpha$  melanotropin in human brain. Neuroendocrinol. 31: 39-45.
- Klee, W.A. and Nirenberg, M. (1976). Mode of action of endogenous opiate peptides. Nature (Lond.) 263: 609-611.
- Knigge, K.M. (1964). Neural regulation of T.S.H. secretion: Effect of diencephalic and intracerebral injections of thyroxine and thyrotropin upon thyroid activity in the cat. In: Major Problems in Neuroendocrinology. E. Bujusz and G. Jasmin (Eds.), pp. 261-265, Basel, Karger.
- Kobayashi, R.M., Brown, M. and Vale, W. (1977). Regional distribution of neurotensin and somatostatin in rat brain. Brain Res. 126: 584-588.
- Kobayashi, R.M., Palkovits, M., Miller, R.J., Chang, R.J. and Cuatrecasas, P. (1978). Brain enkephalin distribution: Unaltered by hypophysectomy. Life Sci. 22: 527-530.

- Kohle, S.J. and Vanucci, R.C. (1977). Glycogen metabolism in fetal and postnatal rat brain: Influence of birth. J. Neurochem. 28: 441-443.
- Korner, A. (1965). Growth hormone control of protein and ribonucleic acid. Rec. Prog. Horm. Res. 21: 205-240.
- Kostyo, C. and Knobil, E. (1966). The pituitary growth hormone: An adventure in physiology. Physiologist (Bethesda) 9: 25-29.
- Kovács, G.L., Bohus, B. and Versteeg, D.H. (1979a). The effects of vasopressin on memory processes. The role of noradrenergic neurotransmission. Neurosci. 4: 1529-1537.
- Kovács, G.L., Bohus, B., Versteeg, D.H., De Kloet, R.E., DeWied, D. (1979b). Effect of oxytocin and vasopressin on memory. Consolidation: Sites of action and catecholaminergic correlates after local microinjection into limbic-midbrain structures. Brain Res. 175: 303-314.
- Krieger, D.T., Liotta, A. and Brownstein, M.J. (1977a). Presence of corticotropin in the limbic system of normal and hypophysectomised rats. Brain Res. 128: 575-579.
- Krieger, D.T., Liotta, A., Suda, T., Palkovits, M. and Brownstein, M.J. (1977b). Presence of immunoassayable  $\beta$ -lipotropin in bovine brain and spinal cord. Lack of concordance with ACTH concentration. Biochem. Biophys. Res. Comm. 76: 930-936.
- Krieger, D.T. and Liotta, A.S. (1979). Pituitary hormones in brain: Where, how and why? Science 205: 366-372.
- Krieger, D.T. and Liotta, A.S. (1980). Rat anterior pituitary ACTH,  $\beta$ -lipotropin and  $\beta$ -endorphin. Parallel synthesis and release in equimolar amounts. Endoc. Soc. 62nd. Ann. Meeting, Abstract #309, Washington.
- Krivoy, W. and Guillemin, R. (1961). On a possible role of MSH in the spinal cord of the cat. Endocrinol. 69: 170-175.
- Krnejević, K. and Letić, D. (1977). Substance P selectively blocks excitation of Renshaw cells by acetylcholine. Canad. J. Physiol. Pharmacol. 55: 958-961.
- Krulich, L., Giachetti, A., Marchleiska-Koj, A., Hefco, E. and Jameson, H.E. (1977). On the role of the central noradrenergic and dopaminergic systems in the regulation of TSH. Secretion in the rat. Endocrinol. 100: 496-505.



- Kubek, M.J., Wilber, J.F. and Leesthema, J.E. (1979). The identification of gonadotropin releasing hormone (G.N.R.H.) in hypothalamic and extrahypothalamic loci of the human nervous system. Horm. Metab. Res. 11: 26-29.
- Kuhar, M.J., Pert, C.B. and Snyder, S.H. (1973). Regional distribution of opiate receptor binding in monkey and human brain. Nature (Lond.) 245: 447-450.
- Larry, F., Willens, C., Lecocq, R., Delcroix, C. and Dumont, J.E. (1971). Stimulation by thyrotropin in vitro of uridine incorporation into the RNA of thyroid slices. Horm. Metab. Res. 3: 414-422.
- Larsson, L.I. and Rehfeld, J.F. (1979a). Localization and molecular heterogeneity of cholecystokinin in the central and peripheral nervous system. Brain Res. 165: 201-218.
- Larsson, L.I., Childen, S. and Snyder, S.H. (1979b). Met- and leu-enkephalin immunoreactivity in separate neurons. Nature (Lond.) 282: 407-410.
- Lawrence, A.M., Barbato, T. and KIRSTEINS, K. (1974). Cerebral spinal fluid stimulation of pituitary protein synthesis and growth hormone release in vitro. Lancet 1: 599-600.
- Lecocq, R.E. and Dumont, J.E. (1972). Stimulation by thyrotropin of amino acid incorporation into proteins in dog thyroid slices in vitro. Biochem. Biophys. Acta 281: 434-441.
- Lappäluoto, J., Ranta, J. and Tuomisto, J. (1974). Diurnal variation of serum immunoassayable thyrotropin (T.S.H.) concentration in rat. Acta Physiol. Scand. 90: 699-702.
- Leeman, S.E., Mroz, E.A. and Carraway, R.E. (1977). Substance P and neurotensin. In: Peptides in Neurobiology. H. Gainer (Ed.), p. 99, Plenum Press, New York.
- Leonard, B.E., Kafoe, W.F., Thody, A.J. and Shuster, S. (1976). The effect of  $\alpha$  melanocyte stimulating hormone on the metabolism of biogenic amines in the rat brain. J. Neurosci. Res. 2: 39-45.
- Levi-Montalcini, R. and Angeletti, P.V. (1968). Nerve growth factor. Physiol. Rev. 48: 534-569.
- Li, C.H., Evans, H.E. and Simpson, M.E. (1945). Isolation and properties of the anterior hypophyseal growth hormone. J. Biol. Chem. 159: 353-366.
- Li, C.H. and Evans, H.M. (1948). The Hormones. G. Pinus and K. Thimann (Eds.), 1: 631, Academic Press, New York.

- Li, C.H. and Dixon, J.S. (1971). Human growth hormone XXXII. The primary structure of the hormone: Revision. Arch. Biochem. Biophys. 146: 233-236.
- Li, C.H. and Graf, L. (1974). Human pituitary growth hormone. Isolation and properties of two biologically active fragments from plasma digests. Proc. Natl. Acad. Sci. U.S.A. 71: 1197-1201.
- Li, C.H., Yamashiro, D., Tseng, L.F. and Loh, H.H. (1977). Synthesis and analgesic activity of human  $\beta$ -endorphins. J. Med. Chem. 20: 325-328.
- Liggins, G.C., Fairclough, R.J., Grieves, S.A., Kendall, J.Z. and Knox, B.S. (1973). The mechanism of initiation of parturition in the ewe. Rec. Prog. Horm. Res. 29: 111-150.
- Liotta, A.S., Hauser, H., Brownstein, M.J. and Krieger, D.T. (1978). Effect of long term hypophysectomy or adrenalectomy on rat brain ACTH. Fed. Proc. 37: p. 665.
- Liotta, A., Gildersleeve, D., Brownstein, M.J. and Krieger, D.T. (1979a). Biosynthesis in-vitro of immunoreactive 31,000 dalton corticotropin/ $\beta$  endorphin-like material by bovine hypothalamus. Proc. Natl. Acad. Sci. U.S.A. 76: 1448-1452.
- Liotta, A. and Krieger, D.T. (1979b). Preliminary characterization of in-vitro synthesised hypothalamic precursor ACTH/ $\beta$ -endorphin-like material. Mol. Cell. Endocrinol. 16: 221-228.
- Luft, R., Efendic, S., Hökfelt, T., Johansson, O. and Arimura, A. (1974). Immunohistochemical evidence for the localization of somatostatin-like immunoreactivity in a cell population of the pancreatic islet. Med. Biol. 52: 428-430.
- Maclusky, N.J., Lieberburg, J. and McEwen, B.S. (1979). The development of estrogen receptor systems in the rat brain: Perinatal development. Brain Res. 178: 129-142.
- Mains, R.E., Eipper, B.A. and Ling, N. (1977). Common precursor to corticotropins and endorphins. Proc. Natl. Acad. Sci. U.S.A. 74: 3014-3018.
- Malesci, A., Strauss, E. and Yalow, R.S. (1980). Cholecystokinin-converting enzymes in brain. Proc. Natl. Acad. Sci. U.S.A. 77: 597-599.
- Mann, H.B. and Whitney, D.R. (1947). On a test of whether one of two variables is stochastically larger than the other. Ann. Math. Statist. 18: 52-54.

- Martin, J.B. (1973). Neural regulation of growth hormone secretion. New. Eng. J. Med. 288: 1384-1393.
- Martin, J.B., Renaud, L.P. and Brazeau, P. (1975). Hypothalamic peptides: New evidence for "peptidergic" pathways in the CNS. Lancet 2: 393-395.
- Martin, J.M. and Gagliardino, J.J. (1967). Effect of growth hormone on the isolated pancreatic islets of the rat in vitro. Nature (Lond.) 213: 630-631.
- Martin, R., Weber, E. and Voight, K.H. (1979). Localisation of corticotropin and endorphin related peptides in the intermediary lobe of the rat pituitary. Cell Tiss. Res. 196: 307-319.
- Martin, T.E. and Young, F.G. (1965). An in-vitro action of human growth hormone in the presence of actinomycin D. Nature (Lond.) 208: 684-685.
- Marx, J.L. (1979). Brain peptides: Is substance P a transmitter of pain signals? Science 205: 886-889.
- Matsuo, M., Arimura, A. and Nair, R. (1971a). Synthesis of the porcine luteinising hormone and follicle stimulating hormone releasing hormone by the solid phase method. Bioch. Biophys. Res. Comm. 45: 822-831.
- Matsuo, H., Baba, Y., Nair, R.M.G., Arimura, A. and Schally, A.V. (1971b). Structure of the porcine, LH, and FSH releasing hormone. I. The proposed amino acid sequence. Biochem. Biophys. Res. Comm. 43: 1334-1339.
- May, P., Mittler, J., Manougian, A. and Erbel, N. (1979). Thyroid stimulating hormone release inhibitory activity of leucine enkephalin. Horm. Metab. Res. 11: 30-33.
- McCaleb, M.L., Myers, R.D., Singer, S. and Willis, G. (1979). Hypothalamic norepinephrine in the rat during feeding and push pull perfusion with glucose, 2-DG or insulin. Am. J. Physiol. 236: 312-321.
- Merimee, T.J. (1979). Growth hormone: Secretion and action. In: Endocrinology. L.J. Degroot (Ed.), 1: 123-132, Grune and Stratton, New York.
- Metcalf, G. (1974). Thyroid releasing hormone: A possible mediator of thermoregulation. Nature 252: 310-311.

- Mezey, E., Kivovics, P. and Palkovits, M. (1979). Pituitary-brain retrograde transport. T.I.N.S. March, 57-60.
- Moi-Yoi, O., Pinkus, G.S., Spragg, J. and Austen, K.F. (1979). Identification of human glandular kallikrein in the  $\beta$  cell of the pancreas. New Eng. J. Med. 300: 1289-1294.
- Moldow, R. and Yalow, R.S. (1978a). Extrahypophyseal distribution of corticotropin as a function of brain size. Proc. Natl. Acad. Sci. U.S.A. 75: 994-998.
- Moldow, R.L. and Yalow, R.S. (1978b). Extrahypophyseal distribution of thyrotropin as a function of brain size. Life Sci. 22: 1859-1864.
- Montoya, E., Seibel, M.J. and Wilber, J.F. (1975). TRH secretory physiology. Studies by RIA and affinity chromatography. Endocrinol. 96: 1416-1418.
- Moriarty, G.C. (1973). Adenohypophysis ultrastructural cytochemistry. A review. J. Histochem. Cytochem. 21: 855-893.
- Moriarty, G.C., Kupek, M., Cole, D., Tobin, R., Lorinez, M. and Wilber, J. (1978). The pituitary thyrotropin (TSH) secretory granules: A novel location for TRH in rats and man. Endoc. Soc. 60th. Ann. Meeting, Abstract #192.
- Morley, J.E., Levin, S.R., Pehlevanian, A., Adlachi, R., Pekary, A.E. and Hershman, J.M. (1978). Thyrotropin releasing hormone (TRH) and the pancreas. Clin. Res. 26: 160A.
- Morley, J.E., Larners, C., Sharp, B., Poitros, P., Walsh, J.H., Carlson, H.E. and Hershman, J.M. (1980). Evidence for serotonergic and opiate modulation of brain cholecystokinin. Clin. Res. 28: 265A.
- Moss, R.L. (1979). Actions of hypothalamic, hypophysiotropic hormones on the brain. Ann. Rev. Physiol. 41: 617-631.
- Motta, M., Fraschini, F. and Martin, L. (1969). "Short" feedback mechanisms in the control of anterior pituitary function. In: Frontiers of Neuroendocrinology. W. Ganong and L. Martin (Eds.), pp. 211-220, Oxford Univ. Press, New York.
- Müller, E.E., Sawano, S. and Arimura, A. (1967). Mechanism of action of growth hormone in altering its own secretion rate: Comparison with the action of dexamethasone. Acta Endocrinol. (Kbh.) 56: 499-509.

- Müller, E.E. and Pecile, A. (1966). Influence of exogenous growth hormone on endogenous growth hormone release. Proc. Soc. Exp. Biol. Med. 122: 1289-1291.
- Müller, E.E. (1973). Nervous control of growth hormone secretion. Neuroendocrinol. 11: 338-342.
- Murphy, W.R., Daughaday, W.H. and Hartnett, C. (1956). The effect of hypophysectomy and growth hormone on the incorporation of labeled sulphate into tibial epiphyseal and nasal cartilage of the rat. J. Lab. Clin. Med. 47: 715-722.
- Nagatsu, T., Levitt, M. and Udenfriend, S. (1964). Tyrosine hydroxylase. The initial step in norepinephrine biosynthesis. J. Biol. Chem. 239: 2910-2917.
- Nemeroff, C.B., Loosen, P.J., Bissette, G., Manberg, P.J., Wilson, J.C. Lipton, M.A. and Prange, A.J. (1979). Pharmacobehavioural effects of hypothalamic peptides in animals and man. Focus on thyroid releasing hormone and neurotensin. Psychoneuroendocrinol. 3: 272-310.
- Nemeroff, C.B., Osbahr, A.J., Birkeno, L.S., Biselle, G., Mailman, R.B., Frye, G.D., Breese, G.R. and Prange, A.J. (1980). Effects of neurotensin on brain neurotransmitter systems, anterior pituitary hormone secretion and behaviour. Endoc. Soc. 62nd. Ann. Meeting 470, Washington.
- Niall, H.D., Hogan, M.L., Sauer, R., Rosenbloom, I.Y. and Greenwood, T. C. (1971). Sequences of pituitary and placental lactogenic and growth hormones: Evolution from a primordial peptide by gene reduplication. Proc. Natl. Acad. Sci. U.S.A. 68: 866-869.
- Niles, L.P., Brown, G.M. and Grota, L.J. (1979). Role of the pineal gland in diurnal endocrine secretion and rhythm regulation. Neuroendocrinol. 29: 14-21.
- O'Briain, D.S., Delellis, R.A., Wolfe, H.J., Reichlin, S., Bollinger, J. and Tashjian, A.H. (1979). Somatostatin immunoreactive cells in C cell hyperplasia and medullary thyroid carcinoma in the rat. Lab. Invest. 40: 275-276.
- O'Donohue, T.L., Miller, R.L., Pendleton, R.C. and Jacobowitz, D.M. (1979). A diurnal rhythm of immunoreactive  $\alpha$  melanocyte-stimulating hormone in discrete regions of the rat brain. Neuroendocrinol. 29: 281-287.

- Ojeda, S.R. and Jameson, H.E. (1977). Developmental patterns of plasma and pituitary growth hormone (GH) in the female rat. Endocrinol. 100: 881-889.
- Oldendorf, W.H. (1977). The blood brain barrier. Exp. Eye Res. 25: 177-190.
- Oliver, C., Eskay, R.L., Ben-Jonathon, N. and Porter, J.C. (1974). Distribution and concentration of thyroid releasing hormone in rat brain. Endocrinol. 95: 540-546.
- Oliver, C. and Porter, J.C. (1978). Distribution and characterization of  $\alpha$ MSH in the rat brain. Endocrinol. 102: 697-705.
- Oliver, E.J. and Baljard, E.J. (1963). Glycogen metabolism in embryonic chick and neonatal rat liver. Biochem. Biophys. Acta 71: 578-588.
- Oomura, Y. (1976). Significance of glucose, insulin and free fatty acid on the hypothalamic feeding and satiety neurons. In: Hunger: Basic Mechanisms and Clinical Implications. D. Novin, W. Wysocka and G. Bray (Eds.), pp. 145-152, Raven Press, New York.
- Otsuka, M., Konishi, S. and Takahashi, T. (1975). Peptide mediating specific synaptic neurotransmitters. Fed. Proc. Am. Soc. Exp. Biol. 34: p. 1922.
- Pacold, S.T., Lawrence, A.M. and Kirsteins, L. (1976). C.N.S. growth hormone: Secretion of growth hormone-like immunoreactivity from monolayer tissue cultures of the amygdala. Clin. Res. 24: 563A.
- Panksepp, J. and Pilcher, C.W.T. (1973). Evidence for an adipokinetic mechanism in the ventromedial hypothalamus. Experientia 29: 793-794.
- Panksepp, J., Reilly, P., Bishop, P., Meeker, R.B. and Vilbeg, J.R. (1976). Effects of  $\alpha$ MSH on motivation, vigilance, and brain respiration. Pharmacol. Biochem. Behav. 5: 59-64.
- Parker, D.C., Pekary, A.E. and Hershman, J.M. (1976). Effect of normal and reversed sleep wake cycles upon nyctohemeral rhythmicity of plasma thyrotropins. Evidence suggestive of an inhibitory influence of sleep. J. Clin. Endocrinol. Metabol. 43: 318-329.
- Parker, C.R., Neaves, W.B., Barrea, A. and Porter, J.C. (1978). Studies on the subsynaptosomal localization of LHRH and TRH in rat hypothalamus. Endocrinol. 102: 1167-1175.

- Parsons, J., Erlandson, S., Hegre, O., McEvoy, R. and Elde, R.P. (1976). Central and peripheral localization of somatostatin. Immunoenzyme immunocytochemical studies. J. Histochem. Cytochem. 24: 872-882.
- Pasternak, G.W., Childers, S.R. and Snyder, S.H. (1980). Opiate analgesia: Evidence for mediation by a subpopulation of opiate receptors. Science 208: 514-516.
- Patrìtti-Laborde, N., Yoshimoto, Y., Wolfsen, A. and Odell, W.D. (1979). Improved method of purifying some radiolabelled glycopeptide hormones. Clin. Chem. 25: 163-165.
- Pavel, S., Goldstein, R., Ghinea, E. and Calb, M. (1977). Chromatographic evidence for vasotocin biosynthesis by cultured pineal ependymal cells from rat fetuses. Endocrinol. 100: 205-208.
- Pavlovic-Hournac, M., Rapaport, L. and Nunez, J. (1971). Incorporation of labelled amino acids into protein by thyroid glands from hypophysectomised rats. I. In vitro studies. Endocrinol. 89: 1477-1484.
- Pavlovic-Hournac, M. and Delbauffe, D. (1973). Action of TSH on the in vivo incorporation of labelled amino acid into thyroglobulin and other thyroidal proteins. Endocrinol. 92: 1273-1276.
- Pearce, L.A. and Schanberg, S.M. (1969). Histamine and spermidine content in brain during development. Science 166: 1301-1303.
- Pearse, A.G.E. (1966). 5-Hydroxytryptophan uptake by dog thyroid C-cells and its possible significance in polypeptide hormone production. Nature (Lond.) 211: 598-600.
- Pearse, A.G.E. (1977). The diffuse neuroendocrine system and the APVD concept: Related "endocrine" peptides in brain, intestine, pituitary, placenta and anurian cutaneous glands. Med. Biol. 55: 115-124.
- Pearse, A.G.E. and Takor, T. (1979). Embryology of the diffuse neuroendocrine system and its relationship to the common peptides. Peptides of the Brain and Gut Kroc. Foundation Conf. Fed. Proc. 38: 2288-2294.
- Pelletier, G., Leclerc, R., Labrie, F., Cote, J., Chretien, M. and Lis, M. (1977). Immunohistochemical localization of  $\beta$ -lipotropic hormone in the pituitary gland. Endocrinol. 100: 770-776.

- Pelletier, G., Leclerc, R., Saavedra, J.M., Brownstein, M.J., Vaudry, H. Ferland, L. and Labrie, F. (1980). Distribution of  $\beta$ -lipotropin ( $\beta$ -LPH), adrenocorticotropin (ACTH) and  $\alpha$  melanocyte stimulating hormone ( $\alpha$ MSH) in the rat brain. I. Origin of the extrahypothalamic fibres. Brain Res. 192: 433-440.
- Pert, C.G., Kuhar, M.J., Pasternak, G.W. and Snyder, S.H. (1976). Opiate receptor: Autoradiographic localization in rat brain. Proc. Natl. Acad. Sci. U.S.A. 73: 3729-3733.
- Phillips, M.I., Mann, H., Dietz, R., Hoffman, W.E. and Ganten, A. (1977). Lowering of hypertension by central saralasin in the absence of plasma renin. Nature (Lond.) 270: 445-447.
- Phillips, L.S., Orawski, A.T. and Belosky, D.C. (1978). Somatomedin and nutrition. IV. Regulation of somatomedin activity and growth cartilage activity by quantity and composition of diet in rats. Endocrinol. 103: 121-127.
- Pierce, J.G. (1974). Chemistry of thyroid stimulating hormone. In: Handbook of Physiology. E. Knobil and W. Sawyer (Eds.), 4: 79-87, Williams and Wilkins, Baltimore.
- Pimstone, B.L., Wittmann, W. and Hansen, J.D.L. (1966). Growth hormone and Kwashiorkor. Role of protein in growth hormone homeostasis. Lancet 2: 779-780.
- Popa, G.T. and Fielding, V. (1930). A portal circulation from the pituitary to the hypothalamic region. J. Anat. 65: 88-91.
- Poth, M.H., Heath, R.G. and Ward, M. (1975). Angiotensin converting enzyme in human brain. J. Neurochem. 25: 83-85.
- Prange, A.J., Breese, G.R., Cott, J.M., Martin, B.R., Cooper, B.R., Wilson, I.C. and Plotnikoff, N.P. (1974). Thyrotropin releasing hormone: Antagonism of pentobarbital in rodents. Life Sci. 14: 447-455.
- Prange, A., Breese, G. and Wilson, I. (1975). Brain behavioural effects of hypothalamic releasing hormone. In: Anatomical Neuroendocrinology. Int. Conf. of Neurobiology of CNS. Hormone Interactions. W. Stumpf and L. Grant (Eds.), pp. 357-371, Karger, Basel.
- Plotnikoff, N.P., Prange, A.J., Breese, G.R., Anderson, M.S. and Wilson, I.C. (1972). Thyrotropin-releasing hormone: Enhancement of DOPA activity by a hypothalamic hormone. Science 178: 417-418.



- Raben, M.S. and Hollenber, C.H. (1959). Effect of growth hormone on plasma fatty acids. J. Clin. Invest. 38: 484-492.
- Rabié, A., Favre, C., Clavel, M.C. and Legrand, J. (1979). Sequential effects of thyroxine on the developing cerebellum of rats made hypothyroid by propyl thiouracil. Brain Res. 161: 469-480.
- Rapaport, B. (1976). Dog thyroid cells in monolayer tissue culture. Adenosine 3'5' cyclic monophosphate response to thyrotropic hormone. Endocrinol. 98: 1189-1197.
- Rehfeld, J.F., Goltermann, N., Larsson, L.I., Ernson, P.M. and Lee, C. M. (1979). Gastrin and cholecystokinin in central and peripheral neurons. Fed. Proc. 38: 2325-2329.
- Reichlin, S. and Mitnick, M. (1973). Biosynthesis of hypothalamic, hypophysiotropic factors. In: Frontiers in Neuroendocrinology. W.F. Ganong and L. Martini (Eds.), pp. 361-369, Oxford. Univ. Press.
- Reid, I.A. and Day, R.P. (1977). Interaction and properties of some components of the renin-angiotensin system in brains. In: Central Action of Angiotensin and Related Hormones. T. Buckley and F. Ferraro (Eds.), pp. 261-282, Pergamon Press, New York.
- Renaud, L.P. and Martin, J.B. (1975). Thyrotropin releasing hormone (TRH): Depressant action on central neuronal activity. Brain Res. 86: 150-154.
- Rezek, M., Harlicek, V., Hughes, K.R. and Friesen, H. (1977). Behavioural and motor excitation and inhibition induced by the administration of small and large doses of somatostatin into the amygdala. Neuropharmacol. 16: 157-162.
- Riddick, D.H., Luciano, A.A., Kusmick, W.F. and Maslar, I.A. (1979). Evidence for a nonpituitary source of amniotic fluid prolactin. Fertility and Sterility 31: 35-39.
- Rieutort, M. (1972). Dosage radioimmunologique de l'hormone somatotrope de rat à la aide d'une nouvelle technique de séparation. C.R. Acad. Sci. (Paris) 274: 3589-3592.
- Rimoin, D.L., Merimee, T.J., Rabinowitz, D. and McKusik, V.A. (1968). Genetic aspects of clinical endocrinology. Rec. Prog. Horm. Res. 24: 365-437.
- Rivier, C., Brown, M. and Vale, W. (1977a). Effect of neurotensin, substance P, and morphine sulfate on the secretion of prolactin and growth hormone in the rat. Endocrinol. 100: 751-754.

- Rivier, C., Vale, W., Ling, W., Brown, M. and Guillemin, R. (1977b). Stimulation in vivo of the secretion of prolactin and growth hormone by  $\beta$ -endorphin. Endocrinol. 100: 238-241.
- Rose, J.C., McDonald, A.A., Heymann, M.A. and Rudolph, A.M. (1978). Developmental aspects of the pituitary-adrenal axis response to hemorrhagic stress in lamb fetuses in utero. J. Clin. Invest. 61: 424-432.
- Rosenfeld, R., Thorsson, A.V. and Hintz, R.L. (1979). Increased somatomedin receptor sites in newborn circulating mononuclear cells. J. Clin. Endocrinol. Metabol. 48: 456-461.
- Rossier, J., French, E., Rivier, C., Shibasaki, T., Guillemin, R. and Bloom, F.E. (1980). Stress induced release of prolactin. Blockade by dexamethasone and naloxone may indicate  $\beta$ -endorphin mediation. Proc. Natl. Acad. Sci. U.S.A. 77: 666-669.
- Sachs, H., Fawcett, P., Takabatake, Y. and Portanova, R. (1969). Biosynthesis and release of vasopressin and neurophysin. Rec. Prog. Horm. Res. 25: 447-491.
- Said, S.I. and Mutt, V. (1972). Isolation from the porcine intestinal wall of a vasoactive octacosapeptide related to secretion and glucagon. Europ. J. Biochem. 28: 199-204.
- Said, S.I. and Rosenberg, R.N. (1976). Vasoactive intestinal polypeptide: Abundant immunoreactivity in neural cell lines and normal nervous tissue. Science 192: 907-908.
- Salmon, W.D. and Daughaday, W.H. (1957). A hormonally controlled serum factor which stimulates sulfate incorporation by cartilage in vitro. J. Lab. Clin. Med. 49: 825-836.
- Salmon, W.D. and DuVall, M.R. (1970). In vitro stimulation of leucine incorporation into muscle and cartilage protein by a serum fraction with sulfation factor activity: Differentiation of effects from those of growth hormone and insulin. Endocrinol. 87: 1168-1180.
- Samson, W.K., Koenig, J., Reeves, J. and McCann, S.M. (1980). Vasoactive intestinal peptide stimulates LHRH release from hypothalamic synaptosomes. Endoc. Soc. 62nd Ann. Meeting. Abstract #746, Washington.
- Samuels, H.H. and Tsai, J. (1973). Thyroid hormone action in cell culture: Demonstration of nuclear receptors in intact cells and isolated nuclei. Proc. Natl. Acad. Sci. U.S.A. 70: 3488-3492.

- Saunders, A., Terry, L.C., Audet, J., Brazeau, P. and Martin, J.B. (1976). Dynamic studies of growth hormone and prolactin secretion in the female rat. Neuroendocrinol. 21: 193-203.
- Sarne, Y., Azov, R. and Weissman, B.A. (1978). A stable enkephalin-like immunoreactive substance in human CSF. Brain Res. 151: 399-403.
- Schaeffer, J.M., Brownstein, M.J. and Axelrod, J. (1977). Thyrotropin-releasing hormone-like material in the rat retina: Changes due to environmental lighting. Proc. Natl. Acad. Sci. U.S.A. 74: 3579-3581.
- Schaeffer, J.M. and Brownstein, M.J. (1980). Ontogeny of TRH-like material in several regions of the rat brain. Brain Res. 182: 207-210.
- Schalch, D.S. (1969). The influence of physical stress and exercise on growth hormone and insulin secretion in man. J. Lab. Clin. Med. 69: 256-269.
- Schalch, D.S., Heinreich, V.E., Johnson, C.J., Koch, J.G. and Miller, L. L. (1977). Hepatic synthesis of somatomedin NSILA is under hormonal control. Clin. Res. 25: 399A.
- Schally, A., Bowers, C. and Redding, T. (1966). Isolation of thyrotropin releasing factor (TRF) from porcine hypothalamus. Biochem. Biophys. Res. Comm. 25: 165-169.
- Scharrer, E. and Scharrer, B. (1954). Hormones produced by neurosecretory cells. Rec. Prog. Horm. Res. 10: 183-240.
- Scharrer, B. (1978). Peptidergic neurons: Facts and trends. Gen. Comp. Endocrinol. 34: 50-62.
- Schenkel-Hulliger, L., Koella, W.P., Hartmann, A. and Maitre, L. (1976). Tremorogenic effect of thyrotropin-releasing hormone in rats. Experientia 30: 1168-1170.
- Schneider, H.P.G. and McCann, S.M. (1969). Possible role of dopamine as transmitter to promote the discharge of LH-releasing factor. Endocrinol. 85: 121-126.
- Scott, T.W., Freinkel, N., Klein, J.H. and Nitzan, M. (1970). Metabolism of phospholipids, neutral lipids and carbohydrate in dispersed porcine thyroid cells. Comparative effects of pituitary thyrotropin and dibutyryl 3',5' adenosine monophosphate on turnover of individual phospholipids in isolated cells and slices from pig thyroid. Endocrinol. 87: 854-863.

- Segal, D.S. and Mandell, A.J. (1974). Differential behavioural effects of hypothalamic polypeptides. In: The Thyroid Axis, Drugs and Behaviour. A.J. Prange (Ed.), pp. 129-133, Raven Press, New York.
- Segal, D.S. and Mandell, A.J. (1975). Intraventricular infusion decrease the spontaneous motor activity of freely moving rats. In: The Thyroid Axis, Drugs and Behaviour. A.J. Prange (Ed.), pp. 129-137, Raven Press, New York.
- Seljelid, R. (1967). Endocytosis in thyroid follicle cells: V. On the redistribution of cytosomes following stimulation with thyrotropic hormone. J. Ultrastructure Res. 18: 479-488.
- Sétáló, G., Flerko, B., Arimura, A. and Schally, A.V. (1978). Brain cells as producers of releasing and inhibiting hormones. In: Int. Rev. Cytol. H.R. Bourne and S.F. Danielli (Eds.), pp. 93-97, Academic Press, New York.
- Shambaugh, G.E., Kulch, M. and Wilber, J. (1978). The placenta: A new locus for TRH. A.S.C.I. Endocrinol. 495A.
- Shore, V.G. and Shore, B. (1973). Heterogeneity of human plasma very low density lipoproteins. Separation of species differing in protein components. Biochem. 12: 502-507.
- Sigel, M.B., Vanderlaan, W.P., Vanderlaan, E.F. and Lewis, V.J. (1980). Measurement of multiple forms of h.growth hormone cross reactivities in conventional and two chain radioimmunoassays. Endocrinol. 106: 92-97.
- Silberberg, M. (1935). Effects of extract of cattle anterior pituitary gland on endochondral ossification in young guinea pigs. Proc. Soc. Exp. Biol. Med. 32: 1423-1425.
- Simantov, R., Kuhar, M.J., Pasternak, G.W. and Snyder, S.H. (1976a). The regional distribution of a morphine-like factor enkephalin in monkey brain. Brain Res. 106: 189-197.
- Simantov, R. and Snyder, S.H. (1976b). Morphine-like peptides, leucine enkephalin and methionine enkephalin: Interactions with the opiate receptor. Molec. Pharmacol. 12: 987-998.
- Sims, K.B., Hoffman, D.L., Said, S.I. and Zimmerman, E.A. (1980). Vasoactive intestinal polypeptide (VIP) in mouse and rat brain. An immunocytochemical study. Brain Res. 186: 165-183.

- Singh, R.N.P., Seavey, B.K., Rice, V.P., Lindsey, T.T. and Lewis, V.J. (1974). Modified forms of human growth hormone with increased biological activities. Endocrinol. 94: 883-891.
- Siret, N.E., McLean, A.S., Bray, J.J. and Hubbard, J.I. (1977). Distribution of angiotensin II receptors in rat brain. Brain Res. 122: 299-312.
- Slater, E.E., Defendi, R. and Zimmerman, E.A. (1980). Renin in human brain. Clin. Res. 28: 267A.
- Smith, C.J.V. (1972). Hypothalamic glucoreceptors - the influence of gold thioglucose implants in the ventromedial and lateral hypothalamic areas of normal and diabetic rats. Physiol. Behav. 9: 391-396.
- Snyder, A.M., Hull, E.M. and Roth, J.A. (1979). The effect of maternal progesterone injections on fetal development of brain monoamine oxidase of rats. Brain Res. 170: 194-197.
- Spies, H.G., Quadri, S.K., Chappel, S.C. and Norman, R.L. (1980). Dopaminergic and opioid compounds. Effects on prolactin and LH release after electrical stimulation of the hypothalamus in ovariectomized rhesus monkeys. Neuroendocrinol. 30: 249-256.
- Steel, R.G.D. and Torrie, J.H. (1960). Principles and Procedures of Statistics. McGraw Hill Book Co. Inc., New York.
- Sternberger, L.A., Hardy, P.H., Cuculis, J.J. and Meyer, H.G. (1970). The unlabeled antibody enzyme method of immunohistochemistry. Preparation and properties of soluble antigen-antibody complex (horseradish peroxidase - antihorseradish peroxidase) and its use in identification of spirochetes. J. Histochem. Cytochem. 18: 315-333.
- Steiner, R.A., Illner, P., Rolfs, A.D., Toivola, P.T.K. and Gale, C.C. (1978). Noradrenergic and dopaminergic regulation of growth hormone and prolactin in baboons. Neuroendocrinol. 26: 15-31.
- Stern, W.C., Millar, M., Resnick, O. and Morgane, P.J. (1975). Distribution of <sup>125</sup>I-labelled rat growth hormone in regional brain areas and peripheral tissue of the rat. Am. J. Anat. 144: 503-508.
- Strauss, E., Muller, J.E., Choi, H.S., Paronatto, F. and Yalow, R.S. (1977). Immunohistochemical localization in rabbit brain of a peptide resembling the COOH-terminal octapeptide of cholecystokinin. Proc. Natl. Acad. Sci. U.S.A. 74: 3033-3034.

- Stroud, S.W., Hoog, J.M.C. and Bieler, E. (1973). A simple method for the extraction and purification of human growth hormone and its assay by paper chromatography. J. Clin. Endocrinol. Metabol. 37: 860-866.
- Szentágothai, J. (1968). Anatomical considerations. In: Hypothalamic Control of the Anterior Pituitary. An Experimental-Morphological Study. J. Szentágothai, B. Flerkó, B. Mess and B. Halász (Eds.), pp. 22-29, Budapest Akad., Kiadó.
- Takahara, J., Yunokis, L., Hosigi, H., Yakushigi, W., Kageyama, J. and Ofuji, T. (1980). Concomitant increases in serum growth hormone and hypothalamic somatostatin in rats after injection of  $\gamma$ -amino-butyric acid on  $\gamma$  hydroxybutyric acid. Endocrinol. 106: 343-347.
- Tan, A.T., Tsang, D., Renaud, L.P. and Martin, J.B. (1977). Effect of somatostatin on Ca transport in guinea pig cortex synaptosomes. Brain Res. 123: 193-196.
- Taurog, A. (1970). Thyroid peroxidase and thyroxine biosynthesis. Rec. Prog. Horm. Res. 26: 189-247.
- Taylor, D.P. and Pert, C.B. (1979). V.I.P: Specific binding to rat brain membranes. Proc. Natl. Acad. Sci. (Wash.) 76: 660-664.
- Terenius, L., Gispen, W.H. and DeWied, D. (1975). ACTH-like peptides and opiate receptors in the rat brain: Structure activity studies. Eur. J. Pharmac. 33: 395-399.
- Teschemacher, H., Opheim, K.E., Cox, B.M. and Goldstein, A. (1975). A peptide-like substance from pituitary that acts like morphine. Life Sci. 16: 1771-1776.
- Tice, L.W. (1974). Effects of hypophysectomy and TSH replacement on the ultrastructural localization of thyroperoxidase. Endocrinol. 95: 421-433.
- Toubeau, G., Desdin, J., Parmastier, M. and Pasteels, J.L. (1979). Localization of prolactin-like and ACTH immunoactivity within the brain of the rat. Neuroendocrinol. 29: 374-384.
- Truding, R., Shelanski, M.L. and Morell, P. (1975). Glycoproteins released into the culture medium of differentiating murine neuroblastoma cells. J. Biol. Chem. 250: 9348-9354.
- Unger, R.H., Eisentraut, A.M. and Madison, L.L. (1965). Fasting levels of growth hormone in men and woman. Nature (Lond.) 205: 804-805.

- Uthne, K., Reagen, C.R., Gimpel, L.P. and Kostyo, J.L. (1974). Effects of human somatomedin preparations on membrane transport and protein synthesis in the isolated rat diaphragm. J. Clin. Endocrinol. Metabol. 39: 548-554.
- Vaitukaitis, J.L., Ross, G.T. and Pierce, J.G. (1973). Generation of specific antisera with the hormone specific  $\beta$ -sub-unit of h. TSH or h. FSH. J. Clin. Endocrinol. Metabol. 37: 653-659.
- Vale, W., Grant, G. and Amoss, M. (1972). Culture of enzymatically dispersed anterior pituitary cells. Functional validation of a method. Endocrinol. 91: 562-572.
- Vanderaeglan, J.T., Signeau, J.C. and Gepts, W. (1975). New peptide in the vertebrate CNS, reacting with antigastrin antibodies. Nature (Lond.) 257: 604-605.
- Van Houten, M., Kroprina, B. and Brawer, J.R. (1980). Insulin binding sites localized to nerve terminals in rat median eminence and arcuate nucleus. Science 207: 1081-1083.
- Van Wyk, J., Hall, K. and Vandenbandie, J.L. (1972). Partial purification from human plasma of a small peptide with sulfation factor and thymidine activity. In: Proc. of 2nd International Symposium on Growth Hormone. A. Pecile and E.E. Müller (Eds.), pp. 115-121. Excerpta Medica.
- Van Wyk, J.J., Underwood, L.E., Hintz, R.L., Clemmons, D.R., Voina, S.J. and Weaver, R.P. (1974). The somatomedins: A family of insulin-like hormones under growth hormone control. Rec. Prog. Horm. Res. 30: 259-318.
- Vaudry, H., Tonon, M.C., Delarue, R., Vaillant, R. and Kraicer, J. (1978). Biological and radioimmunological evidence for MSH of extrapituitary origin in the rat brain. Neuroendocrin. 27: 9-24.
- Vijayan, E. and McCann, S.M. (1977). Suppression of feeding and drinking activity in rats following intraventricular injection of T.R.H. Endocrinol. 100: 1727-1730.
- Vijayan, E., Samson, W.K. and McCann, S.M. (1978). Effects of intraventricular injection of gastrin on release of luteinising hormone, prolactin, TSH and growth hormone in conscious ovariectomised rats. Life Sci. 23: 2225-2232.
- Vijayan, E., Samson, W.K. and McCann, S.M. (1979). In-vivo and in-vitro effects of cholecystokinin on gonadotropin, prolactin, growth hormone and thyrotropin release in the rat. Brain Res. 172: 295-302.

- Villareal, J.A., Rivier, J. and Brown, M. (1978). Evidence for a bombesin-like substance in hypothalamus. Chemical and immunological characterization. Endoc. Soc. Meet. p. 390.
- Vito, C.C., Wieland, S.S. and Fox, T.O. (1979). Androgen receptors exist throughout the critical period of brain sexual differentiation. Nature 282: 308-310.
- Von Euler, U.S. and Gaddum, J.H. (1931). An unidentified depressor substance in certain tissue extracts. J. Physiol. (Lond.) 72: 74-87.
- Voogt, J.L., Clemens, J.A. and Whitemair, C.K. (1970). Effects of implantation of growth hormone into the median eminence of cycling female rats on pituitary growth hormone and hypothalamic GRF content. Fed. Proc. 29: 377, #749.
- Vzhekov, M.G., Murphy, S. and Rose, S.P. (1979). Ontogenesis of serotonin receptors in different regions of rat brain. Brain Res. 168: 195-199.
- Wadell, W.J. (1956). A simple ultraviolet spectrophotometric method for the determination of protein. J. Lab. Clin. Med. 48: 311-314.
- Wäger, G. (1973). Stimulating effect of dibutyryl cyclic adenosine 3', 5'-monophosphate on protein synthesis in a thyroidal cell-free system at low potassium and sodium concentration. Acta Endocrinol. 74: 711-722.
- Wallis, M. and Davies, R.V. (1976). Studies on the chemistry of bovine and rat growth hormones. In: Growth Hormone and Related Peptides. E. Pecile and E.E. Müller (Eds.), pp. 1-13, Excerpta Medica, American Elsevier.
- Wallis, C.J. and Printz, M.P. (1980). Adrenal regulation of regional brain angiotensinogen content. Endocrinol. 106: 337-342.
- Warburg, J., Oliver, C., Eskay, R.L., Parker, C.R., Barrea, A. and Porter, J.C. (1977). Release of  $\alpha$ MSH from a synaptosome enriched fraction prepared from rat hypothalamic tissue. In: Melanocyte Stimulating Hormone: Control, Chemistry and Effects. Frontiers in Hormone Research. F.J.H. Tilden, D.F. Swaab and Tj.B. Wimersma (Eds.), 4: 167-169, Karger, Basel.
- Watson, S.J., Barchas, J.D. and Li, C.H. (1977).  $\beta$ -lipotropin: localization of cells and axons in rat brain by immunocytochemistry. Proc. Natl. Acad. Sci. (Wash.) 74: 5155-5158.



- Watson, S.J., Richard, C.W. and Barchas, J.D. (1978). Adrenocorticotropin in rat brain: Immunocytochemical localization in cells and axons. Science 200: 1180-1171.
- Weiss, S., Bergland, R., Page, R., Turgen, C. and Aymers, W.C. (1975). Pituitary cell transplants to the central ventricles promote growth in the hypophysectomised rat. Proc. Soc. Exp. Biol. Med. 159: 409-413.
- Weisz, J. and Ward, I.L. (1980). Plasma testosterone and progesterone titres of pregnant rats, their male and female fetusus, and neonatal offspring. Endocrinol. 106: 306-316.
- Weitzman, E.D. (1980). Biologic rhythms and hormone secretion patterns. In: Neuroendocrinology. D.T. Krieger and J.C. Hughes (Eds.), pp. 85-92, Sinaur Assoc., Sunderland, Massachusetts.
- Wharton, J., Polak, J.M., Bloom, S.R. and Pearse, A.G.E. (1978). Bombesin-like immunoreactivity in the lung. Nature (Lond.) 273: 769-770.
- Wikland-Alberson, K., Edén, S. and Isarsson, O. (1980). Analysis of early responses to growth hormone on amino acid transport and protein synthesis in the diaphragm of young normal rats. Endocrinol. 106: 291-297.
- Wilber, J.F. (1971). Stimulation of <sup>14</sup>C glucosamine and <sup>14</sup>C alanine incorporation into thyrotropin by synthetic releasing hormone. Endocrinol. 89: 873-877.
- Wilber, J.F. and Siebel, M.J. (1973a). Thyrotropin releasing hormone interactions with an anterior pituitary membrane receptor. Endocrinol. 92: 888-893.
- Wilber, J.F. (1973b). Thyrotropin releasing hormone. Secretion and actions. Ann. Rev. Med. 24: 353-359.
- Williams, J.A. and Malayan, S.A. (1975). Effects of TSH on iodide transport by mouse thyroid lobes in vitro. Endocrinol. 97: 162-168.
- Willoughby, J.O. and Martin, J.B. (1978). The suprachiasmatic nucleus synchronises growth hormone secretory rhythms with the light-dark cycle. Brain Res. 151: 413-417.
- Wimersma, B.Tj. (1977). Effects of MSH and related peptides on avoidance behaviour in rats. In: Melanocyte Stimulating Hormone: Control, Chemistry and Effects. Frontiers of Hormone Research. F.J. Tilden, D.F. Swaab and Tj.B. Wimersma (Eds.), 4: 129-139, Karger Basel.

- Winokur, A. and Utiger, R.D. (1974). T.R.H. regional distribution in rat brain. Science 185: 265-267.
- Wislocki, G.B. and King, S.L. (1936). The permeability of the hypophysis and hypothalamus to vital dyes, with a study of the hypophyseal vascular supply. Am. J. Anat. 58: 421-472.
- Wurtman, R.J. (1980). The pineal as a neuroendocrine transducer. In: Neuroendocrinology. D.T. Krieger and J.C. Hughes (Eds.), pp. 87-93, Sinauer Assoc., Sunderland, Massachusetts.
- Yaksh, T.L. and Yamamura, H.I. (1975). Blockade by morphine of acetylcholine release from the caudate nucleus in the mid-pontine pretrigeminal cat. Brain Res. 83: 520-524.
- Yalow, R.S. and Berson, S.A. (1959). Assay of plasma insulin in human subjects by immunological methods. Nature (Lond.) 184: 1648-1649.
- Yang, H.T. and Neff, N.H. (1972). Distribution and properties of angiotensin converting enzyme of rat brain. J. Neurochem. 19: 2443-2450.
- Yates, F.E. and Moran, J.W. (1974). Handbook of Physiology. Sect. 7. Endocrinology. Williams and Wilkins, 4: 367-371.
- Yasamura, Y. (1968). Retention of differentiated function in clonal animal cell lines, particularly hormone secreting cultures. Am. Zoologist 8: 285-305.
- Zimmerman, E.A., Liotta, P. and Krieger, D.T. (1978).  $\beta$ -lipotropin in brain: Localization in hypothalamic neurons by immunoperoxidase technique. Cell Tiss. Res. 186: 393-398.
- Zimmerman, E.G. (1979). Peptides of the brain and gut. Fed. Proc. 38: 2286-2287.
- Zucker, I. (1980). Light, behaviour and biological rhythms. In: Neuroendocrinology. D.T. Krieger and J.C. Hughes (Eds.), pp. 93-101, Sinauer Assoc., Sunderland, Massachusetts.
- Zusman, D.R. and Burrow, G.N. (1975). TSH regulation of ornithine decarboxylase activity in thyroid. Endocrinol. 97: 1089-1096.

APPROVAL SHEET.

The dissertation submitted by Sally A. Hojvat  
has been read and approved by the following committee:

Dr. A.M. Lawrence, Thesis Director.  
Professor, Medicine and Biochemistry,  
Loyola.

Dr. M. Manteuffel,  
Assoc. Professor, Biochemistry,  
Loyola.

Dr. A. Frankfater,  
Assoc. Professor, Biochemistry,  
Loyola.

Dr. C.C.C. O'Morchoe,  
Professor and Chairman, Anatomy,  
Loyola.

Dr. R. Miller,  
Asst. Professor, Pharmacology and Physiology,  
University of Chicago.

The final copies have been examined by the director of the  
dissertation and the signature which appears below verifies  
the fact that any necessary changes have been incorporated  
and that the dissertation is now given final approval by the  
Committee with reference to content and form.

The dissertation is therefore accepted in partial fulfillment  
of the requirements for the degree of Doctor of Philosophy.

Dec. 8, 1980  
Date

A. M. Lawrence  
Director's Signature