



2016

Draft Genome for a Urinary Isolate of *Lactobacillus crispatus*

Travis Kyle Price

Loyola University Chicago

Majed Shaheen

Loyola University Chicago

Laurynas Kalesinskas

Loyola University Chicago

Kema Malki

Loyola University Chicago

Evann Elizabeth Hilt

Loyola University Chicago

See next page for additional authors

Follow this and additional works at: https://ecommons.luc.edu/bioinformatics_facpub



Part of the [Bacteriology Commons](#), and the [Genomics Commons](#)

Recommended Citation

Price, Travis Kyle; Shaheen, Majed; Kalesinskas, Laurynas; Malki, Kema; Hilt, Evann Elizabeth; Putonti, Catherine; and Wolfe, Alan J.. Draft Genome for a Urinary Isolate of *Lactobacillus crispatus*. *Genome Announcements*, 4, 6: e01278-16, 2016. Retrieved from Loyola eCommons, Bioinformatics Faculty Publications, <http://dx.doi.org/10.1128/genomeA.01248-16>

This Article is brought to you for free and open access by the Faculty Publications at Loyola eCommons. It has been accepted for inclusion in Bioinformatics Faculty Publications by an authorized administrator of Loyola eCommons. For more information, please contact ecommons@luc.edu.



This work is licensed under a [Creative Commons Attribution 4.0 License](#).

© 2016 Price et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Authors

Travis Kyle Price, Majed Shaheen, Laurynas Kalesinskas, Kema Malki, Evann Elizabeth Hilt, Catherine Putonti, and Alan J. Wolfe

Draft Genome Sequence of a Urinary Isolate of *Lactobacillus crispatus*

Travis K. Price,^a Majed Shaheen,^b Laurynas Kalesinskas,^{b,c} Kema Malki,^c Evann E. Hilt,^a Catherine Putonti,^{b,c,d} Alan J. Wolfe^a

Health Sciences Division, Department of Microbiology and Immunology, Stritch School of Medicine, Loyola University Chicago, Maywood, Illinois, USA^a; Bioinformatics Program, Loyola University Chicago, Chicago, Illinois, USA^b; Department of Biology, Loyola University Chicago, Chicago, Illinois, USA^c; Department of Computer Science, Loyola University Chicago, Chicago, Illinois, USA^d

While *Lactobacillus crispatus* contributes to the stability of normal vaginal microbiota, its role in urinary health remains unclear. As part of an on-going attempt to characterize the female urinary microbiota, we report the genome sequence of an *L. crispatus* strain isolated from a woman displaying no lower urinary tract symptoms.

Received 17 September 2016 Accepted 7 October 2016 Published 1 December 2016

Citation Price TK, Shaheen M, Kalesinskas L, Malki K, Hilt EE, Putonti C, Wolfe AJ. 2016. Draft genome sequence of a urinary isolate of *Lactobacillus crispatus*. *Genome Announc* 4(6):e01278-16. doi:10.1128/genomeA.01278-16.

Copyright © 2016 Price et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Alan J. Wolfe, awolfe@luc.edu.

As part of an attempt to characterize the newly discovered female urinary microbiota (1–8), we report the genome sequence and annotation of a strain of *Lactobacillus crispatus* isolated from the bladder of an adult female. This is the first genome report of a urinary isolate of *L. crispatus*, a species associated with bladder health (8).

Using the expanded spectrum version (9) of the enhanced quantitative urine culture protocol (2), *L. crispatus* strain C037 was isolated from a healthy female not displaying any urinary symptoms. The strain was subcultured to purity, analyzed by matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry, and pure cultures were stored at -80°C in 2 mL CryoSaver Brucella broth with 10% glycerol, no beads, cryovials (Hardy Diagnostics). For genome extraction, the preserved pure culture isolate was grown on 5% sheep blood agar (BDBBL™ prepared plated media) under 5% CO_2 at 35°C for 24 h.

To extract genomic DNA, cells were resuspended in 0.5 mL DNA extraction buffer (20 mM tris-Cl, 2 mM EDTA, 1.2% Triton X-100, pH 8) followed by addition of 50 μL lysozyme (20 mg/mL), 30 μL mutanolysin, and 5 μL RNase (10 mg/mL). After a 1-h incubation at 37°C , 80 μL 10% SDS and 20 μL proteinase K were added followed by a 2-h incubation at 55°C . Then, 210 μL of 6M NaCl and 700 μL phenol-chloroform were added. After a 30-min incubation with rotation, the solution was centrifuged at 13,500 rpm for 10 min, and the aqueous phase extracted. An equivalent volume of isopropanol was added; after a 10-min incubation, the solution was centrifuged at 13,500 rpm for 10 min. The supernatant was decanted and the DNA pellet precipitated using 600 μL 70% ethanol. Following ethanol evaporation, the DNA pellet was resuspended in tris-EDTA and stored at -20°C .

Genomic DNA was diluted in water to a concentration of 0.2 ng/ μL . Library preparation was performed using the Nextera XT DNA library preparation kit (Illumina) according to manufacturer's instructions with 1 ng of input DNA. The isolate was sequenced twice, on two separate runs, using the Illumina MiSeq platform and the MiSeq reagent kit v2 (300-cycles). Sequence assembly was performed using Velvet (10) ($k = 99$) followed by SSPACE (11) for scaffolding. *L. crispatus* C037 was assembled into

96 scaffolds with a genome coverage of $113\times$. The scaffolds include 2.147 Mbp of sequence with a G+C content of 36.6%. Gene annotations were performed using GLIMMER (12) and tRNA-Scan (13) identifying 2,096 protein coding genes, 65 RNA (tRNA and rRNA) genes, and four clustered regularly interspaced short palindromic repeats (CRISPR) (14). The 16S rRNA gene sequence of the urinary isolate C037 was identical to that of the species' type strain *L. crispatus* ST1 (NR_074986), an avian enteric strain. One scaffold (10,704 bp in length) produced a hit to the 16,663 bp *L. crispatus* plasmid pLc17 (KR052811); while a significant proportion of the plasmid sequence was detected, a complete, circularized assembly was not possible.

Accession number(s). The draft whole-genome project for *L. crispatus* C037 has been deposited at DDBJ/EMBL/GenBank under accession number **MAKH00000000**.

ACKNOWLEDGMENTS

We acknowledge Gina Kuffel and Michael Zilliox for sequencing this genome.

FUNDING INFORMATION

This work, including the efforts of Alan J. Wolfe, was funded by HHS | National Institutes of Health (NIH) (R01DK104718). This work, including the efforts of Catherine Putonti, was funded by National Science Foundation (NSF) (1149387). This work, including the efforts of Catherine Putonti and Alan J. Wolfe, was funded by Loyola University Chicago (LUC) (Multidisciplinary Research Award).

REFERENCES

- Fouts DE, Pieper R, Szpakowski S, Pohl H, Knoblach S, Suh MJ, Huang ST, Ljungberg I, Sprague BM, Lucas SK, Torralba M, Nelson KE, Groah SL. 2012. Integrated next-generation sequencing of 16S rDNA and metaproteomics differentiate the healthy urine microbiome from asymptomatic bacteriuria in neuropathic bladder associated with spinal cord injury. *J Transl Med* 10:174. <http://dx.doi.org/10.1186/1479-5876-10-174>.
- Hilt EE, McKinley K, Pearce MM, Rosenfeld AB, Zilliox MJ, Mueller ER, Brubaker L, Gai X, Wolfe AJ, Schreckenberger PC. 2014. Urine is not sterile: use of enhanced urine culture techniques to detect resident bacterial flora in the adult female bladder. *J Clin Microbiol* 52:871–876. <http://dx.doi.org/10.1128/JCM.02876-13>.
- Khasriya R, Sathiananthamoorthy S, Ismail S, Kelsey M, Wilson M,

- Rohn JL, Malone-Lee J. 2013. Spectrum of bacterial colonization associated with urothelial cells from patients with chronic lower urinary tract symptoms. *J Clin Microbiol* 51:2054–2062. <http://dx.doi.org/10.1128/JCM.03314-12>.
4. Lewis DA, Brown R, Williams J, White P, Jacobson SK, Marchesi JR, Drake MJ. 2013. The human urinary microbiome; bacterial DNA in voided urine of asymptomatic adults. *Front Cell Infect Microbiol* 3:41. <http://dx.doi.org/10.3389/fcimb.2013.00041>.
 5. Nienhouse V, Gao X, Dong Q, Nelson DE, Toh E, McKinley K, Schreckenberger P, Shibata N, Fok CS, Mueller ER, Brubaker L, Wolfe AJ, Radek KA. 2014. Interplay between bladder microbiota and urinary antimicrobial peptides: mechanisms for human urinary tract infection risk and symptom severity. *PLoS One* 9:e114185. <http://dx.doi.org/10.1371/journal.pone.0114185>.
 6. Siddiqui H, Nederbragt AJ, Lagesen K, Jeansson SL, Jakobsen KS. 2011. Assessing diversity of the female urine microbiota by high throughput sequencing of 16S rDNA amplicons. *BMC Microbiol* 11:244. <http://dx.doi.org/10.1186/1471-2180-11-244>.
 7. Wolfe AJ, Toh E, Shibata N, Rong R, Kenton K, Fitzgerald M, Mueller ER, Schreckenberger P, Dong Q, Nelson DE, Brubaker L. 2012. Evidence of uncultivated bacteria in the adult female bladder. *J Clin Microbiol* 50:1376–1383. <http://dx.doi.org/10.1128/JCM.05852-11>.
 8. Pearce MM, Hilt EE, Rosenfeld AB, Zilliox MJ, Thomas-White K, Fok C, Kliethermes S, Schreckenberger PC, Brubaker L, Gai X, Wolfe AJ. 2014. The female urinary microbiome: a comparison of women with and without urgency urinary incontinence. *mBio* 5:e01283-14. <http://dx.doi.org/10.1128/mBio.01283-14>.
 9. Price TK, Dune T, Hilt EE, Thomas-White KJ, Kliethermes S, Brincat C, Brubaker L, Wolfe AJ, Mueller ER, Schreckenberger PC. 2016. The clinical urine culture: enhanced techniques improve detection of clinically relevant microorganisms. *J Clin Microbiol* 54:1216–1222. <http://dx.doi.org/10.1128/JCM.00044-16>.
 10. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
 11. Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27:578–579. <http://dx.doi.org/10.1093/bioinformatics/btq683>.
 12. Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with glimmer. *Bioinformatics* 23:673–679. <http://dx.doi.org/10.1093/bioinformatics/btm009>.
 13. Schattner P, Brooks AN, Lowe TM. 2005. The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res* 33:W686–W689. <http://dx.doi.org/10.1093/nar/gki366>.
 14. Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35:W52–W57. <http://dx.doi.org/10.1093/nar/gkm360>.