




Contents lists available at ScienceDirect

Chemical Geology

journal homepage: www.elsevier.com/locate/chemgeo

The large-scale evolution of neodymium isotopic composition in the global modern and Holocene ocean revealed from seawater and archive data

Kazuyo Tachikawa^{a,*}, Thomas Arsouze^{b,c}, Germain Bayon^d, Aloys Bory^e, Christophe Colin^f, Jean-Claude Dutay^g, Norbert Frank^h, Xavier Giraud^a, Alexandra T. Gourlanⁱ, Catherine Jeandel^j, François Lacan^j, Laure Meynadier^k, Paolo Montagna^l, Alexander M. Piotrowski^m, Yves Plancherelⁿ, Emmanuelle Pucéat^o, Matthieu Roy-Barman^g, Claire Waelbroeck^g

^a Aix Marseille Univ, CNRS, IRD, Coll France, CEREGE, Aix-en-Provence, France

^b Laboratoire de Météorologie Dynamique, École Polytechnique, Palaiseau, France

^c ENSTA ParisTech, Université Paris-Saclay, 828 bd des Maréchaux, 91762 Palaiseau cedex, France

^d IFREMER, Unité de Recherche Géosciences Marines, F-29280 Plouzané, France

^e Université de Lille, CNRS, Université du Littoral Côte d'Opale, UMR8187, LOG, Laboratoire d'Océanologie et de Géosciences, F-59000 Lille, France

^f Laboratoire Géosciences Paris-Sud (GEOPS), Université de Paris Sud, Université Paris-Saclay, 91405 Orsay, France

^g Laboratoire des Sciences du Climat et de l'Environnement LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

^h Institute of Environmental Physics, University of Heidelberg, 69120 Heidelberg, Germany

ⁱ Université Grenoble Alpes, ISTerre, CS 40700, 38058 Grenoble Cedex 9, France

^j LEGOS (Université de Toulouse/CNRS/CNES/IRD/UPS), Observatoire Midi-Pyrénées, Toulouse, France

^k Equipe de Géochimie-Cosmochimie, Institut de Physique du Globe de Paris – Sorbonne Paris Cité, Université Paris Diderot, UMR CNRS 7154, 1 rue Jussieu, 75005 Paris, France

^l ISMAR-CNR, via Gobetti 101, 40129 Bologna, Italy

^m Department of Earth Sciences, University of Cambridge, Cambridge, UK

ⁿ Department of Earth Sciences, University of Oxford, Oxford OX1 3AN, UK

^o UMR CNRS 6282 Biogéosciences, 6 Bd. Gabriel, 21000 Dijon, France

ARTICLE INFO

Keywords:

Nd isotopes

Water mass tracer

Data compilation

Predicted seawater ϵ_{Nd}

Seawater-archive comparison

ABSTRACT

Neodymium isotopic compositions ($^{143}\text{Nd}/^{144}\text{Nd}$ or ϵ_{Nd}) have been used as a tracer of water masses and lithogenic inputs to the ocean. To further evaluate the faithfulness of this tracer, we have updated a global seawater ϵ_{Nd} database and combined it with hydrography parameters (temperature, salinity, nutrients and oxygen concentrations), carbon isotopic ratio and radiocarbon of dissolved inorganic carbon. Archive ϵ_{Nd} data are also compiled for leachates, foraminiferal tests, deep-sea corals and fish teeth/debris from the Holocene period (< 10,000 years).

At water depths ≥ 1500 m, property-property plots show clear correlations between seawater ϵ_{Nd} and the other variables, suggesting that large-scale water mass mixing is a primary control of deepwater ϵ_{Nd} distribution. At ≥ 200 m, basin-scale seawater T-S- ϵ_{Nd} diagrams demonstrate the isotopic evolution of different water masses. Seawater and archive ϵ_{Nd} values are compared using property-property plots and T-S- ϵ_{Nd} diagrams. Archive values generally agree with corresponding seawater values although they tend to be at the upper limit in the Pacific. Both positive and negative offsets exist in the northern North Atlantic. Applying multiple regression analysis to deep (≥ 1500 m) seawater data, we established empirical equations that predict the main, large-scale, deepwater ϵ_{Nd} trends from hydrography parameters. Large offsets from the predicted values are interpreted as a sign of significant local/regional influence. Dominant continental influence on seawater and archive ϵ_{Nd} is observed mainly within 1000 km from the continents. Generally, seawater and archive ϵ_{Nd} values form gradual latitudinal trend in the Atlantic and Pacific at depths ≥ 600 m, consistent with the idea that Nd isotopes help distinguish between northern/southern sourced water contributions at intermediate and deep water depths.

* Corresponding author.

E-mail address: kazuyo@cerge.fr (K. Tachikawa).

<http://dx.doi.org/10.1016/j.chemgeo.2017.03.018>

Received 30 December 2016; Received in revised form 12 March 2017; Accepted 18 March 2017

0009-2541/ © 2017 Elsevier B.V. All rights reserved.

1. Introduction

Neodymium is one of the Rare Earth Elements (REE) and consists of seven isotopes. One of the isotopes, ^{143}Nd , is an α -decay product of ^{147}Sm that also belongs to REE with a half-life of 1.06×10^{11} years. Since Sm/Nd ratio in rocks and minerals varies during crystallization from magma, $^{143}\text{Nd}/^{144}\text{Nd}$ of continental materials change as a function of age and lithology. Seawater $^{143}\text{Nd}/^{144}\text{Nd}$ (or $\epsilon_{\text{Nd}} = [(^{143}\text{Nd}/^{144}\text{Nd})_{\text{sample}} / (^{143}\text{Nd}/^{144}\text{Nd})_{\text{CHUR}} - 1] \times 10^4$, CHUR is a Chondritic Uniform Reservoir with $^{143}\text{Nd}/^{144}\text{Nd}$ of 0.512638 (Jacobsen and Wasserburg, 1980) or of 0.512630 (Bouvier et al., 2008)) reflects the isotopic signature of continents surrounding the oceanic basins. In the modern ocean, the lowest seawater ϵ_{Nd} value of -26.6 is observed in the Baffin Bay (Stordal and Wasserburg, 1986) encompassed by old craton whereas the most radiogenic value of $+2.7$ is found in the Eastern Equatorial Pacific where young volcanogenic material is abundant (Grasse et al., 2012). The estimated residence time of Nd in the ocean of 360 to 700 years (Tachikawa et al., 2003; Siddall et al., 2008; Rempfer et al., 2011) is shorter than mean mixing time of deep ocean of about 1500 years (Broecker and Peng, 1982). This explains the heterogeneous distribution of deepwater ϵ_{Nd} values, and Nd isotopes have been used as a tracer of ocean circulation in the present and past ocean (Albarède and Goldstein, 1992; von Blanckenburg, 1999; Frank, 2002; Goldstein and Hemming, 2003; Thomas, 2004; Robinson et al., 2010; Lacan et al., 2012).

This basic idea is however oversimplified, and the knowledge on specific processes/mechanisms controlling oceanic ϵ_{Nd} distribution has been progressing. The net external Nd sources to the ocean have been shown to be partially dissolved dust and river particles as well as river water (Tachikawa et al., 1999a; von Blanckenburg, 1999; Frank, 2002; Goldstein and Hemming, 2003). In addition, the contribution of groundwater is suggested (Johannesson and Burdige, 2007) whereas hydrothermalism functions as sink of oceanic Nd (Zheng et al., 2016), although it was hypothesized that it could modify the local seawater isotopic signature through “Ridge exchange” processes (Jeandel et al., 2013).

It is however impossible to reproduce the observed seawater ϵ_{Nd} contrast between the deep Atlantic and the deep Pacific only accounting for these net continental sources and physical mixing of water masses (Tachikawa et al., 2003; Jones et al., 2008). Reversible scavenging, an interaction of Nd between dissolved and particulate phases, is a fundamental process for the vertical transport of the Nd isotopic signature (Bertram and Elderfield, 1993; Jeandel et al., 1995; Tachikawa et al., 1999b). In particular, the reaction along continental margins called “Boundary Exchange (BE)” is a key process and supported by both field observation (Lacan and Jeandel, 2005; Rickli et al., 2009; Rickli et al., 2010; Carter et al., 2012; Stichel et al., 2012; Wilson et al., 2012; Grenier et al., 2013) and modeling experiments (Arsouze et al., 2007; Siddall et al., 2008; Arsouze et al., 2009; Rempfer et al., 2011). Nonetheless, the exact mechanisms of BE remain to be elucidated (Jeandel, 2016), and its geographical extension should be better constrained since in such areas seawater ϵ_{Nd} values are not a straightforward water mass tracer. Recently, the first measurements of pore water Nd isotopic composition suggested the contribution of radiogenic pore water Nd to shallow and intermediate waters on the Northwestern Pacific margin (Abbott et al., 2015; Abbott et al., 2016). Also, a possible influence of unradiogenic Nd from poorly chemically weathered detrital material on bottom water was proposed for the Northern northwestern Atlantic (Howe et al., 2016b). These findings may explain different aspects of BE.

Over the last several decades, the interest in Nd isotopes has been increasing with a growing body of high quality and highly resolved data, thanks to the GEOTRACES program, and the identification of archives that can record and preserve seawater Nd isotopic signatures. Indeed, Nd isotopes are a rare tracer that allows reconstructing past oceanic circulation on timescale ranging from Jurassic to late

Pleistocene (Puceat et al., 2005; Gourelan et al., 2008; Dera et al., 2009; van de Flierdt and Frank, 2010; Charbonnier et al., 2012; Martin et al., 2012; Moiroud et al., 2013; Thomas et al., 2014; Le Houedec et al., 2016 and Frank, 2002; Goldstein and Hemming, 2003 for review) using various archives.

Biogenic apatite (fish teeth/debris) has been considered one of the most reliable recorders of seawater Nd isotopic compositions (Martin and Haley, 2000; Martin and Scher, 2004; Martin et al., 2010; Horikawa et al., 2011) although the occurrence of this archive is not constant and barren samples are not uncommon. The use of deep-sea corals is relatively recent. The advantages are the possibility of precise U-Th dating and high-resolution reconstruction (Robinson and van de Flierdt, 2009; Colin et al., 2010; Copard et al., 2010; van de Flierdt et al., 2010; Copard et al., 2011; Wilson et al., 2014). However, certain time intervals are bare of this archive under hostile environmental conditions for the organism (Frank et al., 2011). Bulk carbonate fraction (Gourelan et al., 2008; Le Houedec et al., 2016) is also used for carbonate-rich sediments.

Sedimentary dispersive Fe-Mn oxyhydroxides extracted with various leaching techniques have been widely used because of their ubiquitous presence in marine sediments and temporal resolution to possibly resolve glacial-interglacial and millennial changes (Rutberg et al., 2000; Bayon et al., 2002; Piotrowski et al., 2004; Piotrowski et al., 2005; Gutjahr et al., 2007; Gutjahr et al., 2008; Haley et al., 2008a; Haley et al., 2008b; Pahnke et al., 2008; Piotrowski et al., 2009; Crocket et al., 2011; Gutjahr and Lippold, 2011). A potential difficulty of this archive is the possible contamination from labile terrigenous fraction such as preformed continental Fe-Mn oxyhydroxides and volcanogenic material (Bayon et al., 2004; Roberts et al., 2010; Elmore and Wright, 2011; Piotrowski et al., 2012; Ehlert et al., 2013; Kraft et al., 2013; Wilson et al., 2013; Wu et al., 2015b; Abbott et al., 2016; Blaser et al., 2016). To minimize the risk, sedimentary oxyhydroxides attached on foraminiferal tests have been used to reconstruct bottom water Nd isotopic compositions since fine detrital fraction can be mechanically removed from foraminiferal tests (Roberts et al., 2010; Elmore and Wright, 2011; Charbonnier et al., 2012; Pena and Goldstein, 2014; Tachikawa et al., 2014; Wu et al., 2015b). However, it has been recently proposed that sedimentary oxyhydroxides record pore water rather than bottom water Nd isotopic composition, so that the observed discrepancies between archives and seawater isotopic signature could be explained at least partly by the different ϵ_{Nd} values between bottom and pore waters (Du et al., 2016).

The main challenges to further improve the application of Nd isotopes to modern and paleo-oceanography studies are (1) to estimate the spatial extension of areas where seawater ϵ_{Nd} values are highly affected by local/regional detrital inputs and (2) to evaluate the faithfulness of archive ϵ_{Nd} values as an indicator of bottom water masses. For these objectives, we have updated a database of the available seawater Nd isotopic compositions, and compiled ϵ_{Nd} values for foraminiferal coatings, deep-sea corals, fish teeth/debris and various leachates from the Holocene period ($< 10,000$ years) in the framework of the French INSU/LEFE project NEOSYMPA (Workshop NEODYMIUM isotopes in marine environments: SYnergy between Modern, Modeling and PAleo communities). The seawater and archive database contains hydrography parameters (temperature, salinity, silicate, phosphate and oxygen concentrations) from original publications and World Ocean Atlas 2009 (WOA09) (Antonov et al., 2010; Garcia et al., 2010a; Garcia et al., 2010b; Locarnini et al., 2010), carbon isotopic ratio ($^{13}\text{C}/^{12}\text{C}$ or $\delta^{13}\text{C}$) of dissolved inorganic carbon (DIC) from the global $\delta^{13}\text{C}$ compilation (Schmittner et al., 2013) and natural (background or pre-bomb) $\Delta^{14}\text{C}$ values from GLODAP database (Key et al., 2004). The database contains also the distance of each data station from the nearest continental margin. Using this new database, we provide novel insight into oceanic Nd cycle.




Fig. 1. Location map of all stations with ϵ_{Nd} data together with vertical distribution of seawater and archives data stations. Black dots indicate seawater stations whereas open circles show various archive stations with color code. The map is visualized using the software Ocean Data View (ODV) (Schlitzer, 2015).

2. Construction of NEOSYMPA global database of seawater and archives Nd isotopes

The seawater Nd database of Lacan et al. (2012) has been updated by integrating all seawater ϵ_{Nd} data published up to September 2016 to our knowledge (Fig. 1) (Bayon et al., 2011; Grenier et al., 2011; Carter et al., 2012; Grasse et al., 2012; Singh et al., 2012; Stichel et al., 2012; Amakawa et al., 2013; Chen et al., 2013b; Ehlert et al., 2013; Jeandel et al., 2013; Freslon et al., 2014; Garcia-Solsona et al., 2014; Goswami et al., 2014; Haley et al., 2014; Huang et al., 2014; Werner et al., 2014; Abbott et al., 2015; Basak et al., 2015; Stichel et al., 2015; Wu et al., 2015a; Du et al., 2016; Dubois-Dauphin et al., 2016a; Fröllje et al., 2016; Lambelet et al., 2016; Dubois-Dauphin et al., 2017). Following the approach of the previous compilation (Lacan et al., 2012), we focus on seawater Nd isotopic compositions. Dissolved (filtered and unfiltered) Nd concentrations are included in the database if they are reported with corresponding isotopic results. Considering the general agreement of seawater Nd isotopic compositions obtained by various analytical techniques during different time spans (Lacan et al., 2012; van de Flierdt et al., 2012), no distinction was made between analytical methods. Some Nd data from low salinity waters (salinity 5–30 psu) are included in the database although the major focus of this study is on waters with salinity ≥ 30 psu. The vertical distribution of seawater stations is heterogeneous with more abundant data for surface/subsurface (Fig. 1).

In contrast to the previous compilations (Lacan et al., 2012; van de Flierdt et al., 2016) and an on-going construction of GEOTRACES database, NEOSYMPA database includes $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ values of DIC extracted from global databases (Key et al., 2004; Schmittner et al., 2013) as well as lateral distances from the data stations to the nearest continental margins at the ocean surface, in addition to hydrography parameters (temperature, salinity, dissolved silica, phosphate and oxygen concentrations). If the hydrographical parameters are not available in original publications, they were extracted from the closest station in WOA09 database (Antonov et al., 2010; Garcia et al., 2010a; Garcia et al., 2010b; Locarnini et al., 2010). The hydrography para-

meters from WOA09 and original publications are in good agreement except for temperature and salinity at some coastal stations such as the Amazon river mouth, Bay of Bengal, Gulf of Alaska, Greenland coast and Japan Sea, as well as dissolved silicate and phosphate at two sites in the Southeast Pacific at 650 m and 1500 m (Fig. S1).

NEOSYMPA database (Table S1) contains ϵ_{Nd} values obtained on sedimentary archives and deep-sea corals (referred as to “archival ϵ_{Nd} values” below) from the following publications (Palmer and Elderfield, 1985; Vance and Burton, 1999; Burton and Vance, 2000; Rutberg et al., 2000; Freydier et al., 2001; Bayon et al., 2002; Bayon et al., 2004; Piotrowski et al., 2004; Scrivner et al., 2004; Tachikawa et al., 2004; Vance et al., 2004; Gutjahr et al., 2007; Stoll et al., 2007; Gutjahr et al., 2008; Haley et al., 2008a; Haley et al., 2008b; Klevenz et al., 2008; Pahnke et al., 2008; Piotrowski et al., 2008; Yu et al., 2008; Piotrowski et al., 2009; Robinson and van de Flierdt, 2009; Basak et al., 2010; Colin et al., 2010; Copard et al., 2010; Gourelan et al., 2010; Murphy and Thomas, 2010; Roberts et al., 2010; Stumpf et al., 2010; van de Flierdt et al., 2010; Bayon et al., 2011; Copard et al., 2011; Elmore et al., 2011; Horikawa et al., 2011; Asahara et al., 2012; Carter et al., 2012; Chen et al., 2012; López Correa et al., 2012; Piotrowski et al., 2012; Wilson et al., 2012; Xie et al., 2012; Haley and Polyak, 2013; Jang et al., 2013; Kraft et al., 2013; Maccali et al., 2013; Montero-Serrano et al., 2013; Noble et al., 2013; Pena et al., 2013; Skinner et al., 2013; Achyuthan et al., 2014; Huang et al., 2014; Molina-Kescher et al., 2014; Werner et al., 2014; Jiménez-Amat and Zahn, 2015; Jonkers et al., 2015; Roberts and Piotrowski, 2015; Abbott et al., 2016; Blaser et al., 2016; Du et al., 2016; Dubois-Dauphin et al., 2016b; Howe et al., 2016a; Hu et al., 2016; Lang et al., 2016; Lippold et al., 2016; Meinhardt et al., 2016; Wei et al., 2016) (Fig. 1). Each archival value is accompanied by all the above-mentioned auxiliary data from various sources (Key et al., 2004; Antonov et al., 2010; Garcia et al., 2010a; Garcia et al., 2010b; Locarnini et al., 2010; Schmittner et al., 2013) and distances from the continents.

We classified the archives into seven categories: sedimentary foraminiferal coatings (referred as to “foraminifera” below), dispersive Fe-Mn oxyhydroxides obtained with hydroxylamine hydrochloride

(HH) leaching after decarbonation, HH leachate obtained from bulk sediments, acetic acid (AA) leachate from bulk sediments, other leachates (for example, HCl leaching), deep-sea corals, and fish teeth/debris. Regarding to the HH leachates, the presence and absence of decarbonation is distinguished because this step induces significant differences in Nd isotopic composition, and recent studies recommend omitting decarbonation (Wilson et al., 2012; Kraft et al., 2013; Wu et al., 2015b; Blaser et al., 2016). To improve the spatial coverage of archive data, we gathered all data with age < 10,000 years, even if we are aware that seawater ϵ_{Nd} values could vary during Holocene as shown by deep-sea corals from the North Atlantic (Colin et al., 2010; Montero-Serrano et al., 2011; Copard et al., 2012). When the evidence of age < 10,000 years is provided, the data are included in the database in spite of the absence of absolute age. The archive data are better distributed vertically compared to seawaters without a predominant surface maximum (Fig. 1).

Both seawater and archive data are classified into seven oceanic basins (Atlantic, Southern Ocean, Indian, Pacific, Arctic and Mediterranean Sea). Our focus of this paper is on the open oceans that create the global general trends. Because of this reason, we treat essentially the data from the Atlantic, the Pacific, the Indian and the Southern Oceans.

3. Results

3.1. General trends of seawater Nd variability

Depth profiles of seawater Nd concentration, seawater ϵ_{Nd} and archive ϵ_{Nd} values are shown for the Atlantic, Southern Ocean, Indian and Pacific with latitudinal trend by color code (Fig. 2). Marked surface water Nd enrichment up to 150 pmol/kg is observed in the North Atlantic, North Indian and North Pacific (Fig. 2). In the North Pacific, seawater Nd concentration is systematically higher than the values in the South Pacific at corresponding latitudes and water depths (Fig. 2). Seawater ϵ_{Nd} depth-profiles present a larger variability at shallower water depths in the Atlantic, the Indian and the Pacific (Fig. 2). Except very radiogenic values ($\epsilon_{Nd} > -5$) at water depths shallower than 500 m (referred as to “< 500 m” below) in the North Atlantic, the Atlantic waters indicate the latitudinal ϵ_{Nd} trend with lower values in the northern North Atlantic (ϵ_{Nd} centered at around -13 at 50°N to 75°N), middle values at mi-latitudes (ϵ_{Nd} of around -12 at 0 to 30°N), and higher values of around -9 at southern hemisphere (Fig. 2). In contrast, the Pacific ϵ_{Nd} values are lower in the southern hemisphere with a mean ϵ_{Nd} value of around -8 at ≥ 500 m at 60°S to 20°S than in equatorial and northern hemisphere with ϵ_{Nd} of about -4 at ≥ 500 m




Fig. 2. Depth profiles of the compiled data in different oceanic basins. Seawater Nd concentrations (upper panels), seawater ϵ_{Nd} values (middle panels), and archive ϵ_{Nd} values (lower panels) for the Atlantic, Southern Ocean, Indian and Pacific. The color code indicates the latitudes of the data stations. The profiles are created using ODV (Schlitzer, 2015).

at 0 to 75°N (Fig. 2).

These observed features are consistent with the conventional view that the general increase of deepwater ϵ_{Nd} along the global thermohaline circulation pathway is caused by a mixing between young North Atlantic waters with low ϵ_{Nd} values and old Pacific waters with high ϵ_{Nd} values. Deepwater ϵ_{Nd} values (≥ 1500 m) correlate well with dissolved silica and phosphate concentrations (Fig. 3). The South Atlantic, the Southern Ocean, the South Indian and the South Pacific Ocean waters present intermediate values (Fig. 3). However, a conventional binary mixing trend is not evident in seawater ϵ_{Nd} versus $1/[Nd]$ plot (here $[Nd]$ is dissolved Nd concentration of seawater) and the highest ϵ_{Nd} values are observed not only for the deep water in the north Pacific but also in the equatorial Pacific (Fig. 3). Taking into account general covariation between seawater ϵ_{Nd} values and other water mass tracers at water depths deeper than 1500 m (Figs. 3), deepwater Nd isotopic composition generally reflects water mass mixing but significant local/regional effects exist for some cases. The lack of the conventional binary mixing (ϵ_{Nd} vs. $1/[Nd]$) thus indicates principally non-conservative nature of dissolved Nd concentration.

The property-property plots between ϵ_{Nd} and other water mass indicators (multi-scatter plots) allow us to visually identify the data that are offset from the general mixing envelopes, such as the values from the Caribbean Sea (number 1 in Fig. 3), Baffin Bay (number 2 in Fig. 3) and Bay of Bengal (number 3 in Fig. 3). This is consistent with the statement of original papers suggesting influence of local detrital inputs (Stordal and Wasserburg, 1986; Singh et al., 2012; Osborne et al., 2014). In the following, we use basin-scale multi-scatter plots to examine the regional variance in each oceanic basin in detail.

3.2. Basin-scale seawater and archive Nd isotopic variability

3.2.1. Seawater multi-scatter plots for deep and intermediate waters

We test the relationships of water masses at ≥ 1500 m to avoid difficulties related to definition of water mass end-members. For the first approximation, offsets from the general trend are evaluated visually and considered to be significant when more than two property-property plots support the departure.

In the Atlantic and the Atlantic sector of the Southern Ocean (Fig. 4a), the general mixing envelopes are formed between younger, warmer, saltier nutrient-depleted and oxygen-rich northern-sourced deep waters with lower ϵ_{Nd} , and older, cooler, fresher nutrient-rich and oxygen-depleted southern-sourced waters with higher ϵ_{Nd} . Offsets from the general mixing are observed for the stations in the Caribbean Sea (number 1 in Fig. 4a), the Labrador Sea (number 2, Fig. 4a), along the eastern Greenland coast (number 4, Fig. 4a), off Amazon River mouth (number 3, Fig. 4a) and a site in Southeastern Atlantic (number 5, Fig. 4a), as well as warm and saline Mediterranean Outflow Water (MOW). In contrast, silicate and phosphate data of these sites are on the general mixing trend (Fig. 4a), suggesting that the observed seawater ϵ_{Nd} offsets are related to specific behavior of Nd. Clear mixing trend is not expressed in the relationship with $1/[Nd]$ and with potential density anomaly (not shown in figure).

The spatial data coverage in the Indian Ocean is not sufficient to determine the mixing envelopes (Fig. 4b). We observe regional trends with the most radiogenic seawater ϵ_{Nd} values of -6 to -4 in the easternmost part of the Equatorial Indian Ocean originating from the Sunda Arch Slope (Jeandel et al., 1998) that shows properties close to




Fig. 3. Seawater multi-scatter plots for data at water depths deeper than 1500 m in the Atlantic, Southern Ocean, Indian and Pacific. All the data to create the plots are compiled in NEOSYMPA database. The numbers on the map show the area where seawater ϵ_{Nd} values are offset from the general trend (see text for detail). The figures and the map are created using ODV (Schlitzer, 2015).

those of Pacific waters (Fig. 3) whereas the waters of ϵ_{Nd} as low as -12 are found in the Bay of Bengal that receive unradiogenic Nd from the Ganga–Brahmaputra river system (Singh et al., 2012) (Fig. 4b). The Arabian Sea waters at 1500–1600 m (numbers 1 and 2 in Fig. 4b) are warmer and saltier than other Indian deep waters.

In the Pacific Ocean and the Pacific sector of the Southern Ocean, deepwater ϵ_{Nd} values correlate well with other water mass indicators, forming mixing envelopes between the younger and colder southern-sourced waters with lower ϵ_{Nd} values of around -8 , and the older and warmer northern and equatorial-sourced waters with higher ϵ_{Nd} values centered at around -4 (Fig. 4c). Positive ϵ_{Nd} offsets from the general trend are observed close to Hawaii (number 1, Fig. 4c), along the eastern Japan coast (number 2, Fig. 4c), in the eastern equatorial Pacific (number 3, Fig. 4c) and Haxby seamount in the Southern Ocean (number 6, Fig. 4c) whereas negative offsets are found east of New Zealand (number 4, Fig. 4c) and in the central North Pacific bottom water (number 5, Fig. 4c).

Basin-scale multi-scatter plots for the intermediate waters (600–1500 m) present weaker relationships in all basins (Fig. S2abc). The geographical distribution of sites where seawater ϵ_{Nd} values are away from the general tendencies is similar to the deepwater cases with some additional sites including lower ϵ_{Nd} values in the Angola Basin (number 2, Fig. S2a) and the Bay of Bengal (white circle in Fig. S2S2b) and higher ϵ_{Nd} values in the Western equatorial Pacific (number 5, Fig. S2c).

3.2.2. Comparison between archive and seawater Nd isotopic compositions

Basin-scale mixing trends defined by seawater multi-scatter plots allows visual comparison between archive and seawater ϵ_{Nd} values at ≥ 600 m even if the sampling locations and water depths are not identical for the two types of samples (Fig. 5abc and Fig. S4abc).

For the deep (≥ 1500 m) Atlantic and the Atlantic sector of the Southern Ocean, the majority of archival ϵ_{Nd} values fall within the domains defined by seawater data except the northern North Atlantic where both positive and negative offsets as large as 10 ϵ -units are observed (the large circle on the plots and the red and bleu stations on the map of Fig. 5a). The most radiogenic archival ϵ_{Nd} values are found around Iceland and downstream the Iceland-Scotland Overflow Waters at about 40°N (Elmore et al., 2011) whereas an about 2 ϵ -unit higher ϵ_{Nd} value is observed at a southernmost site at around 55°S on the Mid-Ocean Ridge (number 5 in Fig. 5a). Three sites from 50°N to 34°N along Deep Western Boundary Current (numbers 1, 2 and 4 in Fig. 5a), and a site in the Angola Basin (number 3 in Fig. 5a) present ϵ_{Nd} values lower than expected from the seawater domain (Fig. 5a). The archive types showing the offsets are diverse including foraminifera, HH leachate with decarbonation, fish teeth/debris and AA leachate (Fig. 5a).

In the deep Indian Ocean, more negative ϵ_{Nd} values (about 2 to 4 ϵ -units) than the expected seawater signals are found on the northern margin of Madagascar (number 1 in Fig. 5b) for foraminifera, HH leachate with and without decarbonation, being consistent with the original publication (Wilson et al., 2012).

In the deep Pacific and the Pacific sector of the Southern Ocean, the archive ϵ_{Nd} values generally follow the mixing envelopes although they




Fig. 4. Seawater property-property plots for water depths deeper than 1500 m. The color code shows latitudes (a and c) or longitude (b) of data stations. a) Atlantic and Atlantic sector of the Southern Ocean, b) Indian, and c) Pacific and Pacific sector of the Southern Ocean. MOW = Mediterranean Outflow Water. The numbers on the maps show the sites where seawater ϵ_{Nd} values are offset from the general trend (see text for detail). The figures and the maps are created using ODV (Schlitzer, 2015).




Fig. 4. (continued)

are prone to occupy the domains at upper limits of seawater ϵ_{Nd} values (Fig. 5c), as shown by the recent study (Hu et al., 2016). Clear positive ϵ_{Nd} offsets up to 4 ϵ -units are observed for the Gulf of Alaska (numbers 5 to 10 in Fig. 5c), the Eastern Equatorial Pacific (numbers 1 to 3 in Fig. 5c), near Hawaii (number 4 in Fig. 5c), for foraminifera and HH leachate with and without decarbonation.

The archival ϵ_{Nd} values at intermediate water depths (600–1500 m) present a larger range of variability than for the deeper depths (Fig. S3abc). In the Atlantic, one HH leachate with decarbonation from the eastern Greenland coast where ϵ_{Nd} of continental margin is very high (Jeandel et al., 2007) presents Nd isotopic composition about 10 ϵ -units higher than the water value (number 1 in Fig. S3a). In the Indian, only one archive ϵ_{Nd} value is available and it is within the seawater ϵ_{Nd} range (Fig. S3b). In the Pacific and the Pacific sector of the Southern Ocean, archive ϵ_{Nd} values are generally within the seawater domain except about 2 ϵ -units more radiogenic data obtained in the Eastern Equatorial Pacific (numbers 1 to 3 in Fig. S3S3c), Amundsen Sea embayment (number 4 in Fig. S3S3c) and Bering Sea (number 5 in Fig. S3S3c).

3.2.3. T-S- ϵ_{Nd} diagrams for seawater and archives

In order to associate ϵ_{Nd} values with different water masses and evaluate the basin-scale isotopic evolution at ≥ 200 m, we use temperature/salinity diagram combined with seawater and archive ϵ_{Nd} values (T-S- ϵ_{Nd} diagram). In T-S diagram, the ϵ_{Nd} values are shown with color code (Fig. 6). The identification of water masses is based on temperature and salinity ranges and the geographical extension (Emery and Meincke, 1986; Amakawa et al., 2009; Bostock et al., 2013). The water masses described below are listed in Table 1 and Table S2.

In the Atlantic, the T-S- ϵ_{Nd} diagram undoubtedly indicates a

latitudinal variation of seawater Nd isotopic compositions with unradiogenic northern water masses including NADW and various central waters, and more radiogenic southern-sourced waters such as AABW and AAIW (Fig. 6a). The mean seawater Nd isotopic composition and the mean Nd concentration of each water mass (Table 1) can be estimated at a given temperature and salinity range, and the geographical extension of the water mass (Table S2). The ϵ_{Nd} value and the dissolved Nd concentration of NADW are estimated at -12.3 ± 0.9 (1σ , $n = 173$) and 20.3 ± 3.2 (pmol/kg) (1σ , $n = 135$), respectively (Table 1). This NADW ϵ_{Nd} value is in an excellent agreement with a recent estimation of -12.4 ± 0.4 (2σ) (Lambelet et al., 2016) that focuses on the North western Atlantic data. AABW and AAIW in the South Atlantic are characterized by the ϵ_{Nd} values of -8.6 ± 0.6 (1σ , $n = 39$, Table 1) and -8.6 ± 1.4 (1σ , $n = 28$), respectively (Table 1). The archive data are concentrated on the mixing line between AABW and NADW, and ϵ_{Nd} values close to the core of NADW are often higher than corresponding seawater data (Fig. 6b). This is consistent with the observation based on the property-property plots (Fig. 5a).

In the Indian Ocean (Fig. 6c), the most radiogenic seawater Nd isotopic compositions of -5 to -4 are observed for Indonesian Intermediate Water (IIW) at intermediate water depths that occupies easternmost equatorial basin (Emery and Meincke, 1986). An array of unradiogenic ϵ_{Nd} values of about -12 at salinity of around 35 (Fig. 6c) is found in the Bay of Bengal. The mean ϵ_{Nd} values for CDW and AAIW in the South Indian are -7.9 ± 1.3 (1σ , $n = 10$, Table 1) and -7.6 ± 0.9 (1σ , $n = 4$, Table 1), respectively. Archive ϵ_{Nd} data are very limited in the Indian Ocean and cover only a few water masses (Fig. 6d).

The most striking feature in the Pacific Ocean is very high seawater




Fig. 4. (continued)

ϵ_{Nd} values for EqPIW (Bostock et al., 2013) at 500–1500 m and PEW at 0–500 m (Emery and Meincke, 1986) (Table 1 and Fig. 6e). NPIW is also characterized by an elevated Nd isotopic composition (Table 1 and Fig. 6e). The mean ϵ_{Nd} values of EqPIW, PEW and NPIW are -3.0 ± 1.4 (1σ , $n = 31$), -2.1 ± 0.9 (1σ , $n = 72$) and -3.4 ± 1.3 (1σ , $n = 12$), respectively (Table 1). The deeper water and southern-sourced waters have lower mean ϵ_{Nd} values with -4.3 ± 1.2 (1σ , $n = 84$), -7.7 ± 1.1 (1σ , $n = 21$) and -7.8 ± 1.2 (1σ , $n = 92$) for NPDW, AAIW and CDW in the Pacific, respectively (Table 1). EqPIW is a mixture of AAIW and upwelled old NPDW (Bostock et al., 2013). Since the Nd isotopic composition of AAIW and NPDW are lower than the value of EqPIW (Fig. 6e, Table 1), additional radiogenic Nd supply is required (Lacan and Jeandel, 2001; Grenier et al., 2013; Grenier et al., 2014). The NPIW formation is mainly density driven by sinking in the Okhotsk Sea (Yasuda et al., 1996), and its high ϵ_{Nd} value is explained by a large contribution of radiogenic Nd supply from the Kuril Islands (Amakawa et al., 2004). Consequently EqPIW, PEW and NPIW are influenced by detrital inputs. The archive T-S- ϵ_{Nd} diagram presents ϵ_{Nd} trends similar to the seawater T-S- ϵ_{Nd} diagram with radiogenic EqPIW and PEW although the mean Nd isotopic compositions are about 1 ϵ -unit higher than the corresponding seawater values (Fig. 6f).

4. Discussion

The deepwater Nd isotopic compositions correlate with other water mass properties (Figs. 3 and 4abc) whereas shallower waters show the weaker relationships (Fig. S2abc) with larger ϵ_{Nd} variability (Fig. 2). Linear correlations between tracers indicate mixing if the tracer is

conservative. Linear relationships with non-conservative tracers (i.e. nutrients) suggest that whatever mechanisms control one variable are also likely the mechanisms that control the other. In modern deep waters, nutrient concentrations vary essentially with water mass mixing (Broecker and Peng, 1982). Almost linear relationships between deepwater ϵ_{Nd} values with hydrography parameters were already reported (Goldstein and Hemming, 2003; Piotrowski et al., 2008; Hu et al., 2016), which is consistent with our observation (Figs. 3 and 4abc). Our hypothesis here is that deepwater ϵ_{Nd} variability that is explained by linear relationships with hydrography parameters corresponds to the mixing of water masses of which ϵ_{Nd} values are distinct. Large offsets from the predicted values indicate that other processes than the ones governing the large-scale tracer evolution are at play. For the case of Nd isotopic compositions, we consider this to be a sign of significant local/regional influence.

4.1. Empirical equations that predict deepwater ϵ_{Nd} trends and evaluation of local/regional influence

Applying multiple regression analysis to deep (≥ 1500 m) seawater data, we establish empirical equations that predict the main, large-scale, deepwater ϵ_{Nd} trends from hydrography parameters. We use hydrography data WOA09 (Antonov et al., 2010; Garcia et al., 2010a; Garcia et al., 2010b; Locarnini et al., 2010) because they are available for the major oceans with 1° resolution and indicate a good agreement with values compiled in NEOSYMPA database (Fig. S1).

For the modern global deep waters, nutrients and oxygen concentrations present a bimodal distribution with a peak corresponding to a nutrient-poor and oxygen-rich Atlantic water (except for the southern-

most Atlantic) and another peak corresponding to deep waters of other oceanic regions with higher nutrient contents and more depleted oxygen (Fig. S4S4). As a consequence, the Atlantic (50°S to 75°N) is treated separately from the global ocean for multiple regression analysis. Only the variables showing a normal distribution (examined mainly with Kolmogorov-Smirnov/Lilliefors Test) are used for regression analysis. The number of variables is optimized based on Akaike's Information Criterion (Motulsky and Christopoulos, 2004) (Table S3ab). Data from restricted basins such as Caribbean Sea, Labrador Sea and Bay of Bengal are not considered for the regression analysis because of the ϵ_{Nd} values decoupled from the general trend (Fig. 3).

Established empirical relationships for the Atlantic (50°S to 75°N) and other global oceans are expressed by Eqs. (1) and (2), respectively:

$$\epsilon_{Nd} = -0.7372 T [^{\circ}\text{C}] - 0.9385 O_2 [\text{ml/l}] - 4.5207 \quad (r = 0.705) \quad (1)$$

$$\begin{aligned} \epsilon_{Nd} = & 2.252 * PO_4 [\mu\text{mol/l}] + 0.04635 * SiO_2 [\mu\text{mol/l}] - 9.958 \\ & * Salinity + 0.2861 * T [^{\circ}\text{C}] + 326.95 \quad (r = 0.845) \quad (2) \end{aligned}$$

They are the first equations that allow calculating deepwater Nd isotopic compositions at ≥ 1500 m. Fig. 7 compares the predicted and observed deepwater ϵ_{Nd} values in the Atlantic and the Atlantic sector of the Southern Ocean (Fig. 7ab), as well as the Indian, the Pacific and the Pacific sector of the Southern Ocean (Fig. 7cd). Large-scale isotopic

variation including the penetration and the extension of northern- and southern-sourced waters are well demonstrated in zonal-average meridional transect in both Atlantic and Pacific (Fig. 7bd). These results confirm that the major deepwater Nd isotopic variation can be explained by the mixing of preformed tracers. However, there are some significant offsets between predicted and observed seawater Nd isotopic signatures in the eastern and western equatorial Pacific where the observed values are higher than the predicted Nd isotopic compositions (Fig. 7cd). Also, predicted values for the MOW is anomaly low because of warmer temperature of this water mass (Eq. (1)) relative to other Atlantic waters (not shown in figure). Except for this specific artifact, any systematic bias is noticed. The mean difference between observed and predicted ϵ_{Nd} values is almost zero (Fig. 8) and residual values present a normal distribution (Fig. 8). The global performance of the multiple regression analysis is shown in Fig. 8.

We now define that significant local/regional detrital influence by a difference between observed and predicted ϵ_{Nd} values, $\epsilon_{Nd}(\text{observed-predicted})$. The values out of range of 2σ (Fig. 8) is considered as a sign of local/regional detrital influence, which corresponds roughly to $\epsilon_{Nd}(\text{observed-predicted})$ values $< +2$ ϵ -units or > -2 ϵ -units (Fig. 8). The predicted ϵ_{Nd} values, the difference between measured and predicted isotopic composition, as well as judgment about significant local/regional influence are all listed in Table S1.




Fig. 5. Basin-scale archive multi-scatter plots at ≥ 1500 m. a) Atlantic, b) Indian, and c) Pacific and Pacific sector of the Southern Ocean. The red and blue stations on the maps indicate the sites with positive and negative offsets, respectively. The color code shows different archive types and gray dots indicate seawater values shown in Fig. 4abc. There is no archive data from deepwater depths (≥ 1500 m) in the Atlantic and Indian sectors of the Southern Ocean. The numbers on the map show the area where seawater ϵ_{Nd} values are offset from the general trend (see text for detail). The figures and the maps are created using ODV (Schlitzer, 2015).




Fig. 5. (continued)

4.2. Spatial distribution of sites affected by local/regional detrital inputs

The distribution of sites with significant local/regional influence on seawater ϵ_{Nd} values (Fig. 9a) is generally consistent with the estimation based on multi-scatter plots (Fig. 4abc). The regression results allow identifying more detail, in particular for the Indian Ocean where the poor spatial coverage of data prevents from establishing general mixing trends (Fig. 4b). Applying the same criteria of local/regional effects as the seawater samples to archive ϵ_{Nd} data, the distribution of locally affected archive sites is determined (Fig. 9c). Again, it generally agrees with the results of multi-scatter plots (Fig. 5abc). It is worth noting that there exists certain similarity of locations between the seawater and archive sites affected by local/regional influence, for instance the Eastern Equatorial Pacific (labeled “a” in Fig. 9ac), around Hawaii (“b” in Fig. 9ac) and close to Iceland (“c” in Fig. 9ac) with positive ϵ_{Nd} offsets, and Bay of Bengal (“d” in Fig. 9ac) and central North Pacific (“e” in Fig. 9ac) with negative bias. The positive offsets observed for the western equatorial Pacific seawater in Fig. 7c do not appear in Fig. 8a because the nutrient data for these coastal sites are not available, which prevents the application of Eq. (2).

The relationships between the water depths and the distance from the margins reveal that the main detrital impact occurs roughly within 1000 km from the margins (Fig. 9bd). For seawater samples, 61 data are evaluated to be affected by local/regional effects, and 57 data are from the sites within 1000 km from the margins, which corresponds to 93%. Totally 195 archive data are estimated to be influenced by detrital inputs, and 179 data (92%) are from sites within 1000 km from the

margins. Several seawater samples (labeled “f” and “g” in Fig. 9ac) show the offsets far from the continental margins. At station GYR in the Southeastern Pacific (“f” in Fig. 9ab), the radiogenic seawater Nd signature was explained by ridge exchange above the East Pacific Rise (Jeandel et al., 2013). Similarly, the positive offset in the South Atlantic (station S4, “g” in Fig. 9ac) may reflect the influence of radiogenic Mid Atlantic Ridge although the contribution of the South American shelf sediments and of the Antarctic Peninsula sediments could not be discarded (Garcia-Solsona et al., 2014).

The archives present more frequently positive offsets than negative ones (Fig. 9cd). There are two ways of explaining this feature. Firstly, the dissolution of labile detrital Nd during leaching procedures could produce such biases (Elmore et al., 2011; Piotrowski et al., 2012; Ehlert et al., 2013; Kraft et al., 2013; Wu et al., 2015b) and consequently the observed offsets could be an artifact. Secondly the archive ϵ_{Nd} values reflect pore water Nd signals that are more radiogenic than the overlying bottom waters (Wilson et al., 2013; Abbott et al., 2015; Molina-Kescher et al., 2014; Wu et al., 2015b; Blaser et al., 2016; Du et al., 2016). Both explanations are plausible, and the prevailing reason may vary with regions. It is also possible that some archive ϵ_{Nd} offsets reflect the real temporal variation of deepwater ϵ_{Nd} values. In fact, some mid- or early Holocene data indicate significant difference from the predicted modern seawater values (Fig. S5). At this stage it is difficult to compare the faithfulness of different archives since data number and sampling locations strongly vary with archive types (Fig. 1 and Fig. S5). Nonetheless, we tentatively suggest that the HH leachate with decarbonation tends to be more frequently biased (Fig. S5, see also Section 4.4).




Fig. 5. (continued)

4.3. Possible processes modulating local/regional detrital inputs

Possible provenance of detrital inputs can be deduced from the Nd isotopic composition. In general, the difference between observed and predicted seawater ϵ_{Nd} values is positive for the circum Pacific and negative in the Labrador Sea and the Bay of Bengal (Fig. 8). This geographical trends is consistent with the Nd isotopic signature of continental margins (Jeandel et al., 2007). In addition, the influence of remote dust source may have influence as suggested by negative offsets at around 30°N in the Pacific (Fig. 9a) that underlies loess dust plume (Maher et al., 2010). The effective influence of detrital inputs depends on relative strength between physical water mass mixing and dissolved/particulate interaction (Roy-Barman et al., 2009; Osborne et al., 2015), stock of exchangeable Nd in particulate phases, as well as the contrast of Nd isotopic signature between dissolved and particulate phases.

The strong local effects close to the continental margins can be explained by the proximity of the sources as well as high detrital and biogenic particle concentrations in the water column that can promote reversible scavenging. A leading role of particulate phases as a vertical vector is further confirmed by another particle reactive tracer, Th isotopes with a recent GEOTRACES study in the North Atlantic (Lerner et al., 2016 and references therein). Indeed, coupled measurements of Ra and Nd isotopes as well as shale-normalized REE patterns on seawater samples from sites close to Hawaii demonstrate that radiogenic Nd from volcanic islands has a strong influence on surface water isotopic signature, and that the radiogenic Nd signal is vertically

transported by dissolved/particulate interaction (Fröllje et al., 2016). In contrast, seasonally variable dust contribution is of secondary importance but affects significantly the surface waters (Fröllje et al., 2016). This observation is consistent with the results of the North-eastern Atlantic under the influence of Saharan dust (Stichel et al., 2015). Seasonal or annual variability of surface water Nd isotopic compositions may exist in relation to variable dissolved and particulate river and dust inputs as well as current reversal according to monsoon regime. Massive particle inputs could release exchangeable Nd in the water column and on seafloor (e.g. Bay of Bengal and at around 30°N in the North Pacific). Such processes could explain partly why certain margins show more important local/regional influences than others do (Fig. 8a).

In addition to the Nd from the surface layer, benthic flux of Nd from coastal regions may contribute to the observed local influences (Abbott et al., 2015; Du et al., 2016; Abbott et al., 2016). The redox cycle during early diagenesis is active in shallow margin sediments because high organic matter content induces reducing pore water conditions. Pore water Nd concentration is tightly linked to Fe and Mn cycle (Elderfield and Sholkovitz, 1987; Haley et al., 2004). Under reduced pore water conditions, scavenged Nd can be released and labile detrital fractions can be dissolved, leading to discrepancy between bottom and pore water Nd isotopic signature if the isotopic composition of the reactive phases in sediments is different from the bottom water value. Indeed, large positive offsets from the predicted values are observed for both seawater and archive Nd isotopic signatures in the Eastern Equatorial




Fig. 6. Atlantic, Indian and Pacific T-S- ϵ_{Nd} diagrams for seawater (a, c, e) and archives (b, d, f) at water depths from 200 m to bottom. Mean seawater ϵ_{Nd} values of major water masses are summarized in Table 1, and the temperature and salinity ranges and the geographical extension of the water masses are shown in Table S2 (Emery and Meincke, 1986; Amakawa et al., 2009; Bostock et al., 2013). The figures and the maps are created using ODV (Schlitzer, 2015). Atlantic, Indian and Pacific T-S- ϵ_{Nd} diagrams for seawater (a, c, e) and archives (b, d, f) at water depths from 200 m to bottom. Mean seawater ϵ_{Nd} values of major water masses are summarized in Table 1, and the temperature and salinity ranges and the geographical extension of the water masses are shown in Table S2 (Emery and Meincke, 1986; Amakawa et al., 2009; Bostock et al., 2013). The figures and the maps are created using ODV (Schlitzer, 2015).

Pacific, one of the regions where total organic carbon content in sediments is the highest in the modern ocean (Seiter et al., 2004).

Fresh river particles and poorly chemically weathered detrital material could be an additional Nd source to the oceans because they may contain exchangeable Nd (Sholkovitz, 1992; Singh et al., 2012; Roberts and Piotrowski, 2015; Rousseau et al., 2015; Howe et al., 2016b). The impact of poorly chemically weathered detrital material

was originally proposed for unradiogenic ϵ_{Nd} values extracted from foraminiferal tests that reflects weathered detrital material due to the retreat of Laurentide ice sheet during deglaciation (Howe et al., 2016b). In addition to the redox changes that create a benthic flux, transformation of meta-stable minerals containing Nd into stable minerals in contact with seawater produces isotopic exchange on time scales from weeks to months (Jeandel et al., 2011; Jones et al., 2012; Jeandel and

Table 1

Mean seawater ϵ_{Nd} values and Nd concentrations for major water masses estimated from NEOSYMPA database.

Water mass	Basin	ϵ_{Nd}	1 σ	n	Nd (pmol/kg)	1 σ	n	Salinity	1 σ	Temp (°C)	1 σ	Water depth (m)	1 σ	n
AABW	S. Atlantic	-8.6	0.6	39	25.5	3.1	37	34.680	0.020	-0.110	0.44	3059	979	39
NADW	N. Atlantic	-12.3	0.9	173	20.3	3.2	135	34.933	0.035	2.567	0.632	2749	836	173
AAIW	S. Atlantic	-8.6	1.4	28	12.1	1.9	28	34.430	0.159	3.270	1.02	901	270	28
CDW	S. Indian	-7.9	1.3	10	27.6	1.6	9	34.723	0.007	1.058	0.28	3776	744	10
AAIW	S. Indian	-7.6	0.9	4	12.9	2.1	4	34.703	0.096	6.205	1.48	874	248	4
CDW	S. Pacific	-7.8	1.2	92	24.0	4.8	87	34.702	0.019	0.624	0.551	3089	1005	92
NPDW	N. Pacific	-4.3	1.2	84	39.2	9.6	45	34.672	0.019	1.006	0.371	4142	1384	84
AAIW	S. Pacific	-7.7	1.1	21	9.6	2.1	19	34.306	0.081	4.708	1.35	827	254	21
EqPIW	Pacific	-3.0	1.4	31	11.6	5.9	30	34.537	0.029	5.346	0.863	816	125	31
NPIW	N. Pacific	-3.4	1.3	12	14.3	2.2	6	34.122	0.075	6.671	0.754	860	176	35
PEW	Pacific	-2.1	0.9	72	6.5	2.5	64	35.017	0.307	14.33	4.192	586	82	11

AABW: Antarctic Bottom Water.
 NADW: North Atlantic Deep Water.
 AAIW: Antarctic Intermediate Water.
 CDW: Circumpolar Deep Water.
 NPDW: North Pacific Deep Water.
 EqPIW: Equatorial Pacific Intermediate Water.
 NPIW: North Pacific Intermediate Water.
 PEW: Pacific Equatorial Water.




Fig. 7. Comparison between predicted deepwater ϵ_{Nd} values (background color) with Eq. (1) (Atlantic at 50°S to 75°N) and Eq. (2) (global ocean except for the Atlantic at 50°S to 75°N) with observed seawater ϵ_{Nd} values (color dots). The predicted deepwater ϵ_{Nd} values at 1500 m are compared with observed seawater ϵ_{Nd} values at ≥ 1500 m in the Atlantic and Atlantic sector of the Southern Ocean (a) and the other oceanic regions (c). Zonally averaged latitudinal transect at 1500–6000 m of the Atlantic (b) and the Pacific (d). Note that the ϵ_{Nd} scales are different with basins. The Atlantic sector of the Southern Ocean and the southernmost Atlantic (75°S to 50°S) is treated as a buffer zone where deepwater ϵ_{Nd} values are calculated by both Eqs. (1) and (2), and weighted-averaged values considering latitudinal position are shown. All the figures and maps are created using ODV (Schlitzer, 2015).

Oelkers, 2015).

All these processes participate in the determination of local/regional effects in a distinct manner according to different regions, which makes it difficult to provide precise meaning for the observed

distance and water depths. Further insight would be obtained by coupling Nd isotopes, REE patterns and modeling approaches (Oka et al., 2009; Chen et al., 2013a; Zheng et al., 2016). Our results do not provide clear evidence for the impact of oxygen minimum zones (OMZ)




Fig. 8. Comparison between the predicted and observed global seawater ϵ_{Nd} values. Map showing the distribution of difference between observed and predicted ϵ_{Nd} values that correspond to $-(\text{residual } \epsilon_{Nd})$ values at ≥ 1500 m (upper left panel). Histogram of $-(\text{residual } \epsilon_{Nd})$ values with 2σ interval (dashed lines, upper right panel). Scatter plots showing the correlation between observed and predicted ϵ_{Nd} values with latitude (lower left panel) or with oceanic basins (lower right panel). The figures and the map are created using ODV (Schlitzer, 2015).




Fig. 9. Distribution of sites where the seawater and archive ϵ_{Nd} values are significantly affected by local/regional detrital inputs. Maps showing the sites at ≥ 1500 m for seawater data (a) and archive data (c). Relationship between the distance from margin and water depth for seawater data (b) and archive data (d). Labels “a” to “g” show sites discussed in the text. All the figures and maps are created using ODV (Schlitzer, 2015).




Fig. 10. Latitudinal trends of seawater (a and c) and archive (b and d) ϵ_{Nd} values in the Atlantic and Atlantic sector of the Southern Ocean at ≥ 600 m without deep water data significantly affected by local/regional effects. The color code indicates water depths. The major water masses are also shown (Table 1). The gray dots on archive figures (b and d) show seawater values. Labels (i) to (iv) indicate the archive data that are offset from the seawater trends (see text for detail). All the figures and maps are created using ODV (Schlitzer, 2015).

on Nd isotopic variability, similarly to the previous study (Jeandel et al., 2013). Subtle variability related to the reduction of Fe-Mn oxides and consequent release of Nd in OMZ might be quantified for different OMZ by coupled Nd isotopes and REE patterns (Zheng et al., 2016).

4.4. Nd isotopic ratio as a paleo-circulation tracer

The global thermohaline circulation controls the transport of heat, water, salt and nutrients, and affects the segregation of carbon between the atmosphere and the ocean. The Nd isotopes can evaluate the variable contribution between northern/southern-sourced water masses in relation to global thermohaline circulation states (Friedrich et al., 2014). We examine latitudinal seawater and archive ϵ_{Nd} variability for the Atlantic, the Pacific and the Southern Ocean at intermediate and deep water depths (≥ 600 m) after the removal of deep water data significantly influenced by local/regional effects (Fig. 10abcd). Since quantitative evaluation of the detrital influence on waters at < 1500 m is not available, the whole intermediate water data are considered.

The Atlantic, Pacific and corresponding sectors in the Southern Ocean indicate clear latitudinal trends for seawater and archive Nd isotopic compositions. We note that the ϵ_{Nd} values of four archive data sets at intermediate water depths (labeled “i” to “iv” in Fig. 10bd) are > 2 ϵ -units different from the corresponding seawater values. They are all HH leachates with decarbonation. The lower ϵ_{Nd} values off Niger River mouth (labeled “i” in Fig. 10b) possibly reflect unradiogenic signals of preformed Fe-Mn oxides (Kraft et al., 2013). Since the higher ϵ_{Nd} values for Brazil margin correspond to early Holocene (labeled “ii” in Fig. 10b), they could correspond to the real temporal variability (Pahnke et al., 2008). The HH leachate of Antarctic sediment of which the detrital fraction is as high as -2.7 shows a strong positive offset (labeled “iii” in Fig. 10d) (Carter et al., 2012). Finally, the archive ϵ_{Nd} data from the South-eastern Pacific (labeled “iv” in Fig. 10d) allow comparing the performance of different archives because foraminiferal authigenic oxides, HH leachates with or without decarbonation were obtained from the same sediments (Molina-Kescher et al., 2014). The Nd isotopic compositions of foraminiferal oxides and HH leachate without decarbonation fall on the general latitudinal trend formed by seawater data, further confirm the preference to omit the decarbonation step.

The clear latitudinal gradients attest the usefulness of Nd isotopes to evaluate the variability in northern and southern origin water proportion in relation to global circulation changes when the sites are not under strong local/regional continental inputs, supporting the application of this tracer to modern and paleo-oceanography studies.

5. Conclusions

A new global database of seawater and archive Nd isotopic compositions was constructed and combined with hydrography parameters (temperature, salinity, nutrient and oxygen concentrations), $\delta^{13}C$ and $\Delta^{14}C$ values of dissolved inorganic carbon. Archive ϵ_{Nd} data correspond to various leachates, foraminiferal tests, deep-sea corals and fish teeth/debris from the Holocene period ($< 10,000$ years). Using property-property (multi-scatter) plots and T-S- ϵ_{Nd} diagrams, we examined the correspondence between the seawater and archive Nd isotopic compositions.

At water depths ≥ 1500 m, multi-scatter plots present clear correlations between seawater ϵ_{Nd} and the other water mass indicators whereas at 600–1500 m water depths, the relationships are weaker. Basin-scale seawater vertical ϵ_{Nd} profiles demonstrate larger variability at surface and subsurface depths, suggesting local/regional detrital influence at shallower water depths.

Basin-scale seawater T-S- ϵ_{Nd} diagrams allow associating Nd isotopic compositions with water masses. One of the most striking features is radiogenic seawater Nd isotopic composition for surface and intermediate water masses in tropical Pacific that cannot be explained by

only water mass mixing.

Archival Nd isotopic compositions generally agree with corresponding seawater values although they tend to be at the upper limit in the Pacific and both positive and negative offsets exist in the northern North Atlantic.

Applying multiple regression analysis to deep (≥ 1500 m) seawater data, we established empirical equations that predict the main, large-scale, deepwater ϵ_{Nd} trends from hydrography parameters. Large offsets from the predicted values are interpreted as a sign of significant local/regional influence. Dominant detrital influences are observed mainly within 1000 km from the continents. Archive data show more frequently positive offsets that may indicate radiogenic Nd contribution from labile detrital fraction to pore water and/or by chemical extraction of authigenic phases. HH leachate with decarbonation at certain sites present larger offsets from the predicted seawater values than foraminiferal oxides and HH leachate without decarbonation.

Except data influenced by local/regional detrital effects, seawater and archive ϵ_{Nd} values present clear latitudinal trends in the Atlantic and Pacific at water depths ≥ 600 m, consistent with the idea that Nd isotopes help distinguish between northern/southern sourced water contributions at intermediate and deep water depths in the present and past ocean.

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.chemgeo.2017.03.018>.

Acknowledgement

This work was supported by NEOSYMPA project (CYBER/LEFE) granted to KT. We thank l'OSU-Institut Pythéas for technical support of data extraction program. Norbert Frank further received support from the German Science Foundation through the project DFG-NEOGLACIAL (FR-1341/3-1 and 3-2). We thank Brian Haley for helpful suggestions. Thibault de Garidel-Thoron is thanked for the initiation of ODV for KT.

References

- Abbott, A.N., Haley, B.A., McManus, J., 2015. Bottoms up: sedimentary control of the deep North Pacific Ocean's ϵ_{Nd} signature. *Geology* 43 (11), 1035.
- Abbott, A.N., Haley, B.A., McManus, J., 2016. The impact of sedimentary coatings on the diagenetic Nd flux. *Earth Planet. Sci. Lett.* 449, 217–227.
- Achyuthan, H., Nagasundaram, M., Gourlan, A.T., Eastoe, C., Ahmad, S.M., Padmakumari, V.M., 2014. Mid-holocene Indian summer monsoon variability off the Andaman Islands, bay of Bengal. *Quat. Int.* 349, 232–244.
- Albarède, F., Goldstein, S., 1992. A world map of Nd isotopes in seafloor ferromanganese deposits. *Geology* 20, 761–763.
- Amakawa, H., Alibo, D.S., Nozaki, Y., 2004. Nd concentration and isotopic composition distributions in surface waters of Northwest Pacific Ocean and its adjacent seas. *Geochem. J.* 38, 493–504.
- Amakawa, H., Sasaki, K., Ebihara, M., 2009. Nd isotopic composition in the central North Pacific. *Geochim. Cosmochim. Acta* 73 (16), 4705–4719.
- Amakawa, H., Tazoe, H., Obata, H., Gamo, T., Sano, Y., Shen, C.C., 2013. Neodymium isotopic composition and concentration in the Southwest Pacific Ocean. *Geochem. J.* 47, 409–422.
- Antonov, J.I., Seidov, D., Boyer, T.P., Locarnini, R.A., Mishonov, A.V., Garcia, H.E., Baranova, O.K., Zweng, M.M., Johnson, D.R., 2010. World Ocean Atlas 2009, volume 2: salinity. In: Levitus, S. (Ed.), NOAA Atlas NESDIS 68. U.S. Government Printing Office, Washington, D.C., pp. 184.
- Arsouze, T., Dutay, J.C., Lacan, F., Jeandel, C., 2007. Modeling the neodymium isotopic composition with a global ocean circulation model. *Chem. Geol.* 239 (1–2), 165–177.
- Arsouze, T., Dutay, J.C., Lacan, F., Jeandel, C., 2009. Reconstructing the Nd oceanic cycle using a coupled dynamical. *Biogeochem. Model. Biogeosci.* 6 (12), 2829–2846.
- Asahara, Y., Takeuchi, F., Nagashima, K., Harada, N., Yamamoto, K., Oguri, K., Tada, O., 2012. Provenance of terrigenous detritus of the surface sediments in the Bering and Chukchi Seas as derived from Sr and Nd isotopes: implications for recent climate change in the Arctic regions. *Deep-Sea Res. II Top. Stud. Oceanogr.* 61–64 (0), 155–171.
- Basak, C., Martin, E.E., Horikawa, K., Marchitto, T.M., 2010. Southern Ocean source of ^{14}C -depleted carbon in the North Pacific Ocean during the last deglaciation. *Nat. Geosci.* 3 (11), 770–773.
- Basak, C., Pahnke, K., Frank, M., Lamy, F., Gersonde, R., 2015. Neodymium isotopic characterization of Ross Sea Bottom Water and its advection through the southern South Pacific. *Earth Planet. Sci. Lett.* 419 (0), 211–221.
- Bayon, G., German, C.R., Boella, R.M., Milton, J.A., Taylor, R.N., Nesbitt, R.W., 2002. An improved method for extracting marine sediment fractions and its application to Sr and Nd isotopic analysis. *Chem. Geol.* 187, 179–199.

- Bayon, G., German, C.R., Burton, K.W., Nesbitt, R.W., Rogers, N., 2004. Sedimentary Fe-Mn oxyhydroxides as paleoceanographic archives and the role of aeolian flux in regulating oceanic dissolved REE. *Earth Planet. Sci. Lett.* 224 (3–4), 477–492.
- Bayon, G., Birot, D., Ruffine, L., Caprais, J.C., Ponzevera, E., Bollinger, C., Donval, J.P., Charlou, J.L., Voisset, M., Grimaud, S., 2011. Evidence for intense REE scavenging at cold seeps from the Niger Delta margin. *Earth Planet. Sci. Lett.* 312 (3–4), 443–452.
- Bertram, C.J., Elderfield, H., 1993. The geochemical balance of the rare earth elements and Nd isotopes in the oceans. *Geochim. Cosmochim. Acta* 57, 1957–1986.
- von Blanckenburg, F., 1999. Tracing past ocean circulation? *Science* 286 (5446), 1862–1863.
- Blaser, P., Lippold, J., Gutjahr, M., Frank, N., Link, J.M., Frank, M., 2016. Extracting foraminiferal seawater Nd isotope signatures from bulk deep sea sediment by chemical leaching. *Chem. Geol.* 439, 189–204.
- Bostock, H.C., Sutton, P.J., Williams, M.J.M., Opydye, B.N., 2013. Reviewing the circulation and mixing of Antarctic Intermediate Water in the South Pacific using evidence from geochemical tracers and Argo float trajectories. *Deep-Sea Res. I Oceanogr. Res. Pap.* 73 (0), 84–98.
- Bouvier, A., Vervoort, J.D., Patchett, P.J., 2008. The Lu–Hf and Sm–Nd isotopic composition of CHUR: constraints from unequilibrated chondrites and implications for the bulk composition of terrestrial planets. *Earth Planet. Sci. Lett.* 273 (1–2), 48–57.
- Broecker, W.S., Peng, T.H., 1982. *Tracers in the Sea*. Columbia University, Palisades, Lamont-Doherty Geological Observatory (243 pp).
- Burton, K.W., Vance, D., 2000. Glacial-interglacial variations in the neodymium isotope composition of seawater in the Bay of Bengal recorded by planktonic foraminifera. *Earth Planet. Sci. Lett.* 176 (3–4), 425–441.
- Carter, P., Vance, D., Hillenbrand, C.D., Smith, J.A., Shoosmith, D.R., 2012. The neodymium isotopic composition of water masses in the eastern Pacific sector of the Southern Ocean. *Geochim. Cosmochim. Acta* 79 (0), 41–59.
- Charbonnier, G., Puceat, E., Bayon, G., Desmares, D., Dera, G., Durlet, C., Deconinck, J.-F., Amedro, F., Gourlan, A.T., Pellenard, P., Bomou, B., 2012. Reconstruction of the Nd isotope composition of seawater on epicontinental seas: testing the potential of Fe-Mn oxyhydroxide coatings on foraminifera tests for deep-time investigations. *Geochim. Cosmochim. Acta* 99 (0), 39–56.
- Chen, T.-Y., Frank, M., Haley, B.A., Gutjahr, M., Spielhagen, R.F., 2012. Variations of North Atlantic inflow to the central Arctic Ocean over the last 14 million years inferred from hafnium and neodymium isotopes. *Earth Planet. Sci. Lett.* 353–354 (0), 82–92.
- Chen, T.-Y., Rempfer, J., Frank, M., Stumpf, R., Molina-Kescher, M., 2013a. Upper ocean vertical supply: a neglected primary factor controlling the distribution of neodymium concentrations of open ocean surface waters? *J. Geophys. Res. Oceans* 118 (8), 3887–3894.
- Chen, T.-Y., Stumpf, R., Frank, M., Beldowski, J., Staubwasser, M., 2013b. Contrasting geochemical cycling of hafnium and neodymium in the central Baltic Sea. *Geochim. Cosmochim. Acta* 123 (0), 166–180.
- Colin, C., Frank, N., Copard, K., Douville, E., 2010. Neodymium isotopic composition of deep-sea corals from the NE Atlantic: implications for past hydrological changes during the Holocene. *Quat. Sci. Rev.* 29 (19–20), 2509–2517.
- Copard, K., Colin, C., Douville, E., Freiwald, A., Gudmundsson, G., De Mol, B., Frank, N., 2010. Nd isotopes in deep-sea corals in the North-eastern Atlantic. *Quat. Sci. Rev.* 29 (19–20), 2499–2508.
- Copard, K., Colin, C., Frank, N., Jeandel, C., Montero-Serrano, J.C., Reverdin, G., Ferron, B., 2011. Nd isotopic composition of water masses and dilution of the Mediterranean outflow along the southwest European margin. *Geochim. Geophys. Geosyst.* 12 (6), Q06020.
- Copard, K., Colin, C., Henderson, G.M., Scholten, J., Douville, E., Sicre, M.A., Frank, N., 2012. Late Holocene intermediate water variability in the northeastern Atlantic as recorded by deep-sea corals. *Earth Planet. Sci. Lett.* 313–314 (0), 34–44.
- Crockett, K.C., Vance, D., Gutjahr, M., Foster, G.L., Richards, D.A., 2011. Persistent Nordic deep-water overflow to the glacial North Atlantic. *Geology* 39 (6), 515–518.
- Dera, G., Puceat, E., Pellenard, P., Neige, P., Delsate, D., Joachimski, M.M., Reisinger, L., Martinez, M., 2009. Water mass exchange and variations in seawater temperature in the NW Tethys during the Early Jurassic: evidence from neodymium and oxygen isotopes of fish teeth and belemnites. *Earth Planet. Sci. Lett.* 286 (1–2), 198–207.
- Du, J., Haley, B.A., Mix, A.C., 2016. Neodymium isotopes in authigenic phases, bottom waters and detrital sediments in the Gulf of Alaska and their implications for paleo-circulation reconstruction. *Geochim. Cosmochim. Acta* 193, 14–35.
- Dubois-Dauphin, Q., Bonneau, L., Colin, C., Montero-Serrano, J.-C., Montagna, P., Blamart, D., Hebbeln, D., Van Rooij, D., Pons-Branchu, E., Hemsing, F., Wefing, A.-M., Frank, N., 2016a. South Atlantic intermediate water advances into the North-east Atlantic with reduced Atlantic meridional overturning circulation during the last glacial period. *Geochim. Geophys. Geosyst.* 17 (6), 2336–2353.
- Dubois-Dauphin, Q., Montagna, P., Siani, G., Douville, E., Wienberg, C., Hebbeln, D., Liu, Z., Kallel, N., Dapoigny, A., Revel, M., Pons-Branchu, E., Colin, C., 2016b. Hydrological variations of the intermediate water masses of the western Mediterranean Sea during the past 20 ka inferred from neodymium isotopic composition in foraminifera and cold-water corals. *Clim. Past Discuss.* 2016, 1–36.
- Dubois-Dauphin, Q., Colin, C., Bonneau, L., Montagna, P., Wu, Q., Van Rooij, D., Reverdin, G., Douville, E., Thil, F., Waldner, A., Frank, N., 2017. Fingerprinting Northeast Atlantic water masses using Neodymium isotopes. *Geochim. Cosmochim. Acta*.
- Ehlert, C., Grasse, P., Frank, M., 2013. Changes in silicate utilisation and upwelling intensity off Peru since the Last Glacial Maximum -insights from silicon and neodymium isotopes. *Quat. Sci. Rev.* 72 (0), 18–35.
- Elderfield, H., Sholkovitz, E.R., 1987. Rare earth elements in the pore waters of reducing nearshore sediments. *Earth Planet. Sci. Lett.* 82, 280–288.
- Elmore, A.C., Wright, J.D., 2011. North Atlantic Deep Water and climate variability during the Younger Dryas cold period. *Geology* 39 (2), 107–110.
- Elmore, A.C., Piotrowski, A.M., Wright, J.D., Scrivner, A.E., 2011. Testing the extraction of past seawater Nd isotopic composition from North Atlantic deep sea sediments and foraminifera. *Geochim. Geophys. Geosyst.* 12 (9), Q09008.
- Emery, W.J., Meincke, J., 1986. Global water masses: summary and review. *Oceanol. Acta* 9 (4), 383–391.
- van de Fliedert, T., Frank, M., 2010. Neodymium isotopes in paleoceanography. *Quat. Sci. Rev.* 29 (19–20), 2439–2441.
- van de Fliedert, T., Robinson, L.F., Adkins, J.F., 2010. Deep-sea coral aragonite as a recorder for the neodymium isotopic composition of seawater. *Geochim. Cosmochim. Acta* 74 (21), 6014–6032.
- van de Fliedert, T., Pahnke, K., Participants, G.I., 2012. GEOTRACES intercalibration of neodymium isotopes and rare earth element concentrations in seawater and suspended particles. Part 1: reproducibility of results for the international intercomparison. *Limnol. Oceanogr. Methods* 10, 234–251.
- van de Fliedert, T., Griffiths, A.M., Lambelet, M., Little, S.H., Stichel, T., Wilson, D.J., 2016. Neodymium in the oceans: a global database, a regional comparison and implications for paleoceanographic research. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 374 (2081).
- Frank, M., 2002. Radiogenic isotopes: tracers of past ocean circulation and erosional input. *Rev. Geophys.* 40 (1), 1001. <http://dx.doi.org/10.1029/2000RG000094>.
- Frank, N., Freiwald, A., Correa, M.L., Wienberg, C., Eisele, M., Hebbeln, D., Van Rooij, D., Henriot, J.-P., Colin, C., van Weering, T., de Haas, H., Buhl-Mortensen, P., Roberts, J.M., De Mol, B., Douville, E., Blamart, D., Hatté, C., 2011. Northeastern Atlantic cold-water coral reefs and climate. *Geology* 39 (8), 743–746.
- Freslon, N., Bayon, G., Toucanne, S., Bermell, S., Bollinger, C., Chéron, S., Etoubleau, J., Germain, Y., Khripounoff, A., Ponzevera, E., Rouget, M.-L., 2014. Rare earth elements and neodymium isotopes in sedimentary organic matter. *Geochim. Cosmochim. Acta* 140 (0), 177–198.
- Freydier, R., Michard, A., De Lange, G.J., Thomson, J., 2001. Nd isotopic compositions of Eastern Mediterranean sediments: tracers of the Nile influence on sapropel S1 formation. *Mar. Geol.* 177 (1–2), 45–62.
- Friedrich, T., Timmermann, A., Stichel, T., Pahnke, K., 2014. Ocean circulation reconstructions from ϵNd : a model-based feasibility study. *Paleoceanography* 29 (11), 1003–1023.
- Fröllje, H., Pahnke, K., Schnetger, B., Brumsack, H.-J., Dulai, H., Fitzsimmons, J.N., 2016. Hawaiian imprint on dissolved Nd and Ra isotopes and rare earth elements in the central North Pacific: local survey and seasonal variability. *Geochim. Cosmochim. Acta* 189, 110–131.
- Garcia, H.E., Locarnini, R.A., Boyer, T.P., Antonov, J.I., Baranova, O.K., Zweng, M.M., Johnson, D.R., 2010a. World Ocean Atlas 2009, volume 3: dissolved oxygen, apparent oxygen utilization, and oxygen saturation. In: Levitus, S. (Ed.), NOAA Atlas NESDIS 70. U.S. Government Printing Office, Washington, D.C., pp. 344.
- Garcia, H.E., Locarnini, R.A., Boyer, T.P., Antonov, J.I., Zweng, M.M., Baranova, O.K., Johnson, D.R., 2010b. World Ocean Atlas 2009, volume 4: nutrients (phosphate, nitrate, silicate). In: Levitus, S. (Ed.), NOAA Atlas NESDIS 70. U.S. Government Printing Office, Washington, D.C., pp. 398.
- Garcia-Solsona, E., Jeandel, C., Labatut, M., Lacan, F., Vance, D., Chavagnac, V., Pradoux, C., 2014. Rare earth elements and Nd isotopes tracing water mass mixing and particle-seawater interactions in the SE Atlantic. *Geochim. Cosmochim. Acta* 125 (0), 351–372.
- Goldstein, S., Hemming, S.R., 2003. *Long Lived Isotopic Tracers in Oceanography, Paleoceanography, and Ice Sheet Dynamics*. Treatise on Geochemistry. Elsevier Pergamon press, Amsterdam.
- Goswami, V., Singh, S.K., Bhushan, R., 2014. Impact of water mass mixing and dust deposition on Nd concentration and ϵNd of the Arabian Sea water column. *Geochim. Cosmochim. Acta* 145, 30–49.
- Gourlan, A.T., Meynadier, L., Allègre, C.J., 2008. Tectonically driven changes in the Indian Ocean circulation over the last 25 Ma: neodymium isotope evidence. *Earth Planet. Sci. Lett.* 267, 353–364.
- Gourlan, A.T., Meynadier, L., Allegre, C.J., Tapponnier, P., Birc, J.-L., Joron, J.-L., 2010. Northern Hemisphere climate control of the Bengali rivers discharge during the past 4 Ma. *Quat. Sci. Rev.* 29 (19–20), 2484–2498.
- Grasse, P., Stichel, T., Stumpf, R., Stramma, L., Frank, M., 2012. The distribution of neodymium isotopes and concentrations in the Eastern Equatorial Pacific: water mass advection versus particle exchange. *Earth Planet. Sci. Lett.* 353–354 (0), 198–207.
- Grenier, M., Cravatte, S., Blanck, B., Menkes, C., Koch-Larouy, A., Durand, F., Melet, A., Jeandel, C., 2011. From the western boundary currents to the Pacific Equatorial Undercurrent: modeled pathways and water mass evolutions. *J. Geophys. Res.* 116 (C12), C12044.
- Grenier, M., Jeandel, C., Lacan, F., Vance, D., Venchiarutti, C., Cros, A., Cravatte, S., 2013. From the subtropics to the central equatorial Pacific Ocean: neodymium isotopic composition and rare earth element concentration variations. *J. Geophys. Res. Oceans* 118 (2), 592–618.
- Grenier, M., Jeandel, C., Cravatte, S., 2014. From the subtropics to the equator in the southwest Pacific: continental material fluxes quantified using neodymium data along modeled thermocline water pathways. *J. Geophys. Res. Oceans* 119 (6), 3948–3966.
- Gutjahr, M., Lippold, J., 2011. Early arrival of Southern Source Water in the deep North Atlantic prior to Heinrich event 2. *Paleoceanography* 26 (2), PA2101.
- Gutjahr, M., Frank, M., Stirling, C.H., Klemm, V., van de Fliedert, T., Halliday, A.N., 2007. Reliable extraction of a deepwater trace metal isotope signal from Fe-Mn oxyhydroxide coatings of marine sediments. *Chem. Geol.* 242 (3–4), 351–370.
- Gutjahr, M., Frank, M., Stirling, C.H., Keigwin, L.D., Halliday, A.N., 2008. Tracing the Nd isotope evolution of North Atlantic Deep and Intermediate Waters in the western

- North Atlantic since the Last Glacial Maximum from Blake Ridge sediments. *Earth Planet. Sci. Lett.* 266 (1–2), 61–77.
- Haley, B.A., Polyak, L., 2013. Pre-modern Arctic Ocean circulation from surface sediment neodymium isotopes. *Geophys. Res. Lett.* 40 (5), 893–897.
- Haley, B.A., Klinkhammer, G.P., McManus, J., 2004. Rare earth elements in pore waters of marine sediments. *Geochim. Cosmochim. Acta* 68 (6), 1265–1279.
- Haley, B.A., Frank, M., Spielhagen, R.F., Eisenhauer, A., 2008a. Influence of brine formation on Arctic Ocean circulation over the past 15 million years. *Nat. Geosci.* 1 (1), 68–72.
- Haley, B.A., Frank, M., Spielhagen, R.F., Fietzke, J., 2008b. Radiogenic isotope record of Arctic Ocean circulation and weathering inputs of the past 15 million years. *Paleoceanography* 23 (1), PA1S13.
- Haley, B.A., Frank, M., Hathorne, E., Piasias, N., 2014. Biogeochemical implications from dissolved rare earth element and Nd isotope distributions in the Gulf of Alaska. *Geochim. Cosmochim. Acta* 126 (0), 455–474.
- Horikawa, K., Martin, E.E., Asahara, Y., Sagawa, T., 2011. Limits on conservative behavior of Nd isotopes in seawater assessed from analysis of fish teeth from Pacific core tops. *Earth Planet. Sci. Lett.* 310 (1–2), 119–130.
- Howe, J.N.W., Piotrowski, A.M., Noble, T.L., Mulitza, S., Chiessi, C.M., Bayon, G., 2016a. North Atlantic Deep Water production during the last glacial maximum. *Nat. Commun.* 7, 11765.
- Howe, J.N.W., Piotrowski, A.M., Rennie, V.C.F., 2016b. Abyssal origin for the early Holocene pulse of unradiogenic neodymium isotopes in Atlantic seawater. *Geology*.
- Hu, R., Piotrowski, A.M., Bostock, H.C., Crowhurst, S., Rennie, V., 2016. Variability of neodymium isotopes associated with planktonic foraminifera in the Pacific Ocean during the Holocene and Last Glacial Maximum. *Earth Planet. Sci. Lett.* 447, 130–138.
- Huang, K.-F., Oppo, D.W., Curry, W.B., 2014. Decreased influence of Antarctic intermediate water in the tropical Atlantic during North Atlantic cold events. *Earth Planet. Sci. Lett.* 389 (0), 200–208.
- Jacobsen, S.B., Wasserburg, G.J., 1980. Sm–Nd isotopic evolution of chondrites. *Earth Planet. Sci. Lett.* 50, 139–155.
- Jang, K., Han, Y., Huh, Y., Nam, S.-I., Stein, R., Mackensen, A., Matthiessen, J., 2013. Glacial freshwater discharge events recorded by authigenic neodymium isotopes in sediments from the Mendeleev Ridge, western Arctic Ocean. *Earth Planet. Sci. Lett.* 369–370 (0), 148–157.
- Jeandel, C., 2016. Overview of the mechanisms that could explain the ‘Boundary Exchange’ at the land–ocean contact. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 374 (2081).
- Jeandel, C., Oelkers, E.H., 2015. The influence of terrigenous particulate material dissolution on ocean chemistry and global element cycles. *Chem. Geol.* 395 (0), 50–66.
- Jeandel, C., Bishop, J.K., Zindler, A., 1995. Exchange of Nd and its isotopes between seawater small and large particles in the Sargasso Sea. *Geochim. Cosmochim. Acta* 59, 535–547.
- Jeandel, C., Thourou, D., Fieux, M., 1998. Concentrations and isotopic compositions of Nd in the eastern Indian Ocean and Indonesian Straits. *Geochim. Cosmochim. Acta* 62, 2597–2607.
- Jeandel, C., Arsouze, T., Lacan, F., Téchine, P., Dutay, J.-C., 2007. Isotopic Nd compositions and concentrations of the lithogenic inputs into the ocean: a compilation, with an emphasis on the margins. *Chem. Geol.* 239, 156–164.
- Jeandel, C., Peucker Ehrenbrink, B., Jones, M., Pearce, C., Oelkers, E., Godderis, Y., Lacan, F., Aumont, O., Arsouze, T., 2011. Ocean margins: the missing term for oceanic element budgets? *Eos* 92, 217–219.
- Jeandel, C., Delattre, H., Grenier, M., Pradoux, C., Lacan, F., 2013. Rare earth element concentrations and Nd isotopes in the Southeast Pacific Ocean. *Geochem. Geophys. Geosyst.* 14 (2), 328–341.
- Jiménez-Amat, P., Zahn, R., 2015. Offset timing of climate oscillations during the last two glacial-interglacial transitions connected with large-scale freshwater perturbation. *Paleoceanography* 30 (6), 768–788.
- Johannesson, K.H., Burdige, D.J., 2007. Balancing the global oceanic neodymium budget: evaluating the role of groundwater. *Earth Planet. Sci. Lett.* 253 (1–2), 129–142.
- Jones, K.M., Khattiwala, S.P., Goldstein, S.L., Hemming, S.R., van de Fliert, T., 2008. Modeling the distribution of Nd isotopes in the oceans using an ocean general circulation model. *Earth Planet. Sci. Lett.* 272 (3–4), 610–619.
- Jones, M.T., Pearce, C.R., Jeandel, C., Gislason, S.R., Eiriksdottir, E.S., Mavromatis, V., Oelkers, E.H., 2012. Riverine particulate material dissolution as a significant flux of strontium to the oceans. *Earth Planet. Sci. Lett.* 355–356 (0), 51–59.
- Jonkers, L., Zahn, R., Thomas, A., Henderson, G., Abouchami, W., François, R., Masque, P., Hall, I.R., Bickert, T., 2015. Deep circulation changes in the central South Atlantic during the past 145 kyrs reflected in a combined $^{231}\text{Pa}/^{230}\text{Th}$, Neodymium isotope and benthic record. *Earth Planet. Sci. Lett.* 419 (0), 14–21.
- Key, R.M., Kozyr, A., Sabine, C.L., Lee, K., Wanninkhof, R., Bullister, J.L., Feely, R.A., Millero, F.J., Mordy, C., Peng, T.H., 2004. A global ocean carbon climatology: results from Global Data Analysis Project (GLODAP). *Glob. Biogeochem. Cycles* 18 (4), GB4031.
- Klevenz, V., Vance, D., Schmidt, D.N., Mezger, K., 2008. Neodymium isotopes in benthic foraminifera: core-top systematics and a down-core record from the Neogene south Atlantic. *Earth Planet. Sci. Lett.* 265 (3–4), 571–587.
- Kraft, S., Frank, M., Hathorne, E.C., Weldeab, S., 2013. Assessment of seawater Nd isotope signatures extracted from foraminiferal shells and authigenic phases of Gulf of Guinea sediments. *Geochim. Cosmochim. Acta* 121 (0), 414–435.
- Lacan, F., Jeandel, C., 2001. Tracing Papua New Guinea imprint on the central Equatorial Pacific Ocean using neodymium isotopic compositions and Rare Earth Elements concentrations. *Earth Planet. Sci. Lett.* 186, 497–512.
- Lacan, F., Jeandel, C., 2005. Neodymium isotopes as a new tool for quantifying exchange fluxes at the continent–ocean interface. *Earth Planet. Sci. Lett.* 232 (3–4), 245–257.
- Lacan, F., Tachikawa, K., Jeandel, C., 2012. Neodymium isotopic composition of the oceans: a compilation of seawater data. *Chem. Geol.* 300–301 (0), 177–184.
- Lambelet, M., van de Fliert, T., Crockett, K., Rehkämper, M., Kreissig, K., Coles, B., Rijkenberg, M.J.A., Gerringa, L.J.A., de Baar, H.J.W., Steinfeldt, R., 2016. Neodymium isotopic composition and concentration in the western North Atlantic Ocean: results from the GEOTRACES GA02 section. *Geochim. Cosmochim. Acta* 177, 1–29.
- Lang, D.C., Bailey, I., Wilson, P.A., Chalk, T.B., Foster, G.L., Gutjahr, M., 2016. Incursions of southern-sourced water into the deep North Atlantic during late Pliocene glacial intensification. *Nat. Geosci.* 9 (5), 375–379.
- Le Houedec, S., Meynadier, L., Allègre, C.J., 2016. Seawater Nd isotope variation in the Western Pacific Ocean since 80 Ma (ODP 807, Ontong Java Plateau). *Mar. Geol.* 380, 138–147.
- Lerner, P., Marchal, O., Lam, P.J., Anderson, R.F., Buesseler, K., Charette, M.A., Edwards, R.L., Hayes, C.T., Huang, K.-F., Lu, Y., Robinson, L.F., Solow, A., 2016. Testing models of thorium and particle cycling in the ocean using data from station GT11-22 of the U.S. GEOTRACES North Atlantic section. *Deep-Sea Res. I Oceanogr. Res. Pap.* 113, 57–79.
- Lippold, J., Gutjahr, M., Blaser, P., Christner, E., de Carvalho Ferreira, M.L., Mulitza, S., Christl, M., Wombacher, F., Böhm, E., Antz, B., Cartapanis, O., Vogel, H., Jaccard, S.L., 2016. Deep water provenance and dynamics of the (de)glacial Atlantic meridional overturning circulation. *Earth Planet. Sci. Lett.* 445, 68–78.
- Locarnini, R.A., Mishonov, A.V., Antonov, J.L., Boyer, T.P., Garcia, H.E., Baranova, O.K., Zweng, M.M., Johnson, D.R., 2010. World Ocean Atlas 2009, volume 1: temperature. In: Levitus, S. (Ed.), NOAA Atlas NESDIS 68. U.S. Government Printing Office, Washington, D.C., pp. 184.
- López Correa, M., Montagna, P., Joseph, N., Rüggeberg, A., Fietzke, J., Flögel, S., Dorschel, B., Goldstein, S.L., Wheeler, A., Freiwald, A., 2012. Preboreal onset of cold-water coral growth beyond the Arctic Circle revealed by coupled radiocarbon and U-series dating and neodymium isotopes. *Quat. Sci. Rev.* 34 (0), 24–43.
- Maccali, J., Hillaire-Marcel, C., Carignan, J., Reisberg, L.C., 2013. Geochemical signatures of sediments documenting Arctic sea-ice and water mass export through Fram Strait since the Last Glacial Maximum. *Quat. Sci. Rev.* 64 (0), 136–151.
- Maher, B.A., Prospero, J.M., Mackie, D., Gaiero, D., Hesse, P.P., Balkanski, Y., 2010. Global connections between aeolian dust, climate and ocean biogeochemistry at the present day and at the last glacial maximum. *Earth Sci. Res.* 99 (1–2), 61–97.
- Martin, E.E., Haley, B.A., 2000. Fossil fish teeth as proxies for seawater Sr and Nd isotopes. *Geochim. Cosmochim. Acta* 64 (5), 835–847.
- Martin, E.E., Scher, H.D., 2004. Preservation of seawater Sr and Nd isotopes in fossil fish teeth: bad news and good news. *Earth Planet. Sci. Lett.* 220 (1–2), 25–39.
- Martin, E.E., Blair, S.W., Kamenov, G.D., Scher, H.D., Bourbon, E., Basak, C., Newkirk, D.N., 2010. Extraction of Nd isotopes from bulk deep sea sediments for paleoceanographic studies on Cenozoic time scales. *Chem. Geol.* 269 (3–4), 414–431.
- Martin, E.E., MacLeod, K.G., Jiménez Berrocoso, A., Bourbon, E., 2012. Water mass circulation on Demerara Rise during the Late Cretaceous based on Nd isotopes. *Earth Planet. Sci. Lett.* 327–328, 111–120.
- Meinhardt, A.K., Pahnke, K., Böning, P., Schnetger, B., Brumsack, H.J., 2016. Climate change and response in bottom water circulation and sediment provenance in the Central Arctic Ocean since the Last Glacial. *Chem. Geol.* 427, 98–108.
- Moiroud, M., Pucéat, E., Donnadié, Y., Bayon, G., Moriya, K., Deconinck, J.-F., Boyet, M., 2013. Evolution of the neodymium isotopic signature of neritic seawater on a northwestern Pacific margin: new constraints on possible end-members for the composition of deep-water masses in the late Cretaceous ocean. *Chem. Geol.* 356 (0), 160–170.
- Molina-Kescher, M., Frank, M., Hathorne, E.C., 2014. Nd and Sr isotope compositions of different phases of surface sediments in the South Pacific: extraction of seawater signatures, boundary exchange, and detrital/dust provenance. *Geochem. Geophys. Geosyst.* 15 (9), 3502–3520.
- Montero-Serrano, J.-C., Bout-Roumzeilles, V., Carlson, A.E., Tribouillard, N., Bory, A., Meunier, G., Sionneau, T., Flower, B.P., Martínez, P., Billy, I., Riboulleau, A., 2011. Contrasting rainfall patterns over North America during the Holocene and Last Interglacial as recorded by sediments of the northern Gulf of Mexico. *Geophys. Res. Lett.* 38, 6.
- Montero-Serrano, J.-C., Frank, N., Tisnerat-Laborde, N., Colin, C., Wu, C.-C., Lin, K., Shen, C.-C., Copard, K., Orejas, C., Gori, A., De Mol, L., Van Rooij, D., Reverdin, G., Douville, E., 2013. Decadal changes in the mid-depth water mass dynamic of the Northeastern Atlantic margin (Bay of Biscay). *Earth Planet. Sci. Lett.* 364 (0), 134–144.
- Motulsky, H., Christopoulos, A., 2004. Fitting Models to Biological Data Using Linear and Nonlinear Regression. Oxford University Press, New York.
- Murphy, D.P., Thomas, D.J., 2010. The negligible role of intermediate water circulation in stadial-interstadial oxygenation variations along the southern California margin: evidence from Nd isotopes. *Quat. Sci. Rev.* 29 (19–20), 2442–2450.
- Noble, T.L., Piotrowski, A.M., McCave, I.N., 2013. Neodymium isotopic composition of intermediate and deep waters in the glacial southwest Pacific. *Earth Planet. Sci. Lett.* 384 (0), 27–36.
- Oka, A., Hasumi, H., Obata, H., Gamo, T., Yamanaka, Y., 2009. Study on vertical profiles of rare earth elements by using an ocean general circulation model. *Glob. Biogeochem. Cycles* 23 (4), GB4025.
- Osborne, A.H., Haley, B.A., Hathorne, E.C., Flögel, S., Frank, M., 2014. Neodymium isotopes and concentrations in Caribbean seawater: tracing water mass mixing and continental input in a semi-enclosed ocean basin. *Earth Planet. Sci. Lett.* 406 (0), 174–186.
- Osborne, A.H., Haley, B.A., Hathorne, E.C., Plancherel, Y., Frank, M., 2015. Rare earth element distribution in Caribbean seawater: continental inputs versus lateral

- transport of distinct REE compositions in subsurface water masses. *Mar. Chem.* 177 (Part 1), 172–183.
- Pahnke, K., Goldstein, S.L., Hemming, S.R., 2008. Abrupt changes in Antarctic Intermediate Water circulation over the past 25,000 years. *Nat. Geosci.* 1, 870–874.
- Palmer, M.R., Elderfield, H., 1985. Variations in the Nd isotopic composition of foraminifera from Atlantic Ocean sediments. *Earth Planet. Sci. Lett.* 73, 299–305.
- Pena, L.D., Goldstein, S.L., 2014. Thermohaline circulation crisis and impacts during the mid-Pleistocene transition. *Science* 345 (6194), 318–322.
- Pena, L.D., Goldstein, S.L., Hemming, S.R., Jones, K.M., Calvo, E., Pelejero, C., Cacho, I., 2013. Rapid changes in meridional advection of Southern Ocean intermediate waters to the tropical Pacific during the last 30 kyr. *Earth Planet. Sci. Lett.* 368 (0), 20–32.
- Piotrowski, A.M., Goldstein, S.L., Hemming, S.R., Fairbanks, R.G., 2004. Intensification and variability of ocean thermohaline circulation through the last deglaciation. *Earth Planet. Sci. Lett.* 225 (1–2), 205–220.
- Piotrowski, A.M., Goldstein, S.L., Hemming, S.R., Fairbanks, R.G., 2005. Temporal relationships of carbon cycling and ocean circulation at glacial boundaries. *Science* 307, 1933–1938.
- Piotrowski, A.M., Goldstein, S.L.R.H.S., Fairbanks, R.G., Zylberberg, D.R., 2008. Oscillating glacial northern and southern deep water formation from combined neodymium and carbon isotopes. *Earth Planet. Sci. Lett.* 272 (1–2), 394–405.
- Piotrowski, A.M., Banakar, V.K., Scrivner, A.E., Elderfield, H., Galy, A., Dennis, A., 2009. Indian Ocean circulation and productivity during the last glacial cycle. *Earth Planet. Sci. Lett.* 285 (1–2), 179–189.
- Piotrowski, A.M., Galy, A., Nicholl, J.A.L., Roberts, N., Wilson, D.J., Clegg, J.A., Yu, J., 2012. Reconstructing deglacial North and South Atlantic deep water sourcing using foraminiferal Nd isotopes. *Earth Planet. Sci. Lett.* 357–358 (0), 289–297.
- Puceat, E., Lecuyer, C., Reissberg, L., 2005. Neodymium isotope evolution of NW Tethyan upper ocean waters throughout the Cretaceous. *Earth Planet. Sci. Lett.* 236 (3–4), 705–720.
- Rempfer, J., Stocker, T.F., Joos, F., Dutay, J.-C., Siddall, M., 2011. Modelling Nd-isotopes with a coarse resolution ocean circulation model: sensitivities to model parameters and source/sink distributions. *Geochim. Cosmochim. Acta* 75 (20), 5927–5950.
- Rickli, J., Frank, M., Halliday, A.N., 2009. The hafnium-neodymium isotopic composition of Atlantic seawater. *Earth Planet. Sci. Lett.* 280 (1–4), 118–127.
- Rickli, J., Frank, M., Baker, A.R., Aciego, S., de Souza, G., Georg, R.B., Halliday, A.N., 2010. Hafnium and neodymium isotopes in surface waters of the eastern Atlantic Ocean: implications for sources and inputs of trace metals to the ocean. *Geochim. Cosmochim. Acta* 74 (2), 540–557.
- Roberts, N.L., Piotrowski, A.M., 2015. Radiogenic Nd isotope labeling of the northern NE Atlantic during MIS 2. *Earth Planet. Sci. Lett.* 423 (0), 125–133.
- Roberts, N.L., Piotrowski, A.M., McManus, J.F., Keigwin, L.D., 2010. Synchronous deglacial overturning and water mass source changes. *Science* 327 (5961), 75–78.
- Robinson, L.F., van de Fliedert, T., 2009. Southern Ocean evidence for reduced export of North Atlantic Deep Water during Heinrich event 1. *Geology* 37 (3), 195–198.
- Robinson, S.A., Murphy, D.P., Vance, D., Thomas, D.J., 2010. Formation of “southern component water” in the late Cretaceous: evidence from Nd-isotopes. *Geology* 38 (10), 871–874.
- Rousseau, T.C.C., Sonke, J.E., Chmeleff, J., van Beek, P., Souhaut, M., Boaventura, G., Seyler, P., Jeandel, C., 2015. Rapid neodymium release to marine waters from lithogenic sediments in the Amazon estuary. *Nat. Commun.* 6.
- Roy-Barman, M., Lemaire, C., Ayrault, S., Jeandel, C., Souhaut, M., Miquel, J.C., 2009. The influence of particle composition on Thorium scavenging in the Mediterranean Sea. *Earth Planet. Sci. Lett.* 286 (3–4), 526–534.
- Rutberg, R.L., Hemming, S.R., Goldstein, S.L., 2000. Reduced North Atlantic Deep Water flux to the glacial Southern Ocean inferred from neodymium isotopic ratios. *Nature* 405, 935–938.
- Schlitzer, R., 2015. *Ocean Data View*. <http://odv.awi.de> (Editor).
- Schmittner, A., Gruber, N., Mix, A.C., Key, R.M., Tagliabue, A., Westberry, T.K., 2013. Biology and air-sea gas exchange controls on the distribution of carbon isotope ratios ($\delta^{13}\text{C}$) in the ocean. *Biogeosciences* 10 (9), 5793–5816.
- Scrivner, A.E., Vance, D., Rohling, E.J., 2004. New neodymium isotope data quantify Nile involvement in Mediterranean anoxic episodes. *Geology* 32 (7), 565–568.
- Seiter, K., Hensen, C., Schröter, J., Zabel, M., 2004. Organic carbon content in surface sediments—defining regional provinces. *Deep-Sea Res. I Oceanogr. Res. Pap.* 51 (12), 2001–2026.
- Sholkovitz, E.R., 1992. Chemical evolution of rare earth elements: fractionation between colloidal and solution phases of filtered river water. *Earth Planet. Sci. Lett.* 114, 77–84.
- Siddall, M., Khatiwala, S., van de Fliedert, T., Jones, K., Goldstein, S.L., Hemming, S., Anderson, R.F., 2008. Towards explaining the Nd paradox using reversible scavenging in an ocean general circulation model. *Earth Planet. Sci. Lett.* 274 (3–4), 448–461.
- Singh, S.P., Singh, S.K., Goswami, V., Bhushan, R., Rai, V.K., 2012. Spatial distribution of dissolved neodymium and ϵ_{Nd} in the Bay of Bengal: role of particulate matter and mixing of water masses. *Geochim. Cosmochim. Acta* 94 (0), 38–56.
- Skinner, L.C., Scrivner, A.E., Vance, D., Barker, S., Fallon, S., Waelbroeck, C., 2013. North Atlantic versus Southern Ocean contributions to a deglacial surge in deep ocean ventilation. *Geology* 41 (6), 667–670.
- Stichel, T., Frank, M., Rickli, J., Haley, B.A., 2012. The hafnium and neodymium isotope composition of seawater in the Atlantic sector of the Southern Ocean. *Earth Planet. Sci. Lett.* 317–318 (0), 282–294.
- Stichel, T., Hartman, A.E., Duggan, B., Goldstein, S.L., Scher, H., Pahnke, K., 2015. Separating biogeochemical cycling of neodymium from water mass mixing in the Eastern North Atlantic. *Earth Planet. Sci. Lett.* 412 (0), 245–260.
- Stoll, H.M., Vance, D., Arealos, A., 2007. Records of the Nd isotope composition of seawater from the Bay of Bengal: implications for the impact of Northern Hemisphere cooling on ITCZ movement. *Earth Planet. Sci. Lett.* 255 (1–2), 213–228.
- Stordal, M.C., Wasserburg, G.J., 1986. Neodymium isotopic study of Baffin Bay water: sources of REE from very old terranes. *Earth Planet. Sci. Lett.* 77, 259–272.
- Stumpf, R., Frank, M., Schonfeld, J., Haley, B.A., 2010. Late Quaternary variability of Mediterranean Outflow Water from radiogenic Nd and Pb isotopes. *Quat. Sci. Rev.* 29 (19–20), 2462–2472.
- Tachikawa, K., Jeandel, C., Roy-Barman, M., 1999a. A new approach to the Nd residence time in the ocean: the role of atmospheric inputs. *Earth Planet. Sci. Lett.* 170 (4), 433–446.
- Tachikawa, K., Jeandel, C., Vangriesheim, A., Dupre, B., 1999b. Distribution of rare earth elements and neodymium isotopes in suspended particles of the tropical Atlantic Ocean (EUMELI site). *Deep-Sea Res. I Oceanogr. Res. Pap.* 46 (5), 733–755.
- Tachikawa, K., Athias, V., Jeandel, C., 2003. Neodymium budget in the modern ocean and paleo-oceanographic implications. *J. Geophys. Res.* 108 (C8), 3254.
- Tachikawa, K., Roy-Barman, M., Michard, A., Thouron, D., Yeghicheyan, D., Jeandel, C., 2004. Neodymium isotopes in the Mediterranean Sea: comparison between seawater and sediment signals. *Geochim. Cosmochim. Acta* 68 (14), 3095–3106.
- Tachikawa, K., Piotrowski, A.M., Bayon, G., 2014. Neodymium associated with foraminiferal carbonate as a recorder of seawater isotopic signatures. *Quat. Sci. Rev.* 88 (0), 1–13.
- Thomas, D.J., 2004. Evidence for deep-water production in the North Pacific Ocean during the early Cenozoic warm interval. *Nature* 430 (6995), 65–68.
- Thomas, D.J., Korty, R., Huber, M., Schubert, J.A., Haines, B., 2014. Nd isotopic structure of the Pacific Ocean 70–30 Ma and numerical evidence for vigorous ocean circulation and ocean heat transport in a greenhouse world. *Paleoceanography* 2013PA002535.
- Vance, D., Burton, K., 1999. Neodymium isotopes in planktonic foraminifera: a record of the response of continental weathering and ocean circulation rates to climate change. *Earth Planet. Sci. Lett.* 173, 365–379.
- Vance, D., Scrivner, A.E., Beney, P., Staubwasser, M., Henderson, G.M., 2004. The use of foraminifera as a record of the past neodymium isotope composition of seawater. *Paleoceanography* 19 (PA2009). <http://dx.doi.org/10.1029/2003PA000957>.
- Wei, R., Abouchami, W., Zahn, R., Masque, P., 2016. Deep circulation changes in the South Atlantic since the Last Glacial Maximum from Nd isotope and multi-proxy records. *Earth Planet. Sci. Lett.* 434, 18–29.
- Werner, K., Frank, M., Teschner, C., Müller, J., Spielhagen, R.F., 2014. Neoglacial change in deep water exchange and increase of sea-ice transport through eastern Fram Strait: evidence from radiogenic isotopes. *Quat. Sci. Rev.* 92 (0), 190–207.
- Wilson, D.J., Piotrowski, A.M., Galy, A., McCave, I.N., 2012. A boundary exchange influence on deglacial neodymium isotope records from the deep western Indian Ocean. *Earth Planet. Sci. Lett.* 341–344 (0), 35–47.
- Wilson, D.J., Piotrowski, A.M., Galy, A., Clegg, J.A., 2013. Reactivity of neodymium carriers in deep sea sediments: implications for boundary exchange and paleoceanography. *Geochim. Cosmochim. Acta* 109 (0), 197–221.
- Wilson, D.J., Crocket, K.C., van de Fliedert, T., Robinson, L.F., Adkins, J.F., 2014. Dynamic intermediate ocean circulation in the North Atlantic during Heinrich Stadial 1: a radiocarbon and neodymium isotope perspective. *Paleoceanography* 29 (11) 2014PA002674.
- Wu, Q., Colin, C., Liu, Z., Douville, E., Dubois-Dauphin, Q., Frank, N., 2015a. New insights into hydrological exchange between the South China Sea and the Western Pacific Ocean based on the Nd isotopic composition of seawater. *Deep-Sea Res. II Top. Stud. Oceanogr.* 122, 25–40.
- Wu, Q., Colin, C., Liu, Z., Thil, F., Dubois-Dauphin, Q., Frank, N., Tachikawa, K., Bordier, L., Douville, E., 2015b. Neodymium isotopic composition in foraminifera and authigenic phases of the South China Sea sediments: implications for the hydrology of the North Pacific Ocean over the past 25 kyr. *Geochem. Geophys. Geosyst.* 16 (11), 3883–3904.
- Xie, R.C., Marcantonio, F., Schmidt, M.W., 2012. Deglacial variability of Antarctic Intermediate Water penetration into the North Atlantic from authigenic neodymium isotope ratios. *Paleoceanography* 27 (3), PA3221.
- Yasuda, I., Okuda, K., Shimizu, Y., 1996. Distribution and modification of north pacific intermediate water in the Kuroshio-Oyashio interfrontal zone. *J. Phys. Oceanogr.* 26 (4), 448–465.
- Yu, J., Elderfield, H., Piotrowski, A.M., 2008. Seawater carbonate ion- $\delta^{13}\text{C}$ systematics and application to glacial-interglacial North Atlantic ocean circulation. *Earth Planet. Sci. Lett.* 271 (1–4), 209–220.
- Zheng, X.-Y., Plancherel, Y., Saito, M.A., Scott, P.M., Henderson, G.M., 2016. Rare earth elements (REEs) in the tropical South Atlantic and quantitative deconvolution of their non-conservative behavior. *Geochim. Cosmochim. Acta* 177, 217–237.