



Swansea University  
Prifysgol Abertawe



## Cronfa - Swansea University Open Access Repository

---

This is an author produced version of a paper published in :

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa32984>

---

### Conference contribution :

Banerjee, D., Biraben, F., Charlton, M., Clade, P., Comini, P., Crivelli, P., Dalkarov, O., Debu, P., Dodd, L., Douillet, A., Dufour, G., Dupre, P., Eriksson, S., Froelich, P., Grandemange, P., Guellati, S., Guerout, R., Heinrich, J., Hervieux, P., Hilico, L., Husson, A., Indelicato, P., Jonsell, S., Karr, J., Khabarova, K., Kim, S., Kim, Y., Kolachevsky, N., Kuroda, N., Lambrecht, A., Leite, A., Liskay, L., Lotrus, P., Lunney, D., Madsen, N., Manfredi, G., Mansouli, B., Matsuda, Y., Mohri, A., Mornacchi, G., Nesvizhevsky, V., Nez, F., Perez, P., Regenfus, C., Rey, J., Reymond, J., Rousse, J., Reynaud, S., Rubbia, A., Sacquin, Y., Schmidt-Kaler, F., Sillitoe, N., Staszczak, M., Torii, H., Heinrich, J., Vallage, B., Valdes, M., van der Werf, D., Voronin, A., Walz, J., Wolf, S., Wronka, S. & Yamazaki, Y. (2016). *Towards a test of the weak equivalence principle of gravity using anti-hydrogen at CERN.*(pp. 1-2). 2016 Conference on Precision Electromagnetic Measurements (CPEM 2016).

<http://dx.doi.org/10.1109/CPEM.2016.7540781>

---

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

# Towards a test of the Weak Equivalence Principle of gravity using anti-hydrogen at CERN

D. Banerjee<sup>2</sup>, F. Biraben<sup>7</sup>, M. Charlton<sup>11</sup>, P. Cladé<sup>7</sup>, P. Comini<sup>1,2</sup>, P. Crivelli<sup>2</sup>, O. Dalkarov<sup>6</sup>, P. Debu<sup>5</sup>, L. Dodd<sup>11</sup>, A. Douillet<sup>7#</sup>, G. Dufour<sup>7</sup>, P. Dupré<sup>10</sup>, S. Eriksson<sup>11</sup>, P. Froelich<sup>14</sup>, P. Grandemange<sup>\*\*</sup>, S. Guellati<sup>7</sup>, R. Guérout<sup>7</sup>, J. M. Heinrich<sup>7</sup>, P.-A. Hervieux<sup>4</sup>, L. Hilico<sup>7#</sup>, A. Husson<sup>1</sup>, P. Indelicato<sup>7</sup>, S. Jonsell<sup>15</sup>, J.-P. Karr<sup>7#</sup>, K. Khabarova<sup>6</sup>, S.K. Kim<sup>16</sup>, Y. Kim<sup>17</sup>, N. Kolachevsky<sup>6</sup>, N. Kuroda<sup>12</sup>, A. Lambrecht<sup>7</sup>, A.M.M. Leite<sup>5</sup>, L. Liskay<sup>5</sup>, P. Lotrus<sup>5</sup>, D. Lunney<sup>1</sup>, N. Madsen<sup>11</sup>, G. Manfredi<sup>4</sup>, B. Mansouli<sup>5</sup>, Y. Matsuda<sup>12</sup>, A. Mohri<sup>13</sup>, G. Mornacchi<sup>\*\*</sup>, V. Nesvizhevsky<sup>3</sup>, F. Nez<sup>7</sup>, P. Pérez<sup>5</sup>, C. Regenfus<sup>2</sup>, J.-M. Rey<sup>5</sup>, J.-M. Reymond<sup>5</sup>, J.-Y. Roussé<sup>5</sup>, S. Reynaud<sup>7</sup>, A. Rubbia<sup>2</sup>, Y. Sacquin<sup>5</sup>, F. Schmidt-Kaler<sup>8</sup>, N. Sillitoe<sup>7</sup>, M. Staszczak<sup>9</sup>, H. Torii<sup>12</sup>, J. M. Heinrich<sup>7</sup>, B. Vallage<sup>5</sup>, M. Valdes<sup>4</sup>, D.P. van der Werf<sup>5,11</sup>, A. Voronin<sup>6</sup>, J. Walz<sup>8</sup>, S. Wolf<sup>8</sup>, S. Wronka<sup>9</sup> and Y. Yamazaki<sup>10</sup>

<sup>1</sup> Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, Orsay, France,

<sup>2</sup>Eidgenössische Tech. Hochschule Zürich (ETH Zürich), Zürich, Switzerland,

<sup>3</sup>Institut Laue Langevin, Grenoble, France,

<sup>4</sup> Institut de Physique et Chimie des Matériaux de Strasbourg, Strasbourg, France,

<sup>5</sup> Institut de recherche sur les lois fondamentales de l'Univers (IRFU), CEA Saclay, France

<sup>6</sup> P.N. Lebedev Institute of Physics, Moscow, Russia,

<sup>7</sup> Laboratoire Kastler Brossel, UPMC-Sorbonne Universités, CNRS, ENS-PSL Research University, Collège de France, Paris, France,

<sup>8</sup> Johannes-Gutenberg-Universität, Mainz, Germany,

<sup>9</sup> National Center for Nuclear Research, Otwock-Swierk, Poland,

<sup>10</sup> Institute of Physical and Chemical Research, Wako, Japan,

<sup>11</sup> Swansea University, Swansea, United Kingdom,

<sup>12</sup> Institute of Physics, University of Tokyo, Komaba, Japan,

<sup>13</sup> Kyoto University, Kyoto, Japan,

<sup>14</sup> Uppsala University, Uppsala, Sweden,

<sup>15</sup> Stockholm University, Stockholm, Sweden,

<sup>16</sup> Seoul National University, Seoul, Korea,

<sup>17</sup> Institute for Basic Science, Daejeon, Korea,

<sup>\*\*</sup> CERN, Switzerland,

<sup>#</sup> also at Université d'Evry, Comue Paris-Saclay, Evry, France  
francois.nez@lkb.upmc.fr

**Abstract**—The aim of the GBAR (Gravitational Behavior of Antimatter at Rest) experiment is to measure the free fall acceleration of an antihydrogen atom, in the terrestrial gravitational field at CERN and therefore test the Weak Equivalence Principle with antimatter. The aim is to measure the local gravity with a 1% uncertainty which can be reduced to few parts of  $10^{-3}$ .

**Index Terms**—Gravitation, Antimatter.

## I. INTRODUCTION

To date, the controversial behavior of antimatter in the gravity field (see [1] and references therein) has never been accurately tested. To avoid stray effects from electric or magnetic fields, the antimatter species has to be neutral. The possible candidates are antineutrons ( $\bar{n}$ ) [2], antihydrogen ( $\bar{H}$ ) [3] [4] and positronium (Ps) (i.e.  $e^+ e^-$ ) atoms [5]. One attractive feature of  $\bar{H}$ , which is made of one antiproton ( $\bar{p}$ )


and one positron ( $e^+$ ), is the possibility of laser cooling. However, cooling  $\bar{H}$  in the 1S ground state is still challenging as there is no powerful laser available at 121nm to drive the 1S-2P transition [6]. The originality of GBAR experiment, which has been approved by the CERN committee [7], is to circumvent this difficulty by producing  $\bar{H}$  at rest from the photo-detachment of one of the positrons of the anti-ion ( $\bar{H}^+$ ) (that is  $\bar{p} e^+ e^+$ ) as presented in [8]. Indeed  $\bar{H}^+$  ions can be efficiently cooled in electromagnetic traps .

## II. PRINCIPLE OF THE EXPERIMENT

This complex experiment can be divided into three challenging parts :

- production of  $\bar{H}^+$  ions,
- cooling of  $\bar{H}^+$  ions,
- free-fall of an anti-hydrogen atom.

The  $\bar{H}^+$  ions are produced by two consecutive reactions using excited Positronium atoms ( $Ps^*$ ):



in which  $\bar{H}^*$  stands for anti-hydrogen in an excited state.

Nowadays, efficient production of Ps atoms from a positron ( $e^+$ ) beam sent on a nanoporous silica sample can be achieved [9]. As the total efficiency of the two reactions is not very high, large quantities of  $e^+$  and  $\bar{p}$  are still needed. The experiment is carried out at CERN, on the Antiproton Decelerator and ELENA ring, using an intense linac based positron source.

### III. DEVELOPMENTS

There is an intense activity on the GBAR project both from the theoretical and experimental sides. A careful theoretical study has been done on the systematic effects arising from the anti-hydrogen detector which can interfere with the free fall signal [10][11]. Possible schemes for a determination of the local gravity at the level of  $10^{-3}$  with the GBAR apparatus have also been proposed [12][13][14][15].

The cooling of  $\bar{H}^+$  will be challenging as its temperature has to be decreased over eleven decades, from 200 eV to the neV range. It will be done in several steps, the last one being ground-state cooling of a  $Be^+/\bar{H}^+$  ion pair. Simulations of the cooling have been performed, and the lasers needed for ions pair cooling are under development [16].

The beam line is also under construction see [17] and references therein. The cross sections of the two consecutive reactions (1) and (2) have been calculated to optimize the production of  $\bar{H}^+$  [18] and more recently, the cross section of the first reaction has been recalculated [19].

It was shown in [18] that, depending on the antiproton energy, the production of  $\bar{H}^+$  should be optimized with Ps atoms excited in 3D state or in 2P state. The velocity distributions of the Ps atoms cloud produced in nano-porous silica is very wide [20] and references therein. Therefore, in order to maximize the Ps excitation efficiency, one possibility is to use a two-photon transition to get rid of the first-order Doppler effect, as proposed for the 1S-3D transition [22]. Alternatively one can build a large line-width laser [20] or an optical frequency comb to excite the 1S-2P transition at 243 nm [21].

Currently, a laser chain for the two-photon excitation of the 1S-3D line of Ps, at 410 nm, has been built and characterized [22]. It is based on the same principle as the laser used for muonic hydrogen spectroscopy [23] excepted that the oscillator cavity is a ring cavity instead of a linear one. The cross section of reaction (1) will be measured in the near future.

A general overview of the project and of recent advances in the development of the experiment will be given at the conference.

### IV. ACKNOWLEDGEMENTS

This work is supported in part by ANTION Project (ANR-14-CE33-0008) operated by the French National Research Agency (ANR).

### REFERENCES

- [1] M. M Nieto and T. Goldman, "the argument against anti-gravity and gravitational acceleration of anti-matter", *Phys. Rept.* **205**, 221-281, (1991).
- [2] F. S. Dzheparov, Y.G. Abov and L.B. Okun, "Test of the equality of the particle and anti particle masses in neutron-antineutron oscillations", *JETP Lett.* **30** 599-600, (1984).
- [3] A. Kellerbauer et al, "Proposed antimatter gravity measurement with antihydrogen beam", *Nucl. Instr. Meth. Phys. Res. B*, **266**, 561 (2008).
- [4] The ALPHA collaboration and A.E. Charman, "Description and first application of a new technique to measure the gravitational mass of antihydrogen", *Nature Comm.* **4**, 1785, (2013).
- [5] P. Crivelli, D.A Cooke and S. Friedreich, "Experimental considerations for testing ant-matter antigravity using positronium 1S-2S spectroscopy", *International Journal of Modern Physics : Conference Series*, **20**, 1460257, (2014).
- [6] A. Müllers et al, "A semiconductor laser system for the production of antihydrogen" *New J. Phys.* **14** 055009 (2012).
- [7] G. Chardin et al., "GBAR, Proposal to Measure the Gravitational Behaviour of Antihydrogen at Rest", CERN-SPSC-P-342, 30/09/2011.
- [8] J. Waltz and T.W. Hänsch, "A proposal to measure antimatter gravity using ultracold antihydrogen atoms", *Gen. Relat. Gravit.* **36**, 561 (2008).
- [9] L. Liszky et al, "Positronium reemission yield from mesostructured silica films", *App. Phys. Lett.* **92**, 063114 (2008).
- [10] G. Dufour et al, "Quantum reflection of antihydrogen from the Casimir potential above matter slabs", *Phys Rev A* **87**, 012901 (2013).
- [11] G. Dufour, R. Guérout, A. Lambrecht, S. Reynaud, "Quantum reflection of antihydrogen from nanoporous media", *Phys Rev A* **87**, 022506 (2013).
- [12] G. Dufour et al, "Shaping the distribution of vertical velocities of antihydrogen in GBAR", *Eur. Phys. J. C*, **74**, 2731 (2014).
- [13] A. Yu. Voronin et al, "A spectroscopy approach to measure the gravitational mass of antihydrogen", arXiv:1403.4783
- [14] A. Yu. Voronin et al, "Quantum ballistic experiment on antihydrogen fall", arXiv:1510.00866;
- [15] G. Dufour et al., "Prospects for Studies of the Free Fall and Gravitational Quantum States of Antimatter", *Advances in High Energy Physics* (2015) 379642.
- [16] L. Hilico et al, "Preparing single ultra-cold antihydrogen atoms for the free-fall in GBAR", *International Journal of Modern Physics, Conference Series* **30**,1460269 (2014).
- [17] D. Lunney et al., "Beam preparation for studying the gravitational behaviour of antimatter at rest", LEAP 2013 Proceedings, *Hyp. Interact.* **229**, 1 (2014)
- [18] P. Comini and P-A. Hervieux, "H<sup>+</sup> ion production from collisions between antiprotons and excited positronium: cross sections calculations in the framework of the GBAR experiment", *New J. Phys.* **15**, 095022 (2013).
- [19] A. S. Kadyrov et al., "Antihydrogen Formation via Antiproton Scattering with Excited Positronium", *Phys. Rev. Lett.* **114**, 183201 (2015).
- [20] A. Deller et al, "Exciting positronium with a solid-state UV laser: the Doppler-broadened Lyman- $\alpha$  transition", *J. Phys. B: At. Mol. Opt. Phys.* **48**, 175001 (2015).
- [21] F. Biraben, GBAR collaboration meeting communication (2014).
- [22] P. Comini thesis, <https://tel.archives-ouvertes.fr/tel-01150446>
- [23] A. Antognini et al, "Powerful fast triggerable 6  $\mu$ m laser for muonic hydrogen 2S-Lamb shift experiment", *Opt. Comm.* **253**, 362-374 (2005).