

Protocol 1. Total RNA preparation

Jürg Bähler and Jo Ann Wise.

Correspondence: j.bahler@ucl.ac.uk

ABSTRACT

This method is widely used by fission yeast researchers to prepare RNA for analysis of individual species (e.g., Helmlinger et al. 2008) or for genome-wide studies (e.g., Bitton et al. 2011; Wilhelm et al. 2008). Treatment with hot phenol breaks open the cells and begins to strip away bound proteins from the RNA. Deproteinization is completed by multiple extractions with chloroform/isoamyl alcohol and separation of the aqueous and organic phases using Phase Lock Gel, an inert material that acts as a physical barrier between them. The final step is concentration of the RNA by ethanol precipitation. The protocol can be used to prepare RNA from several cultures grown in parallel but it is important not to process too many samples at once as delays can be detrimental to RNA quality.

Materials

Reagents: Cells grown in YES or other appropriate media (25-50 mL at $0.5 - 1 \times 10^7$ cells/mL)
Chloroform
Ethanol
ddH₂O (double-distilled)
Isoamyl alcohol
Phenol (water-saturated, acidic-equilibrated, e.g. Sigma P1944)
Sodium acetate (NaOAc) (3 M pH 5.2)
TES buffer <R>
Agarose
TBE Agarose Gel running buffer

Equipment: Bench top centrifuge with swing-out rotor for 50 mL tubes
Dry ice/ethanol bath
Microcentrifuge
Microcentrifuge tubes
Phase Lock Gel tubes (Eppendorf)

Vortex Mixer

Water bath or heat block (65°C)

Agarose Gel electrophoresis tank and casting sets to generate gels

Note: Due to the widespread use of RNase in small-scale DNA preparations, some labs use electrophoresis tanks that are specifically dedicated to running RNA gels to avoid degradation.

Method

1. Prepare the work area by cleaning racks, pipettes and work surfaces, e.g. using RNase-ZAP wipes (Ambion or Sigma). Prepare all solutions including gels and gel buffers using RNase-free technique and wear gloves throughout the procedure.
2. Transfer cells to 50 mL screw-cap tubes and harvest by centrifugation for 5 min at 1,000 x g or 3 min at 2,000 x g and discard the culture fluid by carefully decanting. Snap-freeze the cell pellet by placing the tube in a dry ice/ethanol bath. Cells may be stored at -80°C prior to extracting RNA.
3. Thaw cells by placing the tube on ice (5 min). Add 1 mL of pre-chilled ddH₂O to resuspend the cells (adjust volume accordingly if using more or less than the specified number of cells). Transfer the cell suspension to a 2 mL microfuge tube and spin for 10 sec at 20,000 x g; remove the supernatant (SN) and discard.
4. To the cell pellet add 750 µL of TES buffer, resuspend the cells with a pipette, immediately add 750 µL of acidic phenol (stored at 4°C), vortex vigorously, and incubate in a 65°C water bath or heat block. Treat the next sample in the same way.
5. Incubate all samples at 65°C for 1 hr, vortexing for 10 sec every 10 min.
Note: A similar protocol for budding yeast (Rio et al. 2011, Chapter 2, Protocol 7, p. 38-40) uses a shorter (10 min) incubation at high temperature, which is recommended if problems with degradation are encountered (see step 15).
6. Place the samples on ice for 1 min, vortex for 20 sec, and spin in a micro-centrifuge for 15 min at 20,000 x g at 4°C.
Note: Some labs use room temperature as the phenol should prevent RNA degradation.
7. Pre-spin the appropriate number of yellow Phase Lock tubes for 10 sec to return the gel to the bottom and add 700 µL of acidic phenol-chloroform to each.
8. Take 700 µL of the aqueous phase from step 6 and add to the tubes from step 7, thoroughly mix by inverting (do not vortex as this will fragment the gel), and centrifuge for 5

- min at 20,000 g at 4°C.
9. Add 700 μ L of chloroform:isoamyl alcohol (24:1) to the appropriate number of new 2 mL microfuge tubes.
 10. Take 700 μ L of the aqueous (top) phase from step 8 and add to the tubes from step 9, thoroughly mix by inverting (not vortexing), and centrifuge for 5 min at 20,000 g at 4°C.
 11. To the appropriate number of standard 2 mL microfuge tubes, add 1.5 mL of 100% EtOH (4°C) and 50 μ L of 3 M NaAc pH 5.2.
 12. Transfer 500 μ L of the aqueous phase from step 10 to the tubes from step 11, vortex for 10 sec. The samples can be precipitated at -20°C overnight (or at -70°C in a dry ice-ethanol bath for 30 min).
 13. Spin in a micro-centrifuge for 10 min at 20,000 g at RT. Discard SUPERNATANT, add 500 μ L 70% EtOH (V/V, diluted with water & stored at 4°C), flick the tube gently with a finger but don't vortex, then spin for 1 min using the same tube orientation to avoid dislodging the pellet. Aspirate most of the SUPERNATANT, spin for 5 sec, and remove the remaining liquid with a pipette. Air dry for 5 min at RT.
 14. Add 100 μ L of ddH₂O (adjust volume to pellet size), and incubate for 1 min at 65°C (or 10 min at RT). Dissolve the pellet by first pipetting up and down (~30x) until no particles are left, then gently vortex for 10 sec.
 15. To assess the integrity of the RNA, run 2 μ L of the sample on a 1% agarose gel under RNase-free conditions. An even better option if available is to use a Bioanalyzer, which can generate a "virtual gel" from the electropherogram. If the preparation is free of degradation, the two large ribosomal RNAs will form two single sharp bands, whereas a partially degraded sample will show decreased intensity of the large subunit band and a ladder of more rapidly migrating bands below the major bands (for an example, see Rio et al. 2011, Chapter 2, Protocol 12, Fig. 2-7). On a stained agarose gel, the degradation products may appear as a smear rather than discrete bands. In addition to assessing integrity, such gels can also be used to normalize the amounts of RNA in different preps.
 16. To assess the purity of the RNA, use a spectrophotometer to determine A_{260} , A_{270} and A_{280} . If the sample is uncontaminated, the $A_{260}:A_{280}$ ratio should be between 1.8 and 2.2. An $A_{270} \geq A_{260}$ may indicate phenol contamination (see Rio et al. 2011, Chapter 1, p. 8).
 17. The yield of RNA from this protocol is generally ~200 μ g but cells grown under some conditions may yield less RNA. Quantitate the yield of RNA using a standard or nanoscale

spectrophotometer (see Rio et al. 2011, Chapter 2, Protocol 13, p. 67-68). Adjust different samples to be analyzed in parallel to the same concentration, generally 1-2 $\mu\text{g}/\mu\text{L}$, by adding ddH₂O.

Reagents:

TES buffer: 10 mM Tris, pH 7.5
10 mM EDTA, pH 8.0
0.5% SDS

Additions and variations:

- To eliminate RNase activity, some labs treat glassware with diethyl pyrocarbonate (DEPC) and/or use DEPC-treated water to make solutions. Others avoid this practice, as residual DEPC not destroyed by autoclaving can react with adenines in the RNA. Do NOT treat buffers containing Tris with DEPC.
- To avoid stress responses during harvesting, cells may be collected by filtration rather than centrifugation (Chapter 2) and the filter disc can be snap-frozen in liquid nitrogen (step 2).
- Some labs use standard Eppendorf microfuge tubes rather than Phase Lock tubes to separate the organic and aqueous phases (step 6).
- Some labs add an additional purification step using Qiagen RNeasy columns, but it should be noted that this introduces a bias towards large (> 200 nt) RNAs (see Rio et al. 2011, Chapter 2, Protocol 4, p. 30-32 for additional information about RNA purification kits). If you elect to include Qiagen purification, postpone analysis of purity and integrity until after it is completed. After quantitation and adjusting concentration, proceed as follows:
 18. Use 100 μg of each RNA preparation for Qiagen purification. Measure the volume of the remaining RNA, add 1/10 volume of 3 M NaOAc (pH 5.2) and 3 volumes of 100% EtOH, mix by inversion, and store at -70°C as a backup.
 19. Purify 100 μg of each of your RNAs over an RNeasy mini-spin column as described in the RNeasy Mini Handbook supplied by the manufacturer. Elute twice with 30 μL RNase-free water. Hold on ice.

20. Determine the yield, integrity and purity of the RNA as described above (steps 15-17) and concentrate by ethanol precipitation if necessary.

REFERENCES

- Bitton DA, Grallert A, Scutt PJ, Yates T, Li Y, Bradford JR, Hey Y, Pepper SD, Hagan IM, Miller CJ. 2011. Programmed fluctuations in sense/antisense transcript ratios drive sexual differentiation in fission yeast. *Mol Syst Biol* **20**: 559.
- Helmlinger D, Marguerat S, Villèn J, Gygi SP, Bähler J, Winston F. 2008. The *S. pombe* SAGA complex controls the switch from proliferation to sexual differentiation through opposing roles of its subunits Gcn5 and Spt8. *Genes Dev* **22**: 3184-3195.
- Wilhelm BT, Marguerat S, Watt S, Schubert F, Wood V, Goodhead I, Penkett CJ, Rogers J, Bähler J. 2008. Dynamic repertoire of a eukaryotic transcriptome surveyed at single-nucleotide resolution. *Nature* **26**: 1239-1243.

Protocol 2. Polysome profile analysis and RNA purification

Dieter Wolf, Jürg Bähler and Jo Ann Wise.

Correspondence: dwolf@sanfordburnham.org or j.bahler@ucl.ac.uk

ABSTRACT

Polysome profile analysis is widely used by investigators studying the mechanism and regulation of translation. The method described below uses high-velocity centrifugation of whole cell extracts on linear sucrose gradients (Pospisek and Valasek 2013) to separate 40S and 60S ribosomal subunits from 80S monosomes and polysomes. Cycloheximide is included in the lysis buffer to “freeze” polysomes by blocking translation elongation. After centrifugation, the gradient is fractionated and RNA (and/or protein) is prepared from each fraction for subsequent analysis of individual species on blots, or the entire population for identification by hybridization to microarrays or by high throughput RNA sequencing.

Materials

Reagents: Cells (100 mL at $0.5 - 1 \times 10^7$ cells/mL)
Cycloheximide
Protease inhibitors (2 tablets “Complete, EDTA-free, Roche” for 100 mL culture)
Zirconia beads (Biospec)
Polysomal lysis buffer <R>

Equipment: FastPrep device (MP Biomedicals)
Bench top centrifuge with swing-out rotor for 50 mL tubes
Microcentrifuge
Nanodrop spectrophotometer
Phase contrast/bright field microscope with 40 x objective and 10 x eyepieces

Method

1. Harvest $0.5 - 1 \times 10^9$ mid-log phase cells by decanting the culture directly into 50 mL screw cap centrifuge tubes on ice containing cycloheximide to a final concentration of 100 μ g/mL.

Note: Some protocols advise adding cycloheximide 5 min before harvesting to allow time for the inhibitor to work, in which case cells can be pelleted by centrifugation immediately upon harvesting.

2. For lysis, cells are suspended in 200 μL of ice-cold polysomal lysis buffer. Protease and RNase inhibitors are optional but highly recommended.

Note: In one variation of the protocol, cells are washed in 0.9 mL of polysomal lysis buffer.

3. Cell suspensions are transferred to a pre-chilled 1.5-mL screw cap microfuge tube containing approximately 600 μL of 0.5 mm Zirconia beads (Biospec) and lysed in a FastPrep device at level 6.0 for 18 seconds (shorter times down to 13 sec may also be used).

Note (optional but recommended): An aliquot can be checked using a microscope to ensure that at least 50-70% of the cells have been lysed.

4. After lysis, a hole is poked in the bottom of the tube and the lysate is separated from the beads by centrifugation at 5,000 x g for 5 min at 4°C, collecting the liquid into a 1.5 mL microfuge tube.
5. The flow through is combined with 250-350 μL of lysis buffer (depending on desired final concentration and the density of cells at the time of harvest).
6. The lysate is cleared by one centrifugation step at 20,000 x g for 15 min at 4°C.
7. 2 μL of the lysate is added to 98 μL of dH_2O and the OD_{260} is measured in a Nanodrop spectrophotometer.

8. Aliquots corresponding to 20 OD_{260} units (diluted to 350 μL with lysis buffer with or without the protease inhibitor cocktail and RNasin) are layered carefully onto an 11 mL linear sucrose gradient (linear 10 to 50% w/v; prepared using polysomal lysis buffer).

Note: Sucrose gradients can be stored for up to 4 hours at 4°C, with or without lysate layered on top.

9. The sucrose gradients are centrifuged in a Beckman SW 40Ti rotor at 39,000 rpm (~277,000 x g) for 150 min at 4°C.

Note: for a Beckman SW-41Ti rotor, use 35,000 rpm (~246,000 x g) for 160 min.

10. Remove gradients carefully to avoid disturbing and fractionate each one by upward displacement with 55% (w/v) sucrose, prepared using polysomal lysis buffer, using a gradient fractionator (Brendel or ISCO). The precise manipulations will vary depending on the exact combination of fractionator, UV monitor and fraction collector. For the Brendel machine, the flow rate should be set to 800 μL per min; for the ISCO machine, use a pump speed of 50%, 10x (~1.25 mL/ min).
11. Collect 12-13 fractions of 800-900 μL from each gradient. For protein isolation, fractionate into empty glass tubes. For RNA isolation, tubes should be pre-filled with

100% isopropanol to give a final concentration of 70%, mix and store overnight at -20°C.
[Variation: pre-fill tubes with 2 mL of 8 M guanidinium-HCl (final concentration = 5.5 M);
add 1 volume of 100% ethanol, mix by inversion and precipitate overnight at -20°C.]

12. After precipitation, centrifuge for 20 min at >10,000 x g at 4°C and discard the supernatant.

Note: Alternatively, centrifuge at 3,000 x g for 90 min.

13. Optional: Wash pellets with 1 mL 85% EtOH (V/V, diluted with water; see Protocol 1 for tips to avoid losing the pellet).

14. Air-dry RNA pellets for 20 min, then re-suspend in 100 µL of RNase-free water and proceed with RNA purification according to the instructions provided with the Qiagen RNeasy kit (see Protocol 1 for details).

Reagents:

Polysomal Lysis Buffer* 20 mM Tris-HCl (pH 7.5)
50 mM KCl
10 mM MgCl₂

Just before use, add 1 mM DTT (final concentration)
100 µg/mL cycloheximide (freshly prepared)

Optional 0.2 mg/mL heparin (to combat RNase activity; store at -20 °C)
RNasin (Promega; 1 µL/mL according to manufacturer's instructions)
Protease inhibitor cocktail (Roche; 1 tablet "Complete, EDTA-free")

For meiotic cells, add 1 mM PMSF (from a 10 mM stock; general protease inhibitor)

Note: A 10x salt solution containing 200 mM Tris-HCl, 500 mM KCl and 100 mM MgCl₂ can be prepared and stored on a long-term basis if desired.

Treatment with agents that disrupt polysomes:

Puromycin:

Puromycin is a tRNA mimic that causes polypeptide chain termination.

1. Treat cells growing at 32°C with 1 mM puromycin for 15 min.
2. Include 1 mM puromycin and eliminate cycloheximide from the polysome lysis buffer while increasing KCl to 500 mM and decreasing MgCl₂ to 2 mM.

RNase:

RNase treatment cleaves RNA between individual ribosomes.

Add RNase after disrupting cells (25 μ L/500 μ L cells) and incubate for 15 min at RT.
Do not add cycloheximide

EDTA:

EDTA treatment dissociates ribosomal subunits.

Add EDTA to 25 mM (from a 500 mM stock) after disrupting cells (25 μ L/500 μ L cells) and incubate for 15 min at RT.

Do not add cycloheximide

REFERENCES

- Lackner DH, Schmidt MW, Wu S, Wolf DA, Bähler J. 2012. Regulation of transcriptome, translation and proteome in response to environmental stress in fission yeast. *Genome Biol* **13**: R25.
- Pospisek M, Valasek L. 2013. Polysome profile analysis – yeast. *Meth Enzymol* **530**: 173-181.

Protocol 3. 4-thiouridine labeling to analyze mRNA turnover

Juan Mata and Jo Ann Wise.

Correspondence: jm593@cam.ac.uk

ABSTRACT

In general, mRNA half-lives have been measured after inhibition of transcription to allow decay of the pre-existing population. The protocol presented here is a more recently developed strategy in which mRNA turnover is analyzed by measuring the decline in levels of newly synthesized RNA labeled with 4-thiouridine (4sU) during a brief pulse. After RNA extraction, the 4sU is biotinylated and the labeled species are purified using streptavidin beads. Finally, DNA microarrays can be used to compare this population with total RNA, allowing half-lives to be calculated.

Materials

Reagents: Cells (50 - 100 mL at $0.5 - 0.7 \times 10^7$ cells/mL)
0.5 M EDTA
EZ-Link® Biotin-HPDP (Thermo 21341) <R>
Isopropanol
5 M NaCl
MPG buffer <R>
1 M Tris, pH 7.6
4-thiouridine (Sigma-Aldrich # T4509-100MG; 100 X stock) <R>
Dynabeads® M-280 Streptavidin (Invitrogen catalog # 112-05D)
Yeast tRNA (10 mg/mL)
100 mM dithiothreitol (DTT) or β -mercaptoethanol
RNAqueous columns (Ambion # 1931)
RNAqueous kit (Ambion)
Superscript Direct Plus cDNA labeling system (Invitrogen)

Equipment: Bench top centrifuge with swing-out rotor for 50 mL tubes.
Microcentrifuge
Nanodrop spectrophotometer
Magnetic rack for 1.5 mL tubes

Method

In vivo labeling of wild-type cells

1. Grow *ura*⁺ cells to mid-log phase in EMM or other appropriate media
Note: For wild-type cells, growth is carried out at 32°C but another temperature may be more appropriate for conditional mutants.
2. Add 4-thiouridine (4sU) to a final concentration of 75 µg/mL.
3. Incubate culture at 32°C for the duration of the pulse (30-40 min).
Note: Similar results were obtained with shorter pulses (Amorim et al. 2010; Hasan et al. 2014).
4. Centrifuge cells for 3 min at 2,000 x g; discard the supernatant and freeze the pellet at -80°C.

RNA extraction

Prepare total RNA from the 4sU labeled culture according to Protocol 1, scaling up or down as necessary. Use two separate tubes to extract RNA from 5×10^8 cells. The yield of RNA from 2.5×10^8 cells should be 300 – 400 µg. Stop after step 17 and freeze the RNA at -80°C.

RNA biotinylation

The protocol below is for 200 µg of total RNA, which should yield ~2 µg of biotinylated RNA. 100 µg should be sufficient to allow accurate half-life determination, for example in a time course experiment.

1. Prepare the following reaction:

	Volume (µL)
Biotin-HPDP 1mg/mL in DMSO	400
1 M Tris pH7.6	20
0.5 M EDTA	4
200 µg RNA in water	x
ddH ₂ O	2 mL final
2. Incubate at room temperature for 90 min.
3. During the incubation, purify total RNA to be used as the reference and prepare the streptavidin beads as described below.
4. Transfer the biotinylation reaction to phase-lock tubes (500 µL into each).

5. Add 500 μ L of chloroform:isoamyl alcohol (24:1) and mix by inversion.
6. Spin for 15 min in a microcentrifuge at 20,000 x g at RT.
7. Transfer the upper phases (400 μ L from each tube) to 2 mL Eppendorf tubes (800 μ L in each).
Note: Avoid taking the interphase; sometimes the intermediate phase in the phase-lock tubes may not separate well, possibly due to the DMSO in the reaction.
8. Add 1 volume of isopropanol and 1/10 volume of 5 M NaCl.
9. Incubate at -80°C for 30 min (can be left overnight).
Note: This is a good point to stop if you do not want to complete the entire purification in one day.
10. Centrifuge for 15 min at 25,000 g at RT.
11. Discard supernatant, add 500 μ L 75% Ethanol and spin for 1 min. Aspirate most of the supernatant, spin for 5 sec, and remove remaining liquid with a pipette. Air-dry the pellet for 2 min at RT.
12. Resuspend in 20 μ L of water (each tube), pool into a single tube and proceed to the purification step (see below).

Total RNA purification

During the biotinylation reaction, further purify the total thiouridine-labeled RNA as in steps 18-20 of Protocol 1 for use as a reference. Briefly, pass 200 μ g of the RNA preparation through a Qiagen RNeasy column. Elute in 30 μ L of water and use the same water for a second elution. Quantify using a nanodrop spectrophotometer and prepare aliquots containing 20 μ g of RNA in 12 μ L to use in labeling reactions.

Preparation of streptavidin beads

Prepare ~3.5 mL of MPG buffer for the beads and ~3.5 mL for each sample

1. Use 100 μ L beads/sample.
2. Wash 3x with 500 μ L MPG buffer at room temperature (all washes are quick).
3. Re-suspend in 400 μ L MPG buffer.
4. Add 20 μ L tRNA to block the beads.
5. Incubate at RT (with rocking) for at least 30 min.
6. Wash 3x with 300 μ L MPG buffer.
7. Resuspend in 160 μ L MPG buffer.

Purification of biotinylated RNA

1. Heat RNA at 65°C for 5 min before cooling to RT for 1 min.
2. Add 160 μ L of streptavidin beads equilibrated in MPG buffer as described above to 40 μ L of biotinylated RNA.
3. Incubate at RT for 40 min with rocking.
4. Place tube in magnetic rack and remove the buffer once all the beads have been attracted to the side. Add 500 μ L MPG preheated to 65°C (added hot, then incubated at RT for 5 min with rocking).
5. Repeat step 3 twice more.
Note: If there are problems with bead retrieval, the best strategy is to find a stronger magnet. Alternatively, tubes can be centrifuged for 3 sec before placing in the rack.
6. Wash 3x with 500 μ L MPG at RT (5 min with rocking).
7. Elute RNA from beads by reducing the linker with 5% β -mercaptoethanol (BMETH). Resuspend the beads in 50 μ L of BMETH and incubate at RT for 5 min.
Note: 100 mM DTT can be used in place of 5% BMETH but the latter is preferred.
8. Recover the supernatant.
9. Repeat steps 6 and 7.
10. Pool the supernatants from steps 8 and 9.
11. Further purify RNA from the pooled supernatants using RNAqueous micro columns.
12. Add 5 volumes of RNAqueous lysis buffer (500 μ L) to the sample, mix and add 2.5 volumes of ethanol (250 μ L).
13. Purify as described in the RNAqueous manufacturer's protocol. Load mixture onto columns in two rounds (the volume is too large to fit into the column all at once).
14. Wash 3 times with 180 μ L of wash solution I, then twice with solution 2/3. In between washes, centrifuge at 20,000 x g for 1 min.
15. Switch column to a new eppendorf tube and centrifuge for 1 min at 20,000 x g to dry the column.
16. Elute twice with 8 μ L of RNase free water at 85 °C in the same microfuge tube. For elution, incubate the column and microfuge tube at 85°C for 1 min. Use 1 μ L to quantify the RNA in a spectrophotometer.
17. Use the remainder of the RNA (should be ~12 μ L) for half-life analysis by microarray analysis.

Half-life determination:

Fission yeast mRNA half-lives can be determined by analyzing data from 4sU labeling experiments using equations developed for studies in other organisms (Dolken et al 2008). Briefly, microarray analysis is used to produce an unnormalized measure of the relative abundance of a given 4sU-labeled (nascent) transcript compared with its abundance in total mRNA. See Amorim et al. (2010) and Hasan et al. (2014) for additional details.

Reagents: EZ-Link® Biotin-HPDP
4-thiouridine Prepare a 7.5 mg/mL solution (100X stock) in ddH₂O; can be stored at -20°C but generally better to prepare fresh

MPG buffer:
100 mM Tris pH7.6
1 M NaCl
10 mM EDTA
Prepare using Ambion nuclease free H₂O.

4-thiouridine:
Prepare a 1 mg/mL solution in DMSO (store at 4°C and avoid excessive freezing and thawing).

REFERENCES

- Amorim MJ, Cotobal C, Duncan C, Mata J. 2010. Global coordination of transcriptional control and mRNA decay during cellular differentiation. *Mol Syst Biol* **6**: 380.
- Dolken L, Ruzsics Z, Radle B, Friedel CC, Zimmer R, Mages J, Hoffmann R, Dickinson P, Forster T, Ghazal P, Koszinowski UH. 2008. High-resolution gene expression profiling for simultaneous kinetic parameter analysis of RNA synthesis and decay. *RNA* **14**: 1959-1972.
- Hasan A, Cotobal C, Duncan CD, Mata J. 2014. Systematic analysis of the role of RNA-binding proteins in the regulation of RNA stability. *PLOS Genetics* **10**: e1004684.

ACKNOWLEDGMENTS

Protocol 1 (total RNA isolation) originated in Jürg Bähler's laboratory (University College London, UK); variations noted here were provided by Dom Helmlinger (Centre de Recherche de Biochimie Macromoléculaire, Montpellier FR) and Crispin Miller (Cancer Research UK, Manchester Unit). Protocol 2 (polysome profiling) is an amalgamation of procedures provided by Dieter Wolf (Sanford-Burnham Medical Research Institute, La Jolla CA USA) and Jürg Bähler. Protocol 3 (mRNA half-life analysis) was contributed by Juan Mata (University of Cambridge, UK). The authors would also like to thank the following investigators for supplying protocols that could not be included due to space constraints but were useful in compiling the introductory material: Weijun Chen and Melissa Moore (University of Massachusetts Worcester, affinity purification of large RNA-protein complexes); Dom Helmlinger (qPCR); Juan Mata (RIP-chip); and Crispin Miller (high throughput RNA sequencing). We also thank Charalampos Rallis (University College London, UK) for comments on Protocols 1 and 2.