

ILC2017-RS-3314

***Cirrhosis and its complications: Clinical aspects*****Portosystemic shunts in cirrhosis are associated to more complications and deteriorated quality of life. An international cohort study.**

Macarena Simón-Talero<sup>1</sup>, Davide Roccarina<sup>2</sup>, Avik Majumdar<sup>2</sup>, Emmanuel A. Tsochatzis<sup>2</sup>, Javier Martínez<sup>3</sup>, Carmen Picón<sup>4</sup>, Agustín Albillos<sup>3</sup>, Katharina Lampichler<sup>5</sup>, Daniel Toth<sup>5</sup>, Thomas Reiberger<sup>5</sup>, Anna Baiges<sup>6</sup>, Anna Darnell<sup>7</sup>, Virginia Hernandez-Gea<sup>6</sup>, Gavin Low<sup>8</sup>, Juan G. Abraldes<sup>9</sup>, Puneeta Tandon<sup>9</sup>, Elba Llop<sup>10</sup>, Marta Lopez<sup>10</sup>, Jose Luis Calleja<sup>10</sup>, Michael Praktijn<sup>11</sup>, Guido Kukulj<sup>12</sup>, Jonel Trebicka<sup>11</sup>, Martin Maurer<sup>13</sup>, Annalisa Berzigotti<sup>14</sup>, Alexander Zipprich<sup>15</sup>, Cristina Ripoll<sup>15</sup>, Michela Triolo<sup>16</sup>, Vincenzo La Mura<sup>16</sup>, Guillaume Vangrinsven<sup>17</sup>, Wim Laleman<sup>17</sup>, Rita Garcia-Martinez<sup>18</sup>, Rafael Bañares<sup>18</sup>, Annette Dam<sup>19</sup>, Aleksander Krag<sup>19</sup>, Salvador Augustin<sup>1</sup>, Joan Genescà<sup>1</sup> and the Baveno VI-SPSS group

<sup>1</sup>Liver Unit, Department of Internal Medicine, Hospital Universitari Vall d'Hebron, VHIR, Universitat Autònoma de Barcelona, CIBERehd, Barcelona, Spain, <sup>2</sup>Sheila Sherlock Liver Unit and UCL Institute for Liver and Digestive Health, Royal Free Hospital and UCL, London, United Kingdom, <sup>3</sup>Department of Gastroenterology and Hepatology, Hospital Universitario Ramón y Cajal, IRICYS, Universidad de Alcalá, CIBERehd, <sup>4</sup>Department of Radiology, Hospital Universitario Ramón y Cajal, IRICYS, Universidad de Alcalá, Madrid, Spain, <sup>5</sup>Division of Gastroenterology and Hepatology, Vienna Hepatic Hemodynamic Lab, Medical University of Vienna, Vienna, Austria, <sup>6</sup>Hepatic Hemodynamic Laboratory, Liver Unit, Hospital Clinic, IDIBAPS, Universitat de Barcelona, CIBERehd, <sup>7</sup>Department of Radiology, Hospital Clinic, IDIBAPS, Universitat de Barcelona, Barcelona, Spain, <sup>8</sup>Department of Radiology, University of Alberta, <sup>9</sup>Cirrhosis Care Clinic, University of Alberta, Edmonton, Canada, <sup>10</sup>Liver Unit, Hospital U. Puerta de Hierro, Universidad Autónoma de Madrid, CIBERehd, Madrid, Spain, <sup>11</sup>Department of Internal Medicine I, University of Bonn, <sup>12</sup>Department of Radiology, University of Bonn, Bonn, Germany, <sup>13</sup>Department of Radiology, Inselspital, University of Berne, <sup>14</sup>Hepatology, Inselspital, University of Berne, Berne, Switzerland, <sup>15</sup>First Department of Internal Medicine, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany, <sup>16</sup>First Division of Internal Medicine, IRCCS Policlinico San Donato, University of Milan, Milan, Italy, <sup>17</sup>Department of Gastroenterology and Hepatology, University Hospitals Leuven, Leuven, Belgium, <sup>18</sup>Liver Unit, Hospital General Universitario Gregorio Marañón, Universidad Complutense, CIBERehd, Madrid, Spain, <sup>19</sup>Department of Gastroenterology and Hepatology, Odense University Hospital, Odense, Denmark

Submitting author's email: [msimon@vhebron.net](mailto:msimon@vhebron.net)

**<p><b> Young Investigator Bursary</b></p><br>Do you want to apply for a Young Investigator Bursary?: Yes  
<p><b>Young Investigator Award 2017</b></p><br>Do you want to apply to the YI Award?: No**

**Background and Aims:** Spontaneous portosystemic shunts (SPSS) have been reported in hepatic encephalopathy (HE). However, their prevalence in cirrhosis and association to decompensating events have not been assessed. The aims of this study were (i) to determine the prevalence of SPSS in cirrhosis and (ii) to assess the impact of SPSS on clinical course.

**Methods:** Consecutive cirrhotic patients submitted to abdominal angioCT/angioMR (2010-2015) were included in a retrospective, multicenter, international study. Advanced hepatocellular carcinoma, neurological, psychiatric or terminal illness were excluded. Patients were classified into 3 groups: large SPSS (L-SPSS,  $\geq 8$  mm), small SPSS (S-SPSS,  $< 8$  mm) or without SPSS (W-SPSS).

**Results:** 2978 patients were assessed, 1785 included. The median follow-up was 19.4 months (IR 7.9-34.9). Child-Pugh A 42%, B 38% and C 20%. Prevalence of shunts: 26% L-SPSS, 30% S-SPSS and 44% W-SPSS. Type of L-SPSS more often identified: splenorenal (47%), umbilical (26%). Distribution by Child was different (L-SPSS, S-SPSS, W-SPSS): Child A 24%, 30%, 46%; Child B 35%, 33%, 32%; Child C 41%, 38%, 21% ( $p < 0.001$ ).

Patients with L-SPSS had more HE (prior to CT, at the time of CT and during follow-up) compared to S-SPSS, and S-SPSS compared to W-SPSS (L-SPSS 58%, S-SPSS 42%, W-SPSS 27%,  $p < 0.001$ ). They also had a higher percentage

of recurrent/persistent HE (L-SPSS 50%, S-SPSS 40%, W-SPSS 31%,  $p<0.001$ ). There were no differences in HE grade (West-Haven criteria); however, differences in quality of life were found (autonomy according mRS: L-SPSS 76%, S-SPSS 81%, W-SPSS 91%,  $p<0.001$ ).

The stratified analysis by liver function showed that HE was more frequent in Child A (L-SPSS 35%, S-SPSS 17%, W-SPSS 10%,  $p<0.001$ ) and B (L-SPSS 62%, S-SPSS 45%, W-SPSS 39%,  $p<0.001$ ); recurrent/persistent HE was also related to SPSS in each Child class. The effect on quality of life of SPSS was observed in Child B and C. Child A patients with SPSS had also more ascites and bleeding ( $p=0.03$ ). Transplant-free-survival was not different according to the presence of SPSS in the 3 stages of Child.

**Conclusions:** SPSS are very common in cirrhosis and their prevalence increases with the deterioration of liver function. The most frequent SPSS found is splenorenal. The presence of SPSS is associated with frequent and severe HE-episodes and poorer quality of life, and more portal hypertension complications (ascites/bleeding) in Child A. No association with mortality was observed.

**Disclosure of Interest:** None Declared