



Open Research Online

The Open University's repository of research publications and other research outputs

Hambleton – a new sulphur rich pallasite

Conference or Workshop Item

How to cite:

Johnson, D.; Hutchison, R.; Kirk, C. and Grady, M. M. (2006). Hambleton – a new sulphur rich pallasite. In: 69th Annual Meeting of the Meteoritical Society, 6-11 Aug 2006, Zurich, Switzerland.

For guidance on citations see [FAQs](#).

© [not recorded]

Version: [not recorded]

Link(s) to article on publisher's website:

<http://www.lpi.usra.edu/meetings/metsoc2006/pdf/5216.pdf>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

HAMBLETON - A NEW SULPHUR RICH PALLASITE

D.Johnson¹, R.Hutchison², C Kirk² and M.M.Grady^{1,2}. ¹Open University, Walton Hall, Milton Keynes, MK76AA, United Kingdom. E-mail: D.Johnson@open.ac.uk. ²Natural History Museum, London, SW7 5BD, UK.

Introduction: A new pallasite, a single mass of 17.6 kg, was found south of Hambleton, North Yorkshire, by R and I Elliott in August 2005. The mass is composed of ~60 vol % olivine, ~25 vol% metal and ~15 vol% sulphide. The phases are irregularly distributed and highly weathered. There follow the results of a study by optical and analytical scanning electron microscopy.

Observations: Olivine occurs as cm-sized sub-rounded crystals in a granular mosaic. Many contain sub-parallel sets of fractures, some of which are annealed, while others are filled with metal or sulphide. In metal-rich or sulphide-rich areas olivines are fragmented and angular to sub-angular and veined by metal or sulphide respectively. Some regions <5cm in size are composed entirely of olivine crystals enclosed within troilite. Olivine is Fo_{88.3}, and together with the oxygen isotopic ratios: $\delta^{17}\text{O} = +1.383\text{‰}$; $\delta^{18}\text{O} = +3.029\text{‰}$; $\delta^{17}\text{O} = -0.187\text{‰}$, indicate that the meteorite is a main group pallasite. From the olivine-rich exterior, weathering has penetrated for 4-5 cm towards the interior of the mass. The weathered, olivine-rich outer portion is brittle and prone to disintegration. A blue secondary mineral rich in Mg, P and Fe was shown by XRD to be baricite (Mg, Fe)₃(PO₄)₂.8H₂O. Much of the metal has succumbed to terrestrial oxidation, especially low-Ni phases such as kamacite, cloudy taenite or plessite. The sulphide is more susceptible to terrestrial alteration than the metal.

Discussion: Metal rich regions are consistent with the view of Scott [1] that pallasites formed by the injection of metallic liquid into dunite. Evolved metallic melts, related to IIIAB irons, should be sulphur-rich. Paucity of sulphide in pallasites led Uff-Møller et al [2] to suggest that either FeS-rich liquid was lost or formed pallasites that are underrepresented in our samples.

Conclusion: With Phillips County (pallasite), Hambleton is a rare FeS-rich pallasite.

References:

- [1] Scott E.R.D. 1977. *Geochimica et Cosmochimica Acta* 47: 693-710. [2] Uff-Møller et al. 1998. *Meteoritics & Planetary Science* 33: 221-227.