

Muscle strength can better differentiate between gradations of functional performance than muscle quality in healthy 50 – 70y women.

Short Title: Muscle health and functional performance

Peter Francis^{1,2*}, William Mc Cormack², Clodagh Toomey^{2,3}, Mark Lyons², Philip Jakeman²

¹Musculoskeletal Health Research Group, School of Clinical and Applied Sciences, Leeds Beckett University, LS13HE, United Kingdom, ²Department of Physical Education and Sport Sciences, Faculty of Education and Health Sciences, University of Limerick, Ireland. ³Faculty of Kinesiology, University of Calgary, Alberta, Canada.

***Corresponding author:** Peter Francis, School of Clinical and Applied Sciences, Leeds Beckett University, Leeds, LS13HE, UK. E-mail: peter.francis@leedsbeckett.ac.uk

ABSTRACT

OBJECTIVE: It is not known which laboratory indices of muscle mass, strength or quality can distinguish functional performance in healthy middle aged women. The aim of this study was to a) examine the association between upper leg lean tissue mass (LTM), knee extensor strength, muscle quality (strength per unit LTM) and functional performance and b) to determine the utility of tertiles of muscle strength and muscle quality to distinguish gradations of functional capacity in healthy 50 – 70y women. **METHOD:** Using a cross-sectional study design, one hundred and twenty-eight healthy 50 – 70y women (mean age: 60.4 SD=5.1 years) underwent body composition assessment (dual X-ray absorptiometry) and performed a maximal voluntary isometric contraction of the knee extensors (Con-Trex Dynamometer). Functional performance was assessed using a 5 repetition and 30 second chair rise test and 900m gait speed test. **RESULTS:** Ordered by muscle strength or muscle quality, those in the highest tertile (T1) demonstrated greater functional performance than those in lowest tertile (T3). There was no association between upper leg LTM and functional performance ($r \leq 0.06$). Muscle strength explained a greater proportion of the variance in all functional performance measures relative to muscle quality ($R^2=0.13$ to 0.36 vs. $R^2=0.11$ to 0.16) **CONCLUSION:** Upper leg LTM is not associated with physical performance in

healthy 50 – 70y women. These results suggest strength relative to the body mass being accelerated distinguishes gradations in functional performance better than muscle quality healthy 50-70y women.

Keywords: healthy ageing; muscle function; knee extensor; physical performance; older adult.

Bullet Points

- Muscle strength can better differentiate between gradations of functional performance than muscle quality.
- Muscle strength normalised to body mass may provide clinicians with an objective measure of muscle health that does not require imaging technology.
- Extended gait speed may be a clinically useful measure of functional performance in healthy 50 – 70y women.
- Functional performance measures that allow healthy adults perform close to maximum may be useful for clinicians attempting to distinguish gradations of capacity.

Introduction

Expert working groups^{1,2} have recommended older adult muscle health is evaluated in terms of mass, strength and functional performance. Laboratory measures of muscle mass, strength and quality which are considered to be below 'normal' are associated with functional impairment in the elderly³⁻⁵. Low relative muscle mass, knee extensor strength and muscle quality (strength per unit tissue) have been associated with self-reported and objectively measured functional limitation in older adults³⁻⁷. These associations have been studied extensively in older adults >65y^{6,8-10} and as a result tend to use simple tasks related to activities of daily living (ADL) such as gait speed tests <6m¹¹⁻¹⁵.

However, it is well documented that age-related change in muscle characteristics which precede functional limitation become noticeably different from a young adult aged ~50y^{16,17}. By contrast comparatively little is known about the time course and transition to functional limitation after the age of ~50y. We previously reported age-related difference in lean tissue mass (LTM), muscle strength and quality (strength per unit tissue) in healthy older women¹⁸. Furthermore, our research group¹⁹ and others²⁰ identified short gait speed tests (<10m) as unsuitable measures of functional performance for use in healthy older adults. This is primarily due to short performance tests inability to detect age-related difference where expected in healthy adults. Instead, we proposed that extended gait speed and chair rise tests are used due to their greater construct validity in detecting age-related difference and greater association with maximal voluntary knee extensor strength¹⁹.

Ideally, laboratory measures of muscle mass, strength and quality which can distinguish gradations of capacity with aging would also reflect functional performance. Lower body strength is thought to undergo a preferential age-related decline relative to the upper body. Specifically, quadriceps strength declines at a greater rate than grip strength in healthy volunteers²¹. This preferential decline is thought to be due to a change in activity patterns^{22,23} and is greatest in women²⁴. Hicks et al.¹¹, have suggested that establishing cut points from strength at one time point is predictive of future mobility decline in a population based sample of older (~73y) men and women. Researchers have recently^{12,25} classified healthy older (65 – 84y) adults into tertiles based on body composition, muscle function and functional performance. This approach may be more appropriate for attempting to distinguish gradations of capacity in healthy adults whereby cut-points below 'normal' in muscle health may not be able to identify individuals of lower physical capability²⁴.

At present, there are two indices of lower extremity muscle function most commonly reported in the literature. The first is knee extensor strength which is normally assessed by a commercially available dynamometer and the second is muscle quality normally expressed as knee extensor strength per unit muscle or LTM mass. Both indices have been associated with functional performance in older (>65y) adults. Hairi et al.⁵ reported that muscle quality had a stronger association than muscle mass with both self-reported and objectively measured physical disability in community dwelling older (≥ 70 y) men (n=1,612). However, the authors also identified muscle strength (grip and quadriceps strength) as having stronger associations with all outcome measures than either muscle mass or muscle quality. Similarly, Visser et al.²⁶ reported lower muscle strength and greater fatty infiltration as stronger predictors of self-reported mobility limitation than muscle mass in well-functioning older (70 – 79y) adults (n=2,631). More recently, Hayashida et al.²⁷ reported knee extension strength but not appendicular LTM to be associated with maximal gait speed (11m) in 318 community dwelling Japanese men and women (≥ 65 y).

Whilst muscle strength can be determined using a variety of cost effective techniques which are often portable, muscle quality requires imaging technology, often at significant time, financial and training cost. The association between both indices and functional performance has received considerably less attention in adults <70y of age. Furthermore, it is not yet known whether one index is superior to the other in distinguishing gradations of capacity in healthy middle aged adults.

The aim of this study was to examine the association between upper leg LTM, knee extensor strength, muscle quality (strength per unit upper leg LTM) and functional performance in healthy 50 -70y women. The secondary aim of this investigation was to determine the utility of tertiles of muscle strength and muscle quality to distinguish gradations of functional capacity in healthy 50 – 70y women.

Methods

Participants

A convenience sample (n=204) of healthy older (50 – 70y) adults were recruited via email and word of mouth from the University of Limerick campus community and surrounding area to take part in the University of Limerick Aging Study^{18,19,28,29}. For the present investigation, one hundred and twenty-eight women participated in a cross-sectional study design.

Participants received a full medical screening and physical examination prior to the assessment of body composition, maximal voluntary knee extensor strength and functional performance. Those defined as healthy, i.e. disease free based on Greig et al.³⁰ and independent living were invited to participate. Disease free included the absence of clinical, cardiovascular or musculoskeletal abnormality as determined by a medical doctor. Participants were required to be healthy but not masters athletes as defined by Drey et al.³¹. After receiving a complete explanation of the procedures, benefits and risks of the study, all participants gave their written informed consent. This study was approved by the Research Ethics Committee of the University of Limerick (EHSREC 10/45).

Measures

Whole body and regional body composition analysis was conducted using dual X-ray absorptiometry (iDXATM; GE Healthcare, Chalfont St Giles, Buckinghamshire, UK), in accordance with procedures used in the University of Limerick Body Composition Study^{28,32}. To standardise test conditions and tissue hydration, participants were instructed to refrain from strenuous exercise in the 12-hour period before testing and to attend after an overnight fast. Participants consumed 500ml of water one hour prior to the scan and were instructed to void and defecate, if required, immediately prior to measurement. Height was measured to the nearest 0.1 cm (SECA stadiometer) and body mass to the nearest 0.1 kg (Tanita MC-180MA Body Composition Analyzer, Tanita UK Ltd.). Daily calibration of the scanner employed a phantom spine containing composites of bone, fat and lean tissue mass. Participants wore light clothing and removed all metal objects such as belts and jewellery. The thigh, representing upper leg LTM was measured from the inferior side of the lesser trochanter until the tibio-femoral joint.

Reliability of estimate was calculated using a convenience sub-sample of participants who were asked to take part in the reliability study when attending a 6-month or 1 year follow up as part of the healthy aging study. Reliability of estimate was calculated in accordance with recommendations from the International Society of Clinical Densitometry (Baim et al., 2006). The root mean square coefficient of variance (RMSCV) of the iDXA for repeated measures of whole body composition analysis was 0.6%. Repeat scans on a sub-sample of 87 participants between 18-70y (mean age: 35.0 SD=17.0 years) was used to assess reliability of whole body LTM and total leg LTM. The RMSCV was found to be 0.6% and 1.4% respectively. Of the 87 participants 22 were between the ages of 50 – 70y. The older adult

scans (n=22) were further segmented to provide the additional measurement of upper leg LTM. The RMSCV was found to be 0.7%, 1.4% and 2.3% respectively for whole body, total and upper leg LTM respectively.

Maximal voluntary isometric contractions of the knee extensors of the dominant limb were measured using a Con-Trex MJ Dynamometer (Con-Trex MJ; CMV AG, Dubendorf, Switzerland) in accordance with procedures used by Francis et al.¹⁹. The participants were tested during two identical sessions held 7 days apart. Warm up consisted of 5 minutes on a bicycle ergometer (Monark Ergomedic; 828E), at a workload of 40 watts in accordance with the methodology of Lynch et al.¹⁷. The knee extensors and flexors of the dominant limb (the limb used to kick a ball) were tested. Participants were seated with a hip flexion angle of 110°. The back of the knee joint was on the edge of the seat with a knee angle of -60° from anatomical zero (180°), which has been demonstrated to be the angle of maximal isometric force generation³³. The distal shin pad of the dynamometer was attached 4-5cm proximal to the medial malleolus using a velcro strap. Lever length (mean length: 22.7 SD=2.3 cm) was recorded as in Lynch et al.¹⁷. Two seatbelts were applied across the chest and pelvis while velcro straps were applied to the mid-thigh to reduce extraneous movement during contractions. The dynamometer rotational axis was aligned with the lateral femoral condyle (knee joint axis of rotation). The absolute zero (point at which gravity is acting vertically) was recorded prior to each trial. Participants were instructed to perform 2 submaximal voluntary isometric contractions (50 and 75% of perceived maximum) prior to each test series as in Maffiuletti et al.³⁴, with a 1 minute rest period in between. The participant then performed 3 MVC's of the knee extensors separated by 2 minutes of stationary rest. The participant was instructed to consistently produce their maximal force rapidly (as hard and as fast as possible in the sagittal plane) and to maintain that force for 3-4 seconds. Participants received a 5 second count down with a distinct emphasis on "Go". No overt verbal encouragement was provided due to the difficulty in standardising it for all participants³⁵. Visual feedback of the instantaneous dynamometer torque was provided to the participants on a computer screen. This was used to show the participant the type of contraction required and also to provide encouragement to reach their maximum as in Bazzucchi et al.³⁶. The participant then received 2-3 minutes of stationary rest to allow set up of the protocol to measure MVC of the knee flexors. To assess the knee flexors, the distal shin pad was removed and replaced along the posterior portion of the triceps surae; 4 -5cm above the

medial malleolus. The participant was then instructed to repeat the procedure used for extension above whilst kicking back (flexion) in the sagittal plane.

The coefficient of variance (CV) for the knee extensors and flexors was 3 % in both cases with a range of 0-9%. There was an increase in the number of participants with repeated measures which satisfied the criteria for a MVC (knee extensors (n=9) and flexors (n=7)) on day 2 (CV 3%; range 0-6%). A 2-way mixed model intra class correlation coefficient (ICC) was used to assess absolute agreement. The ICC on day 1 and day 2 for the knee extensors (0.99 (95% CI 0.99 to 1.00) and 1.00 (95% CI 1.00 to 1.00) respectively) and flexors (0.99 (95% CI 0.99 to 1.00) and 0.99 (95% CI 0.99 to 1.00) respectively) demonstrate excellent within day reliability. The intra-class correlations between testing sessions for knee extensors and flexors were 0.85 (95% CI 0.79 to 0.90) and 0.887 (95% CI 0.83 to 0.92) respectively. The highest value for both days was used for analysis in this study. Muscle quality was expressed as knee extensor torque per kilogram upper leg LTM as this has been reported as the most appropriate index of muscle quality when using DXA and isokinetic dynamometry as the measurement tools¹⁹.

The ability to rise from a chair was assessed using a timed 5 repetition chair rise test³⁷ and by counting the number of chair rises completed in 30 seconds³⁸. Maximal extended gait speed, using a one or a combination of walking, jogging or running, was assessed by the time taken to complete 900m (4 laps of a 225m indoor track)^{19,29}. All measures were repeated separated by 7 days in order to reduce the potential for a learning effect. The highest measures from both days are used to report tertiles of functional performance and to examine associations with LTM, muscle strength and muscle quality. All measurements were carried out by the same exercise scientist to exclude issues with inter-tester reliability.

Analysis

Physical characteristics, body-composition, muscle strength and functional performance data were checked for normality of distribution by using a Kolmogorov-Smirnov test or Shapiro-Wilk test and expressed as means and SDs for normally distributed variables and medians (IQRs) for asymmetric distributed variables. Functional performance was classified per tertile of muscle strength and muscle quality. A one-way ANOVA or Kruskal Wallis test was used to determine if differences existed between tertiles. Pairwise comparison resulting from a Bonferroni post hoc test were used to determine differences between tertiles. The association between the independent variables (upper leg LTM, muscle strength and muscle quality) and

the dependent variables (functional performance) were assessed using a Pearson's or Spearman's correlation for normal and non-normal data respectively. Correlations were classified based on British Medical Journal recommendations³⁹: r : 0-0.19 very weak, 0.2-0.39 weak, 0.40-0.59 moderate, 0.6-0.79 as strong and 0.8-1 very strong. Linear regression was used to explain the proportion of variance in functional performance explained by muscle strength and muscle quality. The data were analysed using SPSS 22.0 for Windows (SPSS, Inc., Chicago, IL, USA).

Results

Physical characteristics, body composition and functional performance of 50 -70y women are displayed in Table 1. Functional performance classified per tertile of muscle strength and muscle quality are displayed in Table 2. Classified by muscle strength or muscle quality, those in the highest tertile (T1) had a higher functional performance across all tests than those in lowest tertile (T3) ($P < 0.05$). Classified by tertile of muscle strength extended gait speed was different between those in the highest tertile, middle tertile and lowest tertile. Classified by tertile of muscle quality, those in the lowest tertile were different to those in the middle and highest tertile for extended gait speed and time taken to complete 5 chair rises. Those in the middle and highest tertile of muscle quality did not differ in functional capability.

There was no association between upper leg LTM and functional performance ($r = 0.007$ to 0.061 , $P > .05$). Muscle strength demonstrated a moderate association with extended gait speed (Figure 1) and the number of chair rises undertaken in 30 seconds and a weak association with the time taken to complete 5 chair rises (Table 3). Muscle strength was associated with all functional performance measures in the highest tertile and with gait speed and the number of chair rises completed in 30 seconds in the lowest tertile but was not associated with functional performance in the middle tertile.

Muscle quality demonstrated a moderate association with extended gait speed and a weak association with both chair rise tests. Muscle quality was associated with the time taken to complete 5 chair rises in the highest tertile but was not associated with any other functional performance measure in any other tertile. Compared to muscle quality, muscle strength explained a greater proportion of the variance in extended gait speed (36% vs. 16%), the number of chair rises completed in 30 seconds (22% vs. 15%) and the time taken to complete 5 chair rises (13% vs. 11%).

Discussion

Compared to muscle quality, knee extensor strength per kg body mass had a stronger association with functional performance and could better differentiate between those of high, intermediate and lower functional performance in healthy 50 – 70y women.

Interestingly, classified by tertile of muscle strength or muscle quality those in the highest tertile could be differentiated from those in the lowest tertile in terms of functional performance. It became more challenging to separate those in the middle tertile from those in the highest or lowest; in fact, only gait speed classified by muscle strength could provide this differentiation. Furthermore, knee extensor strength was associated with functional performance in the highest and lowest tertiles of muscle strength but muscle quality was generally not associated with functional performance in any tertile. The absence of an association between strength and functional performance in the middle tertile perhaps suggests a non-linear relationship between strength and functional performance. Buchner et al.⁴⁰ reported an association between strength and gait speed (15.2 m) in weaker participants but not in stronger participants. The authors suggested that small changes in physiological capacity of frail older adults, may lead to large changes in functional performance whereas small changes in physiological capacity of stronger adults, may lead to little or no change in functional performance. Our results suggest that this may be the case in the lowest tertile but also that small changes in strength for healthy adults in the highest tertile may lead to significant changes in functional performance. This has recently been demonstrated in well-trained distance runners who experienced an increase in running economy as a result of a strength training intervention⁴¹. Buchner et al.³⁸ may not have seen this effect at the higher end due to using a population based sample that allowed for participants with musculoskeletal conditions. These explanations whilst plausible need to be interpreted cognisant of the sample size used to derive the associations in each tertile (n=42-43).

As reported by Visser et al.²⁶, Hairi et al.⁵ and Hayashida et al.²⁷ in older (≥ 65 y) adults, we report muscle strength and quality to have a greater association with functional performance than muscle mass in healthy 50 – 70y women. However, compared to muscle quality, we report knee extensor strength corrected for body mass to have a stronger association with all functional performance measures. This suggestion has been supported by Hairi et al.⁵ when investigating the association between knee extensor strength and 6m gait speed and by Vilaça et al.²⁵ using the 6 minute walk test in older (≥ 65 y) adults. In our study, it is perhaps expected

that as the difficulty of the test increases towards maximum so does the association between performance of the test and maximal voluntary knee extensor strength. Strength per unit LTM is considered to be an accurate representation of a muscles force generating capacity. The lower association between muscle quality and functional performance is perhaps because the upper leg LTM does not represent the body mass required to be elevated from a chair or accelerated during a gait speed test. Furthermore, we previously reported no difference in LTM between women in the 6th and 7th decade of life¹⁹ which may suggest that changes in muscle quality are a function of the underlying strength rather than mass^{25,26}. However, it must be acknowledged that we have used DXA to estimate LTM which has been shown to underestimate age-related change in skeletal mass compared to gold standard methods such as MRI and CT⁴².

Previously, Buchner et al.⁴⁰ have reported lower extremity strength to account for ~17 – 20% of short (<16m) gait speed tests in older adults (50 – 96y). To the authors knowledge, the 900m gait speed test is the first gait speed test reported in which >30% of the variance in performance among healthy older women can be explained by knee extensor strength. Our test which requires a similar time (~6 – 7 minutes) to complete as the 6 minute walk test is perhaps a step in the right direction toward the call for the generation of maximal gait speed reference values in healthy adults²⁵. The fact that lower extremity strength has a strong association with a maximal gait speed test which has a range of 1.3 – 3.9 m/s is encouraging. This suggests that the relative muscular effort for those with the mean gait speed (2.3 m/s) is considerably less when performing a habitual gait speed (~1.4 m/s) for extended periods of time. The relative effort required to perform tasks related to activities of daily living (ADL) could be a key determinant as to how long quality of live and independent living are preserved.

Our study makes a contribution to knowledge about the association between laboratory measures of LTM, muscle strength, muscle quality and functional limitation in healthy late middle aged ²⁰ adults who are not yet frail. The applicability of our study is limited by our small (n=128) convenience sample of healthy older women recruited from the University campus community and surrounding area. This may be especially important as education and socioeconomic status have been reported to influence the health status of a population⁴³. The mean habitual gait speed (1.4 m/s) and dietary protein intake (1.2g ·kg⁻¹ BW·day⁻¹) of participants in the University of Limerick Healthy Aging Study demonstrate the relative physical and nutritional health of this cohort compared to more representative cohorts

(dietary protein intake $0.8 - 1.1 \cdot \text{kg}^{-1} \text{BW} \cdot \text{day}^{-1}$ and habitual gait speed 0.8m/s)^{44,45} which may introduce a health bias. Furthermore, we did not assess cognitive function, stage of the menopause or control for habitual physical activity and as such it is unknown how these confounding variables may have affected the associations reported.

Although 900m gait speed had the strongest association with muscle strength and quality, the self-selected mode of maximal gait speed which included walking, jogging or running may have over or underestimated our gait speed and therefore influenced the strength of the associations reported. Furthermore, endurance performance is also dependent on cardiorespiratory and muscle oxidative capacity which we did not control for. The associations reported for all muscle and functional performance parameters must be interpreted cognisant of the cross-sectional study design and therefore survival bias may have influenced the results.

Conclusion

The fact that cross-sectional strength measurement has been shown to be predictive of mobility decline¹¹, that the lower limbs are most affected^{21,24} should make lower limb strength assessment a priority for researchers generating reference values for muscle health and clinicians in practice. As Dulac et al.⁴⁶ have recently reported using handgrip strength, we report knee extensor strength normalised to body mass as an index that could provide a cost effective and simplistic measure of muscle health which could be used in a clinical setting and can represent a significant proportion of the variance in functional capacity. This may be useful to clinicians attempting to assess the functional capability of older adults in the community. By contrast our results suggest that muscle quality assessment, requiring expensive imaging technology may be of lesser value in relation to distinguishing functional performance. Further work is required to determine whether these differences remain when using MRI or CT to determine skeletal muscle mass.

Acknowledgements

The authors would like to thank the University of Limerick for funding this study, specifically the 'Roadbridge Medical Research Scholarship' which funded the authors PhD research. This study was also supported by Food for Health Ireland and Enterprise Ireland grant CC20080001.

Conflict of Interest On behalf of all authors, the corresponding author states that there is no conflict of interest aside from grant funding stated above.

Statement of human and animal rights All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent Informed consent was obtained from all individual participants included in the study.

References

1. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing*. July 1, 2010 2010;39(4):412-423.
2. Fielding RA, Vellas B, Evans WJ, et al. Sarcopenia: An Undiagnosed Condition in Older Adults. Current Consensus Definition: Prevalence, Etiology, and Consequences. International Working Group on Sarcopenia. *Journal of the American Medical Directors Association*. 2011;12(4):249-256.
3. Janssen I, Heymsfield SB, Ross R. Low Relative Skeletal Muscle Mass (Sarcopenia) in Older Persons Is Associated with Functional Impairment and Physical Disability. *J. Am. Geriatr. Soc.* 2002;50(5):889-896.
4. Ostchega Y, Dillon CF, Lindle R, Carroll M, Hurley BF. Isokinetic Leg Muscle Strength in Older Americans and Its Relationship to a Standardized Walk Test: Data from the National Health and Nutrition Examination Survey 1999–2000. *J. Am. Geriatr. Soc.* 2004;52(6):977-982.
5. Hairi NN, Cumming RG, Naganathan V, et al. Loss of Muscle Strength, Mass (Sarcopenia), and Quality (Specific Force) and Its Relationship with Functional Limitation and Physical Disability: The Concord Health and Ageing in Men Project. *J. Am. Geriatr. Soc.* 2010;58(11):2055-2062.
6. Foldvari M, Clark M, Laviolette LC, et al. Association of Muscle Power With Functional Status in Community-Dwelling Elderly Women. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. April 1, 2000 2000;55(4):M192-M199.
7. Bean J, Kiely DK, Herman S, et al. The Relationship Between Leg Power and Physical Performance in Mobility-Limited Older People. *J. Am. Geriatr. Soc.* 2002;50(3):461-467.
8. Pahor M. Effects of a physical activity intervention on measures of physical performance: results of the Lifestyle Interventions and Independence for Elders Pilot (LIFE-P) study. *The Journals of Gerontology: Series A: Biological Sciences and Medical Sciences*. 2006.
9. Vasunilashorn S, Coppin AK, Patel KV, et al. Use of the Short Physical Performance Battery Score to predict loss of ability to walk 400 meters: analysis from the InCHIANTI study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2009;64(2):223-229.
10. Volpato S, Cavalieri M, Sioulis F, et al. Predictive value of the Short Physical Performance Battery following hospitalization in older patients. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2011;66(1):89-96.

11. Hicks GE, Shardell M, Alley DE, et al. Absolute strength and loss of strength as predictors of mobility decline in older adults: the InCHIANTI study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2011;glr055.
12. Choquette S, Bouchard D, Doyon C, Sénéchal M, Brochu M, Dionne IJ. Relative strength as a determinant of mobility in elders 67–84 years of age. A nuage study: Nutrition as a determinant of successful aging. *J. Nutr. Health Aging*. 2010;14(3):190-195.
13. Bouchard DR, Beliaeff S, Dionne IJ, Brochu M. Fat mass but not fat-free mass is related to physical capacity in well-functioning older individuals: nutrition as a determinant of successful aging (NuAge)—the Quebec Longitudinal Study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2007;62(12):1382-1388.
14. Barbat-Artigas S, Rolland Y, Vellas B, Aubertin-Leheudre M. Muscle quantity is not synonymous with muscle quality. *J. Am. Med. Dir. Assoc*. 2013;14(11):852. e851-852. e857.
15. Visser M, Kritchevsky SB, Goodpaster BH, et al. Leg muscle mass and composition in relation to lower extremity performance in men and women aged 70 to 79: the health, aging and body composition study. *J. Am. Geriatr. Soc*. 2002;50(5):897-904.
16. Lindle RS, Metter EJ, Lynch NA, et al. Age and gender comparisons of muscle strength in 654 women and men aged 20–93 yr. *J. Appl. Physiol*. November 1, 1997 1997;83(5):1581-1587.
17. Lynch N, Metter E, Lindle R, et al. Muscle quality. I. Age-associated differences between arm and leg muscle groups. *J. Appl. Physiol*. 1999;86(1):188-194.
18. Francis P, Toomey C, Mc Cormack W, Lyons M, Jakeman P. Measurement of maximal isometric torque and muscle quality of the knee extensors and flexors in healthy 50- to 70-year-old women. *Clin. Physiol. Funct. Imaging*. Jan 07 2016.
19. Francis P, Mc Cormack W, Lyons M, Jakeman P. Age-group Differences in the Performance of Selected Tests of Physical Function and Association with Lower Extremity Strength. *J. Geriatr. Phys. Ther*. 2017 (accepted).
20. Glenn J, Vincenzo J, Canella C, Binns A, Gray M. Habitual and Maximal Dual-task Gait Speeds Among Sedentary, Recreationally Active, and Masters Athlete Late-middle Aged Adults. *Journal of aging and physical activity*. 2014.
21. Samuel D, Wilson K, Martin HJ, Allen R, Sayer AA, Stokes M. Age-associated changes in hand grip and quadriceps muscle strength ratios in healthy adults. *Aging Clin. Exp. Res*. 2012;24(3):245-250.
22. Ferreira L, Gobbi S, Gobbi LTB. An explanatory mechanism for the different decline in limb strength in older women. *Arch. Gerontol. Geriatr*. 2009;49(3):373-377.
23. Theou O, Jones GR, Vandervoort AA, Jakobi JM. Daily muscle activity and quiescence in non-frail, pre-frail, and frail older women. *Exp. Gerontol*. 2010;45(12):909-917.
24. Amaral JF, Alvim FC, Castro EA, Doimo LA, Silva MV, Novo Júnior JM. Influence of aging on isometric muscle strength, fat-free mass and electromyographic signal power of the upper and lower limbs in women. *Brazilian journal of physical therapy*. 2014;18(2):183-190.
25. Vilaca KH, Alves N, Carneiro JA, Ferriolli E, Lima NK, Moriguti JC. Body composition, muscle strength and quality of active elderly women according to the distance covered in the 6-minute walk test. *Brazilian journal of physical therapy*. 2013;17(3):289-296.
26. Visser M, Goodpaster BH, Kritchevsky SB, et al. Muscle mass, muscle strength, and muscle fat infiltration as predictors of incident mobility limitations in well-functioning older persons. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*. 2005;60(3):324-333.
27. Hayashida I, Tanimoto Y, Takahashi Y, Kusabiraki T, Tamaki J. Correlation between muscle strength and muscle mass, and their association with walking speed, in community-dwelling elderly Japanese individuals. *PLoS One*. 2014;9(11):e111810.
28. Norton C, Toomey C, McCormack WG, et al. Protein Supplementation at Breakfast and Lunch for 24 Weeks beyond Habitual Intakes Increases Whole-Body Lean Tissue Mass in Healthy Older Adults. *The Journal of Nutrition*. 2016;146(1):65-69.

29. Francis P. Twelve weeks' progressive resistance training combined with protein supplementation beyond habitual intakes increases upper leg lean tissue mass, muscle strength and extended gait speed in healthy older women. *Biogerontology*. 2016.
30. Greig CA, Young A, Skelton DA, Pippet E, Butler FMM, Mahmud SM. Exercise Studies with Elderly Volunteers. *Age Ageing*. May 1, 1994 1994;23(3):185-189.
31. Drey M, Sieber CC, Degens H, et al. Relation between muscle mass, motor units and type of training in master athletes. *Clin. Physiol. Funct. Imaging*. 2016;36(1):70-76.
32. Leahy S, O'Neill C, Sohun R, Toomey C, Jakeman P. Generalised equations for the prediction of percentage body fat by anthropometry in adult men and women aged 18–81 years. *Br. J. Nutr.* 2013;109(04):678-685.
33. Thorstensson A, Grimby G, Karlsson J. Force-velocity relations and fiber composition in human knee extensor muscles. *J. Appl. Physiol.* January 1, 1976 1976;40(1):12-16.
34. Maffiuletti NA, Bizzini M, Desbrosses K, Babault N, Munzinger U. Reliability of knee extension and flexion measurements using the Con-Trex isokinetic dynamometer. *Clin. Physiol. Funct. Imaging*. 2007;27(6):346-353.
35. Perrin DH, Costill DL. *Isokinetic exercise and assessment*. Vol 2: Human Kinetics Publishers Champaign (IL); 1993.
36. Bazzucchi I, Felici F, Macaluso A, De Vito G. Differences between young and older women in maximal force, force fluctuations, and surface EMG during isometric knee extension and elbow flexion. *Muscle Nerve*. 2004;30(5):626-635.
37. Guralnik JM, Simonsick EM, Ferrucci L, et al. A Short Physical Performance Battery Assessing Lower Extremity Function: Association With Self-Reported Disability and Prediction of Mortality and Nursing Home Admission. *J. Gerontol.* March 1, 1994 1994;49(2):M85-M94.
38. Jones CJ, Rikli RE, Beam WC. A 30-s chair-stand test as a measure of lower body strength in community-residing older adults. *Res. Q. Exerc. Sport*. 1999;70(2):113-119.
39. Swinscow T, Campbell M. Correlation and regression. *Statistics at square one*. 1997;9.
40. Buchner DM, Larson EB, Wagner EH, Koepsell TD, De Lateur BJ. Evidence for a non-linear relationship between leg strength and gait speed. *Age Ageing*. 1996;25(5):386-391.
41. Beattie K, Carson BP, Lyons M, Rossiter A, Kenny IC. The Effect of Strength Training on Performance Indicators in Distance Runners. *Journal of strength and conditioning research/National Strength & Conditioning Association*. 2016.
42. Maden-Wilkinson T, Degens H, Jones D, McPhee J. Comparison of MRI and DXA to measure muscle size and age-related atrophy in thigh muscles. *J. Musculoskelet. Neuronal Interact*. 2013;13(3):282-290.
43. Packard CJ, Bezlyak V, McLean JS, et al. Early life socioeconomic adversity is associated in adult life with chronic inflammation, carotid atherosclerosis, poorer lung function and decreased cognitive performance: a cross-sectional, population-based study. *BMC Public Health*. 2011;11(1):1.
44. Tieland M, Borgonjen-Van den Berg KJ, van Loon LJ, de Groot LC. Dietary protein intake in community-dwelling, frail, and institutionalized elderly people: scope for improvement. *Eur. J. Nutr.* 2012;51(2):173-179.
45. de Almeida Busch T, Duarte YA, Nunes DP, et al. Factors associated with lower gait speed among the elderly living in a developing country: a cross-sectional population-based study. *BMC Geriatr*. 2015;15(1):1.
46. Dulac M, Boutros GEH, Pion C, Barbat-Artigas S, Gouspillou G, Aubertin-Leheudre M. Is handgrip strength normalized to body weight a useful tool to identify dynapenia and functional incapacity in post-menopausal women? *Brazilian Journal of Physical Therapy*. 2016(AHEAD):0-0.

Table 1: Physical characteristics, body composition, muscle strength and functional performance

In healthy 50 – 70y women.

	Mean \pm SD or Median (IQR)	95% CI
Age (years)	60.4 \pm 5.1	59.5 to 61.3
Height (cm)	162.7 \pm 5.5	161.8 to 163.7
Body mass (kg)	66.2 (13.5)	64.8 to 68.1
Lean tissue mass (kg)	40.0 \pm 3.9	39.3 to 40.6
Body fat (%)	37.5 \pm 6.8	36.4 to 38.7
Knee Extensor Torque (N·m/kg)	1.3 \pm 0.4	1.2 to 1.4
Muscle Quality (N·m/kg)	23.7 \pm 5.9	22.6 to 24.7
900m Gait Speed (m/s)	2.3 \pm 0.5	2.2 to 2.4
5 repetition chair rise time (s)	8.4 (2.5)	8.0 to 8.7
Chair rises in 30 seconds (n)	17.2 \pm 4.1	16.2 to 18.2

Values are displayed as mean \pm SD, median (IQR) and 95% or Bootstrap 95% confidence interval (CI).

Table 2: Functional performance classified by tertile of knee extensor torque and muscle quality.

	T1 (n=42)	T2 (n=43)	T3 (n=43)	P
Knee Extensor Torque (N·m /kg)				
	Mean ± SD or Median (IQR)	Mean ± SD or Median (IQR)	Mean ± SD or Median (IQR)	
Gait Speed (m/s) ¹	2.6 ± 0.5	2.3 ± 0.4	2.1 ± 0.4	<0.001
95% CI	2.5 to 2.8	2.2 to 2.4	1.9 to 2.2	
5 Chair Repetitions (s) ²	8.1 ± 1.8	8.3 (2.1)	9.4 ± 2.3	0.006
95% CI	7.5 to 8.6	7.6 to 8.7	8.8 to 10.2	
30 s Chair Repetitions (n) ^{2*}	19.0 ± 4.6 17.0 to 21.0	16.8 ± 3.3 15.4 to 18.2	15.6 ± 3.6 13.8 to 17.4	0.020
Muscle Quality (N·m/kg)				
	Mean ± SD or Median (IQR)	Mean ± SD or Median (IQR)	Mean ± SD or Median (IQR)	
Gait Speed (m/s) ³	2.4 (0.7)	2.4 ± 0.4	2.1 ± 0.2	0.001
95% CI	2.3 to 2.7	2.2 to 2.5	2.0 to 2.3	
5 Chair Repetitions (s) ³	8.0 (2.4)	8.2 ± 1.7	9.6 ± 2.2	0.003
95% CI	7.5 to 8.5	7.7 to 8.7	8.9 to 10.2	
30 s Chair Repetitions (n) ^{2*}	18.3 ± 4.8 16.4 to 20.3	17.8 ± 3.6 16.2 to 19.3	15.2 ± 3.2 13.7 to 16.7	0.027
95% CI				

Values are displayed as mean ± SD or median (IQR) and 95% or Bootstrap 95% CI. P = significance value for one-way ANOVA or Kruskal Wallis test. * = T1 (n=25); T2 (n=23); T3 = (n=20). Post hoc analysis: ¹= differences between T1, T2 and T3 P=<0.05; ²=difference between T1 and T3 P=<0.05; ³=T1 and T2 different to T3.

Table 3: The association between knee extensor torque or muscle quality and functional performance for healthy 50 – 70y women (n=128) and per tertile.

	n=128	T1 (=42)	T2 (n=43)	T3 (n=43)
Knee Extensor Torque (N·m /kg)				
Gait Speed (m/s)	0.60 (<0.01)	0.54 (<0.01)*	0.13 (0.40)	0.46 (<0.01)*
5 Chair Repetitions (s)	-0.35 (<0.01)	0.37 (0.02)*	-0.08 (0.60)	-0.24 (0.12)
30 s Chair Repetitions (n)	0.47 (<0.01) ¹	0.41 (0.05)*	0.20 (0.35)	0.47 (0.04)*
Muscle Quality (N·m/kg)				
Gait Speed (m/s)	0.40 (<0.01)	0.27 (0.08)	0.12 (0.45)	0.24 (0.11)
5 Chair Repetitions (s)	-0.33 (<0.01)	-0.39 (0.01)*	-0.11 (0.48)	-0.21 (0.19)
30 s Chair Repetitions (n)	0.38 (0.01) ¹	0.31 (0.13)	0.04 (0.84)	0.40 (0.08)

Values are displayed as Pearsons or Spearmans r (p-value). *=statistical significance. ¹n=68.




Figure 1: The association between knee extensor torque and 900m gait speed in healthy older women.