

Kada, Sundaran, Satinovic, Milka, Booth, Lisa and Miller, Paul K. (2017) Managing discomfort and involving participation in non-emergency MRI: children's coping strategies during a first procedure. In: UK Radiological Congress, 12-14 June 2017, Manchester Central Convention Centre, Manchester, UK. (Unpublished)

Downloaded from: http://insight.cumbria.ac.uk/2930/

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available <u>here</u>) for educational and not-for-profit activities

provided that

• the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form

- a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found <u>here</u>.

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.

Managing Discomfort and Involving Participation in non-Emergency MRI: The Coping Strategies of Children During a First Procedure Sundaran Kada, Milka Satinovic, Lisa Booth* & Paul K. Miller

Background

For some adults, being inside the MRI scanner's bore can be associated with 'lying as for cremation'^[1,2] and also engender a feeling of being 'buried alive'^[2]. The greatest levels of anxiety are reported to emerge among patients undergoing first time and non-trauma examinations, where no prior referent can be drawn upon and extensive time to deliberate is freely available.^[3] Extant research further demonstrates that uncomfortable or stressful experiences of a medical imaging procedure during childhood can result in negative stances towards pertinent examinations in later life.^[4] It is important to consider the unique concerns that may emerge during MRI work conducted upon pediatric patients. Children are not just 'small adults'; examinations of children require not only understanding of normal variations in anatomy and pediatric pathologies, but also a particular awareness of their cognitions, comprehension and communicative capacities.^[5] It is therefore important that clinicians (a) develop an ability to understand the child's experience during imaging in order to (b) provide high quality care and develop meaningful interactions that might improve the context of the examination for all involved.

With full ethical approval, the study was carried out at a single University Hospital in Western Norway. Grounded Theory (henceforth GT) was adopted as the core investigative approach.^[6] An opportunity sample of N=22 children was recruited, age range = 8 to 16 years, and undergoing a non-emergency MRI brain examination for the first time. The gender split was an even n=11. Semi-structured interviews (mean = 15 minutes) were conducted the children and their parents immediately subsequent to the completion of the procedure itself. Observational field notes were also compiled regarding the activity of children and parents before, during and after the examination. All interviews were transcribed verbatim and translated into English at the point of transcription. All data were incorporatively analyzed using the methods of GT in order to identify key concepts arising from participants' own MRI-

The central concern of the participating children ultimately related to handling the prospective and extant 'discomfort' of their first MRI examination in psychological, sensory and more conventionally physical terms. This was variously expressed in terms of uncertainty/anxiety about what would happen, the handling of restricted space, noise and strong light, and also the demands associated with lying perfectly still and, in some cases, managing nausea. These forms of discomfort were negotiated by the children in an overall process herein termed Involving Participation, which had three chronological phases.

Participation Preparation

- Preparative activities prior to the child's arrival in the Department itself,
- Usually collaboratively achieved with parents.
- Discussion of hospital-produced materials.
- Reference to familiar (comforting) narratives.

Enabling Participation

- Activities undertaken upon arrival in the Department.
- Collaboratively achieved with radiographers and usually parents.
- Humanising staff, encouraging familiarity with light/noise.
- Offering active choices around distraction methods (empowerment).

Sustaining Participation

- Activities embraced during the MRI examination itself.
- Collaboratively achieved with radiographers and sometimes parents.
- Distraction (films, music).
- Adult feedback on opportunities to move between scan sequences, breathing techniques, and handling nausea.

ussion/Conclusions

When the Involving Participation process was successful, the child generally reported a sense of growth and even joy at having 'mastered' the MRI context. The importance of patients actively participating in their own healthcare settings is a common theme in contemporary literature relating to both adults and children.^[8] While the human right to be a participant is often seen a fundamental one for children, they are recurrently not informed of this right, nor significantly involved in discussions about their care. ^[9,10] Given the large, and growing, body of evidence relating not least to the anxiety-allaying impacts of pediatric participation emerging within general medical science, it is perhaps surprising that more attention has not been accorded to it within radiological research fields, and particularly those pertaining to MRI. ^[11] After all, it is well-documented that the noise of the MRI scanner, narrow space and lying still can be acute sources of anxiety, and anxious children have characteristically problematic consequences for the success of MRI examinations. ^[12,13] This study has aimed to illustrate how a child's practical investment in becoming a full and cooperative participant in an MRI procedure is not simply an output of discrete events, nor something that can be managed through one-size-fits-all reasoning, but is rather an ongoing interactive process over three interlinked social contexts, involving the active cooperation of a range of significant others.



- References
- 1. Törnqvist E, Månsson Å, Larsson EM, Hallström I. It's like being in another world–patients' lived experience of magnetic resonance imaging. J Clin Nurs 2006;15:954-961.
- 2. Laidlaw A, Henwood S. Patients with multiple sclerosis: Their experiences and perceptions of the MRI investigation. J Diag Radio Imag 2003;5:19-25.
 3. Heyer CM, Thüring J, Lemburg SP, Kreddig N, Hasenbring M, Dohna M, Nicolas V. Anxiety of patients undergoing CT imaging an underestimated problem?
 Pediatr.Nurs. 2007;33:505-511 7p.
 Acad Radiol 2015;22:105-112.
 D. Rosenbaum P, King S, Law M,
- 4. Bowman S. The radiographer/patient relationship-a short term but vital interaction. Radiogr Today 1993;59:17-18.
- 5. Young SJ, Barnett PL, Oakley EA. Bruising, abrasions and lacerations: Minor injuries in children. Med J Aust 2005;182:588-592.
- 6. Glaser BG. The future of grounded theory. Qual.Health Res. 1999;9:836.
- 7. Glaser BG, Holton J. Remodeling grounded theory. Forum: Qualitative Social Research 2004;5:1-17.

- 8. Eyssen IC, Steultjens MP, Dekker J, Terwee CB. A systematic review of instruments assessing participation: Challenges in defining participation. Arch.Phys.Med.Rehabil. 2011;92:983-997 15p.
- 9. Runeson I, Mårtenson E, Enskär K. Children's knowledge and degree of participation in decision making when undergoing a clinical diagnostic procedure. Pediatr.Nurs. 2007;33:505-511 7p.
- 10. Rosenbaum P, King S, Law M, King G, Evans J. Family-centred service: A conceptual framework and research review. Phys Occup Ther Pediatr 1998;18:1-20.
- Salmela M, Aronen ET, Salanterä S. The experience of hospital-related fears of 4- to 6-year-old children. Child Care Health Dev. 2011;37:719-726 8p.
 Hallowell LM, Stewart SE, de Amorim E Silva CT, Ditchfield MR. Reviewing the process of preparing children for MRI. Pediatr.Radiol. 2008;38:271-279 9p.
 Edwards AD, Arthurs OJ. Paediatric MRI under sedation: Is it necessary? what is the evidence for the alternatives? Pediatr.Radiol. 2011;41:1353-1364 12p.

UK RADIOLOGICAL AND RADIATION ONCOLOGY CONGRESS



www.cumbria.ac.uk *lisa.booth@cumbria.ac.uk



