

## Article (refereed) - postprint

---

Kuhnert, Matthias; Yeluripati, Jagadeesh; Smith, Pete; Hoffmann, Holger; Van Oijen, Marcel; Constantin, Julie; Coucheney, Elsa; Dechow, Rene; Eckersten, Henrik; Gaiser, Thomas; Grosz, Balász; Haas, Edwin; Kersebaum, Kurt-Christian; Kiese, Ralf; Klatt, Steffen; Lewan, Elisabet; Nendel, Claas; Raynal, Helene; Sosa, Carmen; Specka, Xenia; Teixeira, Edmar; Wang, Enli; Weihermüller, Lutz; Zhao, Gang; Zhao, Zhigan; Ogle, Stephen; Ewert, Frank. 2017. **Impact analysis of climate data aggregation at different spatial scales on simulated net primary productivity for croplands.**

© 2016 Elsevier B.V.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>



This version available <http://nora.nerc.ac.uk/516936/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

NOTICE: this is the author's version of a work that was accepted for publication in *European Journal of Agronomy*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *European Journal of Agronomy* (2017), 88. 41-52. [10.1016/j.eja.2016.06.005](https://doi.org/10.1016/j.eja.2016.06.005)

[www.elsevier.com/](http://www.elsevier.com/)

Contact CEH NORA team at  
[noraceh@ceh.ac.uk](mailto:noraceh@ceh.ac.uk)

# Impact analysis of climate data aggregation at different spatial scales on simulated Net Primary Productivity for croplands

*Kuhnert, Matthias<sup>1</sup>; Yeluripati, Jagadeesh<sup>1,13</sup>; Smith, Pete<sup>1</sup>; Hoffmann, Holger<sup>2</sup>; van Oijen, Marcel<sup>3</sup>; Constantin, Julie<sup>4</sup>; Dechow, Rene<sup>5</sup>; Eckersten, Henrik<sup>6</sup>; Gaiser, Thomas<sup>2</sup>; Grosz, Balász<sup>5</sup>; Haas, Edwin<sup>7</sup>; Kersebaum, Kurt-Christian<sup>8</sup>; Kiese, Ralf<sup>7</sup>; Klatt, Steffen<sup>7</sup>; Lewan, Elisabet<sup>9</sup>; Nendel, Claas<sup>8</sup>; Raynal, Helene<sup>4</sup>; Sosa, Carmen<sup>9</sup>; Specka, Xenia<sup>8</sup>; Teixeira, Edmar<sup>10</sup>; Wang, Enli<sup>11</sup>; Weihermüller, Lutz<sup>12</sup>; Zhao, Gang<sup>2</sup>; Zhao, Zhigan<sup>11</sup>; Ogle, Stephen<sup>14</sup>; Ewert, Frank<sup>2</sup>*

<sup>1</sup> Institute of Biological and Environmental Sciences, University of Aberdeen, 23 St Machar Drive, Aberdeen AB24 3UU, UK

<sup>2</sup> Crop Science Group, INRES, University of Bonn, Katzenburgweg 5, 53115 Bonn, DE

<sup>3</sup> Centre for Ecology and Hydrology, CEH-Edinburgh, Bush Estate, Penicuik EH26 0QB, UK

<sup>4</sup> INRA, UMR 1248 AGIR & UR0875 MIA-T, F-31326 Auzeville, France

<sup>5</sup> Thünen-Institute of Climate-Smart-Agriculture, Bundesallee 50, 38116 Braunschweig, DE

<sup>6</sup> Department of Crop Production Ecology, Swedish University of Agricultural Sciences, Ulls väg 16, 750 07 Uppsala, SE

<sup>7</sup> Institute of Meteorology and Climate Research – Atmospheric Environmental Research, Karlsruhe Institute of Technology, Kreuzeckbahnstraße 19, 82467 Garmisch-Partenkirchen, DE

<sup>8</sup> Institute of Landscape Systems Analysis, Leibniz Centre for Agricultural Landscape Research, 15374 Müncheberg, DE

<sup>9</sup> Biogeophysics and water quality, Department of Soil and Environment, Swedish University of Agricultural Sciences, Lennart Hjelm's väg 9, 750 07 Uppsala, SE

<sup>10</sup> Systems Modelling Team (Sustainable Production Group), The New Zealand Institute for Plant and Food Research Limited, Canterbury Agriculture & Science Centre, Gerald St, Lincoln 7608, NZ

<sup>11</sup> CSIRO Land and Water, Clunies Ross Street, Canberra, ACT, AU

<sup>12</sup> Institute of Bio- & Geosciences Agrosphere (IBG-3), Forschungszentrum Jülich, 52425 Jülich, DE

<sup>13</sup> The James Hutton Institute, Craigiebuckler, Aberdeen, AB15 8 QH, UK

<sup>14</sup> Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO 80523, USA

## Abstract

For spatial crop and agro-systems modelling, there is often a discrepancy between the scale of measured driving data and the target resolution. Spatial data aggregation is often necessary, which affects the uncertainty of the simulation results. Previous studies have shown that climate data aggregation has little effect on simulation of phenological stages, but effects on net primary production (NPP) might still be expected through changing the

40 length of the growing season and the period of grain filling. This study investigates the impact  
41 of spatial climate data aggregation on NPP simulation results, applying eleven different  
42 models for the same study region (~34000 km<sup>2</sup>), situated in Western Germany. To isolate  
43 effects of climate, soil data and management were assumed to be constant over the entire  
44 study area and over the entire study period of 29 years. Two crops, winter wheat and silage  
45 maize, were tested as monocultures. The results show only small impacts of climate data  
46 aggregation on averages over the entire simulation period and study region. Maximum  
47 differences between the five scales in the range of 1 to 100 km grid cells show changes of  
48 0.4 – 7.8 % and 0.0 - 4.8 % for wheat and maize, respectively, whereas the simulated  
49 potential NPP averages of the models show a wide range (1.9 - 4.2 g C m<sup>-2</sup> d<sup>-1</sup> and 2.7 - 6.1  
50 g C m<sup>-2</sup> d<sup>-1</sup> for wheat and maize, respectively). The impact of the spatial aggregation was  
51 also tested for shorter periods to test if impacts over shorter periods level out over longer  
52 periods, which shows larger impacts for single years (up to 9.4 % for wheat and up to 13.6  
53 % for maize). An analysis of extreme weather conditions shows an aggregation effect to the  
54 vulnerability up to 12.8 % and 15.5 % between the different resolutions for wheat and maize,  
55 respectively. Simulations of NPP averages over larger areas (e.g. regional scale) and longer  
56 time periods (several years) are relatively insensitive to climate data aggregation, but the  
57 scale of climate data is more relevant for impacts on annual averages of NPP or if the period  
58 is strongly affected or dominated by drought stress. There should be an awareness of the  
59 higher uncertainty for the NPP values if data are not available in a high resolution. On the  
60 other side, the results suggest that there is no need to simulate in high resolution for long  
61 term regional NPP averages based on the simplified assumptions (soil and management  
62 constant in time and space) used in this study.

63

64 Keywords: net primary production, NPP, scaling, extreme events, crop modelling, climate,  
65 data aggregation

66

## 67 **1 Introduction**

68

69 Net primary production (NPP) is a crucial ecosystem variable characterising the condition of  
70 an ecosystem (Pan et al, 2014) and its sensitivity to climate change. Spatial NPP is difficult  
71 to measure and often biased and uncertain (Pan et al., 2014), because measurements show  
72 several limitations (indirect determination, spatially and temporally limited). Spatial  
73 modelling is an important tool for interpolation and extrapolation of measurements or for  
74 providing spatial distributed projections for regional (Reich et al., 1999; Zaehle et al., 2006;  
75 Bandaru et al., 2013; Liu et al, 2015), continental (Ciais et al., 2010) or global scale  
76 (Hemming et al., 2013; Friend et al., 2014). The regional scale is relevant for policy makers  
77 to analyse adaptation and mitigation strategies, but NPP data for this scale are often derived  
78 by extrapolating measured information from the site scale to a region by applying models  
79 developed at site scale (Zhang et al., 2015). This model-based up-scaling requires a balance  
80 between accuracy and simulation time.

81 Spatial modelling of NPP relies on spatially distributed input and driving data like weather  
82 data and information on soil, land use and management characteristics. Depending on  
83 environmental parameters, ecosystem characteristics and the chosen resolution, the  
84 impacts of extrapolation or interpolation may be great or small since there is e.g. a higher  
85 uncertainty for high relief areas compared to relatively flat areas as shown by Pierce and  
86 Running (1995). For this reason, estimates of error and uncertainty arising from data  
87 aggregation across scales needs to be quantified.

88 Several studies have highlighted the impact of data aggregation on simulation results (Cale  
89 et al., 1983; Rastetter et al., 1992; Ewert et al., 2015; Zhao et al., 2015). De Wit et al. (2005)  
90 and Hoffmann et al. (2015) investigated the impact of climate data aggregation on crop  
91 yields. While de Wit et al. (2005) varied precipitation and solar radiation only on the

92 resolutions 10 km and 50 km, Hoffmann et al. (2015) differentiated between five different  
93 resolutions between 1 and 100 km and also considered aggregation effects of temperature  
94 for 13 models. Both studies found only slight impacts of data aggregation on simulated yield  
95 over longer time periods at a regional scale. Van Bussel et al. (2011) investigated the  
96 impacts of climate aggregation on croplands and focused on phenological stages rather than  
97 primary production, but they also found minor effects on simulated average values. The  
98 impacts of climate data aggregation on NPP were tested by Nungesser et al. (1999) and  
99 Pierce and Running (1995), both for American forests. In both studies, the impact was minor  
100 for averages over the entire study area, but showed relevant impacts for smaller areas,  
101 especially areas dominated by strong relief changes (Pierce and Running, 1995). In both  
102 studies, the effects were tested by one model and for two resolutions of 10 and 50 km grid  
103 cells in Nungesser et al. (1999), and 1 km and 110 km in Pierce and Running (1995). The  
104 latter study investigated the effect for different input variables (relief, climate and soil) and  
105 found that climate data aggregation was the dominant variable affecting scale differences of  
106 NPP. They also observed larger scale effects for shorter time periods, which could be an  
107 indication of extreme weather events that average out over larger areas or longer time  
108 periods. Overall, regional simulation results over longer periods seem to be little affected by  
109 climate data aggregation. Over longer periods changes of NPP level out and the impact of  
110 extreme events may be not obvious in a long term average, but relevant for shorter periods.  
111 Reichstein et al. (2013) describe the temporal and spatial scale as very important to detect  
112 impacts of extreme weather conditions on the carbon balance and see a risk of miss out  
113 extreme weather conditions by integration of weather data across scales. Impacts of  
114 extreme weather are also depending on the temporal scale, which is not yet tested for  
115 impacts on NPP. However, there is a lack of studies determining the effect on NPP  
116 simulations of croplands, and no study to date has analysed the relevance of extreme events  
117 during climate data aggregation.

118 Therefore, the objective of this paper is to quantify error and uncertainty of NPP simulations  
119 of croplands caused by climate data aggregation across five resolutions (1, 10, 25, 50 and  
120 100 km grid cell side length). This study addresses the three questions i) what are the  
121 impacts on long term NPP averages over the entire region? ii) how does the aggregation  
122 effect change over shorter time periods? iii) is the aggregation effect more pronounced in  
123 years with extreme weather conditions compared to “normal” years? These questions are  
124 answered by using a simulation approach involving eleven different models. Additionally, a  
125 vulnerability analysis helps to identify the impact of climate data aggregation for years with  
126 extreme weather conditions. Thus, we provide the first systematic analysis considering the  
127 impact of spatial weather data aggregation on NPP using five resolutions and 11 different  
128 models.

129

## 130 **2 Methods**

131

### 132 *2.1 Aggregation effect*

133 Spatial modelling approaches are containing uncertainty, because uncertainty of input data  
134 and limited data availability requires data aggregation, which also contribute to the  
135 uncertainty. In this study we focused on the impact of data aggregation on uncertainty.  
136 Spatial data base on point measurement, small scale measurements or approaches that  
137 averaging the data already during the measurement process. In the data aggregation these  
138 data sets get interpolated, extrapolated and averaged to provide data in its spatial  
139 distribution. This data aggregation increases the uncertainty of the data sets. Beside the  
140 impacts of data aggregation, the chosen format of the model approach adds uncertainty to  
141 the data, too. Spatial model approaches often using data organized in grid maps, while  
142 natural conditions do not follow any symmetric pattern. Therefore, gridded data already  
143 contain uncertainty, which also varies with changing scale. Our focus in this study is on the

144 impact of changing scales of grid map data on simulation results. Because there is a strong  
 145 interaction of different processes, we concentrate on the impact of changing weather data  
 146 as an important driver for plant growth. As we only compare simulated NPP values, we are  
 147 not using the term uncertainty, but aggregation effect ( $E_{aggregation}$ ), which can be formulated  
 148 as:

$$149 \quad E_{aggregation} = \frac{\max(VA_{Res_1}, \dots, VA_{Res_n}) - \min(VA_{Res_1}, \dots, VA_{Res_n})}{VA_{Res_1}} \quad (1)$$

150 In this study the aggregation effect is defined as the maximum difference between the  
 151 simulated NPP averages between the different resolutions and it is quantified by the  
 152 difference between maximum NPP average and minimum NPP average of the five  
 153 resolutions:

$$155 \quad E_{aggregation,model} = \frac{\max(NPP_{Res1,model}, \dots, NPP_{Res100,model}) - \min(NPP_{Res1,model}, \dots, NPP_{Res100,model})}{NPP_{Res1,model}} \quad (2)$$

156  
 157 This allows a model specific calculation of the effect and is independent of any trends  
 158 towards the coarser resolution. The difference describes the maximum expected bias by  
 159 picking one resolution in comparison to the results of another resolution. This calculation is  
 160 applicable on different spatial or temporal averages.

161 The aggregation effect can also be applied on ensemble runs, which is possible in two different  
 162 ways.  $E_{aggregation}$  can be calculated for the resolution specific averages over all models with  
 163 the formulation:

$$164 \quad E_{aggregation,average} = \frac{\max(\overline{NPP_{Res1}}, \dots, \overline{NPP_{Res100}}) - \min(\overline{NPP_{Res1}}, \dots, \overline{NPP_{Res100}})}{\overline{NPP_{Res1}}} \quad (3)$$

165 This allows the quantification of the aggregation effect for ensemble runs.

166

## 167 2.2 Study area

168 The study area is the state North Rhine-Westphalia situated in the West of Germany. The  
 169 state is 34084 km<sup>2</sup> in size with an elevation from 0 to 843 m above sea level, with lower

170 plains in the North West and higher elevations in the South-East. The land-use is dominated  
171 by agriculture (more than 60 % of the area), but in this study the entire area (including the  
172 40 % of forest, urban areas and infrastructure as well as water bodies) was considered to  
173 be cropland. To standardize the simulation runs monocultures of either winter wheat  
174 (*Triticum aestivum* L.) or silage maize (*Zea mays* L.) were assumed for the entire area.  
175 The driving daily weather data are provided at five different resolutions (1, 10, 25, 50 and  
176 100 km grid cells), while soil data (typical soil type) and management (good agricultural  
177 practice) were assumed to be constant during the study period and over the entire study  
178 area. The chosen soil type is a sandy loam, which is typical for this region and the  
179 management includes ploughing, sowing, harvest and three (130, 52, 26 kg ha<sup>-1</sup>) and two  
180 (30, 208 kg ha<sup>-1</sup>) fertilizer applications during spring for wheat and maize, respectively (for  
181 details see Hoffmann et al., 2015). The models were not calibrated for the study area, but  
182 were adjusted based on 30-year yield averages (1982-2011) of about 8 t ha<sup>-1</sup> for winter  
183 wheat and 14 t ha<sup>-1</sup> for silage maize. The weather data, presented and discussed by Zhao  
184 et al. (2015) and Hoffmann et al. (2015), show a 30 year average temperature of 9.7 °C, an  
185 average annual precipitation of 899 mm and mean annual global radiation of 3758 MJ m<sup>-2</sup>  
186 a<sup>-1</sup> (1982-2011) with the standard deviations of 1.2 °C, 214.0 mm a<sup>-1</sup> and 169.4 MJ m<sup>-2</sup> a<sup>-1</sup>,  
187 respectively. The coldest year was 2004 with an average temperature of 8.9 °C and the  
188 warmest year was 1983 (11.2 °C). The driest year was 2001 with 516.2 mm precipitation.  
189 All coarser resolutions of the weather data were based on the grid cells of the 1 km resolution  
190 for daily time steps. The data show a decrease of temperature (from 9.7 °C to 9.4 °C) and  
191 precipitation (from 899 mm to 824 mm) starting from the 1 km resolutions towards the  
192 coarsest resolution of 100 km.

193

194 *2.3 Modelling applications*



195 There are three different approaches using different model settings to analyse the impact of  
196 different processes contributing to the simulation of NPP. In a first approach, no limitations  
197 to growth factors, other than temperature and radiation, are simulated explicitly (switched  
198 off or compensated in all models). We denote this potential, non-limited growth, potential  
199 NPP or PN. The second approach considers only water-limitation (WN), while the third  
200 approach considers nitrogen and water limitation (NN). The way limitations are switched off  
201 differs between the models. Some models switched off the stress factors, other models  
202 compensated the stress by providing additional water and nutrient applications.  
203 The settings for management are presented by Hoffmann et al. (2015). The sowing date is  
204 fixed for all models, while for harvest only a latest date is suggested (if the phenological  
205 model does not determine maturity before this date, there will be an automatic harvest).

206

#### 207 *2.4 Models*

208 Eleven models participated in this study; eight crop models and three biogeochemical  
209 models (Table1). All models provide data on a daily time step (except CENTURY which uses  
210 a monthly time step), consider the complete range of management practices (except AgroC  
211 that does not consider nitrogen limitation) and provide simulations for the two considered  
212 crops, wheat and maize (except COUP that only simulates wheat). The growing season for  
213 the crop models is determined by internal phenological models based on a fixed sowing  
214 date, while the three biogeochemical models CENTURY, DailyDayCent and  
215 LandscapeDNDC and the crop model STICS also used a fixed harvest date (i.e. fixed length  
216 of the growing season).

217 Five of the models determine NPP based on the radiation use efficiency concept (AgroC,  
218 APSIM, APSIMmod., COUP, LINTUL, STICS), while other models determine NPP based on  
219 the difference between gross primary production and respiration (HERMES, MONICA),

220 calculated directly (DailyDayCent) or other approaches (LandscapeDNDC). More details  
221 about the models are provided in Hoffmann et al. (2015).

222

223 Table 1: List of the participating models.

No.	Model	References
1	HERMES	Kersebaum 2007, 2011
2	APSIM	Keating et al. 2003; Holzworth et al. 2014
3	COUP	Conrad & Fohrer 2009; Jansson & Karlberg 2004
4	DailyDayCent	Del Grosso et al. 2001, 2006; Parton et al. 2001; Yeluripati et al. 2009
5	LandscapeDNDC	Haas et al. 2012; Kraus et al., 2014
6	LINTUL	Van Ittersum et al. 2003; Shibu et al. 2010
7	MONICA	Nendel et al. 2011
8	STICS	Bergez et al. 2013; Brisson et al. 1998, 2008
9	APSIMmod	Chen et al. 2010; Keating et al. 2003; Wang et al. 2002
10	CENTURY	Parton et al., 1993, 1995
11	AgroC	Herbst et al., 2008

224

225

## 226 *2.5 Evaluation of aggregation effects over different time periods*

227 The simulation results (NPP averages over the entire study area) were averaged over  
228 different periods (1, 5, 10, 15, 20, 25, 29 years) to examine the maximum differences  
229 between the five resolutions as influenced by the different temporal scales. The number of  
230 averages considered varies for the different time periods (29, 6, 3, 2, 2, 2, 1, respectively).

231 The analysis for the periods 20 and 25 years were applied twice, covering mainly the first  
232 and the last years with some data overlap. The results are presented as the mean  
233 aggregation effect as well as the maximum aggregation effect between the five resolutions.

234

## 235 *2.6 Vulnerability Analysis*

236 Vulnerability and risk are terms that are widely used in different communities and described  
237 in different contexts with different definitions. In this study we use an approach developed  
238 by van Oijen et al. (2014), designed to investigate impacts of extreme weather events on  
239 carbon dynamics. The approach, based on an abiotic definition of extreme periods,

240 compares the impacts on a chosen biotic ecosystem variable on the defined “extreme” or  
241 “hazardous” and “not extreme” or “non-hazardous” conditions. Van Oijen et al. (2014) chose  
242 the standardized precipitation evapotranspiration index (SPEI), developed by Vicente-  
243 Serrano et al. (2010), as the abiotic factor separating hazardous from non-hazardous  
244 conditions. SPEI is a drought index based on the difference between potential  
245 evapotranspiration and precipitation. If the precipitation exceed the potential  
246 evapotranspiration for the given time period, SPEI shows positive values, while negative  
247 values represent a water deficit based on the calculated difference and indicate a drought  
248 impact. There is no fixed threshold, which defines an extreme drought impact or growth  
249 reducing conditions and SPEI can be calculated for any duration. The index is normalized  
250 and normal distributed. The average is about 0 for the considered period of 1982-2011 with  
251 64 % of the values between -1 and +1 and 19 % below -1 for the 1 km resolution. These  
252 statistics stay the same for all resolutions, with the exception of the number of values in the  
253 -1 to +1 interval, which drops down to 63 % for the 100 km resolution. The potential  
254 evapotranspiration can be calculated with different approaches, while in this study the  
255 method developed by Thornthwaite (1948) is used.

256 SPEI is one of the indices that considers both, precipitation and temperature in the  
257 calculation, rather than only precipitation, but is still easy to apply, which makes it an  
258 attractive index to use in this study.. Van Oijen et al. (2014) suggest two thresholds to  
259 separate hazardous from non-hazardous conditions:  $SPEI < -1$  and  $SPEI < -2$ . For the actual  
260 study region there is only a small number of SPEI values below -2, so  $SPEI < -1$  was chosen  
261 as the threshold. Following van Oijen et al. (2014), the period to calculate the SPEI is  
262 restricted to half a year. In contrast to the approach of van Oijen et al. (2014), who suggested  
263 the period April-September, the period February – July was used in this study to better reflect  
264 the crop growth period. The system variable used in this study is NPP.

265 Vulnerability ( $V$ ) describes a possible damage/impact on a system and the risk ( $R$ ) is  
266 described by the product of the probability ( $P$ ) that a hazardous event ( $H$ ) occurs and its  
267 impact on the system.

$$268 \\ 269 R = P(H) \cdot V \quad (2)$$

270  
271 This equation represents the relation between probability, and vulnerability and can be  
272 expressed by using the reduction of the NPP by the hazardous periods, described as risk:

$$273 \\ 274 R = E(NPP|non - hazardous) - E(NPP) \quad (3)$$

275  
276  $E(NPP|non-hazardous)$  is the average value of NPP for all grid cells and years with a SPEI  
277  $\geq -1$  and  $E(NPP)$  is the overall average of NPP (including both, hazardous and non-  
278 hazardous conditions). The vulnerability describes the difference of the NPP averages for  
279 the non-hazardous and the hazardous years and grid cells.

$$280 \\ 281 V = E(NPP|non - hazardous) - E(NPP|hazardous) \quad (4)$$

## 282 283 **3 Results**

### 284 285 *3.1 NPP differences between the models*


286 The different model simulations of NPP are compared for the 1 km grid resolution, which is  
287 considered to be the “best” result for each model separately and thus used as the baseline.

288 The results for wheat vary for the different models with mean values for 29 years of  
289 simulation 1.9 - 4.2 g C m<sup>-2</sup> d<sup>-1</sup>, 1.9 - 4.1 g C m<sup>-2</sup> d<sup>-1</sup> and 1.8 – 4.1 g C m<sup>-2</sup> d<sup>-1</sup> NPP for PN,  
290 WN and NN, respectively (Figure 1). The simulation results of STICS and AgorC show the  
291 highest variation over the 29 years, while the results of COUP and LandscapeDNDC vary  
292 only within a narrow range. Some of the models are sensitive to water limitation (e.g. COUP),

293 which is reflected by the differences between the PN and the WN approach (Figure 1), other  
 294 models show minimal impacts of water limitation (e.g. STICS). The spatial distribution of the  
 295 results shows a dependency on elevation (Figure 2). However, the spatial distribution of high  
 296 and low NPP values is different between the models. While most models show higher NPP  
 297 values for low elevation and lower values in the higher elevations, the other group of models  
 298 (AgroC, COUP, LINTUL, APSIM and APSIMmod) show the opposite spatial separation of high  
 299 and low NPP values.

300 The simulated NPP for maize (7.4 - 12.8 g C m<sup>-2</sup> d<sup>-1</sup>, 7.3 - 12.7 g C m<sup>-2</sup> d<sup>-1</sup> and 7.2 – 10.0 g  
 301 C m<sup>-2</sup> d<sup>-1</sup> for PN, WN and NN, respectively) show higher maximum values than the NPP for  
 302 wheat, and indicate an even lower sensitivity to water limitation, which is represented by a  
 303 comparison of the simulation results of PN and WN (Figure 3). The extreme NPP values for  
 304 APSIMmod and LandscapeDNDC for wheat and maize, respectively, are outside the range  
 305 of the other models, but are included in all analyses.

306



307


308

309 Figure 1: Simulated NPP for winter wheat at 1 km resolution for potential growth (PN), under water  
 310 limitation (WN) and under nutrient and water limitation (NN). The models are in the order HERMES  
 311 (HE), APSIMmod (A2), COUP (CO), DailyDayCent (DA), LandscapeDNDC (DN), LINTUL (LI),  
 312 MONICA (MO), STICS (ST), APSIM (AI), CENTURY (CE) and AgroC (AG).

313


314

315



316  
317  
318  
319  
320

Figure 2: Spatial distribution of the 29 year averages of NPP for the 11 models assuming wheat mono-culture for the PN approach.



321  
322  
323  
324  
325  
326  
327

Figure 3: Simulated NPP values for silage maize at 1 km resolution for potential growth (PN), under water limitation (WN) and under nutrient and water limitation (NN). The models are in the order HERMES (HE), APSIMmod (A2), DailyDayCent (DA), LandscapeDNDC (DN), LINTUL (LI), MONICA (MO), STICS (ST), APSIM (AI), CENTURY (CE) and AgroC (AG).


### 3.2 Model specific aggregation effect

The aggregation effect as described in equation 1 shows a range for all models of 0.4 – 7.8 % and 0 - 4.8 % for wheat and maize, respectively (Table 2). The analysis of the medians shows slightly larger aggregation effects with 0.3 – 11.4 % and 0.0 – 10.0 % for wheat and

332 maize, respectively (Table 2). There are no obvious trends in the changes of NPP from the  
333 1 km resolution to 100 km resolution, neither for the crops nor for the different models  
334 (Figures 4 and 5). However, the models LandscapeDNDC, MONICA, CENTURY and  
335 DailyDayCent show relatively small changes ( $< 1.2\%$ ) for both crops, and HERMES for the  
336 wheat simulations, while APSIM, APSIMmod and AgroC show relatively high aggregation  
337 effects (more than  $4.8\%$ ) between the different scales. The aggregation effect varies  
338 between the models as does the trend. While APSIMmod, LINTUL, AgroC and APSIM show  
339 increasing NPP values towards coarser resolutions for the wheat simulations, HERMES,  
340 MONICA, CENTURY and DailyDayCent show decreasing NPP. The simulation results of  
341 COUP and STICS show no trend, but a minimum NPP averages for the resolutions of 10  
342 km and 50 km, respectively. The median is affected for some models, especially for the  
343 maize simulations, more than the average values and most models show stronger changes  
344 for WN and NN than for PN (Table 2). The results for maize support the findings of the wheat  
345 simulations, but the scale effect is smaller and effect and trends differ for some models  
346 between the two crops. HERMES and DailyDayCent show minimal differences between the  
347 resolutions, while APSIMmod, LINTUL, MONICA, STICS and APSIM show a decreasing  
348 trend with AgroC and LandscapeDNDC showing an increasing trend towards coarser  
349 resolutions.


350 The aggregation effect for the model ensemble is calculated for the resolution specific  
351 average over all models (equation 3). The effect is below  $0.9\%$  for wheat and  $2.0\%$  for  
352 maize, which is below all aggregation effects of the individual models.

353



354  
 355  
 356  
 357  
 358  
 359

Figure 4: Simulated NPP of wheat for the potential growth (PN). These boxplots represent the variability over 29 NPP averages over the growing season for the five resolutions (1, 10, 25, 50 and 100 km).



360



361 Figure 5: Simulated NPP of maize for the potential growth (PN). These boxplots represent the  
 362 variability over 29 NPP averages over the growing season for the five resolutions (1, 10, 25, 50  
 363 and 100 km).

364

365 Table 2: Relative maximum differences of NPP averages (AVG) and median (MED) between the  
 366 five resolutions [%]. The values represent the simulation results of wheat (W) and maize (M) for the  
 367 three approaches (PN, WN, NN). All differences are related to the resolution with the lowest NPP  
 368 average. The models are HERMES (HE), APSIMmod (A2), COUP (CO), DailyDayCent (DA),  
 369 LandscapeDNDC (DN), LINTUL (LI), MONICA (MO), STICS (ST), APSIM (AI), CENTURY (CE)  
 370 and AgroC (AG). Changes of greater 3 % are highlighted by grey boxes.

371


	HE	A2	CO	DA	DN	LI	MO	ST	AI	CE	AG
AVG W PN	0.9	5.0	2.2	1.1	n.s.	4.7	0.8	2.0	4.9	0.9	7.8
AVG W WN	0.8	4.8	2.2	0.7	0.2	2.4	1.1	1.8	4.8	0.8	7.8
AVG W NN	0.5	5.4	2.2	0.5	1.6	3.0	1.2	2.0	5.4	0.7	n.s.
AVG M PN	1.3	2.0	n.s.	0.1	1.3	2.0	2.0	1.5	2.0	0.0	3.2
AVG M WN	1.3	2.0	n.s.	0.3	1.2	2.5	1.5	2.0	2.0	0.5	3.5
AVG M NN	4.8	2.0	n.s.	1.2	2.8	1.7	1.5	2.0	2.0	0.6	n.s.
MED W PN	0.9	4.9	2.2	1.0	n.s.	4.7	0.8	2.0	4.9	0.9	7.8
MED W WN	1.5	7.8	2.3	1.8	0.3	3.2	2.9	2.1	7.8	1.4	11.4
MED W NN	0.9	8.8	2.3	1.6	1.0	3.8	3.0	2.3	8.8	1.3	n.s.
MED M PN	0.7	10.0	n.s.	0.1	2.8	8.7	3.8	3.3	10.0	0.0	3.1
MED M WN	0.7	10.0	n.s.	0.5	2.7	10.0	3.3	3.7	10.0	0.3	3.9
MED M NN	4.0	10.0	n.s.	0.7	5.5	8.8	3.3	3.8	10.0	0.3	n.s.

372 n.s. not simulated


373

### 374 3.3 Aggregation effect over different time periods

375 The impact of scales is also tested for periods shorter than 29 years (Figure 6 and 7). The  
 376 simulation results are averaged for each time step (according to the considered period of 1,  
 377 5, 10, 15, 20, 25 years) over the entire area (for each resolution separately). While the  
 378 maximum aggregation effect is strongest for a single year and does not change greatly for  
 379 periods of 10 years or longer, the average aggregation effect stays almost the same, but  
 380 decreases for the 29 year period. The effects are the same for all models for both crops  
 381 (Figure 6 and 7), but the values differ.



382  
 383 Figure 6: The relative differences between the maximum and minimum NPP (wheat PN) between  
 384 the resolutions for data averaged over different time periods (annual to 29 year averages). On the  
 385 left side the differences are averaged for each period, while the right side shows the maximum  
 386 values for each period.  
 387



388  
 389 Figure 7: The relative differences between the maximum and minimum NPP (maize PN) between  
 390 the resolutions for data averaged over different time periods (annual to 29 year averages). On the  
 391 left side the differences are averaged for each period, while the right side shows the maximum  
 392 values for each period.

393

394 Table 3: Minimum and maximum average length of the growing season of wheat for each  
395 resolution (overall averages) as determined in the different models. The shortest length represents,  
396 in all cases, the length of the growing season for resolution of 1km and longest growing season  
397 was for all models the resolution of 100 km.

398

model	min. length [d]	max. length [d]	change [d]
HERMES	256.1	257.6	1.5
APSIMmod	249.9	253.9	4.0
COUP	239.3	241.7	2.4
LINTUL	243.9	249.1	5.2
MONICA	259.5	261.1	1.6
STICS	236.4	239.0	2.6
APSIM	254.7	257.1	2.4
AgroC	238.2	243.3	5.1

399

400 The NPP values in this study represent NPP during growing season. Length of growing  
401 season varies between the different models and the different years (Table 3), because of  
402 different phenological sub-models and inter-annual variations of temperature. As already  
403 mentioned, the date for the latest possible harvest is fixed and this date is used as actual  
404 harvest by the models CENTURY, DailyDayCent and LandscapeDNDC. The highest  
405 differences between the lengths of the growing season are observed for LINTUL and AgroC,  
406 while there are relative little changes for HERMES and MONICA.


407

### 408 *3.4 Vulnerability analysis*

409 The results of the vulnerability analysis are represented in Figures 8 and 9, showing  
410 vulnerability (grey bars) and the risk (black bars) for each model and resolution. Negative  
411 values indicate a higher NPP average for hazardous conditions than for the non-hazardous  
412 conditions (vulnerability) or a higher NPP for the overall average than for the non-hazardous


413 conditions (risk). The analysis shows positive values for vulnerability and risk for all  
414 simulation results of wheat, except for DailyDayCent and the two APSIM models (Figure 8).  
415 The values between the different resolutions vary for the different models. The vulnerability  
416 analysis for the maize simulations shows a negative risk and vulnerability for  
417 LandscapeDNDC, while LINTUL and DailyDayCent vary between the resolutions (Figure 9).  
418 Overall, vulnerability and risk differ for most models depending on the resolution, but there  
419 is no clear trend for increase or decrease of vulnerability or risk towards coarser resolution.  
420 The average risk for wheat simulations is about  $0.02 \text{ g C m}^{-2} \text{ d}^{-1} \pm 0.02 \text{ g C m}^{-2} \text{ d}^{-1}$ , with no  
421 trend between the different resolutions and the average vulnerability of  $0.13 \text{ g C m}^{-2} \text{ d}^{-1}$  with  
422 a standard deviation of  $0.10 \text{ g C m}^{-2} \text{ d}^{-1}$  shows also no clear trend. The differences of  
423 vulnerability between the five resolutions show the maximum difference between 1.6 % for  
424 CENTURY and 12.8 % for LINTUL (maximum difference relative to the average NPP of  
425 resolution of 1 km) for wheat and between 1.1 % (CENTURY) to 15.5 % (MONICA) for  
426 maize. In these calculations the models AgroC, APSIM, APSIMmod and DailyDayCent for  
427 wheat and APSIMmod, LandscapeDNDC and LINTUL for maize are not considered,  
428 because the results of these models indicate no vulnerability to drought under these  
429 conditions. In contrast to wheat, the vulnerability analysis of maize shows mainly negative  
430 values (Figure 9), except for STICS and AgroC (positive values), and DailyDayCent and  
431 LINTUL (varying values). The number of values (cells and years with a SPEI < -1) may affect  
432 the results, but the number of extreme cells (based on SPEI) is within a narrow range of  
433 17.6 – 18.2 % for the different resolutions, so the relative numbers of hazardous cells stays  
434 about the same.

435



436  
 437 Figure 8: Vulnerability (grey bar) and risk (black bar) for 11 models for water limitation simulation  
 438 results of NPP (WN). The results represent the simulation results for wheat for the period 1983-  
 439 2011. The terms vulnerability and risk are used in the definition by van Oijen et al. (2014) and  
 440 describe the impacts of hazardous in comparison to non-hazardous conditions (see also section  
 441 2.6).

442  
 443



444  
 445 Figure 9: Vulnerability (grey bar) and risk (black bar) for 10 models simulated for water limitation  
 446 considered (WN). The results represent the simulation results for maize for the period 1983-2011.

447 The terms vulnerability and risk are used in the definition by van Oijen et al. (2014) and describe  
448 the impacts of hazardous in comparison to non-hazardous conditions (see also section 2.6).

449

## 450 **4 Discussion**

451

### 452 *4.1 NPP differences between the models*

453 The simulated potential NPP averages of the growing season for the 1 km resolution range  
454 from 1.8 to 4.1 g C m<sup>-2</sup> d<sup>-1</sup> and 7.4-12.8 g C m<sup>-2</sup> d<sup>-1</sup> for wheat and maize, respectively, which  
455 is higher than annual NPP averages of European croplands (550 ± 50 g C m<sup>-2</sup> yr<sup>-1</sup>; Schulze  
456 et al., 2010) and this is expected for crops in the study region. Because the 1 km grid maps  
457 are the highest resolution, we assume these data as the most accurate of the available data  
458 and use these data as baseline, because detailed measurements with crop yields in its  
459 spatial distribution are missing. As mentioned, the results base on simulation runs of  
460 uncalibrated models, but adjusted to proxies for a 30 year average of crop yield. In spatial  
461 modelling data for calibration are rarely available or, if available, often restricted to one or  
462 some point measurements. This makes appropriate calibration for spatial modelling difficult  
463 and adjustment to a 30 year is an appropriate method to set up the model. The two models  
464 with low NPP for wheat (APSIMmod) and high NPP for maize (LandscapeDNDC) are most  
465 likely under- and over-estimates of NPP, because of the lack of calibration. As the results  
466 are not unrealistic for crop yields in central Europe, the results from both models are used  
467 in the analysis. The NPP differs between the models up to 2.3 and 5.4 g C m<sup>-2</sup> yr<sup>-1</sup> for wheat  
468 and maize, respectively, while the range for yield, the target variable of the model settings,  
469 is with 7.6 to 8.7 t ha<sup>-1</sup> and 15.4 to 17.6 t ha<sup>-1</sup> for wheat and maize, respectively, smaller  
470 (Hoffmann et al., 2015). As mentioned above, the approaches for calculating NPP are  
471 different and by grouping the models according to these approaches of radiation use  
472 efficiency, difference between GPP and respiration or direct calculation of NPP reduces the

473 differences within the groups. The two exceptions for wheat (APSIMmod) and for maize  
474 (LandscapeDNDC) are already mentioned above. This means the model structure affects  
475 the NPP and the differences in the structure induce the wide range of NPP averages.  
476 A comparison between the PN, WN and NN approaches enables water limitation and  
477 nitrogen limitation impacts on NPP to be detected. Nitrogen limitations play a minor role for  
478 the study region, because of sufficient fertilization. As Figure 1 shows, the results of the  
479 COUP model indicate a strong sensitivity to water limitation (26 - 30 % decrease of average  
480 NPP), and LINTUL is also sensitive to water limitation (4.7 - 6.1 % decrease of average  
481 NPP), while the other models only show little sensitivity to drought stress on the overall  
482 averages (all < 2 % difference). The differences of sensitivity between the models do not  
483 show a specific impact on the aggregation affect. Neither the strength of the effect nor the  
484 changes between PN and PW are similar to each other or different to the other models.  
485 The contradicting spatial distribution of high and low NPP values reflect different crop  
486 parameters and phenological sub-models applied in the different models. In contrast to the  
487 NPP, the distribution of yield does not necessarily show a similar spatial pattern in the study  
488 area. These differences between NPP and yield are related to the different impacts of  
489 temperature changes on the simulation of phenological stages, which affects the lengths of  
490 the growing season differently to the length of the grain filling period. There is additional  
491 biomass production of wheat from the extension of the growing season, while the grain filling  
492 period does not necessarily benefit from warmer climatic conditions. The example shows  
493 the results for the year 2003, which was a severe drought period (Ciais et al., 2010) starting  
494 from mid-July and was considered by the calculation of the SPEI in the vulnerability analysis.  
495 The harvest at lower elevation started before the drought period, while the primary  
496 production at higher elevation was affected by the drought. The extension of the growing  
497 season allowed an over-compensation of NPP by a growing season that was 53 days longer,  
498 while the yield values were affected by drought, which could not be compensated for by a 7

499 day longer grain filling period. Both day of anthesis and day of maturity are determined based  
500 on temperature sums by the phenological model. For warmer areas, both will be earlier in  
501 the year compared to colder areas during the same period. In contrast, biomass production  
502 benefits from an extension of the growing season, due to additional days of production.  
503 However, the period between anthesis and maturity (both dates are represented by  
504 averages overall years and grid cells per resolution) is up to 5 days shorter when coarser  
505 resolution data are used for all models (Table 3). The models with the higher NPP in the  
506 higher elevated areas show the largest scale effect, which reflects a sensitive reaction of  
507 the phenological models to temperature changes. Van Bussel et al. (2011) reported minimal  
508 scale effects on the phenological stages, but these changes have still larger impacts on  
509 NPP.

510

#### 511 *4.2 Model specific aggregation effect*

512 The differences between the resolutions of the wheat NPP simulations show three groups  
513 of models. APSIM, LINTUL, APSIMmod and AgroC show stronger effects (2.4-7.8 %) than  
514 the other models, and HERMES, CENTURY, DailyDayCent and MONICA show minimal  
515 impacts (<1.2 %), while COUP and STICS lie between (1.8-2.2 %). The groups of models  
516 with medium and strong aggregation effects are all models that determine NPP based on  
517 the radiation use efficiency, while the other models use temperature based approaches.  
518 Hoffmann et al. (2015) investigated the aggregation effect on yields for the same set of  
519 models and suggested that the aggregation effect on radiation may not be much higher than  
520 on temperature, but the models might be sensitive to changes in radiation. Despite trends  
521 of decreasing temperature and solar radiation at coarser resolution (Figure 2 and Table 2 in  
522 Hoffmann et al., 2015), some models show increasing potential NPP values (LINTUL,  
523 APSIMmod and AgroC for wheat and DailyDayCent, LandscapeDNDC and AgroC for  
524 maize). These contradictory trends are also related to an extended growing season caused



525 by different approaches for the calculation of the phenological stages. Therefore, decreasing  
526 temperatures affect an extension of the growing season which compensates, or over-  
527 compensates, the effect of lower temperatures and radiation on crop growth as already  
528 discussed above. The growing season is extended by 1.5 to 5 days on average at coarser  
529 resolutions for wheat, and 1-2 days on average for maize simulations (Table 3),  
530 accompanied by a temperature decrease of 0.3°C (Hoffmann et al., 2015). As the model  
531 structure differs, the models show a different sensitivity to this effect and differ in their trends  
532 through the different resolutions. These results concur with the findings by van Bussel et al.  
533 (2011), who found only small impacts of climate aggregation on determined phenological  
534 stages. The models APSIMmod, LINTUL and AgroC showed the longest (4-5 days)  
535 extension to the average growing season towards the coarser resolutions, while all other  
536 models showed lower extensions (<2.5 days). The models with the larger aggregation  
537 effects are the same models that show large differences in the length of the growing season.  
538 The findings of the overall aggregation effect on NPP agrees to the findings of Nungesser  
539 et al. (1999), who found a mean uncertainty of < 2 % on NPP by modifying precipitation and  
540 solar radiation for resolutions of 10 and 50 km. The average differences for the wheat NN  
541 approach in this study (1.2 %) is also in this range, but includes a large increase for AgroC  
542 of up to 3.3 %. Pierce and Running (1995) determined an aggregation error of 15 – 30 %,  
543 depending on the time step, while the maximum time period is one year and the higher  
544 values are related to impacts on daily NPP. The relative small aggregation effect agrees  
545 also to the findings of de Wit et al. (2005), who found only small impacts of climate data  
546 aggregation on yield. Hoffmann et al. (2015) analysed the aggregation effect for yield and  
547 found an uncertainty of 0.2 t ha<sup>-1</sup> (30 year average, over the entire study area and  
548 considering the simulation results of 13 models), which is about 2.5 % to the determined  
549 yield. A comparison with the maximum aggregation effects on yield for each model shows  
550 4.2 - 8.4 % and 3.6 – 7.6 % for wheat PN and WN, respectively, higher impacts on yield than

551 for NPP, but also only small changes of the aggregation effect between the PN and WN  
552 simulations. Despite the high differences of precipitation between the five resolutions, the  
553 aggregation effect do not change strongly for the approaches WN and NN. For most models  
554 the aggregation effect even decrease. This can be explained by the little relevance of  
555 drought and nitrogen limitation for long term averages in the considered region. In regions  
556 with higher drought or nutrient stress the aggregation effect might be higher. The region is  
557 picked to represent an agricultural managed area in central Europe and the results can be  
558 transferred to comparable regions. Regions of other climate conditions (higher drought  
559 stress) or with different management practices (higher nutrient stress) might show different  
560 aggregation effects and further work needs to be done in upcoming studies to get a  
561 conclusive answer on this.

562 The differences between the aggregation effect for the average and the median NPP reflects  
563 the impacts on the variance. Changes due to aggregation may be more relevant for more  
564 extreme years and show stronger impacts that are levelled out in the average values. It can  
565 be assumed that the extremes are more likely to impact negatively on the NPP values, since  
566 the maximum growth will be restricted. The strongest differences arising from the  
567 aggregation effect on averages and median are detected for models that show a decreasing  
568 aggregation effect towards the coarser resolution. The effect may be compensated by the  
569 general trend of the NPP values for coarser resolutions.

570 If the simulation runs are used as ensembles, the aggregation effect are relatively low. For  
571 the wheat simulation the aggregation effect is almost as low as the minimal effect on an  
572 individual model run and can be explained by the different trends of the individual simulation  
573 approaches of the different models. In contrast, the simulations on maize show similar trends  
574 from the high resolution to the low resolution for models that show a high aggregation affect,  
575 which is reflected in a higher aggregation effect than for the wheat simulation. However, with  
576 2 % is this effect still small.

577

#### 578 *4.3 Aggregation effect over different time periods*

579 The analysis considered 29 year averages, while the impact of scale increases for shorter  
580 time periods (Figures 6 and 7). The scale effect shows the highest differences for single  
581 years, and stabilises to constant differences or only minor changes for periods of 15 years  
582 and longer. This suggests that the resolution of choice depends on the temporal scale as  
583 well as on the research question. The mean uncertainty for longer periods will be below 4  
584 %. The maximum aggregation effect for single years can be more than 9 % for wheat and  
585 more than 13 % for maize, but will decrease to below 4 % for time periods longer than 10  
586 years. While the largest impact on the aggregation for the maximum differences is from 1 to  
587 5 years, the mean aggregation effect shows the maximum change for the step from 25 to  
588 29 years. The NPP is a non-linear process and is especially affected by extreme events  
589 (Reichstein et al., 2013). Extreme weather conditions have an impact for a short period and  
590 affect often only a year and can be reduced by spatial averaging of the climate data. This  
591 spatial averaging is represented by the higher aggregation effect for the annual data, but  
592 this effect is already compensated by averaging over longer time periods. The impact of the  
593 temporal averages is represented in the two graphs in figure 7, which shows lower impacts  
594 on the averaged short term aggregation effect than on the maximum impact for a year.

595 Pierce and Running (1995) also found changes of the aggregation effect depending on  
596 temporal scale. They observed decreasing aggregation errors for increasing periods, which  
597 supports our findings. However, their study considers daily to annual time periods and not  
598 period lasting 1 – 29 years. An error of 15 % is in agreement with the maximum values of  
599 the analysis of the annual aggregation effect, which shows a range of 1.6 – 9.4 % for wheat  
600 and 3.4 - 13.6 % for maize.

601

#### 602 *4.4 Vulnerability analysis*

603 The impact of the the simulated phenological stages is already mentioned earlier in context  
604 of other effects, but is also important in the vulnerability analysis through varying the length  
605 of the growing season. Van Oijen et al., 2014 suggested the period from April to September  
606 as the best period to determine the drought index and found minimal impacts by starting this  
607 calculation of the drought index earlier, while we show a strong impact in this study. Van  
608 Oijen et al. (2014) used a biogeochemical model with fixed harvest dates for crop  
609 simulations or considered simulation results of forests and croplands with a fixed length of  
610 the growing season, but the models in this study are mainly crop models with dynamic  
611 growing season length. Therefore, in some years, the growing season ends before any  
612 drought impacts on crop growth occur, as for the year 2003. Van Bussel et al. (2011)  
613 described the scale impact on the modelling of phenological stages as minor, but these  
614 changes can affect other processes as, in this case, the impact of a drought period on  
615 primary production. In this study the relevant period to define hazardous conditions for the  
616 vulnerability analysis is from February to July, which is similar to the length of the growing  
617 season for most, but not all, models.

618 Not all models show sensitivity to the hazardous conditions defined by the SPEI. For the  
619 model DailyDayCent, the threshold of the SPEI = -1 is not significant in comparison to the  
620 internal drought effects. As the index SPEI is calculated by precipitation and temperature,  
621 there might be a discrepancy in the detection of extreme weather events by the APSIM  
622 models, which use the radiation use efficiency to determine NPP. In contrast to the  
623 vulnerability analysis of wheat, the analysis for maize shows no impact of drought or  
624 negative values for vulnerability, with the exceptions STICS and MONICA, because maize  
625 is a C<sub>4</sub> plant and is more drought tolerant (Lopes et al., 2011). The comparison of the WN  
626 and PN simulation runs support the findings of the vulnerability analysis, as most models  
627 show only small differences. Analysis for higher thresholds is not useful for the considered  
628 study area, because the number of grid cells for hazardous conditions becomes too small

629 for solid statistics. Regarding the resolution, the results show either all positive or negative  
630 values (HERMES, STICS, CENTURY, APSIMmod, LandscapeDNDC, APSIM, AgroC), or  
631 marginal differences to zero vulnerability (DailyDayCent, LINTUL, MONICA). The number  
632 of extreme weather events is expected to decrease at coarser scales, but the overall  
633 averages show little change between the resolutions. The expected pattern of stronger  
634 impacts of extreme events for finer scales with lower impacts at coarser scales is not seen.  
635 Assuming a threshold from SPEI = -1 to define drought conditions affects the same ratio of  
636 hazardous grid cells for the different resolutions. In contrast to the initial assumption of less  
637 extreme events, the number of grid-cells defined as extreme show the same or even higher  
638 ratio for the coarser resolution. One reason is the temporal scale for the NPP calculation,  
639 which is annual. In the vulnerability analysis the extreme events are defined during the first  
640 half of the year, which includes the drought of 2003 which had negative impacts on cropland  
641 NPP (Ciais et al., 2010), but not all model results are affected by this drought, because of  
642 an earlier harvest. Extreme events, therefore, appear to play a minor role in this area for  
643 long term averages.

644 The aggregation effect is stronger for the vulnerability than for the NPP averages. Despite  
645 for AgroC and for the wheat simulation results of APSIMmod the aggregation effect for the  
646 vulnerability at least doubles in comparison to the long term averages. The effect might be  
647 influenced by fewer years considered in the vulnerability analysis, but the maximum values  
648 even exceed the effect for annual averages. This shows that especially periods with extreme  
649 weather conditions get stronger affected by the aggregation of climate input data than other  
650 years. Over long term averages these impacts may level out.

651

## 652 **5 Conclusions**

653

654 NPP differs, depending on spatial resolution of climate input data by up to almost 8 % and  
655 5 % for wheat and maize, respectively. For most models, the overall averages are affected  
656 by only 2 % or less, but for shorter time periods (shorter than 15 years), the aggregation  
657 effect may rise for annual NPP to over 9 or over 13 % for wheat and maize, respectively. It  
658 is concluded that a large part of the aggregation effect is related to the changes in  
659 phenology. The aggregation effect affects the vulnerability stronger than long term averages,  
660 which shows the stronger impact of aggregation effects for periods with extreme weather  
661 conditions. A finer spatial resolution of climate input data will not greatly improve simulations  
662 for long term averages of NPP or vulnerability, but for periods shorter than 15 years, or areas  
663 with extreme conditions finer resolution matters and at most differed by 13 % for averages.  
664 The biggest changes are detected for the steps from 1 km to 10 km resolution and from 50  
665 km to 100 km. The current study suggests that long term NPP averages over large areas  
666 (e.g. regional scale) are relatively insensitive to climate data aggregation, whereas data  
667 aggregation would influence average NPP under extreme weather conditions. Based on  
668 these results there is no need to simulate long term NPP averages for a high resolution, if  
669 soil type and management do not vary in time and space. As this is an unrealistic scenario,  
670 more work is required to investigate the impacts, for heterogenetic soil types and varying  
671 management conditions.

672

## 673 **Acknowledgements**

674 This study was supported by the BMBF/BMELV project on "Modeling European Agriculture with  
675 Climate Change for Food Security (MACSUR)" (grant no. 2812ERA115).

676

## 677 **References**

678 Beguería, S, Vicente-Serrano SM (2012). Calculation of the Standardised Precipitation-  
679 Evapotranspiration Index.  
680

681 Bergez JE, Chabrier P, Gary C, Jeuffroy MH, Makowski D, Quesnel G, Ramat E, Raynal  
682 H, Rousse N, Wallach D, Debaeke P, Durand P, Duru M, Dury J, Faverdin P, Gascuel-  
683 Odoux C, Garcia F (2013) An open platform to build, evaluate and simulate integrated  
684 models of farming and agro-ecosystems. *Environ Modell Softw* 39: 39–49  
685

686 Brisson N, Mary B, Ripoche D, Jeuffroy MH, Ruget F, Nicoullaud B, Gate P, Devienne-  
687 barret F, Recous S, Tayot X, Plenet D, Cellier P, Machet J, Marc J, Delécolle R (1998)  
688 STICS : a generic model for the simulation of crops and their water and nitrogen balances .  
689 1. Theory and parameterization applied to wheat and corn. *Agronomie* 18: 311–346  
690

691 Brisson N, Launay M, Mary B, Beaudoin N (2008) Conceptual basis, formalisations and  
692 parameterization of the STICS crop model, Quae. ed.  
693

694 Cale WG, Oneill RV, Gardner RH (1983) Aggregation error in non-linear ecological  
695 models. *Journal of Theoretical Biology*, 100(3), 539-550  
696

697 Chen C, Wang E, Yu Q (2010) Modeling wheat and maize productivity as affected by  
698 climate variation and irrigation supply in North China Plain. *Agron J* 102: 1037-1049  
699

700 Conrad Y, Fohrer N (2009) Modelling of nitrogen leaching under complex winter wheat and  
701 red clover crop rotation on a drained agricultural field. *Physics and Chemistry of the Earth*  
702 34: 530-540  
703

704 Del Grosso SJ, Parton WJ, Mosier AR, Hartman MD, Brenner J, Ojima DS, Schimel DS  
705 (2001) Simulated interaction of carbon dynamics and nitrogen trace gas fluxes using the  
706 DAYCENT model. In: Schaffer M (Ed.), Modeling Carbon and Nitrogen Dynamics for Soil  
707 Management. CRC Press, Boca Raton, Florida, USA, pp. 303–332.

708

709 Del Grosso SJ, Parton W, Mosier AR, Walsh MK, Ojima D, Thornton PE (2006) DAYCENT  
710 national scale simulations of N<sub>2</sub>O emissions from cropped soils in the USA. J. Environ.  
711 Qual. 35: 1451–1460

712 Ewert F van Bussel LGJ, Zhao G, Hoffmann H, Gaiser T, Specka X, Nendel C, Kersebaum  
713 K-C, Sosa C, Lewan E, Yeluripati J, Kuhnert M, Tao F, Rötter R, Constantin J, Raynal H,  
714 Wallach D, Teixeira E, Grosz B, Bach M, Doro L, Roggero PP, Zhao Z, Wang E, Kiese R,  
715 Haas E, Eckersten H, Trombi G, Bindi M, Klein C, Biernath C, Heinlein F, Priesack E,  
716 Cammarano D, Asseng S, Elliott J, Glotter M, Basso B, Baigorria GA, Romero CC,  
717 Moriondo M (2015). Uncertainties in Scaling-Up Crop Models for Large-Area Climate  
718 Change Impact Assessments. In: Rosenzweig, C; Hillel, D. Handbook of Climate Change  
719 and Agroecosystems. ICP Series on Climate Change Impacts, Adaptation, and Mitigation:  
720 Volume 3

721

722 Haas E, Klatt S, Fröhlich A, Werner C, Kiese R, Grote R, Butterbach-Bahl K (2012)  
723 LandscapeDNDC: A process model for simulation of biosphere-atmosphere-hydrosphere  
724 exchange processes at site and regional scale. Landscape Ecology; DOI:  
725 10.1007/s10980-012-9772-x

726

727 Herbst M, Hellebrand HJ, Bauer J, Huisman JA, Šimůnek J, Weihermüller L, Graf A,  
728 Vanderborght J, Vereecken H (2008). Multiyear heterotrophic soil respiration: Evaluation of  
729 a coupled CO<sub>2</sub> transport and carbon turnover model. Ecological Modelling. 214: 271-283.



730

731 Hoffmann H, Zhao G, van Bussel LGJ, Enders A, Specka X, Sosa C, Yeluripati J, Tao F,  
732 Constantin J, Raynal H, Teixeira E, B. Grosz B, L. Doro L, Zhao Z, Wang E, Nendel C,  
733 Kersebaum KC, Haas E, Kiese R, Klatt S, Eckersten H, Vanuytrecht E, Kuhnert M, Lewan  
734 E, Rötter R, Roggero PP, Wallach D, Cammarano D, Asseng S, Krauss G, Siebert S,  
735 Gaiser T, Ewert F (2015). Variability of effects of spatial climate data aggregation on  
736 regional yield simulation by crop models. *Climate Research*, Vol. 65: 53–69.

737

738 Holzworth DP, Huth NI, deVoil PG, Zurcher EJ, Herrmann NI, McLean G, Chenu K, van  
739 Oosterom EJ, Snow V, Murphy C, Moore AD, Brown H, Whish JPM, Verrall S, Fainges J,  
740 Bell LW, Peake AS, Poulton PL, Hochman Z, Thorburn PJ, Gaydon DS, Dalgliesh NP,  
741 Rodriguez D, Cox H, Chapman S, Doherty A, Teixeira E, Sharp J, Cichota R, Vogeler I, Li  
742 FY, Wang E, Hammer GL, Robertson MJ, Dimes JP, Whitbread AM, Hunt J, van Rees H,  
743 McClelland T, Carberry PS, Hargreaves JNG, MacLeod N, McDonald C, Harsdorf J,  
744 Wedgwood S, Keating BA (2014) APSIM – Evolution towards a new generation of  
745 agricultural systems simulation. *Environmental Modelling & Software*

746

747 Jansson PE, Karlberg L (2004) Coupled heat and mass transfer model for soil-plant-  
748 atmosphere systems, Royal Institute of Technology, Department of Civil and  
749 Environmental Engineering, Stockholm, Sweden, 435 pp. , URL:

750 <http://www2.lwr.kth.se/Vara%20Datorprogram/CoupModel/NetHelp/default.htm> (last fetch:  
751 10.06.2014)

752

753 Keating BA, Carberry PS, Hammer GL, Probert ME, Robertson MJ, Holzworth DP, Huth  
754 NI, Hargreaves G, Meinke H, Hochman Z, Maclean G, Verburg K, Snow V, Dimes JP,  
755 Silburn M, Wang E, Brown S, Bristow KL, Asseng S, Chapman SC, McCown RL,

756 Freebairn DM, Smith CJ (2003) An overview of APSIM, a model designed for farming  
757 systems simulation. *Eur J Agron* 18: 267-288  
758

759 Kersebaum KC (2007) Modelling nitrogen dynamics in soil-crop systems with HERMES.  
760 *Nutrient cycling in agroecosystems* 77: 39-52  
761

762 Kersebaum KC (2011) Special features of the HERMES model and additional procedures  
763 for parameterization, calibration, validation, and applications In: L.R. Ahuja and L. Ma  
764 (ed.): *Advances in Agr Syst Modeling Series 2*. 65-94  
765

766 Li Z, Liu S, Tan Z, Bliss, NB, Young CJ, West TO, Ogle SM (2014). Comparing cropland  
767 net primary production estimates from inventory, a satellite-based model, and a process-  
768 based model in the Midwest of the United States. *Ecological Modelling* 277, pp. 1–12  
769

770 Lopes MS, Araus JL, van Heerden PDR, Foyer CH (2011). Enhancing drought tolerance in  
771 C4 crops. *Journal of Experimental Botany*, Vol. 62, No. 9, pp. 3135–3153,  
772

773 Nendel C, Berg M, Kersebaum KC, Mirschel W, Specka X, Wegehenkel M, Wenkel KO,  
774 Wieland R (2011) The MONICA model: Testing predictability for crop growth, soil moisture  
775 and nitrogen dynamics. *Ecol Model* 222: 1614-1625  
776

777 Nungesser MK, Joyce MLA, David A (1999). Effects of spatial aggregation on predictions  
778 of forest climate change response. *Cimate Research*, Vol. 11, pp. 109–124  
779

780 Pierce LL, Running SW (1995). The effects of aggregating sub-grid land surface variation  
781 on large-scale estimates of net primary production. *Landscape Ecology* vol. 10 no. 4 pp  
782 239-253.

783

784 Hemming D, Betts R, Collins M (2013). Sensitivity and uncertainty of modelled terrestrial  
785 net primary productivity to doubled CO<sub>2</sub> and associated climate change for a relatively  
786 large perturbed physics ensemble. *Agricultural and Forest Meteorology* 170, pp. 79–88

787

788 Pan S, Tian H, Dangal SRS, Ouyang Z, Tao B, Ren W, Lu C, Running S(2014).  
789 Modeling and Monitoring Terrestrial Primary Production in a Changing Global  
790 Environment: Toward a Multiscale Synthesis of Observation and Simulation. *Advances in*  
791 *Meteorology*. Volume 2014, Article ID 965936, 17  
792 pages,<http://dx.doi.org/10.1155/2014/965936>

793

794 Parton WJ, Scurlock JMO, Ojima DS, Gilmanov TG, Scholes RJ, Schimel DS, Kirchner T,  
795 Menaut J-C, Seastedt T, Garcia Moya E, Kamnalrut A, Kinyamario JI (1993). Observations  
796 and modeling of biomass and soil organic matter dynamics for the grassland biome  
797 worldwide. *Global Biogeochem. Cycles* 7, 785–809.

798

799 Parton WJ, Scurlock JMO, Ojima DS, Schimel DS, Hall .O (1995). Impact of climate  
800 change on grassland production and soil carbon worldwide. *Global Change Biol.* 1, 13–22.

801

802 Parton WJ, Holland EA, Del Grosso SJ, Hartman MD, Martin RE, Mosier AR, Ojima DS,  
803 Schimel DS (2001) Generalized model for NO<sub>x</sub> and N<sub>2</sub>O emissions from soils. *J.*  
804 *Geophys. Res.* 106: 17403-17420

805

806 Raes D, Steduto P, Hsiao TC, Fereres E (2009) AquaCrop-the FAO crop model to  
807 simulate yield response to water: ii. Main algorithms and software description. *Agron J*  
808 101: 438-447  
809

810 Raes D, Steduto P, Hsiao T, Fereres E (2012) AquaCrop version 4.0 Reference Manual.  
811 FAO, Rome. 406p.  
812

813 Rastetter EB, King AW, Cosby BJ, Hornberger GM, Oneill RV, Hobbie JE (1992)  
814 Aggregating fine-scale ecological knowledge to model coarser-scale attributes of  
815 ecosystems. *Ecol Appl* 2: 55-70  
816

817 Reich PB, Turner DP, Bolstad P (1999). An Approach to Spatially Distributed Modeling of  
818 Net Primary Production (NPP) at the Landscape Scale and Its Application in Validation of  
819 EOS NPP Products. *Remote Sensing of Environment* 70:69–81  
820

821 Reichstein M, Bahn M, Ciais P, Frank D, Mahecha MD, Seneviratne SI, Zscheischler J,  
822 Beer C, Buchmann N, Frank DC, Papale D, Rammig A, Smith P, Thonicke K, van der  
823 Velde M, Vicca S, Walz A, Wattenbach M (2013). Climate extremes and the carbon cycle.  
824 *Nature* 500, 287-295  
825

826 Schulze ED, Ciais P, Luysaert S, Schrumppf M, Janssen IA, Thiruchittampalam B,  
827 Theloke J, Saurat M, Bringezu S, Lelieveld J, Lohila A, Rebmann C, Jung M, Bastviken D,  
828 Abril G, Grass G, Leip A, Freibauer A, Kutsch W, Don A, Nieschulze J, Börner A, Gash JH,  
829 Dolman AJ (2010). The European carbon balance. Part 4: integration of carbon and other  
830 trace-gas fluxes. *Global Change Biology* 16, 1451–1469, doi: 10.1111/j.1365-  
831 2486.2010.02215.x

832

833 Shibu ME, Leffelaar PA, van Keulen H, Aggarwal PK (2010) LINTUL3, a simulation model  
834 for nitrogen-limited situations: Application to rice. *Eur J Agron* 32: 255-271

835

836 Thornthwaite, CW, 1948: An approach toward a rational classification of climate. *Geogr.*  
837 *Rev.*, 38, 55–94.

838

839 van Bussel LGJ, Ewert F, Leffelaar PA (2011). Effects of data aggregation on simulations  
840 of crop phenology. *Agriculture, Ecosystems and Environment*, 142, 75-84

841

842 Van Ittersum M, Leffelaar P, Van Keulen H, Kropff M, Bastiaans L, Goudriaan J (2003) On  
843 approaches and applications of the Wageningen crop models. *Eur J Agron* 18: 201-234

844

845 Van Oijen M, Thomson A, Ewert F (2009). Spatial upscaling of process-based vegetation  
846 models: An overview of common methods and a case-study for the U.K., *StatGIS*

847 *Geoinformatics for Environmental Surveillance.*, Milos Island, Greece, 17-19 June 2009.,  
848 p. 6.

849

850 Van Oijen M, Balkovi J, Beer C, Cameron DR, Ciais P, Cramer W, Kato T, Kuhnert M,  
851 Martin R, Myneni R, Rammig A, Rolinski S, Soussana JF, Thonicke K, Van der Velde M,  
852 Xu L (2014) Impact of droughts on the carbon cycle in European vegetation: a probabilistic  
853 risk analysis using six vegetation models. *Biogeosciences*, 11, 6357–6375

854

855 Vicente-Serrano SM, Begueria S, Lopez-Moreno JI (2010). A multiscalar drought index  
856 sensitive to global warming: the standardized precipitation evapotranspiration index, *J.*

857 *Clim.*, 23, 1696–1718

858

859 Wallach D, Rivington M, Mearns LO, Antle J (2014). AgMIP White Paper on  
860 Characterizing and Quantifying Uncertainty.

861

862 Wang E, Robertson M, Hammer G, Carberry P, Holzworth D, Meinke H, Chapman S,  
863 Hargreaves J, Huth N, McLean G (2002) Development of a generic crop model template in  
864 the cropping system model APSIM. *Eur J Agron* 18: 121-140

865

866 Yeluripati JB, Van Oijen M, Wattenbach M, Neftel A, Ammann A, Parton WJ, Smith P  
867 (2009) Bayesian calibration as a tool for initialising the carbon pools of dynamic soil  
868 models. *Soil Biology & Biochemistry* 41: 2579–2583

869

870 Zhang X, Izaurrald RC, Manowitz DH, Sahajpal R, West TO, Thomson AM, Xu M, Zhao K,  
871 LeDuc SD, Williams JR (2015). Regional scale cropland carbon budgets: Evaluating a  
872 geospatial agricultural modeling system using inventory data. *Environmental Modelling &*  
873 *Software* 63, 199-216

874

875 Zhao G, Hoffmann H, Van Bussel LGJ, Enders A, Specka X, Sosa C, Yeluripati J, Tao FL  
876 Constantin J, Teixeira E, Grosz B, Doro L, Zhao Z, Nendel C, Kiese R, Raynal H,  
877 Eckersten H, Haas E, Wang E, Kuhnert M, Trombi G, Bindi M, Lewan E, Bach M,  
878 Kersebaum KC, Rötter R, Roggero PP, Wallach D, Krauss G, Siebert S, Gaiser T, Ewert F  
879 (2015) Effect of weather data aggregation on regional crop simulation for different crops,  
880 production conditions, and response variables. *Clim Res*

881

882