

OXFORD
UNIVERSITY PRESSJournal of the
Pediatric Infectious Diseases Society**High reported rates of antimicrobial resistance in Indian neonatal and pediatric blood stream infections.**

Journal:	<i>Journal of the Pediatric Infectious Diseases Society</i>
Manuscript ID	JPIDS-2016-098.R3
Manuscript Type:	Literature Review
Date Submitted by the Author:	n/a
Complete List of Authors:	Dharmapalan, Dhanya; Apollo Hospitals, Navi Mumbai, Pediatrics and Pediatric Infectious Diseases Shet, Anita; Johns Hopkins University Bloomberg School of Public Health, International Vaccine Access Center Yewale, Vijay; Dr Yewale's Multispecialty Hospital for Children, Pediatrics Sharland, Mike; St George's University London, Paediatric Infectious Diseases Unit

SCHOLARONE™
Manuscripts

Only

TITLE PAGE**Manuscript title**

High reported rates of antimicrobial resistance in Indian neonatal and pediatric blood stream infections.

Authors

1. Dhanya Dharmapalan, MD, PGDipPID^{1*}
2. Anita Shet, MD, PhD²
3. Vijay Yewale, MD, DCH³
4. Mike Sharland. MD, FRCPC, DTM&H⁴

¹ Department of Pediatrics and Pediatric Infectious Diseases, Apollo Hospitals, Navi Mumbai, India, Email: drdhanyaroshan@gmail.com

² Associate Scientist. International Vaccine Access Center, Johns Hopkins Bloomberg School of Public Health, Baltimore, USA. Email: ashet1@jhu.edu

³ Department of Pediatrics, Dr Yewale's Multispeciality Hospital for Children, Vashi, Navi Mumbai, India. Email: vnyewale@gmail.com

⁴ Pediatric Infectious Diseases Research Group, St George's University London, UK. Email: mike.sharland@stgeorges.nhs.uk

Corresponding author: Dhanya Dharmapalan

Address for correspondence: Department of Pediatrics, Apollo Hospitals, Parsik Hill Road, Sector - 23, CBD Belapur, Navi Mumbai, Maharashtra 400614, India Email: drdhanyaroshan@gmail.com

1
2
3 **Acknowledgements:** We would like to thank Dr. Asawari Kanade, PhD, for her
4
5 assistance with statistical analysis of the data presented in this review.
6
7

8 **Keywords:** Antimicrobial resistance, India , children, cephalosporins, MRSA
9

10 **Abbreviated title:** Antimicrobial resistance in Indian neonates and children
11

12 **Running head title:** Antimicrobial resistance in Indian neonates and children
13
14

15 **Conflict of interests:** None
16

17 **Source of funding:** None
18
19

20 **Manuscript word count:** 2596
21
22

23
24 **SUMMARY:**
25

26
27 In the absence of a national data on antimicrobial resistance in pediatric population in India,
28
29 this work serves as a first large pooled data review highlighting the extremely high resistance
30
31 in the commonly isolated blood-borne pathogens in Indian neonates and children against all
32
33 routinely used antibiotics.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Background

There is real shortage of national data on antimicrobial resistance rates in Indian neonates and children. A descriptive review was conducted to determine the patterns of antimicrobial resistance in isolates of bloodstream infection among hospitalised children in India

Methods

Published and grey literature on antibiotic resistance in children was searched using “Google Scholar”, “Scopus” and “Pubmed” databases between January 2000 and July 2015. Studies were included if they were original articles that reported a minimum of 10 pathogenic bacterial isolates from the bloodstream within a pediatric population in India, and excluded if they reported studies done during an outbreak or epidemic.

Results

A total of 1,179 studies were screened, and 82 papers were identified as eligible for inclusion. Most studies (78.7%) were reported from neonatal intensive care units. Among a total of 50,545 reported blood cultures, 14,704(29.1%) were positive. *Staphylococcus aureus*, (median:14.7%;IQR7.4%-25.6%) and *Klebsiella pneumoniae* (median:26%;IQR16.7%-35.4%) were the commonest reported gram-positive and gram-negative pathogens respectively. Around half of all *Staphylococcus aureus* isolates were reported as methicillin resistant *Staphylococcus aureus* (median:50%;IQR31.4%-65.1%). After age stratification, the median rate of resistance of common gram-negative pathogens to ampicillin and gentamicin/amikacin were extremely high. (*Klebsiella pneumoniae*/ampicillin 95.9% ; *Klebsiella pneumoniae*/gentamicin 75% ; *Escherichia coli*/ampicillin 92.9% ; *Escherichia coli*/gentamicin 55.6%) .Similarly the median resistance of common gram-negative blood stream isolates to cephalosporins were also high (*Klebsiella pneumoniae*/cefotaxime 62.6% ; *Escherichia coli*/cefotaxime 47.5%).

1
2
3 Conclusions
4

5 High rates of resistance to WHO recommended first-line treatment options for neonates and
6
7 children have been identified in blood stream infections across India. There is an urgent need
8
9 to both enhance antibiotic stewardship and infection prevention and control measures and
10
11 consider urgently how to repurpose older antibiotics back into routine care in India.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

MANUSCRIPT

Introduction

There has been a relentless rise in antimicrobial resistance globally. Concerns about this serious threat have been raised since late 1990s as evident by the Global Strategy for Containment of Antimicrobial Resistance of the World Health Organisation (WHO) in 2001[1]. The WHO in its recent global surveillance report states that this reality of antimicrobial resistance is “far from an apocalyptic fantasy”, is happening right now in every region of the world and has the potential to affect anyone, of any age, in any country[2].

Due to lack of a national robust surveillance system for Antimicrobial Resistance (AMR) in India, there is currently no centrally reported AMR national database. Earlier initiatives to generate AMR data include those carried out by the India clinical epidemiological network (INCLEN) and Indian network for surveillance of antimicrobial resistance (INSAR)[3,4]. In 2011, the Indian Council of Medical Research (ICMR) launched the national policy for containment of antimicrobial resistance which includes sentinel surveillance of AMR phase wise across the country[5]. This surveillance has not defined any age stratification. The Jaipur Declaration in 2011[6] and Chennai declaration[7] in the subsequent year framed consensus roadmaps to tackle AMR.

AMR patterns can significantly differ in adult and children[8]. It is estimated that India has the highest neonatal mortality due to neonatal sepsis caused by bacteria resistant to first-line antibiotics[9]. Nearly one-fifth of neonates with sepsis die in the hospital and the mortality rises to 50% for those with culture-proven sepsis[10]. In the absence of any recent national data of AMR rates in blood stream infections in the Indian neonatal or pediatric population, this study aimed to review the published and unpublished data on AMR during 2000-2015 from India.

Methods:

The published and grey literature on antibiotic resistance in children in English literature was searched using the databases “Google Scholar” “Scopus” and Pubmed using keywords “India” AND “children” OR “pediatrics” OR “neonates” OR “infants” AND “antimicrobial resistance” OR “antibiotic resistance” OR “antibiotic susceptibility” OR “antibiotic sensitivity” AND “blood culture” (see supplementary material 1). Grey literature sources such as pediatric conference proceedings and national reports were examined individually. The period between 1 January 2000 and 1st July 2015 was chosen to access maximum information of published abstracts and full text articles. A single reviewer assessed the study quality and performed data extraction.

Inclusion criteria included original articles describing studies performed in well described health centres in India, published between January 2000 and July 2015 that described blood culture-derived bacterial isolates from children from birth to 18 years, with determination of antimicrobial susceptibility patterns. Excluded studies were those that described less than 10 pathogenic isolates, or those that did not distinguish bloodstream and non-bloodstream sources, and those performed in adult populations. Studies performed during outbreaks or epidemics were also excluded. Bacterial isolates of interest reviewed here included gram-positive bacteria (*Staphylococcus aureus*, coagulase-negative *Staphylococcus*, *Enterococcus fecalis*) and gram-negative bacteria (*Klebsiella pneumoniae*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Enterobacter species*, *Citrobacter species*, *Acinetobacter baumannii*, *Proteus species* and *Salmonella typhi*).

Statistical Analysis

1
2
3 The proportions (%) and ranges of individual blood-borne pathogens in infants and children,
4
5 resistant to various drugs mentioned in Table 1, 2 and 3 were calculated. As the total number
6
7 of reported blood cultures varied markedly between studies, proportions and ranges for
8
9 pathogens were weighted by total sample size of blood cultures and given as weighted
10
11 median of proportions (%) and inter-quartile ranges (IQRs). Similarly, median and inter-
12
13 quartile ranges (IQRs) of resistance rates attributable to a pathogen to recommended
14
15 antibiotics were reported [11].
16
17

18 **Results**

19 **Search Analysis**

20
21
22 Electronic search on 9th July 2015 identified 298 articles in Pubmed, 4,980 articles in the
23
24 database of “Google scholar” and 243 articles in Scopus. Of the 4,980 articles, Google
25
26 scholar engine displays only the first 1,000. Individual search of abstracts from national
27
28 conferences of Indian Academy of Pediatrics identified 8 abstracts. 72 articles were common
29
30 in both search engines. So, in total, 1,179 abstracts were reviewed. Most of the articles were
31
32 excluded because the studies were done in children living outside India or in the adult
33
34 population (Figure 1). Among the remaining 484 articles, those that met the exclusion criteria
35
36 were excluded.
37
38

39
40 Thus, 82 papers (see supplementary material 2) were included for analysis. Among these 82
41
42 papers, some papers had multiple data sets covering multiple years. Since the data set
43
44 belonging to a particular year was considered as a separate study, the total number of studies
45
46 obtained was 89. The positive blood culture rate was calculated using 78 of the 89 studies, as
47
48 the total number of blood cultures analysed were not available in the remaining 11 studies.
49
50

51 **Study characteristics**

52
53 Among the 89 eligible studies that had provided data about positive blood cultures, 78.7%
54
55 data were from neonates, 14.6% from children older than 1 month, and 6.7% studies were
56
57
58
59
60

1
2
3 done in both neonate and pediatric ages. Studies were spread across India, with 36% reports
4 from North India, 31.5% from South, 16.9% from Western India, 12.4% from the East and
5
6 3.4% from Central India. Out of the 89 studies, 98.9% had been conducted at a tertiary level
7
8 health care centre and only one study (i.e. 1.1%) had been reported from a secondary level
9
10 health care centre. The great majority of studies (78.7%) had been reported from NICU and a
11
12 small percentage (<5%) from PICU ward, IPD, or their combinations. One study had
13
14 screened enteric cases using blood cultures from the outpatient department[12].
15
16

17 *Blood Culture positivity*

18
19
20 Data were available from a total of 16,777 positive blood cultures from 89 studies, which
21
22 were used to examine the drug resistance of the isolates. The number of total blood cultures
23
24 reported was available only in 78 of the 89 studies, which summed to a total 50,545 blood
25
26 cultures. The positive blood culture rate was calculated from the 78 studies that, varied from
27
28 7.2% to 88.5% with a median of 35.4% (IQR= 21.2% - 46.6%).
29
30

31 *Pathogens identified*

32
33 Among 72 studies that reported gram-positive bacteria the median of percentage gram-
34
35 positive bacteria was 29.2% (IQR=15.8%-36.8%). Among gram-positive bacteria,
36
37 *Staphylococcus aureus* was found to be the most common isolate (median 14.7%; IQR 7.4%-
38
39 25.6%) followed by *coagulase negative Staphylococcus* (CONS) (median 10.4%; IQR 4.2%-
40
41 15.9%) and *Enterococcus faecalis* (median 0.9%;IQR 0%-4.4%) (Table 1). It was not
42
43 possible to distinguish true pathogens of CONS from contaminants in this review.
44
45

46
47 Gram-negative bacteria were reported in 81 studies. The median percentage gram-negative
48
49 bacteria identified among all reported positive cultures was 61.0% (IQR=34.6% -67.9%).
50

51 Among gram-negative bacteria, *Klebsiella pneumoniae* was found to be the most common
52
53 isolate (median 26%; IQR 16.7%-35.4%) followed by *Escherichia coli* (Median 9.3%; IQR
54
55

1
2
3 4.7%-14.5%), *Acinetobacter baumannii* (median 5.9%; IQR 2.1%-10.8%) and *Pseudomonas*
4
5 *aeruginosa* (median 4.8%; IQR 2.4%-9.4%)

6
7 *Resistance of Gram-positive bacteria:*

8
9 50% of the *Staphylococcus aureus* isolates were methicillin resistant *Staphylococcus aureus*
10 (MRSA). High resistance was also noted for *Staphylococcus aureus* to erythromycin (53%),
11 cefotaxime (57%) and cotrimoxazole (57.7%) (Table 1). Lower resistance pattern was shown
12 by CoNS except to cotrimoxazole (69.9%). However *Enterococcus faecalis* showed very
13 high resistance to antibiotics including penicillins (88.5%), gentamicin (68.5%) and
14 ciprofloxacin (50%). No resistance was found to vancomycin, linezolid and teicoplanin in all
15 the gram-positive bacteria reported.

16
17 *Resistance of Gram-negative bacteria:*

18
19 Details of the gram-negative resistance rates are provided in Table 2. High levels of
20 resistance was reported in *Klebsiella pneumoniae* to ampicillin (95.2%) and cephalosporins
21 (over 60%). The rates of reported resistance to tigecycline were 8.2%, colistin, 3.8% and
22 meropenem, 1%. Reported resistance in *Enterobacter* spp. was 100% to ceftriaxone and
23 aztreonam. Resistance to piperacillin-tazobactam was 16.7%, ciprofloxacin (39%) and
24 meropenem (16.7%). There were very high resistance rates identified in *Escherichia coli*
25 (Table 2) to ampicillin (92.3%), chloramphenicol (63.9%) and ciprofloxacin (40%) and
26 relatively lower resistance to amikacin (22.4%) and piperacillin-tazobactam (16.7%).
27
28 Resistance to meropenem of 9.0% and colistin of 8.8% was noted. As seen in Table 2, high
29 resistance of *Pseudomonas aeruginosa* was noted to ceftazidime (58.7%) which is often used
30 empirically for *Pseudomonas aeruginosa* infections. There was high resistance (16.7%) noted
31 of *Pseudomonas aeruginosa* to meropenem.

32
33 In case of resistance of *Acinetobacter baumannii* and *Citrobacter* spp. (table 2), though the
34 resistance to cephalosporins are high, there was relatively less resistance to ciprofloxacin and
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 beta-lactam antibiotics including piperacillin-tazobactam and cefoperazone-sulbactam.
4
5 *Acinetobacter baumannii* infections were reported to have 11.5% resistance to meropenem.
6
7 *Salmonella typhi* (Table 3) displayed a median resistance of 50% to ampicillin, 35% to
8
9 nalidixic acid and 17.9% to chloramphenicol. However 6.3% resistance to ceftriaxone and
10
11 16% to second-line antibiotics like azithromycin were also noted.

12 13 14 *Age-based stratification for commonly used pathogen-drug combinations* (Table 4)

15
16
17 Using age as criterion, the data was then split by age into neonatal (less than 1 month) and
18
19 pediatric populations (age greater than 1 month) for commonly used pathogen-drug
20
21 combinations.
22

23 24 25 *a. Gram-positive resistance in neonates vs pediatric population*

26
27 The overall resistance was found to be greater in neonatal age than pediatric age. However,
28
29 this difference was not statistically significant. There was no resistance noted of isolates of
30
31 *Staphylococcus aureus* and CONS to vancomycin, and linezolid, in both age groups.
32
33

34 35 36 *b. Gram-negative resistance in neonates vs pediatric population*

37
38 The median resistances of common gram negative pathogens to the WHO recommended
39
40 combination of ampicillin and gentamicin for the empiric treatment of neonatal sepsis were
41
42 extremely high (*Klebsiella pneumoniae* /ampicillin 95.9% ; *Klebsiella pneumoniae*
43
44 /gentamicin 75% ; *Escherichia coli*/ampicillin 92.9% ; *Escherichia coli*/gentamicin 55.6%)
45
46 and median resistance to cephalosporins were also high (*Klebsiella pneumoniae* /cefotaxime
47
48 62.6% ; *Escherichia coli*/cefotaxime 47.5%).
49
50

51 **Discussion**

52 53 54 Main findings

1
2
3 This literature review provides a comprehensive analysis on the antimicrobial resistance
4 pattern of bacterial blood stream infection (BSI) in the pediatric population of India over the
5 past 15 years. This study found higher reports of bacteraemia caused by gram-negative
6 bacteria (53.3%) compared to gram-positive bacteria (30.9%).
7
8

9
10
11 The majority of the BSI identified in this study (78.7%) were reported from tertiary NICU's.
12 These NICUs reported very high rates of MRSA BSI (53.6%). WHO recommends ampicillin
13 plus an aminoglycoside (gentamicin or amikacin) as empirical treatment for community
14 acquired neonatal sepsis[13]. Very high levels of resistance of more than 90% to ampicillin
15 was noted among the majority of the gram-negative isolates and the median resistance to
16 amikacin ranged from 22.4% to 50%. Cephalosporins contribute to over 50% of prescribed
17 antibiotics in paediatrics in India[14,15]. Cephalosporins are recommended as first line
18 antibiotics in various community acquired infections like enteric fever, meningitis and severe
19 pneumonia. The high rates of resistance to cephalosporins in gram-negative BSI isolates of
20 both the neonatal and pediatric population is concerning (*Klebsiella pneumoniae*/cefotaxime
21 62.6% in neonates and 70% in children; *Escherichia coli*/cefotaxime 47.5% in neonates and
22 50% in children). In addition, this study noted an emerging resistance to carbapenems.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 Comparison to other studies

39
40 Despite concern about the high resistance reported from India, it is surprising to note the lack
41 of published data from India. In a study from a mixed population of adults and children
42 across 15 centres in India, the prevalence of MRSA was 42 per cent in 2008 and 40 per cent
43 in 2009. However 60% of the isolates were from skin and soft tissue infections, while the
44 present study reported only on blood culture isolates. No isolate was found to be resistant to
45 vancomycin and linezolid similar to the present study[4].
46
47
48
49
50
51
52
53

54 A high incidence of MRSA BSI in neonates was also reported from a large prospective
55 multisite cohort study by the Delhi Neonatal Infection Study (DeNIS) collaboration (38%)
56
57
58
59
60

1
2
3 [16]. Though the rate of gram-negative sepsis and the most common gram-negative
4 pathogens isolated in this study were similar to ours, the proportion of *Acinetobacter* spp
5 identified (22%) was much higher than ours (5.9%). On comparing our data with a
6 systematic review of gram-negative antimicrobial resistance in sepsis in resource limited
7 countries, the figures from Asian studies show a similar resistance pattern for gram-negative
8 isolates except for a lower resistance to cotrimoxazole and chloramphenicol[11].
9
10 Since 1990s, *Salmonella typhi* had acquired resistance to the former first line drugs
11 recommended by WHO, ampicillin, chloramphenicol and fluoroquinolones[17]. In the
12 present study, one third of invasive *Salmonella* isolates (35%) in children had resistance to
13 nalidixic acid similar to the resistance of 37.3% found in the systematic review of Asian
14 studies[10]. Ceftriaxone and azithromycin are currently recommended as first line and second
15 line therapy for multidrug resistant typhoid fever, (MDRTF). However the resistance rates
16 reported here were significant with ceftriaxone (6.3%) and azithromycin (16%). A recent
17 review on multi drug resistant typhoid fever in India had similarly noted isolated reports of
18 ceftriaxone resistance[18].
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 Study limitations

37 This study has several biases and limitations such as:

38 *Selection bias*

39 Over 80% of studies were reported from tertiary Intensive Care Units where children have
40 prolonged length of stay and underlying medical conditions. Most of the studies in the data
41 were laboratory based, with very limited clinical information, and it was not possible to
42 determine if the infections were community or hospital acquired. Despite the exclusion of any
43 study reporting an outbreak, it is probable that some studies had unrecognised outbreaks.
44
45
46
47
48
49
50
51
52
53

54 *Surveillance bias*

1
2
3 The data was largely contributed by tertiary medical centres. It is noted from CDC's
4
5 surveillance on antimicrobial resistance [19, 20] that expansion of surveillance to include
6
7 smaller community hospitals (<200 beds) resulted in reduction in the pooled mean percent
8
9 resistance for certain resistance phenotypes like MRSA.
10

11 *Reporting bias*

12
13
14 The heterogeneity in data showed the lack of any uniform protocol being followed for
15
16 reporting key drug bug combination data in a standardised manner. There were no formal
17
18 national guidelines on reporting antimicrobial resistance until February 2015 when the Indian
19
20 Council of Medical Research announced the Standard Operating Procedure (SOP)[21].
21

22 *Next steps*

23
24 Although there are many limitations and biases, this study clearly highlights the concerning
25
26 level of antimicrobial resistance in serious infections within the tertiary neonatal and pediatric
27
28 setting in India. It is important to differentiate between hospital acquired and community
29
30 acquired infections to develop appropriate empirical therapy choices. Guidelines on regional
31
32 empirical treatment cannot be made on the basis of the present study as it is not representative
33
34 of the resistance in the community [22]. Improved multi-centre, prospective studies using a
35
36 standardised protocol such as the DeNIS study, are required, with data on underlying disease,
37
38 treatment and clinical outcomes, which are essential to give a more accurate assessment of
39
40 the true burden of disease, especially in relation to the clinical outcome.
41
42
43

44
45 The factors contributing to antimicrobial resistance in India are complex and diverse and
46
47 range from unregulated use of antibiotics in the community and hospital settings, poor
48
49 infection control policies and the unavailability of culture facilities or point of care tests. The
50
51 stewardship programmes recommended by the developed countries require time, personnel
52
53 and resources which are currently completely lacking in the Indian health care system. It is
54
55
56
57
58
59
60

1
2
3 very important to develop simple, easy to implement antimicrobial stewardship programme
4
5 adapted to resource poor settings.
6

7
8 In 2016, the Government of India launched the national treatment guidelines for
9
10 antimicrobial use in infectious diseases [23]. While this is a major step to ensure a uniform
11
12 rational practice in the country, there is a need to formulate a similar guideline for neonates
13
14 and children. In addition, participation of paediatricians should be encouraged in active
15
16 surveillance for antibiotic resistance by bodies like ICMR, Indian Academy of Pediatrics and
17
18 international networks. More importantly, the awareness about the magnitude of antibiotic
19
20 resistance and the essence of rational antibiotic use needs to be highlighted more urgently
21
22 among practicing physicians and families. Resistance rates to current antibiotics are now so
23
24 high, that there is an urgent need to consider how to re-introduce older antibiotics back into
25
26 routine clinical practice and determine how the very few new antibiotics under development
27
28 can be introduced into high risk clinical care in a rational and affordable manner.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgement

We would like to thank Dr. Asawari Kanade, PhD for her assistance with statistical analysis of the data presented in this review.

For Review Only

References:

1. WHO. Global Strategy for Containment of Antimicrobial Resistance.
http://www.who.int/drugresistance/WHO_Global_Strategy_English.pdf. Accessed on August 27, 2015.
2. WHO. Antimicrobial Resistance. Global report on Surveillance. 2014.
http://apps.who.int/iris/bitstream/10665/112642/1/9789241564748_eng.pdf. Accessed August 22, 2015.
3. India clen. The INCLEN India Infectious Disease Initiative (IID) USAID/INCLEN Final Report,2014.
<http://www.inclentrust.org/inclen/uploadedbyfck/file/publication/the%20inclen/USAID%20IID-final%20version.pdf>. Accessed August 26th, 2015.
4. Indian Network for Surveillance of Antimicrobial Resistance (INSAR) group, India. Methicillin resistant Staphylococcus aureus (MRSA) in India: Prevalence & susceptibility pattern. *Indian J Med Res.* 2013;137:363-369.
5. ICMR. National Policy For Containment Of Antimicrobial Resistance, India . Directorate General of Health Services Ministry of Health & Family Welfare Nirman Bhawan, New Delhi, 2011. Available from http://www.ncdc.gov.in/ab_policy.pdf. Accessed 26th August, 2015.
6. WHO. Jaipur Declaration on Antimicrobial Resistance. 2011.
http://www.searo.who.int/entity/world_health_day/media/2011/whd-11_amr_jaipur_declaration_.pdf. Accessed August 27,2015.
7. Ghafur A, Mathai D, Muruganathan A, et al. The Chennai Declaration: a roadmap to tackle the challenge of antimicrobial resistance. *Indian J Cancer.* 2013;Jan-Mar;50(1):71-3.

- 1
2
3 8. Bielikci JA, Lundin R, and Sharland M. Antibiotic Resistance Prevalence in Routine
4
5 Bloodstream Isolates from Children's Hospitals Varies Substantially from Adult
6
7 Surveillance Data in Europe. ARPEC Project. *Pediatr Infect Dis J*. 2015;34(7):734-41.
8
- 9
10 9. Laxminarayan R, Matsoso P, Pant S, Brower C, Røttingen JA, Klugman K, Davies S.
11
12 Access to effective antimicrobials: a worldwide challenge *Lancet* 2016;387:168–75.
13
- 14 10. PHFI, AIIMS, and SC- *State of India's Newborns (SOIN) 2014-* a report. (Eds) Zodpey
15
16 S and Paul VK. Public Health Foundation of India, All India Institute of Medical
17
18 Sciences and Save the Children. New Delhi, India; 2014.
19
- 20
21 11. Le Doare K, Bielikci J, Heath PT, et al. Systematic Review of Antibiotic Resistance
22
23 Rates Among Gram-Negative Bacteria in Children With Sepsis in Resource-Limited
24
25 Countries. *J Pediatric Infect Dis Soc*. 2015;4(1):11-20.
26
27
- 28 12. Chandrashekar et. al. (2013) Study Of Clinical And Laboratory Profile Of Enteric
29
30 Fever In Pediatric Age Group. *International Journal of Basic and Applied Medical*
31
32 *Sciences*, 3(3), pp.16–23.
33
34
- 35 13. WHO. *Pocket book of hospital care for children: guidelines for the management of*
36
37 *common illnesses with limited resources*. Geneva: WHO ; 2005
38
- 39 14. Ramesh L, Samgeeta S and Hussainy SR. Analysis of Antimicrobial Prescriptions in
40
41 Pediatric Patients in a Teaching Hospital. *Asian J Pharm Clin Res*. 2012;5(2):124-128
42
43
- 44 15. Sriram S, Mathew L, Manjula A.S, Rajalingam B, Ramkumar K, Rajeswari R.
45
46 Assessment of antibiotic use in pediatric patients at a tertiary care teaching hospital.
47
48 *IJOPP* 2008;1(1):30-36.
49
- 50
51 16. Investigators of the Delhi Neonatal Infection Study (DeNIS) collaboration.
52
53 Characterisation and antimicrobial resistance of sepsis pathogens in neonates born in
54
55 tertiary care centres in Delhi, India: a cohort study. *Lancet Glob Health*. 2016
56
57 Oct;4(10):e752-60.
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
17. Kundu R, Ganguly N, Ghosh TK, et al; IAP Task Force. IAP Task Force Report: management of enteric fever in children. *Indian Pediatr.* 2006 Oct;43(10):884-7.
 18. Zaki SA and Karande S. Multidrug-resistant typhoid fever: a review. *J Infect Dev Ctries.* 2011;28(5):324-37.
 19. [National Nosocomial Infections Surveillance System](#). National Nosocomial Infections Surveillance (NNIS) System Report, data summary from January 1992 through June 2004, issued October 2004. *Am J Infect Control.* 2004 Dec;32(8):470-85.
 20. Sievert DM, Ricks P, Edwards JR, et al; National Healthcare Safety Network (NHSN) Team and Participating NHSN Facilities. Antimicrobial-resistant pathogens associated with healthcare-associated infections: summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2009-2010.
 21. ICMR Standard Operation Procedures. Bacteriology ICMR, New Delhi. http://icmr.nic.in/Publications/SOP/SOP_Bacteriology.pdf. Accessed on August 28, 2015.
 22. Saha SK, Saha S, Ruhulamin M, Hanif M, Islam M. Decreasing trend of multiresistant *Salmonella typhi* in Bangladesh. *J Antimicrob Chemother.* 1997 Apr;39(4):554-6.
 23. National Centre for Disease Control. National Treatment Guidelines for Antimicrobial Use in Infectious Diseases, version 1.0 (2016). Ministry of Health & Family Welfare India, Nirman Bhawan, New Delhi, 2016. Available from http://www.ncdc.gov.in/writereaddata/linkimages/AMR_guideline7001495889.pdf. Accessed 22nd November, 2016.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legend

Figure 1: Flow Chart of selection of studies based on PRISMA 2009 flow chart

Listing of Supplementary Material

Supplementary material 1

Supplementary material 2

For Review Only

TABLES

Table 1: Median and IQR of percentage resistance of gram-positive bacteria: *Staphylococcus aureus*, *CONS* and *Enterococcus faecalis*

	<i>Staphylococcus aureus</i> 14.7%(7.4%-25.6%) [n=70]	<i>CONS</i> 10.4%(4.2%-15.9%) [n=68]	<i>Enterococcus faecalis</i> 0.9%(0%-4.4%) [n=44]
Penicillin	NS	NS	88.5% (41.7%-100.0%) [n=9]
Ampicillin	NS	NS	100.0% (77.5%-100.0%) [n=4]
Erythromycin	53.0% (39.5%-65.9%) [n=31]	43.3% (30.5%-67.1%) [n=26]	53.2% (44.6%-61.9%) [n=8]
Cloxacillin	50.0% (31.4%-65.1%) [n=33]	42.5% (19.1%-66.7%) [n=24]	NS
Amikacin	25.8% (14.2%-48.7%) [n=40]	28.6% (0.0%-41.0%) [n=35]	NS
Gentamicin	44.9% (24.9%-69.7%) [n=42]	50.0% (29.2%-66.7%) [n=36]	68.5% (41.7%-77.5%) [n=13]
Cephelexin	34.3% (27.3%-66.6%) [n=11]	27.3% (0.0%-51.5%) [n=10]	NS
Cefotaxime	57.1% (25.0%-66.0%) [n=23]	35.4% (18.4%-64.1%) [n=22]	NS
Ceftriaxone	40.0% (21.4%-60.0%) [n=11]	33.0% (9.0%-47.9%) [n=9]	NS
Cotrimoxazole	57.7% (30.0%-72.7%) [n=19]	69.9% (60.6%-87.3%) [n=16]	75.0% (12.5%-100.0%) [n=4]
Ciprofloxacin	40.0% (25.0%-59.0%) [n=39]	38.9% (16.7%-53.6%) [n=31]	50.0% (0.0%-64.4%) [n=10]
Amoxiclav	25.0% (16.0%-53.6%) [n=9]	11.1% (0.0%-40.3%) [n=11]	20.0% (0.0%-40.0%) [n=2]
Clindamycin	29.3% (15.9%-40.2%) [n=14]	27.5% (6.1%-37.1%) [n=14]	NS
Vancomycin	0.0% (0.0%-0.0%) [n=37]	0.0% (0.0%-0.0%) [n=35]	0.0% (0.0%-13.7%) [n=12]
Linezolid	0.0% (0.0%-12.5%) [n=19]	0.0% (0.0%-0.0%) [n=17]	0.0% (0.0%-5.0%) [n=6]
Teicoplanin	0.0% (0.0%-10.5%) [n=5]	0.0% (0.0%-0.0%) [n=5]	0.0% (0.0%-16.6%) [n=3]
Doxycycline	-NS	-NS	40.0% (40.0%-40.0%) [n=1]

* Data is presented in the form [median, (Inter quartile range:Q1-Q3), n=number of studies

NS : Not Studied.

Table 2: Median and IQR of percentage resistance of Gram-negative bacteria: *Klebsiella pneumoniae*, *Enterobacter spp*, *Escherichia coli*, *Acinetobacter baumannii*, *Citrobacter spp* and *Pseudomonas aeruginosa*

	<i>Klebsiella pneumoniae</i> 26%(16.7%-35.4%) [n=74]	<i>Enterobacter spp</i> 3.2%(0%-8.4%) [n =43]	<i>Escherichia coli</i> 9.3%(4.7%-14.5%) [n =72]	<i>Acinetobacter baumannii</i> 5.9%(2.1%-10.8%) [n =62]	<i>Citrobacter spp</i> 1.5%(0%-3%) [n =44]	<i>Pseudomonas aeruginosa</i> 4.8%(2.5%-9.4%) [n =65]
Ampicillin	95.2% (74.0%-100.0%) [n=47]	100.0% (94.7%-100.0%) [n=19]	92.3% (67.7%-100.0%) [n=32]	42.9% (7.0%-63.2%) [n=35]	99.0% (79.2%-100.0%) [n=10]	NS
Cephalexin	73.7% (49.5%-95.2%) [n=4]	100.0% (95.2%-100.0%) [n=3]	58.4% (58.3%-100.0%) [n=3]	NS	NS	NS
Cefazolin	76.5% (53.8%-100.0%) [n=3]	NS	NS	NS	NS	NS
Cefotaxime	63.7% (40.9%-80.5%) [n=52]	70.4% (55.2%-90.3%) [n=22]	50.0% (40.0%-66.0%) [n=37]	60.0% (42.2%-90.6%) [n=29]	60.0% (0.0%-69.1%) [n=13]	NS
Ceftriaxone	65.0% (28.7%-90.4%) [n=31]	100.0% (58.8%-100.0%) [n=11]	50.0% (33.3%-66.7%) [n=19]	68.9% (49.6%-97.1%) [n=14]	50.0% (0.0%-62.5%) [n=11]	NS
Ceftazidime	64.3% (44.4%-88.6%) [n=34]	67.0% (51.9%-97.6%) [n=13]	NS	77.8% (50.0%-100.0%) [n=19]	NS	58.7% (33.3%-73.7%) [n=26]
Amikacin	41.0% (18.8%-50.5%) [n=65]	37.9% (20.6%-66.2%) [n=26]	22.4% (0.0%-40.0%) [n=43]	NS	50.0% (18.6%-63.5%) [n=16]	40.0% (25.0%-50.0%) [n=39]
Gentamicin	75.7% (54.8%-86.8%) [n=64]	83.3% (65.8%-93.6%) [n=25]	55.6% (33.3%-83.3%) [n=43]	64.0% (45.0%-79.6%) [n=37]	50.0% (33.4%-83.3%) [n=15]	65.8% (50.0%-75.0%) [n=36]
Ciprofloxacin	46.2% (28.4%-62.0%) [n=57]	39.0% (8.3%-62.1%) [n=23]	40.0% (22.9%-67.0%) [n=37]	52.8% (32.5%-63.2%) [n=34]	33.4% (0.0%-64.8%) [n=16]	50.0% (32.5%-62.6%) [n=38]
Chloramphenicol	59.2% (9.8%-77.3%) [n=14]	91.1% (51.7%-100.0%) [n=6]	63.9% (28.8%-84.9%) [n=8]	88.9% (65.3%-100.0%) [n=9]	71.9% (37.5%-93.8%) [n=4]	NS
Amoxiclav	64.7% (33.0%-90.0%) [n=19]	NS	74.5% (34.4%-87.8%) [n=12]	NS	52.3% (12.5%-88.6%) [n=4]	NS
Piperacillin-tazobactam	42.0% (1.6%-61.4%) [n=22]	16.7% (0.0%-68.8%) [n=8]	16.7% (0.0%-40.0%) [n=11]	22.3% (0.0%-53.2%) [n=17]	0.0% (0.0%-37.5%) [n=4]	21.0% (1.2%-33.3%) [n=17]
Cefoperazone-sulbactam	23.6% (9.5%-53.9%) [n=12]	23.8% (1.5%-85.4%) [n=4]	NS	20.0% (9.4%-23.2%) [n=5]	1.0% (0.0%-1.9%) [n=2]	NS
Cotrimoxazole	62.5% (50.1%-80.7%) [n=25]	NS	63.2% (50.0%-78.3%) [n=17]	NS	33.4% (0.0%-50.0%) [n=7]	NS
Aztreonam	71.9% (19.0%-84.1%) [n=9]	100.0% (50.0%-100.0%) [n=3]	30.9% (2.9%-57.5%) [n=4]	22.2% (0.0%-86.1%) [n=4]	0.0% (0.0%-100.0%) [n=3]	45.0% (6.3%-83.3%) [n=5]
Meropenem	1.0% (0.0%-17.4%) [n=17]	16.7% (0.0%-83.4%) [n=4]	9.0% (0.0%-27.0%) [n=11]	11.5% (0.0%-47.5%) [n=12]	0.0% (0.0%-0.0%) [n=3]	16.7% (0.0%-30.0%) [n=11]
Colistin	3.8% (0.0%-21.4%) [n=3]	NS	8.8% (0.0%-17.6%) [n=2]	NS	NS	0.0% (0.0%-0.0%) [n=2]
Tigecycline	8.2% (1.0%-18.9%) [n=4]	NS	0.0% (0.0%-0.0%) [n=1]	NS	NS	NS

* Data is presented in the form [median, (Inter quartile range:Q1-Q3), n=number of studies]

NS : Not Studied

Table 3: Median and IQR of percentage resistance of *Salmonella Typhi*

<i>Salmonella typhi</i> 0%(0%-6.8%) [n=37]	
Ampicillin	50.0% (16.6%-87.1%) [n=9]
Cefotaxime	0.0% (0.0%-13.6%) [n=7]
Ceftriaxone	6.3% (0.0%-14.2%) [n=7]
Ceftazidime	0.0% (0.0%-0.0%) [n=2]
Ciprofloxacin	14.3% (0.0%-21.4%) [n=11]
Chloramphenicol	17.9% (0.9%-62.1%) [n=8]
Amoxiclav	0.0% (0.0%-14.3%) [n=3]
Cotrimoxazole	12.2% (0.0%-45.1%) [n=8]
Azithromycin	16.0% (0.0%-31.2%) [n=3]
Nalidixic acid	35.5% (0.0%-81.3%) [n=5]

* Data is presented in the form [median,(Inter quartile range:Q1-Q3), n=number of samples]

NS : Not Studied

Table 4: Age based pathogen resistance in common drug combinations

Pathogen- Antibiotic combination	Neonate [Median(Q1-Q3) number of samples]	Pediatric [Median(Q1-Q3) number of samples]	P ^s
<i>Staphylococcus aureus</i> - Cloxacillin	53.6% (39.8%-66.2%) [#22]	33.3% (29.3%-64.3%) [#11]	0.08
<i>Staphylococcus aureus</i> - Clindamycin	33.0% (20.0%-45.7%) [#11]	27.7% (0.0%-30.0%) [#3]	ns
<i>Staphylococcus aureus</i> - Vancomycin	0.0% (0.0%-0.0%) [#30]	0.0% (0.0%-0.0%) [#7]	ns
<i>Staphylococcus Aureus</i> - Linezolid	0.0% (0.0%-12.5%) [#14]	0.0% (0.0%-13.2%) [#5]	ns
<i>Klebsiella pneumoniae</i> - Ampicillin	95.9% (76.2%-100.0%) [#42]	93.9% (44.6%-100.0%) [#5]	ns
<i>Klebsiella</i> - Gentamicin	75.0% (54.8%-86.2%) [#56]	83.6% (31.3%-98.5%) [#8]	ns
<i>Klebsiella pneumoniae</i> - Cefotaxime	62.6% (42.8%-80.2%) [#46]	70.0% (14.3%-89.0%) [#7]	ns
<i>Klebsiella pneumoniae</i> - Piperacillin- tazobactam	42.0% (5.1%-62.1%) [#20]	25.0% (0.0%-50.0%) [#2]	ns
<i>Klebsiella pneumoniae</i> - Imipenem	0.0% (0.0%-8.0%) [#27]	0.0% (0.0%-0.0%) [#5]	ns
<i>Escherichia coli</i> - Ampicillin	92.9% (66.7%-100.0%) [#27]	83.3% (62.0%-100.0%) [#5]	ns
<i>Escherichia coli</i> - Gentamicin	55.6% (33.3%-83.3%) [#38]	50.0% (0.0%-79.4%) [#5]	ns
<i>Escherichia coli</i> - Amikacin	22.3% (2.3%-40.0%) [#38]	38.8% (0.0%-50.0%) [#5]	ns
<i>Escherichia coli</i> - Cefotaxime	47.5% (40.0%-66.3%) [#32]	50.0% (45.0%-74.3%) [#5]	ns
<i>Pseudomonas</i> - Amikacin	39.4% (23.5%-50.0%) [#32]	40.0% (25.0%-66.3%) [#7]	ns
<i>Pseudomonas</i> - Ceftazidime	50.0% (33.3%-73.3%) [#19]	67.0% (33.3%-75.0%) [#7]	ns
<i>Pseudomonas aeruginosa</i> - Ciprofloxacin	43.0% (30.0%-60.0%) [#31]	63.0% (40.0%-80.0%) [#7]	ns
<i>Enterobacter spp</i> -	100.0% (97.4%-100.0%)	97.1% (94.2%-0.0%)	ns

Ampicillin	[#17]	[#2]	
<i>Enterobacter spp-</i> Gentamicin	88.0% (61.7%-97.4%)	76.5% (67.6%-83.0%)	ns
<i>Citrobacter spp-</i> Ampicillin	95.3% (62.5%-100.0%)	99.0% (87.0%-100.0%)	ns
<i>Citrobacter spp-</i> Gentamicin	52.8% (39.8%-80.3%)	0.0% (0.0%-0.0%)	ns
<i>Acinetobacter baumannii-</i> Gentamicin	63.6% (45.0%-78.9%)	77.1% (18.8%-94.8%)	ns

* Data is presented in the form [Median,(Inter quartile range:Q1-Q3), #number of samples]

§ : significance levels using Mann-Whitney test ; ns – not significant

For Review Only




Figure 1: Flow Chart of selection of studies based on PRISMA 2009 flow chart

eAppendix 1

Search strings used

For Scopus

ALL ("India" AND "Children" OR "Pediatrics" OR "Neonates" OR "Infants" AND
"Antimicrobial resistance" OR "Antibiotic Resistance" OR "Antibiotic Susceptibility" OR
"Antibiotic sensitivity" AND "Blood Culture") AND SUBJAREA (mult OR medi OR
nurs OR vete OR dent OR heal) AND PUBYEAR > 1999 AND PUBYEAR < 2015
AND (EXCLUDE (DOCTYPE , "re") OR EXCLUDE (DOCTYPE , "le") OR
EXCLUDE (DOCTYPE , "ch") OR EXCLUDE (DOCTYPE , "no")) AND (
EXCLUDE (SUBJAREA , "AGRI") OR EXCLUDE (SUBJAREA , "ENVI") OR
EXCLUDE (SUBJAREA , "DENT") OR EXCLUDE (SUBJAREA , "HEAL") OR
EXCLUDE (SUBJAREA , "NURS") OR EXCLUDE (SUBJAREA , "VETE")) AND (
EXCLUDE (LANGUAGE , "French") OR EXCLUDE (LANGUAGE , "Turkish") OR
EXCLUDE (LANGUAGE , "Portuguese") OR EXCLUDE (LANGUAGE , "Spanish"))
AND (EXCLUDE (SRCTYPE , "b") OR EXCLUDE (SRCTYPE , "k")) AND (
LIMIT-TO (AFFILCOUNTRY , "India"))

For pubmed

"India"[All Fields] AND "Children"[All Fields] OR "Pediatrics"[All Fields] OR
"Neonates"[All Fields] OR "Infants"[All Fields] AND "Antimicrobial resistance"[All Fields]
OR "Antibiotic Resistance"[All Fields] OR "Antibiotic Susceptibility"[All Fields] OR
"Antibiotic sensitivity"[All Fields] AND "Blood Culture"[All Fields] AND
("2000/01/01"[PDAT] : "2015/07/01"[PDAT])

eAppendix 2

Included studies

1. Agarwal, A. and S.Bhat, (2015) Clinico-microbiological study of neonatal sepsis. *Journal of International Medicine and Dentistry*, 2(1), pp.22–29.
2. Agnihotri, N., Kaistha, N. and V.Gupta (2004) Antimicrobial susceptibility of isolates from neonatal septicemia. *Japanese Journal of Infectious Diseases*, 57, pp.273–275.
3. Ambade, V.C. et al (2015) Bacteriological and Haematological Profile of Neonatal Sepsis in Rural Teaching Hospital in Central India Authors. *Journal of Medical Science And clinical Research*, 3(2), pp.4358–4364.
4. Bagla, J. et al (2013) Antimicrobial susceptibility patterns following change in antibiotic policy in NICU. *Pediatric Infectious Disease*, 5(2), pp.59–63.
5. Bhat, D.R. and L.P. Baby (2011) Early Onset of Neonatal Sepsis : Analysis of the Risk Factors and the Bacterial Isolates by Using the BacT Alert System. *Journal of Clinical and Diagnostic Research*, 5(7), pp.1385–1388.
6. Bhat, S. and S.Rao, S (2011) Bacteriology of neonatal septicemia. *Indian Journal of Research and Reports in Medical Sciences*, 1(1), pp.18–21.
7. Bhat, Y. Lewis, L.E. and K.Vandana (2011) Bacterial isolates of early-onset neonatal sepsis and their antibiotic susceptibility pattern between 1998 and 2004: an audit from a center in India. *Italian journal of pediatrics*, 37, p.32.
8. Bhatt, K. et al (2012) Bacteriological Profile And Antibigram Of Neonatal Septicemia. *National Journal of Community Medicine*, 3(2), pp.238–241.
9. Bhattacharjee, A. et. al. (2008) Increased prevalence of extended spectrum β lactamase producers in neonatal septicemic cases at a tertiary referral hospital. *Indian Journal of Medical Microbiology*, 26(4), pp.356–360.

- 1
2
3 10. Chandel, D.S. et al(2011) Extended-spectrum β -lactamase-producing Gram-
4 negative bacteria causing neonatal sepsis in India in rural and urban settings.
5
6 *Journal of Medical Microbiology*, 60(4), pp.500–507.
7
8
9
10 11. Chandrashekar et. al. (2013) Study Of Clinical And Laboratory Profile Of Enteric
11 Fever In Pediatric Age Group. *International Journal of Basic and Applied*
12 *Medical Sciences*, 3(3), pp.16–23.
13
14
15 12. Chattopadhyay, T.K. and H.J. Raj (2012) Variations in bacterial pathogens
16 causing early onset neonatal septicaemia, according to birth weights - a 5 year
17 study in a referral hospital of West- Bengal. *Bangladesh Journal of Medical*
18 *Microbiology*, 4(2), pp.9–12.
19
20
21 13. Chelliah, A. et al. (2014) Isolation of MRSA, ESBL and AmpC - β -lactamases
22 from Neonatal Sepsis at a Tertiary Care Hospital. *Journal of clinical and*
23 *diagnostic research : JCDR*, 8(6), pp.DC24–DC27
24
25
26 14. Chopra, S., Kaur, J. and G.Mahajan (2013) Blood culture isolates in paediatric
27 patients and antimicrobial resistance patterns. *Indian Journal for The Practicing*
28 *Doctor*, 8(1), pp.7–14.
29
30
31 15. Dangre-mudey, G., Tankhiwale, N. and A.Mude (2013) Clinico-bacteriological
32 Study of Neonatal Septicaemia in Tertiary Care Rural Hospital of Wardha
33 District. *The Journal of Datta Meghe Institute of Medical Sciences University*,
34 8(3), pp.164–168.
35
36
37 16. Desai, K. and Malek, S (2010) Neonatal Septicemia: Bacterial Isolates and Their
38 Antibiotics Susceptibility Patterns. *National Journal of Integrated Research in*
39 *Medicine*, 1(3), pp.12–15.
40
41
42 17. Dhanalakshmi, V and E.S. Sivakumar (2015) Comparative Study in Early
43 Neonates with Septicemia by Blood Culture , Staining Techniques and C –
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 Reactive Protein (CRP). *Journal of Clinical and Diagnostic Research*, 9(3),
4 pp.12–15.
5
6
7
8 18. Dias, E. and Vigneshwaran, P.(2010) The Bacterial Profile Of Neonatal
9 Septicaemia In A Rural Hospital In South India. *Journal of Clinical and*
10 *Diagnostic Research*, (4), pp.3327–3330.
11
12
13
14 19. Dutta,P.,et.al (2001) Ceftriaxone therapy in ciprofloxacin treatment failure
15 typhoid fever in children.*Indian J Med Res* 113:210-3.
16
17
18
19 20. Gajul, S. et al (2015) Klebsiella pneumoniae in septicemic neonates with special
20 reference to extended spectrum β -lactamase, AmpC, metallo β -lactamase
21 production and multiple drug resistance in tertiary care hospital. *Journal of*
22 *Laboratory Physicians*, 7(1), pp.32–37.
23
24
25
26
27 21. Gandhi, S. et al (2013) Incidence of neonatal sepsis in tertiary care hospital: An
28 overview. *International Journal of Medical Science and Public Health*, 2(3),
29 p.522.
30
31
32
33
34 22. Jain, A. et al(2011) Intravenous device associated blood stream staphylococcal
35 infection in paediatric patients. *Indian Journal of Medical Research*, 134(8),
36 pp.193–199.
37
38
39
40
41 23. Jain, A. et al (2003) Prevalence of extended-spectrum beta-lactamase-producing
42 Gram-negative bacteria in septicemic neonates in a tertiary care hospital. *Journal*
43 *of medical microbiology*, 52, pp.421–425.
44
45
46
47 24. Jain, A., Agarwal, J. and S.Bansal (2004) Prevalence of methicillin-resistant,
48 coagulase-negative staphylococci in neonatal intensive care units: Findings from a
49 tertiary care hospital in India. *Journal of Medical Microbiology*, 53, pp.941–944.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 25. Jain, A. and R.Mondal (2008) TEM & SHV genes in extended spectrum beta-
4 lactamase producing Klebsiella species beta their antimicrobial resistance pattern.
5
6 *The Indian journal of medical research*, 128(6), pp.759–764.
7
8
9
10 26. Javali, N., Banu, N. and S Indi (2014) Incidence and Microbiological Profile of
11 Late Onset Neonatal Sepsis in Preterm and Low Birth Weight Neonates. *Journal*
12 *of Evidence based Medicine and Healthcare*, 1(16), pp.2036–2048.
13
14
15 27. Jeyamurugan, T.,et al (2012)Acinetobacter spp.: An Emerging Pathogen in
16 Neonatal Septicaemia.*J Clin Diagn Res* 6(5):805
17
18
19 28. Jonnala, R.N.R. et al(2013) Clinical And Laboratory Profile Of Neonatal Sepsis.
20 *International Journal of Medical and Applied Sciences*, 2(4), pp.319–331.
21
22
23 29. Kaistha, N. et al (2010) Neonatal septicemia isolates and resistance patterns in a
24 tertiary care hospital of North India. *Journal of Infection in Developing Countries*,
25 4(1), pp.055–057.
26
27
28
29
30 30. Kamble, R. and Ovhal, R (2015) Bacteriological profile of neonatal septicemia.
31 *International Journal of Current Microbiology and Applied Sciences*, 4(2),
32 pp.172–182.
33
34
35
36
37 31. Kapoor, L., Randhawa, V.S. and M.Deb (2005)Antimicrobial resistance of
38 enterococcal blood isolates at a pediatric care hospital in India. *Japanese Journal*
39 *of Infectious Diseases*, 58, pp.101–103.
40
41
42
43
44 32. Karthikeyan, G. and K Premkumar (2001) Neonatal sepsis: Staphylococcus aureus
45 as the predominant pathogen. *Indian journal of pediatrics*, 68(8), pp.715–717.
46
47
48
49 33. Katiyar, R. and S. Bose, (2012) Bacteriological Profile of Neonatal Septicemia in
50 Pravara Rural Hospital. *Pravara Medical Review*, (4(2)).
51
52
53
54 34. Khan, S.N. and S.Joseph (2012) Neonatal Sepsis : Antibiotic Sensitivity &
55 Resistance Pattern Of Commonly Isolated Pathogens In A Neonatal Intensive
56
57
58
59
60

- Care Unit Of A Tertiary Care Hospital , South India. *International Journal of Pharma and Bio Sciences*, 3(4), pp.802–809.
35. Kulkarni, S.R.A.O. et al. (2014) Aerobic Bacteriological Profile Of Neonatal Septicaemia And Study Of Esbl Production In Gram Negative Isolates . *International Journal of Advanced Research*, 2(11), pp.1012–1021.
36. Kumar, A. et al.(2014) Risk factors for carbapenem-resistant *Acinetobacter baumannii* blood stream infections in a neonatal intensive care unit, Delhi, India. *The Journal of Infection in Developing Countries*, 8(08), pp.1049–1054. Available at: <http://www.jidc.org/index.php/journal/article/view/4248>.
37. Kumar, R., Gupta, N. and Shalini (2007) Multidrug — resistant typhoid fever. *The Indian Journal of Pediatrics*, 74(1), pp.39–42.
38. Kumhar, G.D., Ramachandran, V.G. and P.Gupta, (2002) Bacteriological analysis of blood culture isolates from neonates in a tertiary care hospital in India. *Journal of Health Population and Nutrition*, 20(4), pp.343–347.
39. Madavi, D., Aziz, F. and G. Agrawal (2015) Clinico-bacteriological profile and antibiotic sensitivity pattern of neonatal septicaemia- a prospective observational study. *International Journal of Current Research and Review*, 7(5), pp.13–20.
40. Manchanda, V. et al.(2006) Treatment of enteric fever in children on the basis of current trends of antimicrobial susceptibility of *Salmonella enterica* serovar typhi and paratyphi A. *Indian journal of medical microbiology*, 24(2), pp.101–106.
41. Mathur, N.B., Marmu, K. and S. Kumar, (2010) Systemic inflammatory response syndrome in home delivered neonates: A prospective observational study. *Indian Journal of Pediatrics*, 77(10), pp.1109–1113.
42. Marwah,P. et al (2015) Bacteriological profile of neonatal sepsis in a tertiary-care hospital of northern India. *Indian Pediatr*,52:158-159

- 1
2
3 43. Mehar, V. et al.(2013) Neonatal sepsis in a tertiary care center in central India:
4
5 Microbiological profile, antimicrobial sensitivity pattern and outcome. *Journal of*
6
7 *Neonatal-Perinatal Medicine*, 6(2), pp.165–172.
8
9
10 44. Mehta,A.M.,Kaore, N.M. and T.K. Prabhu (2014) Microbial Profile of Neonatal
11
12 septicaemia in a tertiary care hospital of Bhopal *IJBAR* 05(10):459-501
13
14 45. Mehrotra, A. and Mishra, S. (2014) Determination of the Causative Agents of
15
16 Bacteremia in Children under 5 Years and their Susceptibility Pattern to the
17
18 Antibiotics. *Global Journal of Medical Research: C Microbiology and Pathology*,
19
20 14(1), pp.19–24.
21
22
23 46. Modi, J. et al. (2012) A Study Of Extended Spectrum B -Lactamase (Esbl) And
24
25 Ampc B -Lactamase Producing Klebsiella Pneumoniae In Neonatal Intensive Care
26
27 Unit At Tertiary Care Hospital , Ahmedabad. *National Journal of Community*
28
29 *Medicine*, 3(3), pp.523–528.
30
31
32 47. Monica, K.H.S., Devi, K.H.K. and K.H.S. Devi, (2014) Antibigram Of
33
34 Salmonella Typhi Isolated From Enteric Fever Cases In A Tertiary Health Care
35
36 Centre In Imphal. *International Journal Of Pharmacology And Therapeutics*, 4(1),
37
38 pp.15–18.
39
40
41 48. Muley, V., Ghadage, D. and A.Bhore, (2015) Bacteriological profile of neonatal
42
43 septicemia in a tertiary care hospital from Western India. *Journal of Global*
44
45 *Infectious Diseases*, 7(2), pp.75–77.
46
47
48 49. Mundkur, S. (2011) Coagulase Negative Staphylococcus (CONS) Infection in
49
50 Neonatal ICU . *Webmed Central PAEDIATRICALS*, 2(12), pp.1–12. Available at:
51
52 https://www.webmedcentral.com/wmcpdf/Article_WMC002829.pdf.
53
54
55
56
57
58
59
60

- 1
2
3 50. Mustafa, M. and S.Ahmed, (2014) Bacteriological profile and antibiotic
4 susceptibility patterns in neonatal septicemia in view of emerging drug resistance.
5
6 *Journal of Medical and Allied Sciences*, 4(1), pp.2–8.
7
8
9
10 51. Nayak, S. et al.(2014) Distribution of microorganisms in neonatal sepsis and
11 antimicrobial susceptibility patterns in a tertiary care hospital. *Archives of*
12 *Medicine and Health Sciences*, 2(2), p.136. Available at:
13
14 <http://www.amhsjournal.org/text.asp?2014/2/2/136/144304>.
15
16
17
18 52. Pais, M. et al. (2011) A Descriptive Study On The Neonatal Sepsis, Bacterial
19 Isolates And Antibiotic Susceptibility Patterns Among Neonates. *International*
20 *Journal of Community Pharmacy*, 4(2), pp.19–24.
21
22
23
24
25 53. Pal, K. and A.K.Samanta(2013). Bacteriological Profile and Antimicrobial
26 Resistance Pattern in Early Onset Neonatal Sepsis : *Int J Cur Res Rev*, 05(06),
27 pp.51–58.
28
29
30
31
32 54. Patel, D. et al. (2014) Blood culture isolates in neonatal sepsis and their sensitivity
33 in Anand District of India. *Indian Journal of Pediatrics*, 81(8), pp.785–790.
34
35
36
37 55. Ponugoti, M. et al. (2015) Incidence of Early Onset Septicemia, Isolation and
38 Resistant Patterns of Causative Organisms: A Study in a Tertiary Care Hospital
39 *A.P. Journal of Medical Science and Clinical Research*, 3(2), pp.4227–4232.
40
41
42
43 56. Prabhu, K., Bhat, S. and S. Rao(2010) Bacteriologic Profile and Antibigram of
44 Blood Culture Isolates in a Pediatric Care Unit. *J Lab Physicians.*, 2(2), pp.85–88.
45
46
47
48 57. Rajendraprasad, B.P.M., Basavaraj, K. and B.Antony (2013) Bacterial spectrum of
49 neonatal septicemia with their antibiogram with reference to various predisposing
50 factors in a tertiary care hospital in Southern India. *Annals of Tropical Medicine*
51 *and Public Health*, 6(1), pp.96–99. Available at:
52
53
54 <http://www.atmph.org/text.asp?2013/6/1/96/115204>.
55
56
57
58
59
60

- 1
2
3 58. Rao, Y. et al. (2012) Neonatal Septicemia In North India Due To Extended
4
5 Spectrum Beta Lactamase (Esbl) Producing Gram Negative Bacteria.
6
7 *International Journal of Pharma and Bio Sciences*, 3(1), pp.282–290.
8
9
10 59. Roy, I. et al. (2002) Bacteriology of neonatal septicaemia in a tertiary care hospital
11
12 of northern India. *Indian journal of medical microbiology*, 20(3), pp.156–159.
13
14 60. Saravanan, M., Nanda, A. and Tesfaye, T. (2013) Antibiotic Susceptibility Pattern
15
16 of Methicillin Resistant Staphylococcus from Septicemia Suspected Children in
17
18 Tertiary Hospital in Hosur, South India. *American Journal of Microbiological*
19
20 *Research*, 1(2), pp.21–24. Available at:
21
22 <http://pubs.sciepub.com/ajmr/1/2/2/index.html>.
23
24
25 61. Saxena, S. et al. (2013) Detection Of Methicillin-Resistant Staphylococcus
26
27 Species By Multiplex Pcr And Use Of The D Test Method To Detect Inducible
28
29 Clindamycin Resistance. *International Journal of Recent Scientific Research*,
30
31 4(3), pp.231–234.
32
33
34 62. Shah, M.N. and P.B.Desai (2011) Clinical and bacteriological profiles of blood
35
36 culture positive sepsis in newborns. *International Journal Of Pharmacy & Life*
37
38 *Sciences*, 2(9), pp.1041–1045.
39
40
41 63. Sharma CM, et al.(2013)“Neonatal Sepsis”: Bacteria & their Susceptibility Pattern
42
43 towards Antibiotics in Neonatal Intensive Care Unit. *J Clin Diagn Res*.7(11):2511–
44
45 2513.
46
47
48 64. Sharma, M. et al. (2002) Bacteraemia in children. *Indian journal of pediatrics*, 69,
49
50 pp.1029–1032.
51
52
53 65. Sharma, M., Yadav, S. and U. Chaudhary, (2009) Acinetobacter spp as emerging
54
55 pathogen in neonatal septicemia. *Indian Journal of Pediatrics*, 76(8), p.861.
56
57
58
59
60

- 1
2
3 66. Sharma, M., Yadav, S. and U. Chaudhary, (2010) Metallo-beta-lactamase
4 Producing *Pseudomonas aeruginosa* in Neonatal Septicemia. *Journal of*
5 *laboratory physicians*, 2(1), pp.14–16.
6
7
8
9
10 67. Sharma, P., Kaur, P. and A. Aggarwal, (2013) *Staphylococcus aureus*- the
11 predominant pathogen in the neonatal ICU of a tertiary care hospital in Amritsar,
12 India. *Journal of Clinical and Diagnostic Research*, 7(1), pp.66–69.
13
14
15
16 68. Singhi, S. et al. (2008) Nosocomial bloodstream infection in a pediatric intensive
17 care unit. *Indian Journal of Pediatrics*, 75(1), pp.25–30.
18
19
20
21 69. Sood, M., Sharma, S. and Grover, N. (2012) *Staphylococcus aureus* infection:
22 Clinical profile and outcome among children admitted to a tertiary care hospital in
23 developing economy. *Pediatric Infectious Disease*, 4(3), pp.101–106.
24
25
26
27 70. Tallur, S.S. et al. (2000) Clinico-bacteriological study of neonatal septicemia in
28 Hubli. *Indian journal of pediatrics*.
29
30
31
32 71. Tamboli, S. and Nilekar, S. (2011) Neonatal Septicemia Predominant Bacterial
33 Species and Antibiotic Resistance. *Indian Medical Gazette*, (November), pp.421–
34 424.
35
36
37
38 72. Tankhiwale, S. and Agrawal, S. (2013) Bacteriological profile and antibiotic
39 sensitivity pattern of isolates of neonatal septicemia. *Journal of Evolution of*
40 *Medical and Dental Sciences*, 2(31), pp.5021–5029.
41
42
43
44
45 73. Tiwari, D.K. et al. (2013) A study on the bacteriological profile and antibiogram
46 of bacteremia in children below 10 years in a tertiary care hospital in Bangalore,
47 India. *Journal of Clinical and Diagnostic Research*, 7(12), pp.2732–2735.
48
49
50
51 74. Tsering, D.C. et al. (2011) Bacteriological profile of septicemia and the risk
52 factors in neonates and infants in sikkim. *Journal of global infectious diseases*.
53
54
55
56
57
58
59
60

- 1
2
3 75. Tudu, N.K. et al. (2014) a Pilot Study on Bacterial Profile of Neonatal Sepsis in a
4 Tertiary Care Hospital Serving Rural Population. *Journal of Evolution of Medical*
5 *and Dental Sciences*, 3(23), pp.6378–6381.
6
7
8
9
10 76. Venkatnarayan, K., Bej, P. and Thapar, R.,(2014) Neonatal Sepsis : A Profile of a
11 Changing Spectrum. *J. Nepal Paediatr. Soc.*, 34(3), pp.207–214.
12
13
14 77. Vinodkumar, C. et al. (2011) Change in spectrum of microbial aetiology in
15 relation to gestational age and birth weight and emergence of ESBL in tertiary
16 neonatal intensive care units . *International Journal of Biological & Medical*
17 *Research*, 2(3), pp.727–734.
18
19
20
21 78. Vinodkumar, C. et al. (2008) Perinatal risk factors and microbial profile of
22 neonatal septicemia : A multicentred study. *J Obstet Gynecol India*, 58(1), pp.32–
23 40.
24
25
26
27
28
29 79. Viswanathan, R. et al. (2011) Aetiology and antimicrobial resistance of neonatal
30 sepsis at a tertiary care centre in eastern India: A 3 year study. *Indian Journal of*
31 *Pediatrics*, 78(4), pp.409–412.
32
33
34
35
36 80. Viswanathan, R. et al.(2012) Multi-drug resistant gram negative bacilli causing
37 early neonatal sepsis in India. *Archives of Disease in Childhood - Fetal and*
38 *Neonatal Edition*, 97, pp.F182–F187.
39
40
41
42
43 81. Walia, M. et al. (2006) Age-related clinical and microbiological characteristics of
44 enteric fever in India. *Transactions of the Royal Society of Tropical Medicine and*
45 *Hygiene*, 100(10), pp.942–948.
46
47
48
49 82. Zakariya, B. et al. (2012) Risk factors and predictors of mortality in culture
50 proven neonatal sepsis. *Indian Journal of Pediatrics*, 79(3), pp.358–361.
51
52
53
54
55
56
57
58
59
60

Author's Response to Decision Letter for (JPIDS-2016-098.R1)**Extremely high rates of antimicrobial resistance in Indian neonates and children are now a major threat to public health.**

Grateful for your input.

1. The abstract has been structured as suggested.
2. The phrasing of the results changed as advised.
3. Screening and statistical methods updated.
4. Suggested minor edits carried out.
5. Discussion curtailed by removing the repetitions and focusing on trends.
6. Figure 2 deleted as advised

Or Review Only