

How does dynamical downscaling affect model biases and future projections of explosive extratropical cyclones along North America's Atlantic coast?

Article

Accepted Version

Seiler, C., Zwiers, F. W., Hodges, K. I. and Scinocca, J. F. (2018) How does dynamical downscaling affect model biases and future projections of explosive extratropical cyclones along North America's Atlantic coast? *Climate Dynamics*, 50 (1-2). pp. 677-692. ISSN 0930-7575 doi: <https://doi.org/10.1007/s00382-017-3634-9> Available at <http://centaur.reading.ac.uk/70253/>

It is advisable to refer to the publisher's version if you intend to cite from the work.

Published version at: <https://link.springer.com/article/10.1007%2Fs00382-017-3634-9>

To link to this article DOI: <http://dx.doi.org/10.1007/s00382-017-3634-9>

Publisher: Springer

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **How does dynamical downscaling affect model biases and future**
2 **projections of explosive extratropical cyclones along North America's**
3 **Atlantic coast?**

4 **C. Seiler · F.W. Zwiers · K.I. Hodges · J.F. Scinocca**

5 Received: October 14, 2016

6 **Abstract** Explosive extratropical cyclones (EETCs) are rapidly intensifying low pressure systems that generate
7 severe weather along North America's Atlantic coast. Global climate models (GCMs) tend to simulate too few
8 EETCs, which may partly be related to their coarse horizontal resolution and poorly resolved moist diabatic
9 processes. This study explores whether EETC frequency biases can be reduced through dynamical downscaling,
10 and whether this affects climate change projections of storms along North America's Atlantic coast. A regional
11 climate model (CanRCM4) is forced with the CanESM2 GCM for the periods 1981 to 2000 and 2081 to 2100.
12 EETCs are tracked from relative vorticity using an objective feature tracking algorithm. CanESM2 simulates
13 38% fewer EETC tracks compared to reanalysis data, which is consistent with a negative Eady growth rate bias
14 (-0.1 day^{-1}). Downscaling CanESM2 with CanRCM4 increases EETC frequency by one third, which reduces the
15 frequency bias to -22%, and coincides with a 22% increase in maximum EETC precipitation. Anthropogenic

C. Seiler

Pacific Climate Impacts Consortium, University of Victoria, Victoria, BC, Canada, University House 1, PO Box 1700 Stn
CSC, University of Victoria, Victoria, BC, Canada V8W 2Y2

E-mail: cseiler@uvic.ca

F.W. Zwiers

Pacific Climate Impact Consortium, University of Victoria, Victoria, BC, Canada

K.I. Hodges

University of Reading, Reading, United Kingdom

J.F. Scinocca

Canadian Centre for Climate Modelling and Analysis, Victoria, BC, Canada

greenhouse gas forcing is projected to decrease EETC frequency (-15%, -18%) and Eady growth rate (-0.2 day^{-1} , -0.2 day^{-1}), and to increase maximum EETC precipitation (46%, 52%) in CanESM2 and CanRCM4, respectively. The limited effect of dynamical downscaling on EETC frequency projections is consistent with the lacking impact on the maximum Eady growth rate. To conclude, the coarse spatial resolution of GCMs presents an important limitation for simulating extreme ETCs, but biases in the Eady growth rate are likely to be just as relevant.

Keywords Explosive extratropical cyclones · Dynamical downscaling · Model biases · Climate change projections

1 Introduction

Extratropical cyclones (ETCs) affect the general circulation through the exchange of heat, moisture, and momentum, but also impact human activity through the generation of strong surface winds, high waves, extreme precipitation, and associated hazardous conditions. Many of the most violent winter storms along North America's Atlantic coast undergo rapid intensification, with deepening rates typical of "weather bombs" (Sanders and Gyakum, 1980). These explosive ETCs (EETCs) have caused fatalities and billions of \$USD property damage (Kocin et al., 1995), and remain difficult to forecast (Edward B Carroll, personal communication, July 21, 2015). Given the severe weather conditions associated with EETCs, it is of great public interest to better understand how these events may evolve under a warming climate.

Global climate models (GCMs) reproduce the climatology of observed ETC frequency and intensity reasonably well, with a tendency to slightly underestimate both variables and to simulate tracks that are too zonal (Lambert and Fyfe, 2006; Ulbrich et al., 2008; Zappa et al., 2013a). GCMs tend to project an overall decline in ETC frequency as a response to anthropogenic greenhouse gas (GHG) forcing, with a weak polar shift in the Northern Pacific, and a downstream extension of the Atlantic storm track into Europe (Bengtsson et al., 2006; McDonald, 2011; Chang et al., 2012; Collins et al., 2013; Christensen et al., 2013). The latter implies an increase in storm frequency close to the British Isles, and a decrease in storm frequency in the Norwegian and Mediterranean Seas and subtropical central Atlantic (Zappa et al., 2013b). The precipitation rate associated with ETCs is projected to increase along North America's Atlantic coast (Zappa et al., 2013b). GCM performance in simulating EETCs is less impressive; they tend to underestimate EETC frequency by up to two thirds (Seiler and Zwiers, 2016a). The same models project a reduction in EETC frequency in the Northern Atlantic by 17% on average by the end of this century (Seiler and Zwiers, 2016b).

44 Anthropogenic GHG forcing is expected to affect ETCs through multiple competing processes. This includes
45 (i) changes in baroclinicity in the lower and upper troposphere due to polar and tropical amplification, and (ii)
46 an increase in the atmospheric water vapor content that may intensify ETCs through enhanced latent heating
47 (Hall et al., 1994). The latter process has been shown to be sensitive to the horizontal model resolution (Willison
48 et al., 2013). Given the modest spatial resolution of most current GCMs, it is natural to ask if model biases
49 could be reduced by increasing the horizontal model resolution, and whether this would affect the corresponding
50 projections.

51 The impact of horizontal model resolution on ETC biases and projections can be examined in dynamical
52 downscaling experiments, where a regional climate model (RCM) is forced with lateral boundary conditions from
53 a GCM or reanalysis data. Previous studies show that dynamical downscaling has the potential to significantly
54 affect model biases and projections of ETCs. Willison et al. (2013) forced the Weather Research and Forecasting
55 model (WRF) with the NCEP global forecast system (GFS) final analysis, and compared results where WRF was
56 run at lower (120 km) and higher horizontal resolutions (20 km) for a domain covering the North Atlantic. Using
57 an Eulerian approach, the study showed enhancement of the positive feedback between ETC intensification and
58 latent heat release at the higher resolution, resulting in a systematic increase in eddy intensity and a stronger storm
59 track. These results are consistent with findings from Long et al. (2009), who downscaled the Canadian Climate
60 Centre model (CGCM2) with the Canadian Regional Climate Model (CRCM version 3.5) for the Northwest
61 Atlantic and eastern North America. Using a Lagrangian approach in which storms were tracked from 6h mean
62 sea level pressure (MSLP) minima, dynamical downscaling was found to increase the frequency of strong ETCs
63 with $\text{MSLP} < 995$ hPa along North America's Atlantic coast. However, it remains unclear whether this is a
64 response of the ETC or of the background MSLP field. Contrary findings are presented in Colle et al. (2015)
65 who downscaled reanalysis data (NCEP-CFSR) with six different RCMs for North America's Atlantic coast and
66 detected a 5-10% decrease in ETCs tracked from 850 hPa relative vorticity. Côté et al. (2015) related similar
67 findings to the presence of the regional model boundary in the vicinity of the storm track.

68 Downscaling may also significantly affect climate change projections of ETC frequencies and intensities. Willi-
69 son et al. (2015) conducted a pseudo-global warming experiment where projected changes in temperature from
70 five GCMs were applied to reanalysis data, which was then used as the initial and lateral boundary conditions in a
71 regional climate model experiment for a selection of days with storms. The study finds that changes in temperature
72 increases storm activity in the northeastern Atlantic, and that this is further enhanced when increasing the model

73 resolution from a 120-km to a 20-km grid spacing. Long et al. (2009) on the other hand finds that downscaling
74 enhances the reduction in ETC frequency projected by a global model (CGCM2) in the Northwest Atlantic and
75 eastern North America by 10%. Such contrary findings may result from differences in the experimental design and
76 methods employed. Also, it remains unclear to what degree the impacts of dynamical downscaling are affected by
77 differences in physical parameterizations used in the global models compared to the regional models.

78 The studies outlined above show that dynamical downscaling has the potential to significantly affect model
79 ETC biases and projections. The enhanced condensational heating in the high-resolution model runs may play an
80 important role in this context. Only a few studies have assessed the impacts of dynamical downscaling on ETC
81 biases and projections, and their conclusions remain ambiguous. While pseudo-global warming experiments are
82 suitable for exploring mechanisms in controlled experiments (Marciano et al., 2015), the number of tracks that
83 are simulated in such studies is too small to assess the statistical significance of the projected changes. Finally, a
84 focus on EETCs is to our knowledge still missing, despite their potential for severe impacts, and the important
85 role that latent heating can play in rapid intensification (Fink et al., 2012).

86 This study assesses how dynamical downscaling affects model biases and projections of EETCs along North
87 America’s Atlantic coast. This is of particular relevance for our previous work on EETC biases and projections of
88 a multi GCM ensemble (Seiler and Zwiers, 2016a,b). Special attention is paid to the use of different spatial filters
89 required to compute storm tracks from different horizontal grids in a consistent manner that allows intercomparison
90 of results. Section 2 presents our data and approach used for identifying and tracking EETCs in global and
91 regional climate model data from seven experiments. Section 3 assesses biases and projections of all ETCs and of
92 EETCs simulated by the Canadian Earth System Model (CanESM2) and the Canadian Regional Climate Model
93 (CanRCM4). Both models are compatible as they share the exact same physics package. The role of baroclinic
94 instability for EETC biases and projections is explored as well. Section 4 elaborates on the principal findings and
95 discusses opportunities for future research on the competing processes that determine storm biases and projections.

96 **2 Methods**

97 **2.1 Data**

98 We use global data from the ERA-Interim reanalysis (Dee et al., 2011) and the Canadian Earth System Model
99 (CanESM2, Arora et al., 2011). ERA-Interim is produced by the European Centre for Medium-Range Weather
100 Forecasts (ECMWF). The atmospheric component is a spectral model with T255 truncation and 60 vertical levels,

101 and the data are converted to a 480×240 (0.75° , 83 km) horizontal linear grid. CanESM2 is developed by the
102 Canadian Centre for Climate Modelling and Analysis (CCCma) and participated in the fifth phase of the Coupled
103 Model Intercomparison Project (CMIP5). Its atmospheric component has spectral T63 resolution with 35 vertical
104 levels with a lid near 1 hPa, and the data are converted to a 128×64 (2.8125° , 313 km) horizontal linear grid. The
105 CanESM2 simulation (r11i1p1) uses “all” forcings for the historical period ending in 2005 (greenhouse gases, other
106 anthropogenic forcings, solar and volcanic) and follows the Representative Concentration Pathway 8.5 emissions
107 scenario for 2006-2100 (RCP8.5) (Taylor et al., 2012).

108 CanESM2 is dynamically downscaled to a horizontal resolution of 0.22° (24 km) with the Canadian Regional
109 Climate Model (CanRCM4) (Scinocca et al., 2016). CanRCM4 is a hydrostatic model with a hybrid vertical
110 coordinate system and a regular latitude–longitude grid with rotated pole. Vertical model levels match those in
111 CanESM2, except that CanRCM4 has a lower model lid located at 13 hPa. Sea surface temperatures and sea ice
112 extent in CanRCM4 are prescribed by the forcing data. CanESM2 and CanRCM4 are compatible, as they share
113 the same physics package and parameter settings of physical parametrizations. The employed cloud microphysics
114 scheme of both models treat condensation as an instantaneous adjustment of the thermodynamic properties to
115 equilibrium (von Salzen et al., 2013). Spectral nudging is applied to the horizontal wind fields from the model top
116 down to 850 hPa and to temperature in the top three vertical levels of the model domain. The regional model
117 grid is similar to that of the Coordinated Regional Climate Downscaling Experiment (CORDEX) North America
118 experiment (Giorgi et al., 2009), with the difference that it extends further into the Atlantic to include storm
119 tracks that affect North America’s coastal region (NAE22 grid, hereafter) (Figure 1).

120 Variables used for ETC tracking include 6 hourly zonal and meridional wind components at 850 hPa, and
121 6h MSLP. Baroclinic instability is quantified from daily mean values of zonal and meridional wind components,
122 geopotential height, air temperature, and specific humidity at 850 hPa and 500 hPa. Finally, we also use daily
123 precipitation in order to assess the hydrological component of EETCs. Our data covers the historical period from
124 1981 to 2000 and the projected high emission scenario RCP8.5 from 2081 to 2100.

125 2.2 Cyclone tracking

126 ETCs are identified using the objective feature tracking algorithm TRACK (Hodges, 1994; Hodges et al., 1995;
127 Hodges, 1999). The algorithm computes relative vorticity from the horizontal wind velocity at 850 hPa at the
128 resolution of the input data. We spatially filter relative vorticity to T42 resolution prior to tracking in all cases

129 to ensure that results from data sources with different horizontal resolutions are comparable. This is achieved
 130 through spherical harmonic decomposition in the case of global data (Anderson et al., 2003), and by discrete
 131 cosine transform for dynamically downscaled data (Denis et al., 2002). The large scale background is removed
 132 by setting the coefficients in the spectral transform expansion for total wave-numbers < 5 to zero in the global
 133 filtering and the equivalent for the discrete cosine transform. Feature points are detected based on local extreme
 134 values. The algorithm computes tracks by determining the correspondence between feature points for adjacent
 135 time steps. Tracks are first initialized using a nearest neighbor method and then refined by minimizing a cost
 136 function, in spherical geometry, for track smoothness subject to adaptive constraints to produce the optimal set
 137 of smoothest tracks. Post-tracking filters are applied to retain only storms that have a relative vorticity greater
 138 than 10^{-5} s^{-1} , that last longer than two days, and that travel further than 1000 km.

139 MSLP values are assigned to each ETC center by searching for the closest T42 MSLP minimum to the location
 140 of the relative vorticity maxima within a great circle radius of 5° . T42 MSLP is used rather than the full resolution
 141 MSLP to ensure that results from data sources with different horizontal resolutions are comparable. Deepening
 142 rates are computed for each time step by comparing the MSLP values that belong to the same track and that
 143 are 24 h apart. EETCs are identified as ETCs with deepening rates greater than or equal to one bergeron (b)
 144 (Sanders and Gyakum, 1980):

$$b = \frac{24hPa}{24hours} \frac{\sin(\theta)}{\sin(60)}, \quad (1)$$

145 where θ is the latitude of the ETC center. The pressure drop between two cyclone centers is scaled according to
 146 the latitude of the second cyclone center. EETC maximum wind speed and maximum precipitation are computed
 147 by finding the highest T42 wind speed and precipitation within a 6° radius around each ETC center. Daily
 148 precipitation values are assigned to each six hourly ETC center for the day that the ETC center is occurring.
 149 As for MSLP, T42 values rather than full resolution values are used. The projected increase in maximum EETC
 150 precipitation is compared to the projected increase in monthly mean near surface air temperature from October
 151 to March, which is averaged over the coastal region enclosed by the polygon depicted in Figure 1.

152 2.3 Experiments

153 The impacts of dynamical downscaling on model biases and projections of EETC frequency and intensity are
 154 assessed by comparing results from seven experiments (Table 1). The first three experiments consist of ETC

155 tracks that are computed from global grids, where relative vorticity is spatially filtered to T42 resolution through
 156 spherical harmonic decomposition (SH). This includes tracks from ERA-Interim and from CanESM2 for the
 157 historical (H) period 1981-2000 (ERAH-SH and GCMH-SH, respectively), and for the future period (F) 2081-
 158 2100 under RCP8.5 forcing (GCMF-SH). The remaining four experiments consist of ETC tracks that are computed
 159 from the NAE22 grid with relative vorticity spatially filtered to T42 equivalent resolution through discrete cosine
 160 transform (DC). This applies to ERA-Interim (ERAH-DC) and the historical experiment of CanESM2 (GCMH-
 161 DC), which are spatially interpolated to the NAE22 grid by bilinear interpolation, as well as to the historical
 162 and projected simulation of CanRCM4 forced with CanESM2 (RCMH-DC and RCMF-DC, respectively). ETC
 163 statistics are computed for the North American Atlantic coast enclosed by the inner polygon depicted in Figure
 164 1.

165 The impacts of dynamical downscaling on model biases and projections are quantified by comparing runs
 166 that are based on the same grid extension and spatial filter (Table 1). This separates the effects of dynamical
 167 downscaling from the potential impacts of a limited model domain and the use of different grid types and associated
 168 spatial filtering techniques.

169 Model biases in CanESM2 are assessed by contrasting results from GCMH-SH against ERAH-SH. To gain
 170 insight into how spatial filtering affects the bias we compare results from GCMH-DC against ERAH-DC. The
 171 impact of dynamical downscaling is quantified by comparing RCMH-DC against GCMH-DC, and the effects on
 172 model biases are determined by comparing RCMH-DC against ERAH-DC. Finally, we assess whether dynamical
 173 downscaling alters climate change projections by comparing the projected changes from GCMF-SH and GCMH-
 174 SH against RCMF-DC and RCMH-DC.

175 2.4 Clausius-Clapeyron relation

176 The projected change in maximum EETC precipitation per degree warming is compared to the projected change
 177 in the saturation vapor pressure (e_s) per degree warming. The warming is quantified from the climatological
 178 mean cold season (October to March) near-surface air temperature averaged over the polygon that encloses North
 179 America's Atlantic coast (Figure 1). The saturation vapor pressure is estimated from near-surface air temperature
 180 following the Clausius-Clapeyron equation (Stull, 2000):

$$e_s = e_0 \exp \left[\frac{L_v}{R_v} \left(\frac{1}{T_0} - \frac{1}{T} \right) \right], \quad (2)$$

181 where e_0 is the saturation vapor pressure at 273.15 K (0.611 kPa), L_v is the latent heat of vaporization per unit
 182 mass ($2.5 \cdot 10^6$ J kg⁻¹), R_v is the gas constant for water vapor (461 J K⁻¹ kg⁻¹), T_0 is freezing temperature (273.15
 183 K), and T is the near-surface air temperature simulated by CanESM2 and CanRCM4.

184 2.5 Baroclinic instability

185 Changes in EETC frequency as a consequence of dynamical downscaling or anthropogenic GHG forcing are
 186 compared to the corresponding changes in baroclinic instability. Baroclinic instability is measured by the maximum
 187 Eady growth rate (σ_{BI}) with units of s⁻¹ (Hoskins and Valdes, 1990):

$$\sigma_{BI} = 0.31f \left| \frac{\partial \mathbf{v}}{\partial z} \right| N^{-1}, \quad (3)$$

188 where N is the static stability, or Brunt-Väisälä frequency (Stull, 2000):

$$N = \sqrt{\frac{g}{T_v} \left(\frac{\partial T_v}{\partial z} + \Gamma_d \right)}, \quad (4)$$

189 with

$$T_v = T (1 + 0.61r), \quad (5)$$

190 and

$$r = q/(1 - q), \quad (6)$$

191 where f is the Coriolis parameter, \mathbf{v} is the horizontal wind velocity, z is the geopotential height, g is the gravi-
 192 tational acceleration, Γ_d is the dry adiabatic lapse rate, T_v is the virtual temperature, T is the air temperature,
 193 r is the water vapor mixing ratio, and q is the specific humidity. The Eady growth rate is computed for daily
 194 values between 850 and 500 hPa, and is then averaged for each month. The partial derivatives with respect to z
 195 are approximated by computing the differences between the two pressure levels. Our analysis of the Eady growth
 196 rate considers the months October to March, as most EETCs occur during the cold season (Seiler and Zwiers,
 197 2016a). Also, we focus on North America's Atlantic coast, a major region of EETC genesis related to the presence
 198 of the Gulf Stream (Roebber, 1984).

3 Results

3.1 All extratropical cyclones

Looking at all ETCs along North America's Atlantic coast shows that storms in GCMH-SH tend to be too weak compared to ERAH-SH, leading to overestimation of weaker ETCs with maximum vorticities below $6 \cdot 10^{-5} \text{ s}^{-1}$, and too few stronger ETCs above this threshold (GCMH-SH minus ERAH-SH) (Figure 2a and c). The absolute negative bias is largest for strong ETCs that reach vorticities of about $10 \cdot 10^{-5} \text{ s}^{-1}$. The relative negative bias becomes larger as relative vorticity increases, reaching 76% for ETCs with maximum vorticities of $12 \cdot 10^{-5} \text{ s}^{-1}$ (Figure 2e). This is consistent with findings from Zappa et al. (2013a) who showed that CMIP5 models generally underestimate the frequency of extreme ETCs. Regridding data to NAE22 prior to ETC tracking yields similar but not identical values (GCMH-DC minus ERAH-DC) (Figure 2b, c and e). The impact of using two different grids and spatial filtering techniques is sufficiently small to yield comparable results. Nevertheless it is necessary to restrict comparisons to data sets that are based on the same grid and spatial filtering technique as the results are not identical. Downscaling CanESM2 with CanRCM4 decreases the frequency bias of stronger ETCs with vorticities ranging from 6 to $13 \cdot 10^{-5} \text{ s}^{-1}$ (RCMH-DC minus ERAH-DC) (Figure 2c). This is consistent with Long et al. (2009), who also found that dynamical downscaling can increase the frequency of strong ETCs along North America's Atlantic coast.

Anthropogenic GHG forcing is projected to generally weaken ETCs, leading to more events with vorticities below $5 \cdot 10^{-5} \text{ s}^{-1}$ and fewer events with vorticities between 5 and $11 \cdot 10^{-5} \text{ s}^{-1}$ (GCMF-SH minus GCMH-SH) (Figure 2a, d and f). The total number of tracks decreases by 2%, but changes are not statistically significant. This agrees with results from Christensen et al. (2013) who found only a weak reduction of ETC frequency for the CMIP5 climate model ensemble in the same region. Extreme ETCs with relative vorticity exceeding $11 \cdot 10^{-5} \text{ s}^{-1}$ (99th percentile) on the other hand are projected to increase, rising from 79 tracks to 97 tracks per 20-year period (23%; not statistically significant). Dynamical downscaling further reduces the projected number of ETC tracks with vorticities between 4 and $9 \cdot 10^{-5} \text{ s}^{-1}$ (RCMF-DC minus RCMH-DC) (Figure 2d).

To summarize, CanESM2 tends to simulate too many weak and not enough strong ETCs. Dynamical downscaling reduces this bias by increasing the number of stronger storms. Anthropogenic GHG forcing is projected to weaken ETCs, leading to more weak storms and fewer strong events. Dynamical downscaling further reduces

the number of strong storms. The total number of ETCs, and the frequency of extreme ETCs (99th percentile) is not projected to change significantly.

3.2 Explosive extratropical cyclones

3.2.1 Frequency

The most intense ETCs that pass along North America's Atlantic coast undergo rapid intensification (Figure 2a, magenta-colored bars). The annual mean EETC frequency is 45 tracks for ERAH-SH (Figure 3a). GCMH-SH underestimates this number by 38%, or 17 EETC tracks per year (Table 2), which is consistent with the negative frequency bias of stronger ETCs documented in section 3.1. Comparable EETC frequency biases are also found for other CMIP5 models (Seiler and Zwiers, 2016a). Regridding the same data to the NAE22 grid prior to tracking leads to similar results, with 40 tracks for ERAH-DC, and 24 tracks for GCMH-DC, and a negative EETC frequency bias of 41% (17 tracks per year). The slightly lower frequencies for data that are interpolated to the NAE22 grid are consistent with a loss of information that can be expected through the regridding process and somewhat different filtering approach that must be used in non-global domains. Nevertheless, as for ETCs in general, we conclude that the impact of using two different grids and spatial filtering techniques is sufficiently small to yield comparable results, but that comparisons should be restricted to data sets that are based on the same grid and spatial filtering technique as the results are not identical.

Dynamical downscaling increases EETC track frequency by 33% (8 tracks per year) when comparing RCMH-DC against GCMH-DC (Table 2, Figure 3a, and Figure 4). Downscaling therefore reduces the frequency bias from -41% in the global data on the NAE22 grid (GCMH-DC minus ERAH-DC) to -22% (RCMH-DC minus ERAH-DC), which is consistent with the increase in stronger ETCs in the downscaled solution documented in section 3.1.

CanESM2 projects a decrease in EETC frequency of 15% (4 tracks yr^{-1}) (GCMF-SH minus GCMH-SH) (Table 2, Figure 3a). This agrees with the projected decrease in stronger ETCs shown in section 3.1, and is comparable to the reduction in the Northern Atlantic (-17%) found for the CMIP5 climate model ensemble (Seiler and Zwiers, 2016b). Results from dynamically downscaled data are similar, with a projected decrease by 18% (6 tracks yr^{-1}) (RCMF-DC minus RCMH-DC). This result is in agreement with findings from Long et al. (2009) but is in contrast to results from Willison et al. (2015) for reasons discussed in section 4.

253 To summarize, CanESM2 underestimates the frequency of EETCs by 38%, and dynamical downscaling reduces
254 this bias to -22%. The projected relative reduction in EETC frequency is similar for the global model (-15%) and
255 its dynamically downscaled counterpart (-18%).

256 3.2.2 Intensity

257 The average EETC track along North America's Atlantic coast has a maximum relative vorticity of $8 \cdot 10^{-5} \text{ s}^{-1}$, a
258 minimum MSLP of 981 hPa, a maximum deepening rate of 21 hPa 24 h^{-1} , a maximum wind speed of 36 m s^{-1} (130
259 km h^{-1}), and a maximum precipitation rate of 33 mm day^{-1} (ERAH-SH) (Figure 3b-f). Intensities computed from
260 GCMH-SH are similar, with slightly lower values in maximum relative vorticity ($-0.8 \cdot 10^{-5} \text{ s}^{-1}$, -9%), maximum
261 wind speed (-2 m s^{-1} , -5%), and maximum precipitation (-2.8 mm day^{-1} , -8%), and slightly higher values in
262 minimum MSLP (5 hPa, 0.5%) (Table 2, Figure 3b-f). Regridding the data to the NAE22 grid prior to tracking
263 has no major impacts on EETC intensities. The biases of GCMH-SH and GCMH-DC differ by $0.36 \cdot 10^{-5} \text{ s}^{-1}$ (5%)
264 for relative vorticity, 1 hPa (0.07%) for MSLP, 1 m s^{-1} (2%) for wind speed, and 1 mm day^{-1} (3%) for maximum
265 precipitation.

266 Dynamical downscaling increases maximum EETC precipitation by 22% (7 mm day^{-1}) (RCMH-DC minus
267 GCMH-DC) (Table 2, Figure 3f). This is consistent with the increase in EETC frequency described in section 3.2.1,
268 and indicates that the frequency increase is linked to an enhanced positive feedback between ETC intensification
269 and latent heat release at the higher resolution as suggested by Willison et al. (2013). EETC relative vorticity,
270 MSLP, deepening rate, and wind speed are not significantly affected by dynamical downscaling (Table 2).

271 Anthropogenic GHG forcing is projected to increase maximum EETC precipitation by 46%, or by 14 mm
272 day^{-1} (GCMF-SH minus GCMH-SH) (Table 2, Figure 3f). This is consistent with findings from Zappa et al.
273 (2013b) who show that CMIP5 models project an increase in the frequency of ETCs that are associated with
274 strong precipitation in the Northern Atlantic. The projected increase in EETC precipitation is consistent with a
275 projected 35% increase in specific humidity at 850 hPa from October to March (not shown). The maximum EETC
276 precipitation increase per degree warming is $8.9\% \text{ K}^{-1}$, which exceeds the Clausius-Clapeyron rate of $6.6\% \text{ K}^{-1}$
277 for the region (Table 3). While maximum wind speed is projected to increase by 1.4 m s^{-1} (4%), other measures
278 of EETC intensity are not significantly affected by anthropogenic GHG forcing (Table 2, Figure 3b-e).

279 Dynamical downscaling enhances the projected increase in maximum EETC precipitation from 46% (14 mm
280 day^{-1}) to 52% (20 mm day^{-1}) (RCMF-DC minus RCMH-DC) (Table 2, Figure 3f). Maximum precipitation

281 associated with EETCs increases by 10.6% per degree warming for the downscaled data (Table 3). As in CanESM2,
 282 the projected increase is consistent with a 35% increase in specific humidity at 850 hPa for October to March
 283 (not shown).

284 To summarize, EETCs simulated by CanESM2 are slightly weaker compared to ERA-Interim. Dynamical
 285 downscaling increases maximum EETC precipitation by 22%, which suggests that dynamical downscaling may
 286 have enhanced the positive feedback between ETC intensification and latent heat release. Anthropogenic GHG
 287 forcing is projected to increase maximum EETC precipitation by 46% for the global data and by 52% for the
 288 downscaled data. The corresponding precipitation increase per degree warming is 8.9% K⁻¹ and 10.6% K⁻¹ for
 289 CanESM2 and CanRCM4, respectively, which exceeds the Clausius-Clapeyron relation of 6.6% K⁻¹.

290 3.2.3 Life cycle

291 Life cycle composites show that explosive development occurs within a period of two days, with a maximum
 292 deepening rate of 24 hPa 24h⁻¹ (Figure 5, ERAH-SH). MSLP drops from 1011 hPa to 971 hPa, relative vorticity
 293 increases from 3.5 10⁻⁵ s⁻¹ to 9.9 10⁻⁵ s⁻¹, and maximum precipitation increases from 8 mm day⁻¹ to 25 mm
 294 day⁻¹. EETCs reach their minimum MSLP and maximum precipitation values approximately 6 hours after their
 295 maximum vorticity. The rate of MSLP intensification is much stronger compared to the rate of MSLP decay. The
 296 corresponding rates for relative vorticity and maximum precipitation on the other hand are quite similar. Similar
 297 rates are presented in Bengtsson et al. (2009).

298 CanESM2 reproduces these patterns well, with slightly smaller changes in MSLP (from 1011 hPa to 976 hPa),
 299 relative vorticity (from 3.1 10⁻⁵ s⁻¹ to 9.2 10⁻⁵ s⁻¹), and maximum precipitation (from 7 mm day⁻¹ to 23 mm
 300 day⁻¹) (Figure 5, GCMH-SH). The timing of the maximum intensities in GCMH-SH is accurate for MSLP and
 301 relative vorticity, and 12 hours early for maximum precipitation.

302 Regridding ERA-Interim to the NAE22 grid prior to tracking (ERAH-DC) leads to similar but not identical
 303 life cycle composites compared to ERAH-SH, with MSLP dropping from 1011 hPa to 973 hPa, relative vorticity
 304 increasing from 2.8 to 9.0 10⁻⁵ s⁻¹, and precipitation rising from 6 to 26 mm day⁻¹. Life cycle composites of GCMH-
 305 DC are similar to the ones of ERAH-DC for MSLP (from 1012 hPa to 978 hPa) and maximum precipitation
 306 (from 5 mm day⁻¹ to 25 mm day⁻¹). The increase in maximum relative vorticity however is smaller in GCMH-DC
 307 compared to ERAH-DC (from 2.6 10⁻⁵ s⁻¹ to 7.7 10⁻⁵ s⁻¹). Dynamical downscaling slightly improves the result by

308 increasing maximum vorticity to $8.3 \cdot 10^{-5} \text{ s}^{-1}$ (RCMH-DC). However, the MSLP minimum in RCMH-DC is still
309 too high, and the precipitation maximum is still 12 hours too early compared to ERAH-DC.

310 Projections under RCP8.5 forcing exhibit no major impacts on the overall life cycle patterns described above.
311 The most noticeable change is the projected increase in maximum precipitation from 23 mm day^{-1} in the historical
312 experiment (GCMH-SH) to 34 mm day^{-1} under RCP8.5 forcing (GCMF-SH). The corresponding values for the
313 downscaled experiment are slightly higher, with 27 mm day^{-1} for RCMH-DC and 39 mm day^{-1} for RCMF-DC.

314 To summarize, life cycle composites show that explosive development occurs within a period of two days,
315 where a sharp drop in MSLP by $24 \text{ hPa } 24 \text{ h}^{-1}$ coincides with a rise in relative vorticity and maximum precipitation.
316 CanESM2 reproduces these patterns reasonably well, with a tendency for too high MSLP and too low relative
317 vorticity values, and a peak in maximum precipitation that is 12 hours too early. Downscaling CanESM2 slightly
318 improves this result by increasing maximum vorticity. Projections under RCP8.5 forcing exhibit a projected
319 increase in maximum precipitation from 23 mm day^{-1} (GCMH-SH) to 34 mm day^{-1} (GCMF-SH), with slightly
320 higher values for the corresponding downscaled counterparts.

321 3.3 Baroclinic instability

322 The highest density of EETC tracks occurs in a region with strong cold season lower tropospheric Eady growth
323 rate (1.8 to 2.0 day^{-1}) (ERAH-SH) (Figure 6a and b). The highest zonal mean frequency is located at about
324 45° north where the zonal mean Eady growth rate is about 1.6 day^{-1} (Figure 7a and c). This also applies to
325 experiments that are based on discrete cosine transform rather than spherical harmonic decomposition (Figure
326 7b). CanESM2 underestimates the zonal mean Eady growth rate value by 0.1 day^{-1} , which is consistent with
327 the negative EETC frequency bias (Figure 6c and Figure 7c). Similar underestimation along North America's
328 Atlantic coast is also found in the multi-model mean of Eady growth rates in the CMIP5 climate model ensemble
329 (Seiler and Zwiers, 2016a; their Figure 10l). The impact of dynamical downscaling on the Eady growth rate is not
330 statistically significant in the main region of the storm track (Figure 6d and Figure 7d).

331 Projections under RCP8.5 forcing show a decrease of the lower tropospheric Eady growth rate by 0.2 day^{-1}
332 between 40° and 45° latitude (Figure 6e and Figure 7c), which is consistent with the projected decline in EETC
333 frequency in CanESM2, and with Eady growth rate projections in other CMIP5 models (Seiler and Zwiers,
334 2016b). Also, the zonal mean maximum Eady growth rate and the maximum EETC frequency are projected to
335 shift northwards by about two to three degrees latitude (Figure 7a and c). Downscaling does not significantly

336 impact the projected decrease in the Eady growth rate in most of the storm track region (Figure 6f and Figure
337 7c and d). This is in contrast to Willison et al. (2015) who showed that the lower tropospheric Eady growth
338 rate increases with anthropogenic GHG forcing, and that this impact is enhanced when increasing the horizontal
339 model resolution in a pseudo global warming experiment.

340 The low Eady growth rate in CanESM2 is due to underestimation of vertical wind shear rather than overes-
341 timation of the Brunt-Väisälä frequency, as CanESM2 slightly underestimates the latter by $0.25 \cdot 10^{-3}$ to $0.5 \cdot 10^{-3}$
342 s^{-1} (not shown). The Eady growth rate bias coincides with a warm temperature bias in the lower troposphere
343 (850 hPa) over Eastern Canada between 40° and 60° latitude (2-4 K) (Figure 8a and b, and Figure 9a). The
344 resulting meridional temperature gradients are therefore smaller in CanESM2 compared to ERA-Interim, with
345 differences as large as 0.3 K per 100 km at 850 hPa (Figure 8c, yellow shaded area along North America's Atlantic
346 coast). This could lead to underestimation of the Eady growth rate, as vertical wind shear and the meridional
347 temperature gradients are linked through the thermal wind equation.

348 To summarize, the highest density of EETC tracks occurs in a region with strong cold season lower tropospheric
349 Eady growth rate. CanESM2 underestimates the lower tropospheric Eady growth rate by 0.1 day^{-1} , which is
350 consistent with the model's negative EETC frequency bias. Anthropogenic GHG forcing is projected to decrease
351 the lower tropospheric Eady growth rate by 0.2 day^{-1} , which may explain the projected decline in EETC frequency
352 in CanESM2. The negative Eady growth rate bias in CanESM2 is due to an underestimation of vertical wind
353 shear rather than an overestimation of the Brunt-Väisälä frequency. The weak vertical wind shear is consistent
354 with too weak meridional temperature gradients related to a warm temperature bias in the lower troposphere over
355 Eastern Canada. Dynamical downscaling has no significant impacts on Eady growth rate biases and projections
356 in most of the storm track region.

357 4 Discussion

358 This study explores how dynamical downscaling affects model biases and projections of EETCs along North
359 America's Atlantic coast. The regional climate model CanRCM4 is forced with the global climate model CanESM2
360 for the periods 1981 to 2000 and 2081 to 2100. ETCs are tracked from relative vorticity using an objective feature
361 tracking algorithm. Special attention is paid to the impact of different spatial filters used for data with different
362 spatial grids. CanESM2 is shown to simulate too many weak and too few strong ETCs, which is consistent with
363 findings from Zappa et al. (2013a). Dynamical downscaling reduces this bias by increasing the number of stronger

364 storms, consistent with results from Long et al. (2009). Forcing under the RCP8.5 emission scenario is projected
365 to weaken ETCs when comparing the late 21st century with the late 20th century, leading to more weaker and
366 fewer stronger storms. Downscaling further reduces the projected number of stronger storms. The total number of
367 ETCs, and the frequency of extreme ETCs (99th percentile) are not projected to change statistically significantly.

368 CanESM2 underestimates EETC track frequency by 38%, which is comparable to the biases of other CMIP5
369 models (Seiler and Zwiers, 2016a). Dynamical downscaling reduces the CanESM2 bias to -22%. The relatively
370 higher EETC frequency in CanRCM4 coincides with a corresponding 22% increase in maximum EETC precipita-
371 tion. This is consistent with Willison et al. (2013) who show that a positive feedback between ETC intensification
372 and latent heat release is enhanced at higher horizontal model resolution. Dynamical downscaling with CanRCM4
373 does not completely eliminate the EETC frequency bias of CanESM2, possibly due to underestimation of the Eady
374 growth rate, which is not affected by dynamical downscaling. The use of spectral nudging may have constrained
375 the Eady growth rate in CanRCM4 to that of the driving model.

376 The negative Eady growth rate bias in CanESM2 is due to underestimation of vertical wind shear rather than
377 overestimation of the Brunt-Väisälä frequency. The weak vertical wind shear may be caused by weak meridional
378 temperature gradients, as both variables are linked through the thermal wind equation. Meridional temperature
379 gradients are smaller in CanESM2 compared to ERA-Interim, possibly due to a warm temperature bias in the
380 lower troposphere over Eastern Canada. We conclude that the coarse spatial resolution of global climate models
381 presents an important limitation for simulating extreme ETCs, but biases in the large scale circulation that affect
382 baroclinic instability may be just as relevant. Future research on model biases of intense coastal storms should
383 assess the mechanisms that lead to an underestimation of the lower-tropospheric Eady growth rate along North
384 America's Atlantic coast.

385 CanESM2 projects a 15% decrease in EETC frequency (4 tracks yr^{-1}) by the end of this century, which is
386 comparable to CMIP5 model projections (Seiler and Zwiers, 2016b). Dynamical downscaling has little impacts on
387 the relative changes that are projected in EETC track frequency and intensity, suggesting that CMIP5 projections
388 in Seiler and Zwiers (2016b) may not be very sensitive to horizontal model resolution. This is consistent with our
389 finding that dynamical downscaling has no significant impact on the projected reduction in the lower tropospheric
390 Eady growth rate, and consistent with Long et al. (2009) (their Figure 9). In contrast Willison et al. (2015) showed
391 that the lower tropospheric Eady growth rate increases with anthropogenic GHG forcing, and that this impact is
392 enhanced when increasing the horizontal model resolution in a pseudo global warming experiment. Such contrasts

393 may arise due to the different nature of pseudo global warming experiments, and may also be related to the
394 fact that our experiments were spectrally nudged. Future research could compare these results to climate model
395 biases and projections of EETCs in HighResMIP simulations. It is possible that such simulations also reduce the
396 negative Eady growth rate bias, as HighResMIP runs are not constrained by spectral nudging.

397 The projected decline in EETC frequency is consistent with a 0.2 day^{-1} decrease in the lower tropospheric Eady
398 growth rate between 40° and 45° latitude. The Eady growth rate bias (-0.1 day^{-1}) is smaller than its projected
399 reduction (-0.2 day^{-1}), but the EETC track frequency bias (-38%) is larger than its projected decline (-15%). This
400 suggests that the impact of the projected Eady growth rate reduction might be partially offset by a mechanism
401 that favors ETC intensification, such as enhanced baroclinic instability in the upper troposphere related to tropical
402 amplification (Hall et al., 1994), or condensational heating corresponding to the projected increase in maximum
403 EETC precipitation. Changes in the contribution of moist diabatic processes to EETC intensification could be
404 analyzed from the potential vorticity perspective (Davis and Emanuel, 1991).

405 Maximum EETC precipitation is projected to increase by 46% in the global model. The projected precipitation
406 rate per degree warming is estimated as $8.9\% \text{ K}^{-1}$, which exceeds the Clausius-Clapeyron rate of $6.6\% \text{ K}^{-1}$ for
407 the region, reflecting the role of moisture convergence in maximum EETC precipitation. Changes in the spatial
408 distribution of precipitation could be studied in more detail in storm centered composites (*e.g.* Bengtsson et al.,
409 2009).

410 To conclude, this study assesses how dynamical downscaling with a spectrally nudged model affects biases and
411 projections of EETCs along North America's Atlantic coast. Dynamical downscaling significantly reduces EETC
412 biases, likely due to enhanced precipitation and associated condensational heating. The remaining bias is consistent
413 with underestimation of baroclinic instability, which is not affected by dynamical downscaling. Projections under
414 RCP8.5 forcing exhibit a reduction in EETC frequency, and an increase maximum EETCs precipitation. The
415 projected relative changes are not very sensitive to horizontal model resolution. Future research should assess the
416 relative contribution of moist diabatic processes for ETC intensification, and study how these contributions vary
417 with storm intensity under current and projected climatic conditions.

418 **Acknowledgements**

419 The authors gratefully acknowledge the financial support of the Marine Environmental Observation Prediction
420 and Response Network (MEOPAR) for this research. We thank Dr. Yanjun Jiao from the Canadian Centre for

421 Climate Modelling and Analysis (CCCma) for providing us with data from CanRCM4. We acknowledge the
422 World Climate Research Programme’s Working Group on Coupled Modelling, which is responsible for CMIP,
423 and we thank CCCma and ECMWF for producing and making available their model output. For CMIP the
424 U.S. Department of Energy’s Program for Climate Model Diagnosis and Intercomparison provides coordinating
425 support and led development of software infrastructure in partnership with the Global Organization for Earth
426 System Science Portals.

427 References

- 428 Anderson, D., K. I. Hodges, and B. J. Hoskins, 2003: Sensitivity of feature-based analysis methods of storm tracks
429 to the form of background field removal. *Monthly weather review*, **131** (3), 565–573.
- 430 Arora, V. K., and Coauthors, 2011: Carbon emission limits required to satisfy future representative concentration
431 pathways of greenhouse gases. *Geophysical Research Letters*, **38** (5), n/a–n/a, doi:10.1029/2010GL046270,
432 URL <http://dx.doi.org/10.1029/2010GL046270>, 105805.
- 433 Bengtsson, L., K. I. Hodges, and N. Keenlyside, 2009: Will extratropical storms intensify in a warmer climate?
434 *Journal of Climate*, **22** (9), 2276–2301.
- 435 Bengtsson, L., K. I. Hodges, and E. Roeckner, 2006: Storm tracks and climate change. *Journal of Climate*,
436 **19** (15), 3518–3543.
- 437 Chang, E. K. M., Y. Guo, and X. Xia, 2012: CMIP5 multimodel ensemble projection of storm track change
438 under global warming. *Journal of Geophysical Research: Atmospheres*, **117** (D23), n/a–n/a, doi:10.1029/
439 2012JD018578, URL <http://dx.doi.org/10.1029/2012JD018578>, d23118.
- 440 Christensen, J., and Coauthors, 2013: Climate Phenomena and their Relevance for Future Regional Climate
441 Change. In: Climate Change 2013: The Physical Science Basis. Working Group I Contribution to the Fifth
442 Assessment Report of the Intergovernmental Panel on Climate Change. Tech. rep., Groupe d’experts intergou-
443 vernemental sur l’évolution du climat/Intergovernmental Panel on Climate Change-IPCC, C/O World Meteo-
444 rological Organization, 7bis Avenue de la Paix, CP 2300 CH-1211 Geneva 2 (Switzerland).
- 445 Colle, B. A., J. F. Booth, and E. K. Chang, 2015: A Review of Historical and Future Changes of Extratropical
446 Cyclones and Associated Impacts Along the US East Coast. *Current Climate Change Reports*, **1** (3), 125–143.
- 447 Collins, M., and Coauthors, 2013: Long-term Climate Change: Projections, Commitments and Irreversibility. In:
448 Climate Change 2013: The Physical Science Basis. Working Group I Contribution to the Fifth Assessment

- 449 Report of the Intergovernmental Panel on Climate Change. Tech. rep., Groupe d'experts intergouvernemental sur l'évolution du climat/Intergovernmental Panel on Climate Change-IPCC, C/O World Meteorological
450 Organization, 7bis Avenue de la Paix, CP 2300 CH-1211 Geneva 2 (Switzerland).
- 451
- 452 Côté, H., K. M. Grise, S.-W. Son, R. de Elía, and A. Frigon, 2015: Challenges of tracking extratropical cyclones
453 in regional climate models. *Climate Dynamics*, **44** (11-12), 3101–3109.
- 454 Davis, C. A., and K. A. Emanuel, 1991: Potential vorticity diagnostics of cyclogenesis. *Monthly weather review*,
455 **119** (8), 1929–1953.
- 456 Dee, D., and Coauthors, 2011: The ERA-Interim reanalysis: Configuration and performance of the data assimila-
457 tion system. *Quarterly Journal of the Royal Meteorological Society*, **137** (656), 553–597.
- 458 Denis, B., J. Côté, and R. Laprise, 2002: Spectral decomposition of two-dimensional atmospheric fields on limited-
459 area domains using the discrete cosine transform (DCT). *Monthly Weather Review*, **130** (7), 1812–1829.
- 460 Fink, A. H., S. Pohle, J. G. Pinto, and P. Knippertz, 2012: Diagnosing the influence of diabatic processes on the
461 explosive deepening of extratropical cyclones. *Geophysical Research Letters*, **39** (7), n/a–n/a, doi:10.1029/
462 2012GL051025, URL <http://dx.doi.org/10.1029/2012GL051025>, 107803.
- 463 Giorgi, F., C. Jones, G. R. Asrar, and Coauthors, 2009: Addressing climate information needs at the regional
464 level: the CORDEX framework. *World Meteorological Organization (WMO) Bulletin*, **58** (3), 175.
- 465 Hall, N. M., B. J. Hoskins, P. J. Valdes, and C. A. Senior, 1994: Storm tracks in a high-resolution GCM with
466 doubled carbon dioxide. *Quarterly Journal of the Royal Meteorological Society*, **120** (519), 1209–1230.
- 467 Hodges, K., 1994: A general method for tracking analysis and its application to meteorological data. *Monthly
468 Weather Review*, **122** (11), 2573–2586.
- 469 Hodges, K., 1999: Adaptive constraints for feature tracking. *Monthly Weather Review*, **127** (6), 1362–1373.
- 470 Hodges, K., and Coauthors, 1995: Feature tracking on the unit-sphere. *Monthly Weather Review*, **123** (12),
471 3458–3465.
- 472 Hoskins, B. J., and P. J. Valdes, 1990: On the existence of storm-tracks. *Journal of the atmospheric sciences*,
473 **47** (15), 1854–1864.
- 474 Kocin, P. J., P. N. Schumacher, R. F. Morales Jr, and L. W. Uccellini, 1995: Overview of the 12-14 March 1993
475 superstorm. *Bulletin of the American Meteorological Society*, **76** (2), 165–182.
- 476 Lambert, S. J., and J. C. Fyfe, 2006: Changes in winter cyclone frequencies and strengths simulated in enhanced
477 greenhouse warming experiments: results from the models participating in the IPCC diagnostic exercise. *Climate*

- 478 *Dynamics*, **26 (7-8)**, 713–728.
- 479 Long, Z., W. Perrie, J. Gyakum, R. Laprise, and D. Caya, 2009: Scenario changes in the climatology of winter
480 midlatitude cyclone activity over eastern North America and the Northwest Atlantic. *Journal of Geophysical*
481 *Research: Atmospheres*, **114 (D12)**, n/a–n/a, doi:10.1029/2008JD010869, URL [http://dx.doi.org/10.1029/](http://dx.doi.org/10.1029/2008JD010869)
482 2008JD010869, d12111.
- 483 Marciano, C. G., G. M. Lackmann, and W. A. Robinson, 2015: Changes in US East Coast cyclone dynamics with
484 climate change. *Journal of Climate*, **28 (2)**, 468–484.
- 485 McDonald, R. E., 2011: Understanding the impact of climate change on Northern Hemisphere extra-tropical
486 cyclones. *Climate dynamics*, **37 (7-8)**, 1399–1425.
- 487 R Core Team, 2013: *R: A Language and Environment for Statistical Computing*. Vienna, Austria, R Foundation
488 for Statistical Computing, URL <http://www.R-project.org/>.
- 489 Roebber, P. J., 1984: Statistical analysis and updated climatology of explosive cyclones. *Monthly Weather Review*,
490 **112 (8)**, 1577–1589.
- 491 Sanders, F., and J. R. Gyakum, 1980: Synoptic-dynamic climatology of the “bomb”. *Monthly Weather Review*,
492 **108 (10)**, 1589–1606.
- 493 Scinocca, J., and Coauthors, 2016: Coordinated global and regional climate modeling*. *Journal of Climate*,
494 **29 (1)**, 17–35.
- 495 Seiler, C., and F. Zwiers, 2016a: How well do CMIP5 climate models reproduce explosive cyclones in the extrat-
496 ropics of the Northern Hemisphere? *Climate Dynamics*, **46 (3-4)**, 1241–1256.
- 497 Seiler, C., and F. Zwiers, 2016b: How will climate change affect explosive cyclones in the extratropics of the
498 Northern Hemisphere? *Climate Dynamics*, **46 (11)**, 3633–3644.
- 499 Stull, R. B., 2000: *Meteorology for scientists and engineers: a technical companion book with Ahrens’ Meteo-*
500 *rology Today*. Brooks/Cole.
- 501 Taylor, K. E., R. J. Stouffer, and G. A. Meehl, 2012: An overview of CMIP5 and the experiment design. *Bulletin*
502 *of the American Meteorological Society*, **93 (4)**, 485–498.
- 503 Ulbrich, U., J. Pinto, H. Kupfer, G. Leckebusch, T. Spanghel, and M. Reyers, 2008: Changing Northern Hemisphere
504 storm tracks in an ensemble of IPCC climate change simulations. *Journal of climate*, **21 (8)**, 1669–1679.
- 505 von Salzen, K., and Coauthors, 2013: The Canadian fourth generation atmospheric global climate model
506 (CanAM4). Part I: representation of physical processes. *Atmosphere-Ocean*, **51 (1)**, 104–125.

-
- 507 Willison, J., W. A. Robinson, and G. M. Lackmann, 2013: The importance of resolving mesoscale latent heating
508 in the North Atlantic storm track. *Journal of the Atmospheric Sciences*, **70 (7)**, 2234–2250.
- 509 Willison, J., W. A. Robinson, and G. M. Lackmann, 2015: North Atlantic Storm-Track Sensitivity to Warming
510 Increases with Model Resolution. *Journal of Climate*, **28 (11)**, 4513–4524.
- 511 Zappa, G., L. C. Shaffrey, and K. I. Hodges, 2013a: The Ability of CMIP5 Models to Simulate North Atlantic
512 Extratropical Cyclones*. *Journal of Climate*, **26 (15)**, 5379–5396.
- 513 Zappa, G., L. C. Shaffrey, K. I. Hodges, P. G. Sansom, and D. B. Stephenson, 2013b: A multimodel assessment of
514 future projections of north atlantic and european extratropical cyclones in the cmip5 climate models*. *Journal*
515 *of Climate*, **26 (16)**, 5846–5862.




Fig. 1 EETC tracks and their along track MSLP values computed from ERA-Interim for the period 1981-2000. The outer polygon presents the CanRCM4 eastward extended North American model domain, the small polygon shows the area that is used to compute statistics for coastal storms, and the middle-sized polygon encloses the area used to compute zonal mean values in Figure 7.




Fig. 2 Frequency histograms of the maximum relative vorticity that is attained in each ETC track along North America’s Atlantic coast for data on (a) the global grid and (b) the NAE22 grid. Subplots (c) and (d) show the absolute biases and the projected changes, and subplots (e) and (f) show the relative biases and the projected changes, respectively. The relative biases are shown for relative vorticity values below $13 \cdot 10^{-5} \text{ s}^{-1}$ due to the small number of ETC tracks at higher intensities. Only ETC segments that are enclosed by the polygon that encompasses North America’s Atlantic coast in Figure 1 are considered.




Fig. 3 Percentiles (0.05, 0.25, 0.50, 0.75, 0.95) (boxplots) and mean values (asterisks) of (a) annual mean frequency, (b) maximum relative vorticity, (c) minimum MSLP, (d) maximum deepening rate, (e) maximum wind speed, and (f) maximum precipitation of EETC tracks along North America’s Atlantic coast. Only EETC segments that are enclosed by the polygon that encompasses North America’s Atlantic coast in Figure 1 are considered.




Fig. 4 EETC tracks with explosive segments shaded in red for (a) ERAH-DC, (b) GCMH-DC, and (c) RCMH-DC for the period 1981 to 2000. Subplots (d) to (f) show the corresponding annual frequency of 6 hourly EETC centers.




Fig. 5 Life cycle composites of all EETCs that pass through the polygon that encompasses North America’s Atlantic coast in Figure 1. Contrary to Figure 3, this also includes the parts of EETCs that lie outside the polygon. Parameters shown are (a, b) MSLP, (c, d) relative vorticity, and (e, f) maximum precipitation. Subplots (a, c, and e) and (b, d, and f) show results for experiments that are based on spherical harmonic decomposition and discrete cosine transform, respectively. The centered lines show the medians and the outer lines enclose the interquartile ranges. The life cycles of EETC tracks are combined by centering the tracks according to the time step of maximum relative vorticity.




Fig. 6 (a) EETC frequency in ERA-Interim (ERA-SH) for the period 1981 to 2000, (b) lower tropospheric Eady growth rate of ERA-Interim for the same period during the cold season, (c) Eady growth rate bias of CanESM2 (1981-2000), (d) the difference between Eady growth rate in CanESM2 and CanRCM4 (1981-2000), (e) Eady growth rate projections under RCP8.5 by CanESM2 (1981-2000 and 2081-2100), and (f) the difference between Eady growth rate projections from CanESM2 and CanRCM4 (2081-2100). Regions with statistically significant differences at the 5% level are hatched (Wilcox test, R Core Team, 2013). The red lines represent 40° and 45° of northern latitude.




Fig. 7 Zonal mean frequency of EETC centers for experiments based on the (a) spherical harmonic decomposition and (b) discrete cosine transform. Subplots (c) and (d) show the corresponding zonal mean Eady growth rates. Zonal means are computed from 25° to 60° latitude and from -90° to -25° longitude, corresponding to the area enclosed by the middle-sized polygon shown in Figure 1.




Fig. 8 Air temperature biases at (a) 850 hPa and (b) 500 hPa (CanESM2 minus ERA-Interim) from October to March for the historical period 1981-2000. Subplots (c) and (d) show the corresponding meridional temperature gradient biases. Regions with statistically insignificant differences at the 5% level are masked out in white (Wilcoxon test, R Core Team, 2013). The gray box in (a) shows the region that is used to compute zonal mean values in Figure 9.




Fig. 9 (a) Zonal mean isotherms and air temperature biases (CanESM2 minus ERA-Interim), and (b) the corresponding temperatures of isobars for selected pressure levels. Zonal means are computed from 1981 to 2000 for the months October to March for a region enclosed by the longitudes -102° to -30° as denoted by the gray polygon in Figure 8a. Regions with statistically insignificant differences at the 5% level are masked out in white in subplot (a) (Wilcox test, R Core Team, 2013).

Table 1 Experiment ID's, input data, climate model experiments, grids, and spatial filtering techniques. Final grid refers to the grid that is used as input data for TRACK. All data is spatially filtered to T42 or T42 equivalent.

ID	Data	Experiment	Native Grid		Final Grid		Spatial filter (T42 or equivalent)
			Resolution	Extent	Resolution	Extent	
ERAH-SH	ERA-Interim	historical	0.75°	global	0.75°	global	spherical harmonic decomposition
GCMH-SH	CanESM2	historical	2.8125°	global	2.8125°	global	spherical harmonic decomposition
GCMF-SH	CanESM2	RCP8.5	2.8125°	global	2.8125°	global	spherical harmonic decomposition
ERAH-DC	ERA-Interim	historical	0.75°	global	0.22°	NAE22	discrete cosine transform
GCMH-DC	CanESM2	historical	2.8125°	global	0.22°	NAE22	discrete cosine transform
RCMH-DC	CanRCM4[CanESM2]	historical	0.22°	NAE22	0.22°	NAE22	discrete cosine transform
RCMF-DC	CanRCM4[CanESM2]	RCP8.5	0.22°	NAE22	0.22°	NAE22	discrete cosine transform

Table 2 Differences in the frequency and intensity of EETC tracks in proximity of North America’s Atlantic coast for selected experiments. Intensities (*i.e.* relative vorticity, MSLP, deepening rate, wind speed, and precipitation) refer to the strongest intensities that are reached on average by EETC tracks. Only EETC segments that are enclosed by the polygon that encompasses North America’s Atlantic coast in Figure 1 are considered. Statistically significant differences are written in bold (two-sampled Wilcoxon test, 5% level).

Experiments	Track freq.		Vorticity		MSLP		Δ MSLP		Windspeed		Precipitation	
	yr ⁻¹	%	10 ⁻⁵ s ⁻¹	%	hPa	%	hPa/24h	%	m s ⁻¹	%	mm day ⁻¹	%
GCMH-SH minus ERAH-SH	-17	-38	-0.8	-9.4	5.0	0.5	1.6	-7.6	-1.9	-5.4	-2.8	-8.5
GCMH-DC minus ERAH-DC	-17	-41	-1.2	-14.1	4.4	0.4	1.3	-5.7	-1.4	-3.9	-1.7	-5.2
RCMH-DC minus GCMH-DC	8	33	0.3	3.8	-0.2	0.0	1.5	-7.0	0.2	0.7	7.1	22.3
RCMH-DC minus ERAH-DC	-9	-22	-0.9	-10.8	4.2	0.4	2.7	-12.3	-1.2	-3.2	5.4	16.0
GCMF-SH minus GCMH-SH	-4	-15	0.3	4.4	0.1	0.0	-0.4	1.8	1.4	4.2	13.9	45.7
RCMF-DC minus RCMH-DC	-6	-18	0.3	3.8	0.2	0.0	-0.5	2.8	0.0	0.0	20.2	52.0

Table 3 Projected changes in near-surface air temperature (ΔT), and saturated vapor pressure (Δe_s) averaged over the area enclosed by the small polygon in Figure 1, and maximum EETC precipitation (ΔP), and the corresponding rates of change per degree warming for CanESM2 and CanRCM4.

	ΔT (K)	Δe_s (%)	ΔP (%)	$\frac{\Delta e_s}{\Delta T}$ ($\frac{\%}{K}$)	$\frac{\Delta P}{\Delta T}$ ($\frac{\%}{K}$)
CanESM2	5.2	34.3	45.3	6.6	8.9
CanRCM4	4.9	32.5	54.6	6.7	10.6