

Electrical and structural adaptations of the paediatric athlete's heart: A systematic review and meta-analysis

Gavin McClean,^{1,2} Nathan R Riding,¹ Clare L Ardern,^{1,3,4} Abdulaziz Farooq,¹ Guido E Pielles,^{5,6} Victoria Watt,⁷ Carmen Adamuz,⁷ Keith P George,² David Oxborough,² and Mathew G Wilson^{1,2,8}

¹. Athlete Health and Performance Research Centre, ASPETAR Qatar Orthopaedic and Sports Medicine Hospital, Qatar.

². Research Institute for Sport and Exercise Science, Liverpool John Moores University, UK.

³. Division of Physiotherapy, Linköping University, Linköping, Sweden.

⁴. School of Allied Health, La Trobe University, Melbourne, Victoria, Australia.

⁵. National Institute for Health Research (NIHR) Cardiovascular Biomedical Research Unit, Congenital Heart Unit, Bristol Royal Hospital for Children and Bristol Heart Institute, Bristol, UK.

⁶. University of Bristol, UK.

⁷. Department of Sports Medicine, ASPETAR Qatar Orthopaedic and Sports Medicine Hospital, Qatar.

⁸. Research Institute of Sport and Exercise Sciences, University of Canberra, Australia.

Abstract: 244

Word Count: 4664

ABSTRACT

Aim: To describe the electrocardiographic (ECG) and echocardiographic manifestations of the paediatric athlete's heart, and examine the impact of age, race and sex upon cardiac remodelling responses to competitive sport.

Design: Systematic review and meta-analysis

Data sources: Six electronic databases were searched to May 2016: MEDLINE, PubMed, EMBASE, Web of Science, CINAHL and SPORTDiscus.

Inclusion criteria: 1) Male and/or female competitive athletes, 2) participants aged 6-18 years, 3) original research article published in English language.

Results: Data from 14,278 athletes and 1,668 non-athletes were included for qualitative (43 articles) and quantitative synthesis (40 articles). Paediatric athletes demonstrated a greater prevalence of training-related and training-unrelated ECG changes than non-athletes. Athletes ≥ 14 years were 15.8 times more likely to have inferolateral T-wave inversion than athletes < 14 years. Paediatric black athletes had significantly more training-related and training-unrelated ECG changes than Caucasian athletes. Age was a positive predictor of left ventricular (LV) internal diameter during diastole, interventricular septum thickness during diastole, relative wall thickness and LV mass. When age was accounted for, these parameters remained significantly larger in athletes than non-athletes. Paediatric black athletes presented larger posterior wall thickness during diastole (PWTd) than Caucasian athletes. Paediatric male athletes also presented with larger PWTd than females.

Conclusions: The paediatric athlete's heart undergoes significant remodelling both before and during 'maturational years'. Paediatric athletes have a greater prevalence of training related and training-unrelated ECG changes than non-athletes, with age, race and sex mediating factors on cardiac electrical and LV structural remodelling.

Word count: 244

What is already known?

- Chronic training loads are associated with a number of electrophysiological, structural and functional cardiac adaptations in adult athletes.
- Race and sex significantly impact upon the cardiac remodelling of the adult athlete's heart.
- Paediatric athletes undergo significant growth and maturational changes; but unlike known musculoskeletal changes, there is limited information regarding how the heart may adapt to training before, during and after puberty.

What are the new findings?

- Paediatric athletes were up to 13 times more likely to have deep T-wave inversion (TWI) ($\geq 2\text{mm}$) than age-matched non-athletes
- Paediatric athletes ≥ 14 years of age were up to 16 times more likely to have inferolateral TWI (warranting further investigation) than athletes < 14 years
- Paediatric black athletes were up to 36 times more likely to have extended anterior TWI (leads V1-V4) than Caucasians.
- Even after accounting for age, left ventricular structural parameters were larger among paediatric athletes than paediatric non-athletes

INTRODUCTION

Regular and sustained intensive physical activity is associated with a number of electrophysiological¹, structural and functional cardiac adaptations²; collectively referred to as the 'Athlete's Heart'. It is also well documented that race and sex significantly impact these manifestations of the adult athlete's heart^{3,4}. Whilst previous systematic reviews and meta-analyses have detailed the adult athlete's heart phenotype^{2,5}, with some accounting for race and sex^{6,7}, data from paediatric (6-18 years) athletic populations is limited to original research; often restricted by inadequate sample sizes and heterogeneity to assess the impact of age, race and sex in tangent.

Sports academies are increasingly used by clubs and governing bodies alike to develop and nurture talented sports stars of the future. Consequently, there is increasing competitiveness, professionalism and training demands placed upon the paediatric athlete during the maturational period. The International Olympic Committee, amongst others, has called for more diligence to safeguard the physiological development of the paediatric athlete⁸⁻¹⁰. Performing a cardiac pre participation evaluation (PPE) within paediatric populations is controversial due to a lack of international consensus with regards to when, how, and who should undertake such examinations^{11,12}. Whilst data from the USA indicate that paediatric black athletes are particularly susceptible to sudden cardiac death (SCD)¹³, there is a general lack of understanding as to which factors (e.g., physical growth, race and sex) have the potential to increase the likelihood of generating a false-positive diagnosis and unnecessary disqualification from

competitive sport. Consequently, the distinction between paediatric athlete's heart and cardiac pathology associated with SCD is especially important for this population.

Therefore, the primary aim of this systematic review and meta-analysis was to describe the electrocardiographic (ECG), structural and functional manifestations of the paediatric athlete's heart compared to that of age-matched non-athletes. The secondary aims were to determine the impact of an athlete's chronological age, race, and sex on cardiac remodelling responses to intensive competitive sport.

METHODS

This review was conducted and reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines¹⁴.

Literature Searching

A systematic search was conducted using six electronic databases; (1) MEDLINE, (2) PubMed, (3) EMBASE, (4) Web of Science, (5) CINAHL, and (6) SPORTDISCUS. Databases were searched from inception, to May 2016. Search terms were mapped to relevant MeSH terms or subject headings under four concepts:

- 1) 'Paediatric'
- 2) 'Athlete',
- 3) 'Electrocardiography', 'Echocardiography', 'Magnetic Resonance Imaging'

- 4) 'European Society of Cardiology Criteria', 'Seattle Criteria', 'Ventricle', 'Atrium' and 'Septum'.

Terms within each concept were combined with the Boolean operator 'OR', then concepts were combined with the 'AND' operator to, produce the search strategy (Supplementary Appendix A). To supplement the electronic database searching, we hand searched reference lists of eligible articles, ePublication lists of key journals, and undertook citation tracking using Google Scholar (Supplementary Appendix B). All identified articles were imported into Endnote X4 for application of selection criteria (Thomson Reuters, California, USA).

Selection Criteria

Titles and abstracts of potentially eligible articles were independently screened by two authors (GM and NR) against the selection criteria. For articles where it was not immediately clear from the title and/or abstract whether they should be included, we obtained the full text for independent screening. Discrepancies were resolved via consensus discussion, with a third reviewer (MW) consulted if consensus could not be reached.

Inclusion criteria were: 1) data reported for male and/or female competitive athletes, with or without comparison to non-athletes, 2) all participants were aged 6-18 years old at the time of assessment, and 3) an original research article published in English language. We defined a competitive athlete as:

“One who participates in an organised team or individual sport that requires regular competition against others as a central component, places a high premium on excellence and achievement, and requires some form of systematic (and usually intense) training”¹⁵.

Participants not meeting this definition were classified as non-athletes. Articles were limited to English-language owing to translation costs. Articles that did not document athlete age range were excluded because of the risk of including athletes >18 years. If ECG and/or echocardiographic outcome data were not reported, or if professional guidelines for data acquisition were not observed or cited, articles were also excluded.

Risk of Bias Assessment

We developed a 15-item risk of bias assessment checklist (Supplementary Appendix C), comprising items from Downs & Black’s ‘Assessment of Methodological Quality of Randomised and Non-Randomised Studies’ checklist¹⁶, and a previously published athletes heart meta-analysis checklist⁵. The purpose was to identify articles of low methodological quality that could bias results¹⁷; with articles achieving $\leq 50\%$ of total possible appraisal score, excluded from quantitative synthesis. Two reviewers (GM & NR) independently assessed all included articles. Discrepancies were resolved via consensus discussion and consistency was measured using an interclass correlation coefficient (ICC_{2,1}).

Data Extraction

All ECG and echocardiographic data were extracted by one reviewer using a predefined extraction form and reviewed by a second reviewer, with discrepancies resolved by consensus (Supplementary Appendix D). Data extraction included the calculated mean (SD) for continuous data and *n* for dichotomous data. If insufficient data were reported, corresponding authors were contacted to request additional data.

Data management

Demographics

Body surface area (BSA)¹⁸ was extracted or manually calculated from the height and body mass reported in individual articles.

ECG

The 2010 European Society of Cardiology (ESC) recommendations for interpretation of the 12-lead ECG in athletes were applied, dividing ECG patterns into Group 1 training-related and Group 2 training un-related patterns accordingly¹⁹. T-wave inversion (TWI) was classified if ≥ 1 mm and in ≥ 2 contiguous leads, localised as follows: anterior leads (V1–V3), extended anterior leads (V1–V4), inferior leads (Leads II–aVF), lateral leads (V5–V6/I–aVL) and infero-lateral leads (Leads II–aVF/V5–V6/I–aVL). Deep TWI was defined as a negative T wave ≥ 2 mm in ≥ 2 contiguous leads, (excluding leads III and aVR). ECG classification of left ventricular hypertrophy (LVH) was made according to the Sokolow–Lyon criteria²⁰. Early repolarisation was defined as ST segment

elevation (≥ 0.1 mV) and/or J point elevation manifested either as QRS slurring or notching, in ≥ 2 contiguous leads²¹.

Echocardiography

Two-dimensional echocardiography data, where the American Society of Echocardiography paediatric guidelines were followed, were extracted^{22,23}. On the basis of previous publications within the paediatric athlete's heart, participants with an LV wall thickness >12 mm were considered to have left ventricular hypertrophy (LVH)²⁴. Left ventricular mass (LVM) was calculated according to the formula of Devereux²⁵. Relative left ventricular wall thickness (RWT) was calculated and expressed as a fraction: $[\text{PWTd} + \text{IVSd}]/\text{LVIDd}$ posterior wall thickness during diastole (PWTd), interventricular septum thickness during diastole (IVSd) and left ventricular internal diameter during diastole (LVIDd). If IVSd was not reported, it was considered equal to PWTd².

Data synthesis

Data were analysed using StatsDirect (Altrincham, UK) and Stata V.12 (Stata Corp, College Station, Texas, USA). Demographic data were analysed using arithmetic means. Pooled dichotomous data were analysed using random-effects proportion meta-analyses (as we expected significant statistical heterogeneity), and presented as odds ratios (ORs) or risk ratios (RRs) as appropriate. We only pooled data for variables with a minimum of 3 articles reporting on the variable. Pooled

continuous data were presented as standardised mean differences (i.e. effect size). The magnitude of pooled standardised mean differences were interpreted according to Cohen's guidelines; with small medium and large effects interpreted as $\geq 20\%$, $\geq 50\%$ and $\geq 80\%$ respectively²⁶. A p-value of <0.05 was used to denote statistical significance.

Random-effects meta-regression (Kendall's non-parametric statistic) was utilised to explore and account for the impact of the covariates; age, race (black vs. Caucasian) and sex (Male vs. Female) upon ECG and echocardiographic variables. Random-effects meta-regression analysis was deemed inappropriate when <10 articles were available for synthesis²⁷.

Sub-analysis was used to explain the effect of the covariates; age (<14 years vs. ≥ 14 years), race (black vs. Caucasian), sex and where possible, the interaction of age (<14 years vs. ≥ 14 years) was explored within race and sex. With regards to maturational age, 14 years was set according to the attainment of selected development landmarks in boys (Mean age of peak height velocity = 14; Peak weight velocity = 14.3; Peak leg length velocity = 14.4; 90% of adult stature = 13.9; 95% of adult stature = 14.9; Genital stage IV = 14.6; and Pubic hair stage IV = 15.1)²⁸ and the onset of menarche within females (13.2 years)²⁹.

Data were combined as per Cochrane guidelines³⁰. If data were reported for the same participants in more than one article, the data were extracted from the article with the largest cohort size (with corresponding author's confirmation). If an article reported multiple follow-ups, data were extracted from the latest visit (i.e. longest follow-up). When standard deviation (SD)

was not reported, it was imputed from the average SD³⁰, only utilising articles containing ≥ 30 participants. To ensure results were not subsequently biased, sensitivity analysis was conducted omitting imputed SD data. Statistical heterogeneity was examined using the I^2 index³¹.

RESULTS

Literature Search

The literature search identified 2030 potentially eligible articles, of which 972 were duplicates. After application of the selection criteria, 43 articles remained for qualitative analysis and 40 remained for quantitative analysis (Figure 1).

<INSERT FIGURE 1 HERE>

Risk of bias assessment

There was substantial agreement (71% 95% CI (49-84))³² between the reviewers for the risk of bias assessment (Supplementary Appendix E). Most frequently, discrepancies occurred when assessing 'professional guidelines' and 'missing data' (77% 95% CI 61-68)). Risk of bias scores ranged from 4 to 13 out of a maximum possible score of 15. No articles reported 'power analysis' or 'intra-observer reliability', with non-athlete 'activity levels' poorly described in 44%. Three articles were excluded³³⁻³⁵ from quantitative synthesis due to low methodological quality.

Demographic data

Data from 14,278 athletes (mean age 13.8 ± 1.3 years [range: 6-18]) and 1,668 non-athletes (mean age 12.6 ± 0.6 years [7-18]) were extracted from 43 articles. There were no differences in age or BSA between paediatric athletes and non-athletes. Athletes competed in 30 different sports, with football (soccer) predominating (33%). There were more males and Caucasians, but proportionately distributed among both athletes and non-athletes. In 2 articles, sex was not reported^{35,36}. In 23 articles, race was not documented³³⁻⁵⁶ and in 29 articles maturational status was not reported^{24,33-46,49,50,53,55-67} (Table 1).

<INSERT TABLE 1 HERE >

Data management

Within the 40 articles that were quantitatively synthesised; two articles reported overlapping data from a group of 155 athletes^{58,62}, two articles reported overlapping data from a cohort of 158 athletes^{43,44} and two articles reported overlapping data from a cohort of 900 athletes^{24,68}. Four articles presented multiple follow-up data^{40,59,64,69}. Adjustments were made, to account for this in the meta-analysis (Supplementary Appendix F).

Electrocardiographic characteristics

Paediatric athlete vs. paediatric non-athlete

Paediatric athletes had a significantly longer PR interval, and a significantly greater frequency of sinus bradycardia, 1st °AV block, incomplete right bundle branch block (IRBBB), voltage criteria for LVH and early repolarisation when compared to paediatric non-athletes (Table 2). The

prevalence of TWI ≥ 1 mm was similar between athletes and non-athletes (6.7% vs. 5.9%). However, athletes were 12.7 times more likely to have deep TWI ≥ 2 mm in ≥ 2 contiguous leads (except leads III and aVR) than non-athletes (4.7% vs. 0.3%). Athletes were 1.4 times more likely to have anterior TWI (6.5% vs. 5.7%) and 1.5 times more likely to have extended anterior TWI (1.4% vs. 0.9%) than non-athletes. Whilst inferior (0.9%) and lateral (0.2%) TWI was present among athletes, these were not observed in non-athletes. Other training un-related ECG patterns suggestive of underlining cardiac pathology including ST segment depression, abnormal Q waves, complete bundle branch blocks and abnormal QTc measurements were rarely observed in athletes ($\leq 0.6\%$) and were not observed in non-athletes.

<INSERT TABLE 2 HERE>

Impact of paediatric athlete age

Paediatric athletes ≥ 14 years had a significantly longer QRS duration, and a significantly greater frequency of sinus bradycardia and voltage criteria for LVH than athletes < 14 years (Table 3). Athletes ≥ 14 years were 1.3 times more likely to have TWI than athletes < 14 years (6.9% vs 5.4%). Athletes < 14 years were 1.2 times more likely to have anterior TWI than athletes ≥ 14 years (6.7% vs. 5.4%). Athletes ≥ 14 years were 3.1 times more likely to have extended anterior TWI (1.7% vs. 0.5%), and 15.8 times more likely to have inferolateral TWI (2.5% vs. 0.1%) than athletes < 14 years.

<INSERT TABLE 3 HERE>

Impact of paediatric athlete race

Black paediatric athletes had a significantly greater frequency of sinus bradycardia, 1st °AV block, IRBBB, voltage criteria for LVH and early repolarisation compared to Caucasian athletes (Table 4). Black athletes were 4 times more likely to have TWIs (23.4% vs. 5.9%) and 2.6 times more likely to have deep TWIs (10.6 vs. 4.2%), than Caucasian athletes. Further, black athletes were 2.9 times more likely to have anterior TWI (12.2% vs. 4.2%), 36 times more likely to have extended anterior TWI (10.8% vs. 0.3%) and 6.5 times more likely to have inferolateral TWI (8.2% vs. 1.3%) than Caucasian athletes. Finally, black athletes were 5 times more likely to have abnormal Q waves (0.5% vs. 0.1%) and 2.9 times more likely to have LAE (5.7% vs. 2.0%) when compared to Caucasian athletes.

<INSERT TABLE 4 HERE>

Echocardiographic patterns

Paediatric athletes vs paediatric non-athletes

Athletes had a significantly greater LVIDd (+8.2%), LVID during systole (LVIDs) (+14.2%), IVSd (+12.9%), PWTd (+12.2%), relative wall thickness (RWT), (+5.6%) LV mass (LVM) (+27.6%), and left atrial diameter (LAD) (+12.3%) than non-athletes (Table 5). One percent of athletes (95% CI 0.3 - 2.3, 5 articles; n=4460) had LVH (LV wall thickness >12 mm). LVH was not observed in non-athletes. There were no significant differences in cardiac functional parameters between athletes and non-athletes. Using imputed SDs did not influence the results.

<INSERT TABLE 5 HERE>

Impact of age: paediatric athletes vs paediatric non-athletes

Age was a positive predictor of LVIDd, IVSd, PWTd, RWT and LVM in athletes and non-athletes ($P \leq 0.001$). After accounting for age, athletes had greater LVIDd, IVSd, RWT and LVM ($P \leq 0.05$) than non-athletes.

Impact of paediatric athlete age

Paediatric athletes ≥ 14 years had a significantly greater LVIDd (+13.5%), LVIDs (+15.9%), IVSd (+15.2%), PWTd (+21.3%), LVM (+38.7%), Aortic Root (+14.2%), and LAD (+15.6%) than athletes < 14 years (Table 6). With the exception of E/A ratio (+13.6% greater in athlete's ≥ 14 years) there were no statistical differences with regards to left ventricle function.

<INSERT TABLE 6 HERE>

Impact of paediatric athlete race

Black athletes had a significantly greater PWTd (+12.4%) and LAD (+13.4%) than Caucasian athletes (Table 7). Prevalence of LVH (LV wall thickness > 12 mm) was 17.1 times greater among black [2 articles, $n=319$] than Caucasian athletes [3 articles, $n=3318$] (7.1% vs. 0.4%).

<INSERT TABLE 7 HERE>

Impact of paediatric athlete sex

Male athletes had a significantly larger IVSd (+9.2%) than female athletes (Table 8). Prevalence of LVH was 2.6 times greater among male [5 articles; $n=4028$] than female athletes [2 articles; $n=432$] (1.2% vs. 0.4%).

<INSERT TABLE 8 HERE>

DISCUSSION

In the first systematic review and meta-analysis investigating the ECG, structural and functional manifestations of the paediatric athlete's heart, we found that 1) Paediatric athletes had a greater prevalence of training-related and training unrelated ECG changes than non-athletes, 2) Whilst the overall prevalence of TWI remained similar, the distribution and magnitude differed; 3) Paediatric athletes had larger echocardiographic derived LV dimensions than non-athletes, even after accounting for age; 4) Paediatric black athletes had increased levels of training and training unrelated ECG findings (particularly TWI); and finally 5) Paediatric black athletes had a greater prevalence of echocardiographic derived indices of LVH compared to Caucasian athletes.

ECG characteristics of the paediatric athlete

This study confirms that regular and prolonged physical training is associated with a high prevalence of bradycardia, repolarisation changes, atrial enlargement and ventricular hypertrophy in paediatric athletes⁷⁰. However, the magnitude, prevalence and distribution of such changes are dependent on the chronological age of the paediatric athlete. Similar to adult athletes, race impacted ECG remodelling in the paediatric athlete⁷¹. Black paediatric athletes had significantly more training-related changes, anterior, extended anterior, inferolateral and deep TWIs, in addition to Q waves and LAE compared to Caucasian athletes⁶.

T wave inversion in the paediatric athlete: impact of age and race

Inverted T-waves may represent the only sign of an inherited heart muscle disease even in the absence of any other features or before structural changes in the heart can be detected⁶³. Yet, until complete formation of adult ventricular mass, T wave inversions may persist across leads V1-V3 within the paediatric population, owing to right ventricular dominance⁷². Our findings on over 9000 paediatric athletes and over 800 paediatric non-athletes, support this notion, with a relatively high, but similar prevalence of anterior TWI (V1-V3) observed in both athletes and non-athletes (6.5% vs. 5.7%) respectively; suggesting this is a maturational trait largely not resultant upon athletic training. The slightly higher prevalence of anterior TWI in athletes vs. non-athletes also suggests that regular exercise may exacerbate or prolong the presence of juvenile TWI. Nevertheless, paediatric athletes were 12.7-times more likely to present with deep TWI ($\geq 2\text{mm}$) than non-athletes. Deep TWI ($\geq 2\text{mm}$) in the precordial leads are a major concern as these ECG alterations are a recognised manifestation of hypertrophic cardiomyopathy and arrhythmogenic right ventricular cardiomyopathy⁷³.

TWIs are uncommon among adult Caucasian athletes. Conversely, African/Afro–Caribbean black athletes have a higher prevalence of TWI, as well as more striking repolarisation changes and magnitude of voltage criteria for LVH than Caucasian athletes of similar age and size participating in identical sports^{74,75}. Similar to their adult counterparts⁷⁶, we found that black paediatric athletes are 4 times more likely to exhibit any TWI and 36 times more likely to exhibit extended anterior TWI (V1-V4) than Caucasian paediatric athletes⁶³; this likely represents an ethnic

response to physiological adaptation to exercise rather than an effect of race alone, exuberated by right ventricular dominance during pubertal years.

When is anterior T-Wave inversion normal?

Recently updated international consensus standards for 12-lead ECG interpretations in athletes^{77,78} recommends that TWI ≥ 1 mm in depth in two contiguous anterior leads (V2-V4) is abnormal (with the exception of TWI confined to leads V1-V4 in black athletes and leads V1-V3 in all athletes aged <16 years) and should prompt further evaluation for underlying structural heart disease. Our data support this recommendation, demonstrating a significantly reduced prevalence of anterior TWI (V1-V3) in athlete's ≥ 14 years likely as a consequence of maturation. Based on current evidence, TWI in the anterior leads (V1-V3) in paediatric athletes <14 years of age (or pre-pubertal athletes) should not prompt further evaluation in the absence of symptoms, signs or a family history of cardiac disease.

Our data also support the observation that like their adult counterparts, paediatric black athletes were 3 times more likely to have anterior TWI (V1-V3) and 36 times more likely to have extended anterior TWI (V1-V4) when compared to Caucasians. In adult black athletes, it is recognised that anterior TWI is a normal variant when preceded by J-point elevation and convex ST segment elevation⁷⁹, unlike in arrhythmogenic right ventricular cardiomyopathy where the J-point and/or ST segment is usually isoelectric or depressed prior to TWI. Appreciating the J-point and preceding ST segment may help differentiate between physiological adaptation and

cardiomyopathy in athletes with anterior TWI affecting leads V3 and/or V4, and may prove to be especially useful in athletes of mixed race. A recent study compared black and Caucasian healthy athletes against hypertrophic cardiomyopathy and arrhythmogenic right ventricular cardiomyopathy patients, all of whom had anterior TWI. Within athletes, the combination of J-point elevation $\geq 1\text{mm}$ and TWI confined to leads V1-V4 excluded hypertrophic cardiomyopathy or arrhythmogenic right ventricular cardiomyopathy with 100% negative predictive value, regardless of race⁷⁹. Conversely, anterior TWI associated with minimal or absent J-point elevation ($<1\text{ mm}$) may reflect a cardiomyopathy. Such detailed investigations have yet to be extended to the paediatric athletic population.

Inferior and/or lateral TWI warrants investigation

We were surprised by the high prevalence of inferolateral TWI in both black (8.5%) and Caucasian (1.3%) paediatric athletes. It is unlikely that all such athletes harbour a sinister cardiomyopathy and may represent a racial variant in black athletes. Despite this, lateral lead TWI should be viewed with caution. We recently investigated 155 athletes presenting with pathological TWI with clinical examination, ECG, echocardiography, exercise testing, 24h Holter ECG and cardiac magnetic resonance⁸⁰. Cardiac disease was established in 44.5% of athletes (81% hypertrophic cardiomyopathy). Inferior and/or lateral TWI were the most commonly observed ECG abnormalities (83.9%) and were largely isolated findings without other ECG abnormalities (43.2%). In our experience, regardless of an increased frequency after 14-years and a higher prevalence in adolescent black athletes, inferolateral TWI should be considered pathological in

all cases until proven otherwise. While exclusion from competitive sport is not warranted in the asymptomatic paediatric athlete without a family history of SCD and normal secondary examinations, annual follow-up is essential to ascertain possible disease expression.

Left ventricular morphology of the paediatric athlete

While most adult athletes have left ventricular structural changes that are considered physiological, there are a small proportion who develop pronounced morphological changes that overlap with phenotypic expressions of cardiac pathology associated with SCD. Several groups have produced algorithms to aid in this differentiation⁸¹⁻⁸³. Data for these algorithms primarily derives from five large echocardiographic studies^{75,84-87} examining 5053 elite, predominately male adult athletes; 134 (2.7%) had a maximal wall thickness ≥ 12 mm (of which 27 (0.5%) athletes had a maximum wall thickness of ≥ 13 mm). In absolute terms and regardless of an athlete's BSA, the upper limit of physiological hypertrophy for adult male athletes is considered ≥ 13 mm for maximal wall thickness and ≥ 65 mm for LVIDd.

Despite undergoing significant changes in anthropometry during maturation paediatric athletes have significantly larger cardiac diameters, wall thicknesses and LV mass than non-athletes even after adjusting for age. From 4460 paediatric athletes analysed, just 1.1% presented with a maximal wall thickness ≥ 12 mm; although a maximal wall thickness of 15mm was documented in one study. A pooled mean LVIDd of 47mm (<14 years: 44.2mm vs. ≥ 14 years: 51.1 mm) is similar to upper limits previously observed among paediatric hypertrophic cardiomyopathy

patients (48mm)⁸⁸. Thus, such adult upper limit criteria may not be applicable to the paediatric athlete. Regardless of race, values above these should be viewed with suspicion in paediatric athletes, particularly if the athlete also presents with cardiac symptoms, a family history of SCD and/or an abnormal ECG. Given the widely recognised impact of chronological age and somatic growth upon paediatric echocardiographic variables, it is our suggestion that Z scores (which account for the effects of body size and chronological age) are instead used for differential diagnosis when normative data are available^{89–91}, as previously suggested within paediatric specific echocardiographic guidelines²².

Impact of chronological age on LV remodelling

Cardiac enlargement increased with chronological age, as demonstrated by our meta-regression as well as by others⁹², and helps to explain the heterogeneity observed within this dataset. After accounting for age (using meta-regression), paediatric athletes had greater LV morphology than paediatric non-athletes, demonstrating the potent stimulus exercise has upon cardiac structure. These changes appeared to be exaggerated during the pubertal growth stage, suggesting a potential role of hormonal factors in cardiac remodelling⁹³. We recognise that whilst chronological age is a linear factor, growth and maturation are not⁹⁴, and thus maturational status for children of the same chronological age can differ dramatically^{95,96}. Yet, assessment of maturational status was conducted among only 14 of the 43 (33%) articles included for qualitative synthesis, and relied largely on assessment by Tanner Scale (79%), regarded to be inappropriate by many due to obvious child protection concerns. In our experience, clinical interpretation of cardiac PPE data

should be governed by biological age rather than chronological age. According to the International Olympic Committee consensus statement on youth athletic development⁹⁷, skeletal age is the most useful estimate of maturity status and can be used from childhood into late adolescence. However, this can only be confirmed by radiological hand-wrist imaging. Since this is not widely available in most cardiological units, alternative simple measures such as percentage of predicted mature (adult) height at the time of observation may provide an estimate of maturity status⁹⁸. However care is warranted, as 1) predicted mature (adult) height only demonstrates moderate concordance with classifications of maturity status, based on skeletal age^{99,100}, and 2) historical height data of the patient is required to rule out sudden growth spurts.

Impact of race on LV remodelling

Data from the USA indicate that paediatric black athletes are particularly susceptible to SCD¹³, and therefore, the distinction between athlete's heart and cardiac pathology is of particular relevance in this group. Consistent with previous observations in adults^{75,101,102}, we found that paediatric black athletes had increased LVH in response to chronic training loads compared to Caucasian athletes. This change is consistent with a concentric remodelling pattern. Furthermore, the likelihood of LVH was 17.1 times greater among black when compared to Caucasian athletes. We speculate that these ethnic-specific manifestations of the athlete's heart are the result of hemodynamic influences; specifically greater peripheral vascular resistance and a smaller nocturnal decline in BP¹⁰⁰.

Impact of sex on LV remodelling

The last three decades have witnessed an exponential rise in the number of females participating in high-level competitive sport¹⁰⁵. Consistent with observations among adults⁴, we found a reduced LVH response to chronic training loads in female athletes compared to males. This might be due to hormonal differences and lower testosterone concentrations¹⁰⁶. However, the relative differences of sex across maturational years has yet to be fully elucidated among paediatric athletes. Females reach complete pubertal development at an earlier chronological age and thus we may expect such relative differences between female and male athletes to be smaller during the early stages of pubertal development.

Limitations

A high statistical heterogeneity (I^2) was observed; this may be because it was not possible to stratify data according to biological age, race or sex due to inconsistent methodology and designs implemented within the observational studies included. Because of this, a random-effects meta-analysis model was adopted to provide a more conservative pooled estimate. Activity levels of our non-athlete cohort are unknown and thus they may not actually be sedentary, however, in all cases, participants did not meet classification criteria for a competitive athlete.

Whilst we utilised the 2010 ESC recommendations for interpretation of the 12-lead ECG¹⁹, at the time of publication, it was not intended to be used in athletes ≤ 12 years old. We recently observed that the 2014 'Refined Criteria' for ECG interpretation in athletes outperformed both the 2013 Seattle Criteria and the 2010 ESC recommendations, by significantly reducing the number of false-positive ECGs in Arabic, Black and Caucasian adult athletes while maintaining 100% sensitivity for serious cardiac pathologies¹⁰⁷. Again, however, all three ECG criteria are only applicable for adult athletes and not paediatric athletes. Thus for paediatric cardiac PPEs, the attending cardiologist or sports medicine physician is left with the conundrum of which criteria should be used for ECG interpretation. Recently published International consensus standards for ECG interpretation in athletes⁷⁷ do account for age and race respectively. TWI in the anterior leads (V1-V3) in adolescent athletes < 16 years of age (or pre-pubertal athletes) and black adult athletes with J-point elevation and convex ST segment elevation followed by TWI in V2-V4, would now not prompt further evaluation in the absence of symptoms, signs or a family history of cardiac disease. But in most non-black athletes age ≥ 16 years, anterior TWI beyond lead V2 would prompt further evaluation given the potential overlap with arrhythmogenic right ventricular cardiomyopathy.

Finally, echocardiographic data were largely limited to left ventricle structural variables, owing to insufficient data available for synthesis. Such limitations highlight the importance of further research in the paediatric athlete extending to other chambers of the heart, and beyond load dependent measurements of cardiac function (ejection fraction or fractional shortening) towards Tissue Doppler imaging and myocardial deformation (strain) imaging.

Conclusion

Similar to adult athletes, paediatric athletes had a greater prevalence of training related and training unrelated ECG changes than non-athletes. Significant cardiac remodelling in paediatric athlete occurs both before and during their 'maturational years'; with race and sex significantly impacting upon the pattern of remodelling observed. The results demonstrate the importance of adjusting for age when assessing LV morphology in paediatric athletes, whilst consideration for an athletes' race and sex is further required when differentiating between physiological and pathological cardiac remodelling.

Contributions: GM, NRR, CLA and MGW contributed to the conception and design of the review. GM applied the search strategy. GM, NRR and MGW applied the selection criteria. GM and NRR completed the assessment of risk of bias. GM, CLA and AF analysed the data. All authors contributed to data interpretation. GM and MGW wrote the manuscript. NRR, CLA, AF, GEP, VW, CA, KPG and DO critically revised the manuscript for important intellectual content. GM and MGW are responsible for the overall content as guarantors.

Competing interests: None.

Provenance and peer review: Not commissioned; externally peer reviewed.

REFERENCES

1. Drezner J a, Ackerman MJ, Anderson J, et al. Electrocardiographic interpretation in athletes: the “Seattle criteria”. *Br J Sports Med.* 2013;47(3):122-124.
2. Pluim BM, Zwinderman a H, van der Laarse a, van der Wall EE. The athlete’s heart. A meta-analysis of cardiac structure and function. *Circulation.* 2000;101(3):336-344.
3. Sheikh N, Papadakis M, Ghani S, et al. Comparison of electrocardiographic criteria for the detection of cardiac abnormalities in elite black and white athletes. *Circulation.* 2014;129(16):1637-1649.
4. Pelliccia A, Maron BJ, Culasso F, Spataro A, Caselli G. Athlete’s Heart in Women Elite Female Athletes. *J Am Med Assoc.* 1996;276(3):211-215.
5. Utomi V, Oxborough D, Whyte GP, et al. Systematic review and meta-analysis of training mode, imaging modality and body size influences on the morphology and function of the male athlete’s heart. *Heart.* 2013;99(23):1727-1733.
6. Papadakis M, Wilson MG, Ghani S, Kervio G, Carre F, Sharma S. Impact of ethnicity upon cardiovascular adaptation in competitive athletes: relevance to preparticipation screening. *Br J Sports Med.* 2012;46(Suppl_1):i22-i28.
7. Whyte GP, George K, Nevill A, Shave R, Sharma S, McKenna WJ. Left ventricular morphology and function in female athletes: A meta-analysis. *Int J Sports Med.* 2004;25(5):380-383.
8. Bergeron MF, Mountjoy M, Armstrong N, et al. International Olympic Committee

- consensus statement on youth athletic development. *Br J Sports Med.* 2015;49(13):843-851.
9. Mountjoy M, Bergeron M. Youth athletic development: aiming high while keeping it healthy, balanced and fun! *Br J Sports Med.* 2015;49(13):841-842.
 10. Mountjoy M, Rhind DJ a, Tiivas a, Leglise M. Safeguarding the child athlete in sport: a review, a framework and recommendations for the IOC youth athlete development model. *Br J Sports Med.* 2015;49(13):883-886.
 11. Vetter VL. Electrocardiographic Screening of All Infants, Children, and Teenagers Should Be Performed. *Circulation.* 2014;130(8):688-697.
 12. Friedman R a. Electrocardiographic Screening Should Not Be Implemented for Children and Adolescents Between Ages 1 and 19 in the United States. *Circulation.* 2014;130(8):698-702.
 13. Harmon KG, Asif IM, Maleszewski JJ, et al. Incidence and Etiology of Sudden Cardiac Arrest and Death in High School Athletes in the United States. *Mayo Clin Proc.* 2016;91(11):1493-1502.
 14. Moher D, Liberati a, Tetzlaff J, Altman DG, Grp P. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement (Reprinted from *Annals of Internal Medicine*). *Phys Ther.* 2009;89(9):873-880.
 15. Maron BJ, Zipes DP. Introduction: Eligibility recommendations for competitive athletes with cardiovascular abnormalities - General considerations. *J Am Coll Cardiol.*

2005;45(8):1318-1321.

16. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health*. 1998;52(6):377-384.
17. van Tulder MW, Suttorp M, Morton S, Bouter LM, Shekelle P. Empirical evidence of an association between internal validity and effect size in randomized controlled trials of low-back pain. *Spine (Phila Pa 1976)*. 2009;34(16):1685-1692.
18. Du Bois, D., Du Bois EF. A formula to estimate the approximate surface area if height and weight be known. *Nutrition*. 1989;50(5):303.
19. Corrado D, Pelliccia A, Heidbuchel H, et al. Recommendations for interpretation of 12-lead electrocardiogram in the athlete. *Eur Heart J*. 2010;31(2):243-259.
20. Sokolow M, Lyon T. The ventricular complex in left ventricular hypertrophy as obtained by unipolar precordial and limb leads. *Am Heart J*. 1949;37(2):161-186.
21. Miyazaki S, Shah AJ, Haïssaguerre M. Early Repolarization Syndrome. *Circ J*. 2010;74(10):2039-2044.
22. Lopez L, Colan SD, Frommelt PC, et al. Recommendations for Quantification Methods During the Performance of a Pediatric Echocardiogram: A Report From the Pediatric Measurements Writing Group of the American Society of Echocardiography Pediatric and Congenital Heart Disease Council. *J Am Soc Echocardiogr*. 2010;23(5):465-495.
23. Lai WW, Geva T, Shirali GS, et al. Guidelines and Standards for Performance of a Pediatric

- Echocardiogram: A Report from the Task Force of the Pediatric Council of the American Society of Echocardiography. *J Am Soc Echocardiogr.* 2006;19(12):1413-1430.
24. Sharma S, Maron BJ, Whyte G, Firoozi S, Elliott PM, Mckenna WJ. Physiologic Limits of Left Ventricular Hypertrophy in Elite Junior Athletes : Relevance to Differential Diagnosis of Athlete's Heart and Hypertrophic Cardiomyopathy. *J Am Coll Cardiol.* 2002;40(8):1431-1436.
 25. Devereux RB, Alonso DR, Lutas EM, et al. Echocardiographic assessment of left ventricular hypertrophy: comparison to necropsy findings. *Am J Cardiol.* 1986;57(6):450-458.
 26. Cohen J. *Statistical power analysis for the behavioral sciences.* 2nd edn. Hillsdale, NJ: Erlbaum, 1988.
 27. Thompson AM, Baxter-Jones ADG. Endurance training in young female athletes. *Sports Med Arthrosc.* 2002;10(1):33-41.
 28. Bielicki T, Koniarek J, Malina RM. Interrelationships among certain measures of growth and maturation rate in boys during adolescence. *Ann Hum Biol.* 1984;11(3):201-210.
 29. Wellens R, Malina RM, Beunen G, et al. Age at menarche in Flemish girls: current status and secular change in the 20th century. *Ann Hum Biol.* 1990;17(2):145-152.
 30. Higgins JP. *Cochrane Handbook for Systematic Reviews of Interventions.*; 2008.
 31. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ Br Med J.* 2003;327(7414):557-560.

32. Viera AJ, Garrett JM. Understanding interobserver agreement: The kappa statistic. *Fam Med*. 2005;37(5):360-363.
33. Attisani G, Faiola F, Luciani U, Bianchi G, Veicstenas A, Casasco M. Negative T waves in right precordial leads in pre-adolescent subjects . A personal experience. *Medicina Dello Sport*. 2011;64(4):423-434.
34. Morales MC. LV mass in adolescent basketball players. *Cardiovasc Rev reports*. 1992;13(10):60-62.
35. Dinu V, Dutsi S, Lucia C, Gusti A. Aspects of morphofunctional and cardiovascular.pdf. *Arch Balk Med Union*. 2010;45(4):282-285.
36. Medved R, Fabecic-Sabadi V, Medved V. Echocardiographic Findings in Children Participating in Swimming Training. *Int J Sports Med*. 1986;7(2):94-99.
37. Hauser M, Petzuch K, Kühn A, et al. The Munich Triathlon Heart Study: ventricular function, myocardial velocities, and two-dimensional strain in healthy children before and after endurance stress. *Pediatr Cardiol*. 2013;34(3):576-582.
38. Hoogsteen J, Hoogeveen A, Schaffers H, Wijn PFF, Van Der Wall EE. Left atrial and ventricular dimensions in highly trained cyclists. *Int J Cardiovasc Imaging*. 2003;19(3):211-217.
39. Koch S, Cassel M, Linne K, Mayer F, Scharhag J. ECG and echocardiographic findings in 10-15-year-old elite athletes. *Eur J Prev Cardiol*. 2012;21(6):774-781.
40. Meško D, Jurko A, Vrlík M, Novomeská M, Horniak E, Dzurenková D. Development of the

left ventricular hypertrophy and dilation in adolescent ice hockey players evaluated with echocardiography. *Sport Med Train Rehabil.* 1993;4(3):177-188.

41. Obert P, Stecken F, Courteix D, Lecoq a M, Guenon P. Effect of long-term intensive endurance training on left ventricular structure and diastolic function in prepubertal children. *Int J Sports Med.* 1998;19(2):149-154.
42. Ozer S, Cil E, Baltaci G, Ergun N, Ozme S. Left Ventrtricular Structure and Function by Echocardiography in Childhood Swimmers. *Jpn Heart J.* 1994;35(3):295-300.
43. Pelà G, Li Calzi M, Crocamo A, et al. Ethnicity-related variations of left ventricular remodeling in adolescent amateur football players. *Scand J Med Sci Sports.* 2014;25(3):382-389.
44. Pela G, Crocamo A, Li Calzi M, et al. Sex-related differences in left ventricular structure in early adolescent non-professional athletes. *Eur J Prev Cardiol.* 2015;23(7):777-784.
45. Rowland TW, Delaney BC, Siconolfi SF, Rowland, Thomas W.Delaney, Brian C.Siconolfi SF. "Athlete"s heart' in prepubertal children. *Pediatrics.* 1987;79(5):800-804.
46. Rowland TW, Unnithan VB, MacFarlane NG, Gibson NG, Paton JY. Clinical manifestations of the "athlete"s heart' in prepubertal male runners. *Int J Sports Med.* 1994;15(8):515-519.
47. Rowland T, Goff D, DeLuca P, Popowski B. Cardiac effects of a competitive road race in trained child runners. *Pediatrics.* 1997;100(3):E2.
48. Rowland T, Wehnert M, Miller K. Cardiac responses to exercise in competitive child

- cyclists. *Am Coll Sport Med.* 2000;32(4):747-752.
49. Shi JR, Selig S. Cardiac Structure and Function in Young Endurance Athletes and Nonathletes. *J Exerc Sci Fit.* 2005;3(2):74-80.
 50. Sundberg S, Elovainio R. Resting ECG in athletic and non-athletic adolescent boys: correlations with heart volume and cardiorespiratory fitness. *Clin Physiol.* 1982;2(5):419-426.
 51. Telford RD, McDonald IG, Ellis LB, Chennells MH, Sandstrom ER, Fuller PJ. Echocardiographic dimensions in trained and untrained 12-year-old boys and girls. *J Sport Sci.* 1988;6(1):49-57.
 52. Valente-Dos-Santos J, Coelho-e-Silva MJ, Vaz V, et al. Ventricular mass in relation to body size, composition, and skeletal age in adolescent athletes. *Clin J Sport Med.* 2013;23(4):293-299.
 53. Madeira RB, Trabulo M, Alves F, Pereira JG. Effects of chronic exercise training on left ventricular dimensions and function in young athletes. *Port J Cardiol an Off J Port Soc Cardiol.* 2008;27(7-8):909-922.
 54. Menafoglio a., Di Valentino M, Porretta a. P, et al. Cardiovascular evaluation of middle-aged individuals engaged in high-intensity sport activities: implications for workload, yield and economic costs. *Br J Sports Med.* 2014;49(11):757-761.
 55. Yildirim Ş, Binnetoğlu FK, Battal F, Aylanç H, Kaymaz N, Tekin M. Relation between QT Variables and Left Ventricular Geometry in Athletes and Obese Children. *Acta Med Port.*

2016;29(2):95-100.

56. Bartkevičienė A. Echocardiographic characteristics of left ventricular geometry of 12-17 years athletes. *Heal Sci.* 2015;25(4):65-72.
57. Bessem B, de Bruijn MC, Nieuwland W. The ECG of high-level junior soccer players: comparing the ESC vs. the Seattle criteria. *Br J Sports Med.* 2014;49(15):1000-1006.
58. Di Paolo FM, Schmied C, Zerguini Y a., et al. The Athlete's heart in adolescent Africans: An electrocardiographic and echocardiographic study. *J Am Coll Cardiol.* 2012;59(11):1029-1036.
59. Kinoshita N, Katsukawa F, Yamazaki H. Modeling of Longitudinal Changes in Left Ventricular Dimensions among Female Adolescent Runners. *PLoS One.* 2015;10(10):e0140573.
60. Konopka M, Banach M, Burkhard-Jagodzińska K, et al. Echocardiographic evaluation of cardiovascular system in adolescent athletes in view of physiological adaptation to physical training. *Folia Cardiol.* 2015;10(4):233-241.
61. Petridis L, Kneffel Z, Kispéter Z, Horváth P, Sidó Z, Pavlik G. Echocardiographic characteristics in adolescent junior male athletes of different sport events. *Acta Physiol Hung.* 2004;91(2):99-109.
62. Schmied C, Zerguini Y, Junge A, et al. Cardiac findings in the precompetition medical assessment of football players participating in the 2009 African Under-17 Championships in Algeria. *Br J Sports Med.* 2009;43(9):716-721.

63. Sheikh N, Papadakis M, Carre F, et al. Cardiac adaptation to exercise in adolescent athletes of African ethnicity: an emergent elite athletic population. *Br J Sports Med.* 2013;47(9):585-592.
64. Stoner JE. Cardiac dimensions gymnasts and swimmers. *Biol Sport.* 1997;14(2):115-125.
65. Vasiliauskas D, Venckuna T, Marcinkevicius J, Bartkeviciene A. Development of structural cardiac adaptation in basketball players. *Eur J Prev Cardiol.* 2006;13(6):985-989.
66. Zdravkovic M, Perunicic J, Krotin M, et al. Echocardiographic study of early left ventricular remodeling in highly trained preadolescent footballers. *J Sci Med Sport.* 2010;13(6):602-606.
67. Agrebi B, Tkatchuk V, Hlila N, Mouelhi E, Belhani A. Impact of specific training and competition on myocardial structure and function in different age ranges of male handball players. *PLoS One.* 2015;10(12):e0143609.
68. Makan J, Sharma S, Firoozi S, Whyte G, Jackson PG, McKenna WJ. Physiological upper limits of ventricular cavity size in highly trained adolescent athletes. *Heart.* 2005;91(4):495-499.
69. Csajági E, Szauder I, Major Z, Pavlik G. Left Ventricular Morphology in Different Periods of the Training Season in Elite Young Swimmers. *Pediatr Exerc Sci.* 2015;27(2):185-191.
70. Sharma S, Whyte G, Elliott P, et al. Electrocardiographic changes in 1000 highly trained junior elite athletes. *Br J Sports Med.* 1999;33(5):319-324.
71. Papadakis M, Carre F, Kervio G, et al. The prevalence, distribution, and clinical outcomes

- of electrocardiographic repolarization patterns in male athletes of African/Afro-Caribbean origin. *Eur Heart J*. 2011;32(18):2304-2313.
72. Molinari G, Brunetti ND, Biasco L, et al. Electrocardiograms of Children and Adolescents Practicing Non-competitive Sports: Normal Limits and Abnormal Findings in a Large European Cohort Evaluated by Telecardiology. *Sport Med*. 2016:1-9.
 73. Drezner J a, Ackerman MJ, Cannon BC, et al. Abnormal electrocardiographic findings in athletes: recognising changes suggestive of cardiomyopathy. *Br J Sports Med*. 2013;47(3):137-152.
 74. Magalski A, Maron BJ, Main ML, et al. Relation of Race to Electrocardiographic Patterns in Elite American Football Players. *J Am Coll Cardiol*. 2008;51(23):2250-2255.
 75. Basavarajaiah S, Boraita A, Whyte G, et al. Ethnic Differences in Left Ventricular Remodeling in Highly-Trained Athletes. Relevance to Differentiating Physiologic Left Ventricular Hypertrophy From Hypertrophic Cardiomyopathy. *J Am Coll Cardiol*. 2008;51(23):2256-2262.
 76. Papadakis M, Carre F, Kervio G, et al. The prevalence, distribution, and clinical outcomes of electrocardiographic repolarization patterns in male athletes of African/Afro-Caribbean origin. *Eur Heart J*. 2011;32(18):2304-2313.
 77. Sharma S, Drezner J, Baggish A, et al. International Consensus Standards for Electrocardiographic Interpretation in Athletes. *Eur Heart J*. 2017.
 78. Drezner J, Sharma S, Baggish A, et al. International Criteria for Electrocardiographic

- Interpretation in Athletes. *Br J Sports Med*. 2017.
79. Calore C, Zorzi A, Sheikh N, et al. Electrocardiographic anterior T-wave inversion in athletes of different ethnicities: differential diagnosis between athlete's heart and cardiomyopathy. *Eur Heart J*. 2015.
 80. Schnell F, Riding N, O'Hanlon R, et al. Recognition and Significance of Pathological T-Wave Inversions in Athletes. *Circulation*. 2014;131(2):165-173.
 81. Caselli S, Maron MS, Urbano-Moral J a, Pandian NG, Maron BJ, Pelliccia A. Differentiating left ventricular hypertrophy in athletes from that in patients with hypertrophic cardiomyopathy. *Am J Cardiol*. 2014;114(9):1383-1389.
 82. Sheikh N, Papadakis M, Schnell F, et al. Clinical profile of athletes with hypertrophic cardiomyopathy. *Circ Cardiovasc Imaging*. 2015;8(7).
 83. Rudski LG, Lai WW, Afilalo J, et al. Guidelines for the Echocardiographic Assessment of the Right Heart in Adults: A Report from the American Society of Echocardiography. *J Am Soc Echocardiogr*. 2010;23(7):685-713.
 84. Pelliccia A, Maron BJ, Spataro A, Proschan MA, Spirito P. The Upper limit of physiological cardiac hypertrophy in highly trained elite athletes. *N Engl J Med*. 1991;5:295-301.
 85. Whyte GP, George K, Sharma S, et al. The upper limit of physiological cardiac hypertrophy in elite male and female athletes: the British experience. *Eur J Appl Physiol*. 2004;92(4-5):592-597.
 86. Sun B, Ma JZ, Yong YH, Lv YY. The upper limit of physiological cardiac hypertrophy in elite

- male and female athletes in China. *Eur J Appl Physiol*. 2007;101(4):457-463.
87. Basavarajaiah S, Wilson M, Whyte G, Shah A, McKenna W, Sharma S. Prevalence of hypertrophic cardiomyopathy in highly trained athletes: relevance to pre-participation screening. *J Am Coll Cardiol*. 2008;51(10):1033-1039.
88. Maron BJ, Casey S a, Poliac LC, Gohman TE, Almquist a K, Aeppli DM. Clinical course of hypertrophic cardiomyopathy in a regional United States cohort. *JAMA*. 1999;281(7):650-655.
89. Kampmann C, Wiethoff CM, Wenzel A, et al. Normal values of M mode echocardiographic measurements of more than 2000 healthy infants and children in central Europe. *Heart*. 2000;83(6):667-672.
90. Zilberman M V., Khoury PR, Kimball RT. Two-dimensional echocardiographic valve measurements in healthy children: Gender-specific differences. *Pediatr Cardiol*. 2005;26(4):356-360.
91. Daubeney PE, Blackstone EH, Weintraub RG, Slavik Z, Scanlon J, Webber S a. Relationship of the dimension of cardiac structures to body size: an echocardiographic study in normal infants and children. *Cardiol Young*. 1999;9(4):402-410.
92. George K, Sharma S, Batterham A, Whyte G, McKenna W. Allometric analysis of the association between cardiac dimensions and body size variables in 464 junior athletes. *Clin Sci*. 2001;100(1):47-54.
93. Cavasin MA, Sankey SS, Yu A-L, Menon S, Yang X-P. Estrogen and testosterone have

- opposing effects on chronic cardiac remodeling and function in mice with myocardial infarction. *Am J Physiol Heart Circ Physiol*. 2003;284(5):H1560-H1569.
94. Malina RM, Eisenmann JC, Cumming SP, Ribeiro B, Aroso J. Maturity-associated variation in the growth and functional capacities of youth football (soccer) players 13-15 years. *Eur J Appl Physiol*. 2004;91(5-6):555-562.
 95. Cox LA. The biology of bone maturation and ageing. *Acta Paediatr Suppl*. 1997;423(3):107-108.
 96. Mirwald RL, Baxter-Jones ADG, Bailey D a, Beunen GP. An assessment of maturity from anthropometric measurements. *Med Sci Sports Exerc*. 2002;34(4):689-694.
 97. Engebretsen L, Steffen K, Bahr R, et al. The International Olympic Committee Consensus statement on age determination in high-level young athletes. *Br J Sports Med*. 2010;44(7):476-484.
 98. Roche A, Tyleshevski F, Rogers E. Non-invasive measurement of physical maturity in children. *Res Q Exerc Sport*. 1983;54(4):364-371.
 99. Malina RM, Dompier TP, Powell JW, Barron MJ, Moore MT. Validation of a Noninvasive Maturity Estimate Relative to Skeletal Age in Youth Football Players Validation of a Noninvasive Maturity Estimate Relative to Skeletal Age in Youth Football Players. *Clin J Sport Med*. 2007;17(5):362-368.
 100. Malina RM, Silva MJCE, Figueiredo AJ, et al. Interrelationships among invasive and non-invasive indicators of biological maturation in adolescent male soccer players. *J Sport Sci*.

2012;30(15):1705-1717.

101. Rawlins J, Carre F, Kervio G, et al. Ethnic differences in physiological cardiac adaptation to intense physical exercise in highly trained female athletes. *Circulation*. 2010;121(9):1078-1085.
102. Kervio G, Pelliccia A, Nagashima J, et al. Alterations in echocardiographic and electrocardiographic features in Japanese professional soccer players: comparison to African-Caucasian ethnicities. *Eur J Prev Cardiol*. 2013;20(5):880-888.
103. Heffernan KS, Jae SY, Wilund KR, Woods JA, Fernhall B. Racial differences in central blood pressure and vascular function in young men. *Am J Physiol Circ Physiol*. 2008;295(6):h2380-h2387. .
104. DeLoach SS, Daskalakis C, Gidding S, Falkner B. Central blood pressures are associated with left ventricular mass index among African-American adolescents. *Am J Hypertens*. 2012;25(1):41-45.
105. International Olympic Committee. *Factsheet: Women in the Olympic Movement – Update May 2014*. International Olympic Committee, Lausanne, Switzerland (2014).
106. McGill Jr HC, Sheridan PJ. Nuclear uptake of sex steroid hormones in the cardiovascular system of the baboon. *Circ Res*. 1981;48(2):238-244.
107. Riding NR, Sheikh N, Adamuz C, et al. Comparison of three current sets of electrocardiographic interpretation criteria for use in screening athletes. *Heart*. 2014;101(5):384-390.

108. Ayabakan C, Akalin F, Mengütay S, Cotuk B, Odabas I, Ozüak A. Athlete's heart in prepubertal male swimmers. *Cardiol Young*. 2006;16(1):61-66.
109. Calò L, Sperandii F, Martino A, et al. Echocardiographic findings in 2261 peri-pubertal athletes with or without inverted T waves at electrocardiogram. *Heart*. 2015;101:193-200.
110. Migliore F, Zorzi A, Michieli P, et al. Prevalence of cardiomyopathy in italian asymptomatic children with electrocardiographic T-wave inversion at preparticipation screening. *Circulation*. 2012;125(3):529-538.
111. Papadakis M, Basavarajaiah S, Rawlins J, et al. Prevalence and significance of T-wave inversions in predominantly Caucasian adolescent athletes. *Eur Heart J*. 2009;30(14):1728-1735.
112. Pavlik G, Olexó Z, Osváth P, Sidó Z, Frenkl R. Echocardiographic characteristics of male athletes of different age. *Br J Sports Med*. 2001;35(2):95-99.

Table 1 Article characteristics

Author, year	Athletes						Non-Athletes					Outcome data	
	n	Chronological age Mean (range)	Biological age	Race (C:B:O)	Sex (M:F)	Sport	n	Chronological age Mean (range)	Biological age	Race (C:B:O)	Sex (M:F)	ECG	ECHO
Agrebi 2015 ⁶⁷	24	13.9 (11-17)	NR	0:0:24	24:0	Handball							
Attisani 2011 ³³	1865	13.7 (6-18)	NR	NR	1865:0	Soccer/Gymnastics						Y	N
Ayabakan 2006 ¹⁰⁸	22	11.0 (9-12)	Prepubertal	22:0:0	22:0	Swimming	21	10.7 (9-12)	Prepubertal	21:0:0	21:0	N	Y
Bartkevičienė 2015 ⁵⁶	167	14.8 (12-17)	NR	NR	167:0							N	Y
Bessem 2014 ⁵⁷	193	14 (10-19)	NR	134:29:30	193:0	Soccer						Y	N
Calò 2015 ¹⁰⁹	2261	12.4 (8-18)	Peripubertal	2261:0:0	2261:0	Soccer						Y	Y
Csajági 2015 ⁶⁹	18	13.7 (13-15)	Mid pubertal	18:0:0	8:7	Swimming	15	13.8 (13-15)	Mid pubertal	15:0:0	8:7	N	Y
Di Paolo 2012 ⁵⁸	216	16.1 (14-18)	NR	63:153:0	216:0	Soccer						Y	Y
Dinu 2010 ³⁵	40	12.7 (10-17)	NR	NR	NS	Athletics						N	Y
Hauser 2013 ³⁷	26	12.6 (7-17)	NR	NR	18:8	Triathlon						N	Y
Hoogsteen 2003 ³⁸	66	17.5 (17-18)	NR	NR	66:0	Cycling						N	Y
Kinoshita 2015 ⁵⁹	34	16.5 (16-17)	NR	0:0:34	0:34	Middle / long-distance runners						N	Y
Koch 2012 ³⁹	343	13 (10-15)	NR	NS	189:154	High school athletes						Y	Y
Konopka 2015 ⁶⁰	78	14.3 (12-17)	NR	78:0:0	64:14	Soccer, Tennis, Rowing.						N	Y
Madeira 2008 ⁵³	21	15.9 (15-16)	NR	NR	21:0	Soccer; Swimming						N	Y
Makan 2005 ⁶⁸	900	15.7 (14-18)	Post pubertal	882:0:18	693:207	10 sporting disciplines (Invasion games/ racket/ endurance/combat)	250	15.5 (14-18)	Post pubertal	NR	177:48	N	Y
Medved 1986 ³⁶	72	10 (8-14)	NR	NR	NR	Swimming	72	10 (8-14)	NR	NR	NS	N	Y
Meško 1993 ⁴⁰	23	14.5 (14-15)	NR	NR	23:0	Hockey	17	14.5 (14-15)	NR	NR	17:0	N	Y
Migliore 2012 ¹¹⁰	2765	13.9 (8-18)	Peri pubertal	2765:0:0	1914:851	18 sporting disciplines (Invasion games/ gymnastics/winter sports/horse-riding/racket/ endurance/combat)						Y	N
Moarles 1992 ³⁴	9	16.2 (14-17)	NR	NR	9:0	Basketball						N	Y
Obert 1998 ⁴¹	10	10.7 (10-11)	Pre pubertal	NR	4:6	Swimming	11	10.9 (10-11)	Prepubertal	NR	4:7	N	Y
Ozer 1994 ⁴²	82	11.2 (7-14)	NR	NR	41:41	Swimming	41	10.8 (7-15)	NS	NR	22:19	N	Y

Papadakis 2009 ¹¹¹	1710	16 (14-18)	Post-pubertal	1642:0:0	1414:291	15 sporting disciplines (Invasion games/ racket/ endurance/ combat)	400	16 (14-18)	Post pubertal	385:0:0	330:70	Y	N
Pavlik 2001 ¹¹²	165	14.7 (10-18)	NR	165:0:0	165:0	7 sporting disciplines (Endurance/ invasion games/ weightlifting)	22	14.7 (10-18)	NR	22:0:0	22:0	N	Y
Pelà 2014 ⁴³	138	14.3 (11-17)	NR	96:42:0	138:0	Soccer						Y	Y
Pelà 2015 ⁴⁴	206	13.8 (11-17)	NR	206:0:0	158:48	Soccer						Y	Y
Petridis 2004 ⁶¹	137	16.6 (15-18)	NR	NS	137:0	Swimming						N	Y
Rowland 1987 ⁴⁵	14	11 (8-14)	Prepubertal	NR	14:0	Swimming	19	10.4 (8-13)	Prepubertal	NR	19:0	Y	Y
Rowland 1994 ⁴⁶	10	12.2 (11-13)	Prepubertal	NR	10:0	Middle distance runners	18	11.3 (10-14)	Pre pubertal	NR	18:0	Y	Y
Rowland 1997 ⁴⁷	9	12.2 (9-15)	Early pubertal	NR	9:0	Cyclists						N	Y
Rowland 2000 ⁴⁸	8	11.9 (10-13)	Early pubertal	NS	8:0	Cyclists & Triathletes	39	12.2 (10-13)	Early pubertal	NR	39:0	N	Y
Schmied 2009 ⁶²	155	16.4 (14-17)	NR	0:155:0	155:0	Soccer						Y	Y
Sharma 1999 ⁷⁰	1000	15.7 (14-18)	Post-pubertal	998:8:4	730:180	9 Sporting disciplines (Invasion games/ racket/ endurance/ combat)	300	15.6 (14-18)	Post pubertal	293:0:7	210:90	Y	N
Sharma 2002 ²⁴	720	15.7 (14-18)	NR	706:14:0	540:180	10 Sporting disciplines (Invasion games/ racket/ endurance/ combat)						N	Y
Sheikh 2013 ⁶³	1232	16.4 (14-18)	NR	903:329:0	980:252	Swimming/Athletics	134	15.3 (14-18)	NR	0:134:0	88:46	Y	Y
Shi & Selig 2005 ⁴⁹	13	15.3 (14-16)	NR	NR	13:0	Gymnastics/Swimmers						N	Y
Stoner 1997 ⁶⁴	37	9.9 (7-11)	NR	NR	0:37	Athletics	22	9.1 (7-11)	NR	NR	0:22	N	Y
Sundberg & Elovainio 1982 ⁵⁰	59	13.7 (10-17)	NR	NR	59:0	Athletics	81	13.9 (10-17)	NR	NR	81:0	Y	N
Telford 1988 ⁵¹	85	11.9 (11-12)	Pre-mid pubertal	NR	48:37	Hockey	106	12.3 (12-13)	Pre-mid pubertal		60:46	N	Y
Valente-Dos-Santos 2013 ⁵²	73	15.4 (15-17)	Skeletal age 16.4	NR	73:0	Basketball						N	Y
Vasiliauskas 2006 ⁶⁵	62	13.6 (8-17)	NR	62:0:0	62:0	Soccer						N	Y
Yildirim 2016 ⁵⁵	140	14.3 (10-18)	NR	NS	107:33	Basketball, Soccer, Swimmers.	31	14.1 (10-18)	NR	NR	21:10	Y	Y
Zdravkovic 2010 ⁶⁶	94	12.9 (12-14)	NR	94:0:0	94:0	Soccer	47	12.9 (12-14)	NR	47:0:0	47:0	N	Y

Abbreviations: C, Caucasian; B, Black; O, Other; M, Male; F, Female; NR, Not reported; Y, Yes; N, No.

Table 2 ECG characteristics of Paediatric athletes and Paediatric non-athletes

Characteristics	Athletes	Non-athletes	% Difference
PR interval (ms)	148 (142 – 154)* [10; 5671] {98%}	139 (136- 141) [4; 737] {41%}	6.1%
QRS duration (ms)	86 (84 -88) [11; 6938] {97%}	83 (79 -86) [6; 952] {97%}	4%
QTc duration (ms)	396 (391-400) [11; 7018] {97%}	386 (375 - 398) [6; 902] {97%}	3%
QRS axis*, (degrees)	70 (63.1 - 76.1) [8; 5476] {99%}	70 (67.9 - 73.0) [4; 779] {60%}	0%
Group 1 ECG patterns			Odds Ratio
Sinus bradycardia (%)	37.4 (17.6 – 59.7)** [11; 9745] {99%}	19.2 (16.6 21.90) [3; 834] {0%}	2.5 (2.1 - 3.0)
Sinus arrhythmia (%)	45.8 (35.7 -56.0) [3; 2898] {95.9%}		
1 st °AV block (%)	2.2 (0.8 - 4.2)** [8; 9488] {97%}	0.4 (0.1 - 1.1) [3; 834] {26%}	4.6 (1.7 - 12.4)
2 nd °AV block (Morbitz Type I) (%)	0.2 (0.1 - 0.4) [3; 2898] {0%}		
Incomplete RBBB (%)	25.8 (18.2 - 33.7)** [10; 9736] {97%}	7.8 (4.2 - 12.4) [3; 834] {78%}	4.3 (3.5 – 5.6)
LVH (%)	35.2 (26.0 - 45.0)** [11; 9745] {98%}	24.1 (20.3 - 28.1) [3; 834] {41%}	1.7 (1.5 - 2.0)
ER (%)	37.1 (25.6 -49.2)** [10; 9736] {99.3%}	29.2 (17.2 – 43.0) [3; 834] {93.9%}	1.4 (1.2 -1.7)
Group 2 ECG patterns			Risk Ratio
TWI (≥1mm) (%)	6.7 (4.7 - 8.9) [7; 9372] {93.3%}	5.9 (2.2 – 11.2) [3; 834] {86.7%}	1.1 (0.8 – 1.5)
Deep TWI (≥2mm) (%)	4.7 (2.3 - 8.1)** [7; 6514] {95.9%}	0.3 (0.04 - 1.8) [2; 534] {60.3%}	12.7 (3.1 – 50.7)
Anterior (%)	6.5 (2.9 - 11.3) [7; 9372] {98.4%}	5.7 (2.2 - 10.6) [3; 834] {84.9%}	1.2 (0.9 – 1.5)
Extended Anterior (%)	1.4 (0.2 - 3.5) [4; 5391] {95.8%}	0.9 (0.2 - 2.2) [2; 534] {24.1%}	1.5 (0.6 – 3.6)
Inferior (%)	1.0 (0.3 - 2) [5; 7446] {93%}	0.0 [3; 834]	NC
Lateral (%)	0.3 (0.05 - 0.6) [5; 7446] {80%}	0.0 [3; 834]	NC
Infero-lateral (%)	2.0 (1.0 – 3.3) [8; 9256] {93%}	0.0 [3; 834]	NC
ST-segment depression (%)	0.03 (0.003 - 0.08) [6; 7615] {0%}	0.0 [3; 834]	NC
Abnormal Q waves (%)	0.1 (0.03 - 0.2) [10; 9902] {25%}	0.0 [3; 834]	NC
LAE (%)	3.5 (0.4 - 9.5) [8; 5804] {98%}		

RAE (%)	5.9 (0.9 - 14.8) [4; 2575] {98%}		
LAD (%)	0.4 (0.1 - 0.9) [6; 5683] {75%}		
RAD (%)	3.7 (0.1 - 11.8) [5; 4352] {98%}		
RVH (%)	9.8 (7.0 - 13.0) [3; 2420] {81%}		
Ventricular pre- excitation (%)	0.6 (0.2 - 1.1) [6; 7422] {79%}		
Complete RBBB (%)	0.5 (0.3 - 0.7) [8; 9715] {55%}	0.0 [3; 834]	NC
Complete LBBB (%)	0.1 (0.008 - 0.3) [7; 9499] {81%}	0.0 [3; 834]	NC
Long QT interval (%)	0.6 (0.1 - 1.3) [9; 10247] {57%}	0.0 [3; 834]	NC
Short QT interval (%)	0.4 (0.02 - 1.1) [4; 4108] {81%}	0.0 [3; 834]	NC
Brugada-like ER (%)	0.2 (0.03- 0.4) [5; 7079] {0%}		

Data are presented mean or percentage (95%CI) [number of articles; number of participants] {heterogeneity}

*p ≤0.05 Significantly greater or more prevalent in athletes than non-athletes

** p ≤0.001 Significantly greater or more prevalent in athletes than non-athletes

NC: Non-Computable

Abbreviations: AV, atrioventricular; RBBB: right bundle branch block; LVH: left ventricular hypertrophy; ER: early repolarisation; TWI: T-wave inversion; Anterior: V1–V3; Extended Anterior: V1–V4; Inferior: Leads II-aVF; Lateral: V4-V6/I-aVL; Infero-lateral: Leads II-aVF/V4-V6/I-aVL; LAE: left atrial enlargement; RAE: right atrial enlargement; LAD: left axis deviation; RAD: right axis deviation; LBBB: left bundle branch block

Table 3 ECG characteristics of Paediatric athletes: Impact of Age

Characteristics	≥14 years	<14 years	% Difference
PR interval (ms)	151 (140 - 162) [4; 1985] {99%}	142 (137 - 147) [4; 89] {58%}	6%
QRS duration (ms)	92 (91 - 93)* [4; 1991] {59%}	74 (70 - 82) [3; 77] {88%}	9%
QTc duration (ms)	377 (354 - 400) [5; 4205] {99%}	394 (375 - 412) [3; 77] {6%}	-5%
QRS axis*, degree	76 (73 - 78) [4; 2816] {89%}	74.7 (61.4 - 88.0) [2; 63] {75%}	2%
Group 1 ECG patterns			Odds Ratio
Sinus bradycardia (%)	61.3 (46.3 - 75.3)** [5; 4205] {98%}	18.8 (12.1 - 26.7) [2; 109] {0%}	6.6 (4.1-10.7)
LVH (%)	48.0 (36.4 - 59.5)** [5; 4205] {98%}	20.7 (9.3 - 35.1) [2; 109] {30%}	3.4 (2.2 -5.5)
Group 2 ECG patterns			Risk Ratio
TWI (≥1mm) (%)	6.9 (3.7 - 10.9)** [6; 5051] {96%}	5.4 (0.2 - 16.9) [2; 1272] {92%}	1.3 (1.0-1.7)
Anterior (%)	5.4 (1.4 - 11.8) [7; 6575] {98%}	6.7 (4.4 - 9.4)† [3; 2516] {78%}	1.2 (1.0 - 1.5)
Extended Anterior (%)	1.7 (0.4 - 4.0)* [4; 3823] {94%}	0.5 (0.1 - 3.0) [2; 1257] {68%}	3.1 (1.4 - 6.6)
Infero-lateral (%)	2.5 (1.0 - 4.6)** [5; 3710] {89%}	0.1 (0.01 - 0.4) [2; 1272] {0%}	15.8 (3.9 - 63.9)

Data are presented mean or percentage (95%CI) [number of articles; number of participants] {heterogeneity}

*p ≤0.01 Significantly greater or more prevalent in athletes ≥14 years than in athletes <14 years

** p ≤0.001 Significantly greater or more prevalent in athletes ≥14 years than in athletes <14 years

† ≤0.05 Significantly greater or more prevalent in athletes <14 years than in athletes ≥14 years

NC: Non-Computable

Abbreviations: LVH: left ventricular hypertrophy; TWI: T-wave inversion; Anterior: V1–V3; Extended Anterior: V1–V4; Infero-lateral: Leads II-aVF/V4-V6/I-aVL.

Table 4 ECG characteristics of Paediatric athletes: Impact of race			
Characteristics	Black (10-18 years)	Caucasian (8-18 years)	% Difference
PR interval (ms)	161 (146 -177) [2; 196] {91.6%}	141 (135 to 148) [3; 2529] {95%}	12%
QRS duration (ms)	86 (82 - 90)** [3; 525] {94.9%}	92 (88 - 95) [4; 3232] {98.3%}	-7%
QTc duration (ms)	394 (387 - 401) [3; 525] {95%}	398 (392 - 403) [4; 3232] {98.4%}	-1%
Group 1 ECG patterns			Odds Ratio
Sinus bradycardia (%)	38.2 (18.6 - 60.1)*** [3; 525] {95%}	29.3 (10.9 - 52.2) [5; 6197] {99%}	1.5 (1.3 - 1.8)
1st °AV block (%)	11.4 (6.9 - 16.9)*** [2; 483] {65%}	1.1 (0.25 - 2.5) [4; 5991] {92%}	11.6 (8.0 – 17.0)
Incomplete RBBB (%)	22.1 (13.1 - 32.7) [3; 525] {83%}	21.1 (15.0 - 27.9) [5; 6197] {97%}	1.1 (0.9 - 1.3)
LVH (%)	60.3 (11.0 - 98.3)*** [3; 525] {99%}	28.1 (20.2 - 36.7) [5; 6197] {97%}	3.9 (3.3 - 4.7)
ER (%)	74.3 (41.0 - 96.6)*** [3; 525] {98%}	31.0 (17.4 - 46.5) [5; 6197] {99%}	6.4 (5.2 – 7.9)
Group 2 ECG patterns			Risk Ratio
TWI (≥1mm) (%)	23.4 (19.8 – 27.1)*** [3; 512] {69%}	5.9 (5.3 - 6.6) [5; 5263] {71%}	4.0 (3.3 – 4.8)
Deep TWI (≥2mm) (%)	10.6 (5.5 -17.2)*** [3; 525] {73%}	4.2 (0.7 - 10.4) [4; 3936] {97%}	2.6 (1.9 – 3.4)
Anterior (%)	12.2 (8.2 -16.9)*** [3; 512] {43%}	4.2 (3.0 - 5.6) [4; 6063] {25%}	2.9 (2.2 – 3.8)
Anterior Extended (%)	10.8 (7.8 -14.2)*** [2; 358] {0%}	0.3 (0.03 - 0.8) [3; 3298] {66%}	36 (18 – 71)
Infero-lateral (%)	8.2 (6.0 - 10.7)*** [3; 512] {95%}	1.3 (0.3 - 3.1) [4; 6063] {0%}	6.5 (4.5 – 9.3)
Abnormal Q waves (%)	0.5 (0.0 - 2.0)* [3; 526] {52%}	0.1 (0.04 - 0.3) [4; 6135] {19%}	5.0 (1.3 - 19.3)
LAE (%)	5.7 (1.2 - 1.3)*** [4; 680] {91%}	2.0 (0.02 - 7.0) [4; 3936] {97%}	2.9 (2.0 - 4.2)
LAD (%)	0.8 (0.0 - 3.1) [2; 484] {70%}	0.7 (0.4 - 0.9) [2; 3668] {0%}	1.3 (0.4 - 3.6)
Complete RBBB (%)	0.3 (0.02 - 1.1) [2; 483] {0%}	0.3 (0.2 -0.6) [4; 5991] {46%}	1.2 (0.3 – 5.0)
Long QT (%)	1.1 (0.03 - 3.7) [3; 638] {79%}	0.1 (0.0 - 0.2) [4; 5991] {0%}	16.4 (4.8 – 56.0)
Data are presented mean or percentage (95%CI) [number of articles; number of participants] {heterogeneity}			
*p ≤0.05 Significantly greater or more prevalent in black than Caucasian athletes			
**p ≤0.01 Significantly greater or more prevalent in black than Caucasian athletes			
*** p ≤0.001 Significantly greater or more prevalent in black than Caucasian athletes			

Abbreviations: LVH: left ventricular hypertrophy; TWI: T-wave inversion; Anterior: V1–V3; Extended Anterior: V1–V4; Infero-lateral: Leads II-aVF/V4-V6/I-aVL.

Table 5 Echocardiographic patterns of Paediatric Athletes and Paediatric Non-athletes

Parameter	Athletes	Non-athletes	% Difference
LVIDd (mm)	47.3 (46.2 - 48.3)*** [33; 6681] {99%}	43.4 (41.7 - 45.1) [18; 1042] {98%}	8.2
LVIDs (mm)	29.6 (28.4 - 30.8)*** [19; 3354] {98%}	25.4 (24.8 - 26.0) [7; 177] {64%}	14.2
IVSd (mm)	8.5 (8.2 - 8.8)*** [28; 5083] {99%}	7.4 (7.1 - 7.8) [16; 804] {98%}	12.9
PWTd (mm)	8.2 (7.8 - 8.6)** [29; 5168] {99%}	7.2 (6.6 - 7.8) [17; 908] {92%}	12.2
RWT	0.36 (0.34 - 0.37)*** [29; 6315] {99%}	0.34 (0.33 - 0.35) [16; 804] {99%}	5.6
LVM (g)	135.7 (122.2 - 149.1)*** [29; 5086] {99%}	98.2 (84.6 - 111.8) [17; 908] {99%}	27.6
LVEDV (ml)	106.8 (91.8 - 121.8) [6; 494] {98.3%}		
LVESV (ml)	38.3 (35.1 - 41.6) [5; 457] {86%}		
Aortic Root (mm)	26.3 (24.9 - 27.8)* [10; 3055] {99%}	23.5 (20.9 - 26.0) [6; 563] {99%}	10.6
LAD (mm)	30.2 (28.7 - 31.7)* [13; 5324] {99%}	26.5 (24.5 - 28.6) [8; 587] {97%}	12.3
EF (%)	65.6 (61.1 - 70.1) [11; 3150] {99%}	70.9 (63.8 - 77.9) [4; 130] {99%}	-8.1
FS (%)	37.2 (35.5 - 38.9) [14; 1829] {98%}	36.9 (34.7 - 39.1) [11; 666] {96%}	0.8
E Wave (m/s)	0.88 (0.81 - 0.96) [10; 1915] {0%}	0.91 (0.86 - 0.96) [4; 480] {93%}	-3.4
A Wave (m/s)	0.46 (0.43 - 0.49) [10; 1915] {98%}	0.49 (0.46 - 0.52) [4; 480] {84%}	-6.5
E/A ratio	2.1 (2.0 - 2.2) [14; 3634] {96%}	1.9 (1.8 - 2.1) [8; 672] {93%}	9.5
DT (ms)	133 (108 - 157) [4; 201] {97%}		
IVRT (ms)	60 (39 - 82) [3; 168] {99%}		

Data are mean (95% CI), [number of studies; number of participants] {heterogeneity}

*p ≤0.05 Significantly greater in athletes than non-athletes

**p ≤0.01 Significantly greater in athletes than non-athletes

*** p ≤0.001 Significantly greater in athletes than non-athletes

LVIDd, left ventricular cavity diameter in end-diastole; LVIDs, left ventricular cavity diameter in end-systole; IVSd, interventricular septum thickness in end-diastole; PWTd, posterior wall thickness in end-diastole; RWT, relative wall thickness; LVM, left ventricular mass; LVEDV, left ventricular end-diastolic volume; LVESV, left ventricular end-systolic volume; Ao, aortic root diameter; LAD, left atrial diameter; EF, ejection fraction; FS, fractional shortening; DT, Deceleration time; IVRT, interventricular septum relaxation time

Table 6 Echocardiographic patterns of Paediatric Athletes: Impact of Age

Parameter	≥14 years	<14 years	% Difference
LVIDd (mm)	51.2 (50.6 - 51.9)** [14; 2856] {93%}	44.3 (43.3 - 45.3) [14; 872] {94%}	13.5
LVIDs (mm)	32.8 (30.8 - 34.7)** [5; 288] {97%}	27.6 (25.9 - 29.2) [8; 363] {98%}	15.9
IVSd (mm)	9.2 (8.8 - 9.6)** [12; 1366] {97%}	7.8 (7.5 - 8.0) [13; 787] {96%}	15.2
PWTd (mm)	8.9 (8.5 - 9.3)** [12; 1378] {98%}	7 (6.6 - 7.3) [14; 872] {0%}	21.3
RWT	0.36 (0.33 - 0.39) [12; 2857] {99%}	0.35 (0.33 - 0.36) [13; 787] {96%}	2.8
LVM (g)	167 (153.5 - 180.4)** [12; 1378] {96%}	102.3 (91.8 - 112.8) [14; 872] {87%}	38.7
Ao (mm)	28.9 (27.3 - 30.4)** [5; 2396] {98%}	24.8 (23.7 - 25.8) [6 2420] {97%}	14.2
LAD (mm)	33.3 (32.0 - 34.5)** [6; 2462] {95%}	28.1 (27.0 - 29.2) [8; 601] {93%}	15.6
EF (%)	63.7 (59.1 - 68.2) [4; 285] {99%}	67.5 (55.2 - 79.8) [4; 217] {99%}	-6.0
FS (%)	35.8 (33.0 - 38.7) [3; 1052] {98%}	38.1 (37.0 - 39.2) [6; 226] {56%}	-6.4
E Wave (m/s)	0.86 (0.83 - 0.90) [10; 1915] {92%}	0.72 (0.56 - 0.88) [4; 480] {99%}	16.3
A Wave (m/s)	0.43 (0.39 - 0.46) [10; 1915] {95.3%}		
E/A ratio	2.2 (2.1 - 2.3)* [9; 2710] {92%}	1.9 (1.9 – 2.0) [5; 530] {77%}	13.6

Data are mean (95% CI), [number of studies; number of participants] {heterogeneity}

*p ≤0.05 Significantly greater in athletes ≥14 years than in athletes <14 years

** p ≤0.001 Significantly greater in athletes ≥14 years than in athletes <14 years

LVIDd, left ventricular cavity diameter in end-diastole; LVIDs, left ventricular cavity diameter in end-systole; IVSd, interventricular septum thickness in end-diastole; PWTd, posterior wall thickness in end-diastole; RWT, relative wall thickness; LVM, left ventricular mass; Ao, aortic root diameter; LAD, left atrial diameter; EF, ejection fraction; FS, fractional shortening.

Table 7 Echocardiographic patterns of Paediatric Athletes: Impact of race

Parameter	Black (12-18 years)	Caucasian (8-18 years)	% Diff
LVIDd (mm)	49.5 (47.0 - 51.9) [3; 525] {97%}	48.2 (46.3 - 50.0) [10; 3919] {99%}	2.6
IVSd (mm)	9.7 (9.5 - 9.9) [2; 196] {0%}	8.7 (8.0 - 9.3) [9; 3016] {99%}	10.3
PWTd(mm)	9.7 (9.4 - 10.1)* [2; 196] {44%}	8.5 (7.9 - 9.0) [9; 3016] {99%}	12.4
RWT	0.39 (0.38 - 0.40) [4; 680] {90%}	0.36 (0.34 - 0.38) [10; 3919]{ 99%}	7.7
LVM (g)	169.4 (143 - 195.9) [2; 196] {95%}	148.2 (129.0 -167.4) [9; 3016] {99%}	12.5
Aortic Root (mm)	29.7 (28.9 - 30.5) [3; 638] {90%}	26.9 (24.1 - 29.7) [4; 1137] {99%}	9.4
LAD(mm)	35.4 (34.6 - 36.1)** [3; 638] {81%}	30.5 (27.0 - 34.0) [4; 3320] {99%}	13.4
E/A	2.1 (1.9 - 2.3) [2; 483] {88%}	2.1(1.9 - 2.3) [5; 1207] {93%}	0.0

Data are mean (95% CI), [number of studies; number of participants]

*p ≤0.05 Significantly greater in black than Caucasian athletes

**p ≤0.01 Significantly greater in black than Caucasian athletes

LVIDd, left ventricular cavity diameter in end-diastole; IVSd, interventricular septum thickness in end-diastole; PWTd, posterior wall thickness in end-diastole; RWT, relative wall thickness; LVM, left ventricular mass; Ao, aortic root diameter; LAD, left atrial diameter.


Table 8 Echocardiographic patterns of Paediatric Athletes: Impact of sex

Parameter	Male (8-18 years)	Female (10-18 years)	% Diff
LVIDd (mm)	47.8 (46.5 - 49.2) [21; 4294] {99%}	45.3 (43.5 - 47.1) [6; 479] {98%}	5.2
LVIDs (mm)	30.2 (28.5 - 31.8) [12; 2879] {99%}	28.5 (25.2 - 31.9) [3; 92] {98%}	5.6
IVSd (mm)	8.7 (8.3 - 9.1)* [19; 4066] {99%}	7.9 (7.5 - 8.4) [6; 452] {96%}	9.2
PWTd (mm)	8.4 (7.8 - 8.9) [19; 4066] {99%}	7.8 (6.9 - 8.7) [6; 452] {99%}	7.1
RWT	0.36 (0.34 - 0.38) [19; 4066] {99%}	0.34 (0.33 - 0.36) [6; 452] {95%}	5.6
LVM (g)	137.5 (115.3 - 159.6) [14; 3482] {99%}	129.5 (99.8 - 159.2) [5; 298] {98%}	5.8
EDV (ml)	114.3 (108.0 - 120.7) [5; 409] {86%}	82.2 (69.0 - 95.4) [2; 85] {94%}	28

Data are mean (95% CI), [number of studies; number of participants]

*p ≤0.05 Significantly greater in Male than Female athletes

LVIDd, left ventricular cavity diameter in end-diastole; LVIDs, left ventricular cavity diameter in end-systole; IVSd, interventricular septum thickness in end-diastole; PWTd, posterior wall thickness in end-diastole; RLVWT, relative left ventricular wall thickness; LVM, left ventricular mass; LVEDV, left ventricular end-diastolic volume.



ONLINE APPENDIX

Supplementary appendix A

Full search strategy as applied to the MEDLINE electronic database

#	Query	Results
1	Adolescent/ or Child/ or Paediatrics/	2340649
2	(Girl* OR Boy* OR Junior* OR Juvenile* OR Teen* OR Paediatric* OR Prepubescent OR Pubescent OR Peri Pubertal).ti,ab,kw.	316474
3	Athlete/	4416
4	(Athlete* or Players).ti,ab,kw.	52511
5	Electrocardiography/	169256
6	(Electrocardiogram* or 12 Lead Electrocardiogram* or ECG* or 12 Lead ECG* or EKG* or 12 Lead EKG*).ti,ab,kw.	81490
7	Echocardiography, Doppler/ or Echocardiography, Doppler, Color/ or Echocardiography/ or Echocardiography, Transesophageal/ or Echocardiography, Doppler, Pulsed/	103453
8	(Echocardiogram* or Speckle Tracking or STE Resolution or 2D STE or 2DSTE or Speckle or STE).ti,ab,kw.	21310
9	Magnetic Resonance Imaging/	304307
10	(MRI* or CMRI*).ti,ab,kw.	162605
11	Ultrasonography/ or Ultrasonography, Doppler/ or Blood Flow Velocity/	52496
12	Ultrasound.ti,ab,kw.	167633
13	Ventricular Function/ or Hypertrophy, Right Ventricular/ or Ventricular Function, Left/ or Ventricular Function, Right/ or Ventricular Septum/ or Hypertrophy, Left Ventricular/ or Ventricular Remodeling/ or Myocardial Contraction/ or Heart Atria/	130852
14	(Ventric* or Atria* or Atrium or Septum).ti,ab,kw.	443254
15	Arteries/ or Brachial Artery/ or Radial Artery/ or Carotid Artery/ or Femoral Artery/ or Popliteal Artery/ or Vasodilation/ or Vasoconstriction/ or Vascular Resistance/ or Muscle Smooth, Vascular/ or Endothelium, Vascular/ or Arterioles/	282428
16	(Artery Structure or Artery Function or Arteriolar or Conduit Artery or Resistance Artery or Arterial Size or Arterial Wall Thickness or Intima Media Wall Thickness or Arterial Remodeling or Lumen Dimension or Vascular Function or FMD or Flow Mediated Dilatation or Flow Mediated Dilatation or Shear Stress or Shear Pattern or Shear Rate).ti,ab,kw.	36920
17	(Heart rate or HR or PR Interval or QT Interval or QTc or QRS Duration or QRS or LVH or RVH or Sokolow or Cornell or Pediatric Specific or Romhilt Estes or Early Repolarization or ER or ST Elevation or J Point Elevation or J Wave* or ST Segment Elevation or QRS Slurring or Incomplete Right Bundle Branch Block or Incomplete RBBB or Incomplete Left Bundle Branch Block or Incomplete LBBB or T Wave Inversion or TWI or First Degree Atrio Ventricular Block or 1st Degree AV Block or Q Wave* or LAE or RAE or Left Atrial Enlargement or Right Atrial Enlargement or Bradycardia or Arrhythmia or Ectopic Atrial Rhythm or Junctional Rhythm or Mobitz Type I or Mobitz I or Wenckebach or Second Degree AV Block or 2nd Degree AV Block or Premature Ventricular Contraction* or PVC* or ESC Criteria or European Society of Cardiology Recommendation or Seattle Criteria or Refined Criteria or ECG Criteria).ti,ab,kw.	417843
18	OR/1-2	2470231
19	OR/3-4	53440
20	OR/5-12	842415
21	OR/13-17	1096521
22	AND/18-21	433

Supplementary appendix B

ePublication lists of key journals hand searched to supplement electronic database searching.

#	Journal	Yield
1	Journal of the American College of Cardiology	0
2	Circulation	0
3	Circulation: Arrhythmia and Electrophysiology	0
4	Circulation: Cardiovascular Imaging	0
5	European Heart Journal	0
6	European Heart Journal: Cardiovascular Imaging	0
7	American Heart Journal	0
8	Chest	0
9	Heart	0
10	British Journal of Sports Medicine	1
11	Nature Cardiology	0
12	The New England Journal of Medicine	0
13	European Journal of Preventive Cardiology	0
14	Journal of Electrocardiology	0
15	Journal of the American Society of Echocardiography	2
16	Scandinavian Journal of Sports Medicine	2
17	Europace	1
18	European Journal of Applied Physiology	2
19	Pediatric Exercise Science	0
20	Pediatric Cardiology	1
21	Cardiology in the Young	1
22	Pediatrics	0
23	European Journal of Pediatrics	1
24	American Journal of Hypertension	1
25	The Journal of Physiology	0
26	Journal of Applied Physiology	0
Total		12

Supplementary appendix C

15-item risk of bias assessment checklist

Study: _____

#	RISK OF BIAS ASSESSMENT ITEM	YES	NO/UNCLEAR
1	Sufficient power to detect clinically important effect where probability for difference being due to chance < 5% (<i>answer yes if sample size calculated and adequate to detect clinically important effect</i>)		
2	Are the inclusion and exclusion criteria clearly stated?		
Test-control			
3	Are activity levels for the control group reported?		
4	Are the control group matched for age?		
5	If groups are unmatched, have statistical differences been controlled for? (<i>Answer yes if groups matched for age</i>)		
Test-athletes			
6	Are athletes of competitive status? <i>"One who participates in an organised team or individual sport that requires regular competition against others as a central component places a high premium on excellence and achievement, and requires some form of systematic (and usually intense) training"</i>		
7	Are training details available? (<i>years, volume, duration/intensity</i>)		
Data acquisition			
8	Is there detailed information to allow replication? (<i>Answer yes if professional guidelines cited</i>)		
9	Are the observer(s) stated?		
10	Are more than one observer used? If so is interobserver variability stated?		
Measurement technique			
11	Are professional guidelines observed/cited		
Reporting Data			
12	Is an explanation for missing data given? (<i>Score yes if none missing</i>)		
13	Is data clearly and accurately presented? (<i>Simple outcome data, including denominators and numerators, should be reported for all major findings</i>)		
14	Estimates of random variability in data provided for main outcomes? (<i>e.g. interquartile range, standard error, standard deviation, confidence intervals</i>)		
15	Are anthropometrics reported? (Height and weight or BSA (with formula presented))		
Total Score			

Note:

Items 1,2,14 were selected from Downs & Black's Assessment of Methodological Quality of Randomised and Non-Randomised Studies checklist¹.

Items 3-5,7,10-13 were selected from a previously published athletes heart meta-analysis checklist².

Items 6,9,15 were written specifically for the purposes of this review.

Supplementary appendix D

Variables extracted for analysis

Primary Variables	ECG	Characteristics	Group 1: common and training-related ECG changes	Group 2: uncommon and training-unrelated ECG changes
		Heart rate, bpm	Sinus bradycardia (≥ 30 bpm)	T-wave inversion
		PR interval, ms	Sinus arrhythmia	ST-segment depression
		QRS duration, ms	1 st degree AV block (PR interval > 200 ms)	Pathological Q waves
		QTc duration, ms	Morbitz type 1 (Wenckebach) 2nd degree AV block	Left atrial enlargement
		QRS axis, degree	Incomplete RBBB (QRS duration, 120 ms)	Left axis deviation
		S V1 + R V5/6, mm	Early repolarisation (ST elevation, J-point elevation, J waves, notching or terminal QRS slurring)	Right axis deviation
			Isolated QRS voltage criteria for LVH (Sokolow-Lyon)	Complete LBBB or RBBB
				Long QT interval
				Ventricular pre-excitation
				Brugada-like early repolarization
Echo parameters	Structure	Function		
	LV end-diastolic internal diameter, mm	Ejection fraction, %		
	LV end-systolic internal diameter, mm	Fractional shortening, %		
	LV end-diastolic volume, ml	Stroke volume, ml		
	LV end-systolic volume, ml	Cardiac output, lpm		
	Interventricular septal wall thickness, mm	E wave (m/s)		
	Posterior wall thickness, mm	A wave (m/s)		
	Maximal wall thickness, mm	E/A		
	Relative wall thickness.			

Aortic root, mm
RV end-diastolic area, cm²
RV basal dimension, mm
RV mid-ventricular dimension
RV longitudinal dimension
RV outflow tract dimension (parasternal), mm
RV outflow tract dimension (proximal), mm
RV outflow tract dimension (distal), mm
RV free wall thickness, mm
Left atrial diameter, mm
LV mass, grams

**Secondary
Variables**

Contextual Factors

Age range
Gender
Ethnicity
Height, cm
Weight, kg
BSA
Sport
Training Hours/Week, hours
Training Years, years

Bpm: beats per minute; ms: milliseconds; mm; millimeters; cm; centimeters; LBBB: left bundle branch block; RBBB: right bundle branch block; LVH: left ventricle hypertrophy; LV: left ventricle; RV: right ventricle; m/s: meters per second; kg: kilograms; BSA: body surface area.

Supplementary appendix E

Risk of bias assessment

	Criterion 1	Criterion 2	Criterion 3	Criterion 4	Criterion 5	Criterion 6	Criterion 7	Criterion 8
	Power analysis	Selection criteria	Nonathlete activity levels	Control age matched	Statistical differences accounted for	Competitive athletes	Athlete training details	Detailed data acquisition
Article								
Agrebi 2015 ³	N	Y	N	N	N	Y	Y	N
Attisani 2011 ⁴	N	Y	N	N	N	Y	N	N
Ayabakan 2006 ⁵	N	Y	Y	Y	Y	Y	Y	Y
Bartkevičienė 2015 ⁶	N	Y	Y	Y	Y	Y	Y	Y
Bessem 2014 ⁷	N	Y	N	N	N	Y	N	Y
Calò 2015 ⁸	N	Y	N	N	N	Y	Y	Y
Csajági 2015 ⁹	N	Y	Y	Y	Y	Y	Y	Y
Di Paolo 2012 ¹⁰	N	Y	N	N	N	Y	Y	Y
Dinu 2010 ¹¹	N	N	Y	Y	Y	Y	N	N
Hauser 2013 ¹²	N	Y	N	N	N	Y	Y	Y
Hoogsteen 2003 ¹³	N	Y	N	N	N	Y	Y	Y
Kinoshita 2015 ¹⁴	N	Y	N	N	N	Y	Y	Y
Koch 2012 ¹⁵	N	Y	N	N	N	Y	Y	Y
Konopka 2015 ¹⁶	N	Y	N	N	N	Y	Y	Y
Madeira 2008 ¹⁷	N	Y	N	N	N	Y	Y	Y
Makan 2005 ¹⁸	N	Y	Y	Y	Y	Y	Y	Y
Medved 1986 ¹⁹	N	Y	Y	Y	Y	Y	Y	Y
Meško 1993 ²⁰	N	Y	Y	Y	Y	Y	Y	Y
Migliore 2012 ²¹	N	Y	N	N	N	Y	N	Y
Moarles 1992 ²²	N	Y	N	N	N	Y	N	Y
Obert 1998 ²³	N	Y	Y	Y	Y	Y	Y	Y
Ozer 1994 ²⁴	N	Y	Y	Y	Y	Y	Y	Y
Papadakis 2009 ²⁵	N	Y	Y	Y	Y	Y	Y	Y
Pavlik 2001 ²⁶	N	Y	N	N	Y	Y	Y	Y
Pelà 2014 ²⁷	N	Y	N	N	N	Y	Y	Y
Pelà 2015 ²⁸	N	Y	N	N	N	Y	Y	Y
Petridis 2004 ²⁹	N	Y	N	Y	Y	Y	Y	Y
Rowland 1987 ³⁰	N	Y	Y	Y	Y	Y	Y	Y
Rowland 1994 ³¹	N	Y	N	Y	Y	Y	Y	Y
Rowland 1997 ³²	N	Y	N	N	N	Y	Y	Y

Article	Criterion 1	Criterion 2	Criterion 3	Criterion 4	Criterion 5	Criterion 6	Criterion 7	Criterion 8
	Power analysis	Selection criteria	Nonathlete activity levels	Non-athlete age matched	Statistical differences accounted for	Competitive athletes	Athlete training details	Detailed data acquisition
Rowland 2000 ³³	N	Y	Y	Y	Y	Y	Y	Y
Schmied 2009 ³⁴	N	Y	N	N	N	Y	N	Y
Sharma 1999 ³⁵	N	Y	Y	Y	Y	Y	Y	Y
Sharma 2002 ³⁶	N	Y	Y	Y	Y	Y	Y	Y
Sheikh 2013 ³⁷	N	Y	Y	Y	Y	Y	N	Y
Shi & Selig 2005 ³⁸	N	Y	Y	Y	Y	Y	Y	Y
Stoner 1997 ³⁹	N	Y	Y	Y	Y	Y	Y	Y
Sundberg & Elovainio 1982 ⁴⁰	N	Y	N	Y	Y	Y	Y	Y
Telford 1988 ⁴¹	N	Y	Y	Y	Y	Y	Y	Y
Valente-Dos-Santos 2013 ⁴²	N	Y	N	N	N	Y	Y	Y
Vasiliauskas 2006 ⁴³	N	Y	N	N	N	Y	Y	Y
Yildirim 2016 ⁴⁴	N	Y	Y	Y	Y	Y	Y	Y
Zdravkovic 2010 ⁴⁵	N	Y	Y	Y	Y	Y	Y	Y
No. of articles fulfilling each criterion (% of total included studies)	0(0%)	42(98%)	19(44%)	23(53%)	24(56%)	42(100%)	36(84%)	40(93%)
Proportions of agreement	100%(90-100)	88%(74-96)	84%(69-93)	91%(77-97)	86%(71-94)	91%(77-97)	91%(77-97)	95%(83-99)
<i>Y = criterion fulfilled, N = criterion not fulfilled</i>								

Article	Criterion 9	Criterion 10	Criterion 11	Criterion 12	Criterion 13	Criterion 14	Criterion 15	Total
	Observer(s) stated	Interobserver reliability	Professional guidelines	Missing data	Data presentation	Random variability	Anthropometrics	
Agrebi 2015 ³	N	N	Y	Y	Y	Y	Y	8
Attisani 2011 ⁴	N	N	N	Y	Y	N	N	4
Ayabakan 2006 ⁵	Y	N	Y	Y	Y	Y	Y	13
Bartkevičienė 2015 ⁶	N	N	Y	Y	Y	Y	Y	12
Bessem 2014 ⁷	Y	N	Y	Y	Y	Y	Y	9
Calò 2015 ⁸	Y	N	Y	Y	Y	Y	Y	10
Csajági 2015 ⁹	Y	N	Y	Y	Y	Y	Y	13
Di Paolo 2012 ¹⁰	N	N	Y	Y	Y	Y	Y	9
Dinu 2010 ¹¹	N	N	N	N	Y	Y	Y	7
Hauser 2013 ¹²	Y	N	Y	Y	Y	N	Y	9
Hoogsteen 2003 ¹³	Y	N	Y	N	Y	Y	Y	9
Kinoshita 2015 ¹⁴	Y	N	Y	Y	Y	Y	Y	10
Koch 2012 ¹⁵	Y	N	Y	Y	Y	Y	Y	10
Konopka 2015 ¹⁶	N	N	Y	Y	Y	Y	Y	9
Madeira 2008 ¹⁷	Y	N	Y	Y	Y	Y	Y	10
Makan 2005 ¹⁸	Y	N	Y	Y	Y	Y	Y	13
Medved 1986 ¹⁹	N	N	Y	Y	Y	Y	N	11
Meško 1993 ²⁰	N	N	Y	Y	Y	Y	Y	12
Migliore 2012 ²¹	Y	N	Y	Y	Y	Y	N	8
Moarles 1992 ²²	N	N	Y	N	N	N	Y	5
Obert 1998 ²³	Y	N	Y	Y	Y	Y	Y	13
Ozer 1994 ²⁴	Y	N	Y	Y	Y	Y	N	12
Papadakis 2009 ²⁵	Y	N	Y	Y	Y	Y	Y	13
Pavlik 2001 ²⁶	Y	N	Y	Y	Y	Y	N	10
Pelà 2014 ²⁷	Y	N	Y	Y	Y	Y	Y	10
Pelà 2015 ²⁸	Y	N	Y	Y	Y	Y	Y	10
Petridis 2004 ²⁹	N	N	Y	Y	Y	Y	Y	11
Rowland 1987 ³⁰	Y	N	Y	Y	Y	Y	N	12
Rowland 1994 ³¹	Y	N	Y	Y	Y	Y	Y	12
Rowland 1997 ³²	N	N	Y	Y	Y	Y	Y	9

	Criterion 9	Criterion 10	Criterion 11	Criterion 12	Criterion 13	Criterion 14	Criterion 15	Total
	Observer(s) stated	Interobserver reliability	Professional guidelines	Missing data	Data presentation	Random variability	Anthropometrics	
Article								
Rowland 2000 ³³	N	N	Y	Y	Y	Y	Y	12
Schmied 2009 ³⁴	Y	N	Y	Y	Y	Y	Y	9
Sharma 1999 ³⁵	N	N	Y	Y	Y	Y	Y	12
Sharma 2002 ³⁶	Y	N	Y	Y	Y	Y	Y	13
Sheikh 2013 ³⁷	Y	N	Y	Y	Y	Y	Y	12
Shi & Selig 2005 ³⁸	N	N	Y	Y	Y	Y	Y	12
Stoner 1997 ³⁹	N	N	Y	Y	Y	Y	Y	12
Sundberg & Elovainio 1982 ⁴⁰	Y	N	N	Y	Y	Y	Y	11
Telford 1988 ⁴¹	N	N	Y	Y	Y	Y	Y	12
Valente-Dos-Santos 2013 ⁴²	Y	N	Y	Y	Y	Y	Y	10
Vasiliauskas 2006 ⁴³	Y	N	Y	Y	Y	Y	Y	10
Yildirim 2016 ⁴⁴	N	N	Y	Y	Y	Y	Y	12
Zdravkovic 2010 ⁴⁵	Y	N	Y	Y	Y	Y	Y	13
No. of articles fulfilling each criterion (% of total included studies)	26(60%)	0(0%)	40(93%)	40(93%)	43(98%)	40(93%)	37(86%)	452 (70%)
Proportions of agreement	81%(66-91)	100%(90-100)	77%(61-88)	77%(61-88)	84%(69-93)	93%(80-98)	88%(74-96)	70%(47-83)
<i>Y = criterion fulfilled, N = criterion not fulfilled</i>								

Supplementary appendix F

Articles with overlapping electrocardiographic data

First Author	Overlapping participants	Number of participants included
Di Paolo 2012 ¹⁰	Athlete (males, black; n=154)	155/155
Schmied 2009 ³⁴	Athlete (males, black; n=155)	
Pelà 2015 ²⁸	Athlete (males, Caucasian; n=158)	158/158
Pelà 2014 ²⁷	Athlete (males, Caucasian; n=96)	

Articles with overlapping Echocardiographic data

First Author	Overlapping participants	Number of participants included
Di Paolo 2012 ¹⁰	Athletes (males, black; n=154)	155/155
Schmied 2009 ³⁴	Athletes (males, black; n=155)	
Makan 2005 ¹⁸	Athletes (males, mixed race; n=693)	900/900
Sharma 2002 ³⁶	Athletes (females, mixed race; n=207)	
	Athletes (males, mixed race; n=540)	
	Athletes (females, mixed race; n=180)	
Pelà 2015 ²⁸	Athletes (male, Caucasian; n=158)	158/158
Pelà 2014 ²⁷	Athletes (male, Caucasian; n=96)	

Articles reporting repeat echocardiographic measurements (within the same article)

First Author	Follow-up	Participants	Number of participants included
Csajági 2015 ⁹	Six repeat assessments: <ul style="list-style-type: none"> • Start • Endurance (GEP1) • Race 1 (RP1) • Detraining (DT) • Endurance (GEP2) • Race2 (RP2) 	Athletes (mixed gender, Caucasian; n=15)	15/15
Meško 1993 ²⁰	Four repeat assessments: <ul style="list-style-type: none"> • Year 1 • Year 2 • Year 3 • Year 4 	Athletes (male; n=23) Nonathletes (males; n=17)	40/40
Stoner 1997 ³⁹	Two assessments: <ul style="list-style-type: none"> • Pre-onset of training • 1 Year post onset of training 	Athletes (male; n=37) Nonathletes (male; n=20)	57/57
Kinoshita 2015 ¹⁴	Five repeat assessments: <ul style="list-style-type: none"> • Baseline • 0.5 Years post • 1 Year post • 1.5 Years post • 2 Years post • 2.5 Years post • 3 Years post 	Athletes (females, Japanese; n=51)	34/51

REFERENCES

1. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health*. 1998;52(6):377-384. .
2. Utomi V, Oxborough D, Whyte GP, et al. Systematic review and meta-analysis of training mode, imaging modality and body size influences on the morphology and function of the male athlete's heart. *Heart*. 2013;99(23):1727-1733.
3. Agrebi B, Tkatchuk V, Hlila N, Mouelhi E, Belhani A. Impact of specific training and competition on myocardial structure and function in different age ranges of male handball players. *PLoS One*. 2015;10(12):e0143609.
4. Attisani G, Faiola F, Luciani U, Bianchi G, Veicstenas A, Casasco M. Negative T waves in right precordial leads in pre-adolescent subjects . A personal experience. *Medicana Dello Sport*. 2011;64(4):423-434.
5. Ayabakan C, Akalin F, Mengütay S, Cotuk B, Odabas I, Ozüak A. Athlete's heart in prepubertal male swimmers. *Cardiol Young*. 2006;16(1):61-66.
6. Bartkevičienė A. Echocardiographic characteristics of left ventricular geometry of 12-17 years athletes. *Heal Sci*. 2015;25(4):65-72.
7. Bessem B, de Bruijn MC, Nieuwland W. The ECG of high-level junior soccer players: comparing the ESC vs. the Seattle criteria. *Br J Sports Med*. 2014;49(15):1000-1006.
8. Calò L, Sperandii F, Martino A, et al. Echocardiographic findings in 2261 peri-pubertal athletes with or without inverted T waves at electrocardiogram. *Heart*. 2015;101:193-200.
9. Csajági E, Szauder I, Major Z, Pavlik G. Left Ventricular Morphology in Different Periods of the Training Season in Elite Young Swimmers. *Pediatr Exerc Sci*. 2015;27(2):185-191.
10. Di Paolo FM, Schmied C, Zerguini Y a., et al. The Athlete's heart in adolescent Africans: An electrocardiographic and echocardiographic study. *J Am Coll Cardiol*. 2012;59(11):1029-1036.
11. Dinu V, Dutsi S, Lucia C, Gusti A. Aspects of morphofunctional and cardiovascular.pdf. *Arch Balk Med Union*. 2010;45(4):282-285.
12. Hauser M, Petzuch K, Kühn A, et al. The Munich Triathlon Heart Study: ventricular function, myocardial velocities, and two-dimensional strain in healthy children before and after endurance stress. *Pediatr Cardiol*. 2013;34(3):576-582.
13. Hoogsteen J, Hoogeveen A, Schaffers H, Wijn PFF, Van Der Wall EE. Left atrial and ventricular dimensions in highly trained cyclists. *Int J Cardiovasc Imaging*.

2003;19(3):211-217.

14. Kinoshita N, Katsukawa F, Yamazaki H. Modeling of Longitudinal Changes in Left Ventricular Dimensions among Female Adolescent Runners. *PLoS One*. 2015;10(10):e0140573.
15. Koch S, Cassel M, Linne K, Mayer F, Scharhag J. ECG and echocardiographic findings in 10-15-year-old elite athletes. *Eur J Prev Cardiol*. 2012;21(6):774-781. .
16. Konopka M, Banach M, Burkhard-Jagodzińska K, et al. Echocardiographic evaluation of cardiovascular system in adolescent athletes in view of physiological adaptation to physical training. *Folia Cardiol*. 2015;10(4):233-241.
17. Madeira RB, Trabulo M, Alves F, Pereira JG. Effects of chronic exercise training on left ventricular dimensions and function in young athletes. *Port J Cardiol an Off J Port Soc Cardiol*. 2008;27(7-8):909-922.
18. Makan J, Sharma S, Firoozi S, Whyte G, Jackson PG, McKenna WJ. Physiological upper limits of ventricular cavity size in highly trained adolescent athletes. *Heart*. 2005;91(4):495-499. .
19. Medved R, Fabecic-Sabadi V, Medved V. Echocardiographic Findings in Children Participating in Swimming Training. *Int J Sports Med*. 1986;7(2):94-99.
20. Meško D, Jurko A, Vrlík M, Novomeská M, Horniak E, Dzurenková D. Development of the left ventricular hypertrophy and dilation in adolescent ice hockey players evaluated with echocardiography. *Sport Med Train Rehabil*. 1993;4(3):177-188.
21. Migliore F, Zorzi A, Michieli P, et al. Prevalence of cardiomyopathy in italian asymptomatic children with electrocardiographic T-wave inversion at preparticipation screening. *Circulation*. 2012;125(3):529-538.
22. Morales MC. LV mass in adolescent basketball players. *Cardiovasc Rev reports*. 1992;13(10):60-62.
23. Obert P, Stecken F, Courteix D, Lecoq a M, Guenon P. Effect of long-term intensive endurance training on left ventricular structure and diastolic function in prepubertal children. *Int J Sports Med*. 1998;19(2):149-154.
24. Ozer S, Cil E, Baltaci G, Ergun N, Ozme S. Left Ventricular Structure and Function by Echocardiography in Childhood Swimmers. *Jpn Heart J*. 1994;35(3):295-300.
25. Papadakis M, Basavarajaiah S, Rawlins J, et al. Prevalence and significance of T-wave inversions in predominantly Caucasian adolescent athletes. *Eur Heart J*. 2009;30(14):1728-1735.
26. Pavlik G, Olexó Z, Osváth P, Sidó Z, Frenkl R. Echocardiographic characteristics of male athletes of different age. *Br J Sports Med*. 2001;35(2):95-99.
27. Pelà G, Li Calzi M, Crocamo A, et al. Ethnicity-related variations of left ventricular

- remodeling in adolescent amateur football players. *Scand J Med Sci Sports*. 2014;25(3):382-389.
28. Pela G, Crocamo A, Li Calzi M, et al. Sex-related differences in left ventricular structure in early adolescent non-professional athletes. *Eur J Prev Cardiol*. 2015;0(0):1-8.
 29. Petridis L, Kneffel Z, Kispéter Z, Horváth P, Sidó Z, Pavlik G. Echocardiographic characteristics in adolescent junior male athletes of different sport events. *Acta Physiol Hung*. 2004;91(2):99-109.
 30. Rowland TW, Delaney BC, Siconolfi SF, Rowland, Thomas W. Delaney, Brian C. Siconolfi SF. "Athlete's heart" in prepubertal children. *Pediatrics*. 1987;79(5):800-804. .
 31. Rowland TW, Unnithan VB, MacFarlane NG, Gibson NG, Paton JY. Clinical manifestations of the "athlete's heart" in prepubertal male runners. *Int J Sports Med*. 1994;15(8):515-519.
 32. Rowland T, Goff D, DeLuca P, Popowski B. Cardiac effects of a competitive road race in trained child runners. *Pediatrics*. 1997;100(3):E2.
 33. Rowland T, Wehnert M, Miller K. Cardiac responses to exercise in competitive child cyclists. *Am Coll Sport Med*. 2000;32(4):747-752.
 34. Schmied C, Zerguini Y, Junge A, et al. Cardiac findings in the precompetition medical assessment of football players participating in the 2009 African Under-17 Championships in Algeria. *Br J Sports Med*. 2009;43(9):716-721.
 35. Sharma S, Whyte G, Elliott P, et al. Electrocardiographic changes in 1000 highly trained junior elite athletes. *Br J Sports Med*. 1999;33(5):319-324.
 36. Sharma S, Maron BJ, Whyte G, Firoozi S, Elliott PM, McKenna WJ. Physiologic Limits of Left Ventricular Hypertrophy in Elite Junior Athletes : Relevance to Differential Diagnosis of Athlete's Heart and Hypertrophic Cardiomyopathy. *J Am Coll Cardiol*. 2002;40(8):1431-1436.
 37. Sheikh N, Papadakis M, Carre F, et al. Cardiac adaptation to exercise in adolescent athletes of African ethnicity: an emergent elite athletic population. *Br J Sports Med*. 2013;47(9):585-592.
 38. Shi JR, Selig S. Cardiac Structure and Function in Young Endurance Athletes and Nonathletes. *J Exerc Sci Fit*. 2005;3(2):74-80.
 39. Stoner JE. Cardiac dimensions gymnasts and swimmers. *Biol Sport*. 1997;14(2):115-125.
 40. Sundberg S, Elovainio R. Resting ECG in athletic and non-athletic adolescent boys: correlations with heart volume and cardiorespiratory fitness. *Clin Physiol*. 1982;2(5):419-426.
 41. Telford RD, McDonald IG, Ellis LB, Chennells MH, Sandstrom ER, Fuller PJ. Echocardiographic dimensions in trained and untrained 12-year-old boys and girls. *J*

Sport Sci. 1988;6(1):49-57.

42. Valente-Dos-Santos J, Coelho-e-Silva MJ, Vaz V, et al. Ventricular mass in relation to body size, composition, and skeletal age in adolescent athletes. *Clin J Sport Med.* 2013;23(4):293-299.
43. Vasiliauskas D, Venckuna T, Marcinkevicius J, Bartkeviciene A. Development of structural cardiac adaptation in basketball players. *Eur J Prev Cardiol.* 2006;13(6):985-989.
44. Yildirim Ş, Binnetoğlu FK, Battal F, Aylanç H, Kaymaz N, Tekin M. Relation between QT Variables and Left Ventricular Geometry in Athletes and Obese Children. *Acta Med Port.* 2016;29(2):95-100.
45. Zdravkovic M, Perunicic J, Krotin M, et al. Echocardiographic study of early left ventricular remodeling in highly trained preadolescent footballers. *J Sci Med Sport.* 2010;13(6):602-606.