Accepted Manuscript

Variations in the APOE allele or BDNF Val66Met polymorphism are not associated with changes in cognitive function following a tertiary education intervention in older adults: The Tasmanian Healthy Brain Project

Megan E. Thow, Mathew J. Summers, Jeffery J. Summers, Nichole L. Saunders, James C. Vickers

PII: S0197-4580(17)30104-5

DOI: 10.1016/j.neurobiolaging.2017.03.028

Reference: NBA 9889

To appear in: Neurobiology of Aging


Received Date: 10 October 2016

Revised Date: 21 February 2017

Accepted Date: 23 March 2017

Please cite this article as: Thow, M.E., Summers, M.J., Summers, J.J., Saunders, N.L., Vickers, J.C., Variations in the APOE allele or BDNF Val66Met polymorphism are not associated with changes in cognitive function following a tertiary education intervention in older adults: The Tasmanian Healthy Brain Project, *Neurobiology of Aging* (2017), doi: 10.1016/j.neurobiolaging.2017.03.028.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



NEGATIVE RESULT

Variations in the APOE allele or BDNF Val66Met polymorphism are not associated

with changes in cognitive function following a tertiary education intervention in older

adults: The Tasmanian Healthy Brain Project

Megan E. THOW ^{a,b, ‡}, Mathew J. SUMMERS ^{b,c, †}, Jeffery J. SUMMERS ^{a.d}, Nichole L.

SAUNDERS ^b & James C. VICKERS ^b

- a School of Medicine, University of Tasmania, Hobart, Tasmania, AUSTRALIA
- b Wicking Dementia Research & Education Centre, Faculty of Health, University of Tasmania, Hobart, Tasmania, AUSTRALIA
- c Sunshine Coast Mind and Neuroscience Thompson Institute, University of the Sunshine Coast, Birtinya, Queensland, AUSTRALIA
- d Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK

[†] Corresponding author: A/Prof Mathew Summers, Sunshine Coast Mind and Neuroscience -Thompson Institute, University of the Sunshine Coast, Locked Bag 4 (ML59), Maroochydore DC, Queensland, AUSTRALIA, 4558. Tel +61 7 5456 3758, email: <u>msummers@usc.edu.au</u>

‡ Megan E. THOW has previously published under the name Megan E. LENEHAN.

Disclosures

MT received a University of Tasmania Postgraduate Research scholarship as well as supplemental postgraduate scholarships from the Wicking Dementia Research and Education Centre and the Alzheimer's Australia Dementia Research Foundation. This project is funded by National Health and Medical Research Council (NHRMC) Project grants (1003645 and 1108794), as well as the JO and JR Wicking Trust (Equity Trustees). MS reports personal fees from Eli Lily (Australia) Pty Ltd and grants from Novotech Pty Ltd, outside the submitted work. All other authors report nothing to disclose.

5 Supplement files for online access

Abstract

The APOE ε 4 allele and the Met variant of the BDNF Val66Met polymorphism are associated with reduced cognitive function in older adults. The aim of this study was to examine the independent and interactional effect of the APOE ε 4 allele and BDNF Val66Met polymorphism on cognitive function in a cohort of healthy older adults who had undertaken further university level education. Multiple group Latent Growth Curve Modelling revealed no change in cognitive function over time in APOE ε 4-carriers or in BDNF Met-carriers, nor in carriers of both APOE- ε 4 and BDNF-Met alleles. Further, the results indicate that allelic variation in either APOE or BDNF does not modify the beneficial effects of a university based education intervention on cognitive function over a four-year period following the intervention.

Introduction (see Supplement 1 for extended Introduction)

Carriers of the apolipoprotein (APOE) ε 4 allele demonstrate poorer performance compared to non-carriers across a range of cognitive functions, including episodic memory, executive function and general cognition (Wisdom, et al., 2011); with the cognitive decline in ε 4carriers becoming more pronounced with age (Wisdom, et al., 2011). Similarly, brain derived neurotrophic factor (BDNF) is thought to contribute to variance in adult cognitive function, with several studies reporting that healthy older BDNF-Met carriers display reduced cognitive function relative to Val-homozygotes (Egan, et al., 2003). We examined the influence of the APOE ε 4 allele and BDNF Met polymorphism on the longitudinally assessed cognitive performance of healthy older adults who have undertaken an education intervention. This intervention has been shown to result in improved cognitive reserve in this sample (Lenehan, et al., 2016).

Methods (see Supplement 2 for extended Method)

ACCEPTED MANUSCRIPT

A total of 444 healthy adults aged between 50 and 79 years were recruited from the Tasmanian Healthy Brain Project (THBP) participant pool (Summers, et al., 2013). The intervention group (n = 344) had completed a minimum of 12 months part-time or full-time university study comprising a minimum of two units at undergraduate or post graduate levels. The remaining 100 participants were a control reference group who had not undertaken any intervention. The project was approved by the Human Research Ethics Committee (Tasmania) Network and further details of the study protocol have been previously published (Summers, et al., 2013). All participants in the THBP completed a comprehensive testing battery (see Supplement 2; Summers, et al., 2013). DNA was extracted from saliva samples with APOE and BDNF genotype being determined using a one-step ARMS-PCR.

Core Data (see Supplement 2 for extended description of Analyses)

Multiple Group Latent Growth Curve Modelling (LGCM) using maximum likelihood estimation was used to compare group differences in APOE genotype (control non- ϵ 4 carrier; control ϵ 4-carrier; intervention non- ϵ 4 carrier; and, intervention ϵ 4-carrier), BDNF genotype (control Val-homozygote; control Met-carrier; intervention Val-homozygote; and intervention Met-carrier), and APOE/BDNF genotype (ϵ 4/ Met, non- ϵ 4/ Met, ϵ 4/ Val, non- ϵ 4/ Val) in longitudinal performance on composite measures of episodic memory, working memory, executive function and language processing. For APOE (see Supplement 3), the intervention ϵ 4 group displayed higher working memory performance at baseline compared to the control group non- ϵ 4 carriers. In addition, there was a significant difference between the rate of change in language processing over time between the control non- ϵ 4 group (decrease over time) and the intervention non- ϵ 4 group (increase over time). There were no significant APOE group differences in episodic memory or executive function over time. For BDNF (see Supplement 4) the control Met-carrier group displayed lower baseline episodic memory than the intervention Val-carrier group and both intervention groups (Val-

ACCEPTED MANUSCRIPT

homozygote and Met-carrier) displayed significantly higher linear slopes in language processing compared to the control Val-carrier groups. For the interactive effect of APOE/BDNF (see Supplement 5) no differences between groups on any of the cognitive domains were detected at baseline, nor over time.

Discussion of data (see Supplement 1 for extended Conclusions)

No significant differences in baseline score or linear rate of change over time to working memory, episodic memory, executive function or language processing capacity were found between: intervention group ε 4-carriers and non- ε 4 carriers or between control group ε 4carriers and non- ε 4 carriers (Supplement 1); between Val-homozygotes and Met-carriers or between control group Val-homozygotes and Met-carriers (Supplement 2); or between any combination of APOE/BDNF genotype (Supplement 3). Further, there was no evidence of a genotype specific enhanced benefit of education-based intervention on cognitive function, with no differences in the linear slope of the intervention subgroups (APOE or BDNF) across any of the four cognitive domains. These results suggest that in a group of healthy older adults the APOE and BDNF genotypes are unrelated to cognitive performance or to the fouryear trajectory of cognitive change.

References

- Egan, M.F., Kojima, M., Callicott, J.H., Goldberg, T.E., Kolachana, B.S., Bertolino, A., Zaitsev, E., Gold, B., Goldman, D., Dean, M., Lu, B., Weinberger, D.R. 2003. Article: The BDNF val66met Polymorphism Affects Activity-Dependent Secretion of BDNF and Human Memory and Hippocampal Function. Cell 112, 257-69. doi:10.1016/s0092-8674(03)00035-7.
- Lenehan, M.E., Summers, M.J., Saunders, N.L., Summers, J.J., Ward, D.D., Ritchie, K., Vickers, J.C. 2016. Sending your grandparents to university increases cognitive reserve: The Tasmanian Healthy Brain Project. Neuropsychology 30(5), 525-31. doi:10.1037/neu0000249.
- Summers, M.J., Saunders, N.L., Valenzuela, M.J., Summers, J.J., Ritchie, K., Robinson, A., Vickers, J.C. 2013. The Tasmanian Healthy Brain Project (THBP): A prospective longitudinal examination of the effect of university level education in older adults in preventing age-related cognitive decline and reducing the risk of dementia. International Psychogeriatrics 25(7), 1145-55. doi:10.1017/S1041610213000380.
- Wisdom, N.M., Callahan, J.L., Hawkins, K.A. 2011. The effects of apolipoprotein E on nonimpaired cognitive functioning: A meta-analysis. Neurobiol Aging 32(1), 63-74. doi:10.1016/j.neurobiolaging.2009.02.003.