



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Amazonian White-Sand Forests Show Strong Floristic Links with Surrounding Oligotrophic Habitats and the Guiana Shield

Citation for published version:

Garcia-Villacorta, R, Dexter, K & Pennington, RT 2016, 'Amazonian White-Sand Forests Show Strong Floristic Links with Surrounding Oligotrophic Habitats and the Guiana Shield' *Biotropica*, vol. 48, no. 1. DOI: 10.1111/btp.12302

Digital Object Identifier (DOI):

[10.1111/btp.12302](https://doi.org/10.1111/btp.12302)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Biotropica

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 LRH: García-Villacorta *et al.*

2 RRT: Phytogeography of Amazonian white-sand forests

3

4 **Amazonian White-Sand Forests Show Strong Floristic Links with Surrounding**
5 **Oligotrophic Habitats and the Guiana Shield**

6

7 Roosevelt García-Villacorta^{1,2,3,5}, Kyle G. Dexter^{3,4}, and R. Toby Pennington³

8

9 ¹Institute of Molecular Plant Sciences, University of Edinburgh, Edinburgh EH9 3BF, UK

10 ²Centro Peruano para la Biodiversidad & Conservación, PCBC, Iquitos, Peru

11 ³Royal Botanic Garden Edinburgh, 20a Inverleith Row, Edinburgh EH3 5LR, UK

12 ⁴School of GeoSciences, University of Edinburgh, Edinburgh EH9 3FF, UK

13 ⁵corresponding author; Email: roosevelt.garcia@gmail.com

14

15 Received ____; revision accepted ____

16

1 **ABSTRACT**

2 Amazonian white-sand forests occur on quartzitic sandy soils, are distributed as an archipelago
3 of habitat islands across the rainforests of Amazonia and contain many endemic plant species.
4 Surprisingly, we found that only 23% of plant species in western Amazon white-sand forests are
5 white-sand specialists, while the remaining species (77%) also occur in other habitat types.
6 Overall, our analyses revealed (i) somewhat unexpected composition similarity of white-sand
7 forests with nearby non-white-sand forests, (ii) phytogeographical connections among distant
8 white-sand forests, and (iii) a large proportion of western Amazon white-sand specialists
9 occurring in floras of the western and central Guiana Shield region (7-43%). These results
10 suggest that dispersal from both neighbouring oligotrophic non-white-sand habitats and distant
11 white-sand forests is fundamental in shaping western Amazonian white-sand forests' species
12 composition and diversity. While endemism in Amazonian white-sand forests may be lower than
13 previously estimated, conservation of this unique and fragile environment should remain a
14 priority. Such conservation will require the maintenance of regional dispersal processes that
15 connect these archipelagos of habitat islands and other ecologically similar oligotrophic habitats
16 across the Amazon and the Guiana Shield.

17

18 *Key words:* Amazon, arenosol, biogeography, campinarana, ecoregion, floristics, Guiana Shield,
19 habitat specialization, podzol, varillales

20

1 **RESUMEN**

2 Los bosques de arena blanca de la Amazonía ocurren en suelos arenosos quartzíticos, tienen
3 muchas especies endémicas y están distribuidos como un archipiélago de hábitats, dispersos a lo
4 largo de la Amazonía y la región del escudo Guyanés. Inesperadamente, encontramos una baja
5 proporción de plantas restringidas a ellas (23%) mientras que la mayoría de las especies (77%)
6 también ocurren en otros tipos de hábitat de la Amazonía y del Neotrópico. Estos bosques tienen
7 (i) similitudes florísticas con bosques en otro tipo de suelos de áreas cercanas, (ii) conexiones
8 fitogeográficas con otros bosques de arena blanca distantes, (iii) una alta proporción de su flora
9 especialista habitando áreas del centro y oeste del escudo Guyanés (7-43%). Estos resultados
10 sugieren que la dispersión, desde hábitats oligotróficos cercanos, distintos a arena blanca, así
11 como desde bosques de arena blanca distantes es fundamental en estructurar su composición y
12 diversidad. Aun cuando el nivel de endemismo en los bosques de arena blanca de la Amazonía
13 occidental puede ser más bajo de lo que previamente se estimaba, la conservación de estos
14 ecosistemas únicos y frágiles debería continuar siendo una prioridad. Enfatizamos la necesidad
15 de implementar estrategias regionales de conservación que ayuden a conservar los distintos
16 archipiélagos de arena blanca así como hábitats oligotróficos con condiciones ecológicas
17 similares en la Amazonia y el escudo Guyanés.

18

19 *Palabras clave:* Amazonía, arenosol, biogeografía, campinarana, ecoregión, especialización de
20 hábitat, florística, Guiana Shield, podzol, varillales

21

1 QUARTZ-RICH SANDY SOILS ARE FOUND ACROSS AMAZONIA, A BIOGEOGRAPHIC UNIT
2 encompassing the Amazon basin and the Guiana Shield region. These soils support a complex of
3 vegetation types known as white-sand forests, which occupy relatively large extensions in the
4 Guiana Shield region and Rio Negro basin, one of the oldest geological regions in northern South
5 America (Hammond 2005b). Across the rest of Amazonia, white-sand forest is scattered in
6 island-like patches within a matrix of *terra firme*, upland rainforests on clay and sandy-clay soils,
7 with patches varying in size from several to hundreds of hectares (Macedo & Prance 1978,
8 Anderson 1981, Prance 1996).

9
10 There is a sharp physiognomic contrast when one crosses from a multi-layered cathedral-
11 like *terra firme* forest to white-sand forest: a reduction in forest stature, an increase in the density
12 of pole-like stems, and a relatively open canopy, with a large amount of sunlight reaching into
13 the understory (Coomes & Grubb 1998, García-Villacorta *et al.* 2003). Likewise, white-sand
14 forests are substantially distinct floristically from the typical *terra firme* forest, with many local
15 and regional habitat specialists as well as endemic species (Anderson 1981, Prance 1996, Gentry
16 1986, Fine *et al.* 2010). Given their distinctive structure, patchy distribution and floristic
17 composition, it is not surprising that white-sand forests across the Amazon have received distinct
18 local designations such as *varillal*, *chamizal* (in Peru, Colombia), *Amazon caatinga*, *campina*,
19 *campinarana* (Brazil), *caatinga*, *bana* (Venezuela), *wallaba* forest, and *muri bush* (Guyana,
20 French Guiana, Surinam) (Richards 1941, Revilla 1974, Cooper 1979, Anderson 1981).

21

1 Fundamental to the existence of these forests is the presence of nutrient-poor, sandy soils.
2 Pedological and geological evidence about the origin of these soils (García-Villacorta 2015)
3 suggests that they may have at least four different origins: (i) the product of deep *in situ*
4 weathering of quartzitic sandstones (Kubitzki 1989, Potter 1994, Hammond 2005a); (ii)
5 deposition by eolian transport (Ab'Saber 1982, Clapperton 1993, Horbe *et al.* 2004); (iii) as
6 fluvial deposits of paleo-channels (Klinge 1965, Anderson 1981, Ab'Saber 1982, Hoorn 1994,
7 Räsänen & Linna 1998, Hermoza *et al.* 2005, Rossetti *et al.* 2012); and (iv) the final product of
8 on-going Ferralsol/Acrisol to Podzol transformation (Lucas *et al.* 1984, Dubroeuq & Volkoff
9 1998, Lucas *et al.* 2012, Mendonça *et al.* 2014).

10

11 Taxonomic revisions and local floristic studies in Amazonian white-sand forests have
12 emphasized the existence of plant species and genera disjunctly distributed between the Guiana
13 Shield region and western Amazonian white-sand forests (e.g. Spruce 1908, Gentry & Ortiz
14 1993, Berry *et al.* 1995, Cortés & Franco 1997, Silveira 2003, Arbeláez & Duivenvoorden 2004,
15 García-Villacorta & Hammel 2004, Struwe & Albert 2004, Fine *et al.* 2010). To date, there has
16 been no attempt to study species distribution and compositional patterns of these floras at pan-
17 Amazonian scales. To shed light onto the phytogeography of western Amazon white-sand
18 forests, we addressed three main questions: (1) Are western Amazonian white-sand forests
19 comprised primarily of white-sand specialist species?; (2) What are the phytogeographic
20 connections of western Amazon white-sand species?; and (3) Are white-sand forests of the
21 western Amazon floristically more similar to floras on adjacent areas of non-white-sand soils or
22 to white-sand floras of the Guiana Shield region?

1 **METHODS**

2 STUDY AREA AND FLORISTIC DATASETS.—The study area encompasses the Amazon and Guiana
3 Shield region (Fig. 1). The border of the Amazon and Guiana regions was extracted from the
4 ecoregions map of the world (Olson *et al.* 2001), following closely the limits of the Guiana
5 Shield (Hammond 2005b) and HYBAM’s Amazon basin watershed limits (Seyler *et al.* 2009).
6 We term this entire area “Amazonia”.

7

8 White-sand forests in the western Amazon occur patchily dispersed in the southwest of
9 the Colombian Amazon, northern Peruvian Amazon, and around the area of Cruzeiro do Sul in
10 the state of Acre, Brazil. The white-sand flora in all these three areas has been studied intensively
11 in the last few years, making them amenable to a floristic assessment. To assess the floristic
12 affinities of the white-sand forests from the western Amazon we compiled a list of all vascular
13 plant species known to occur in the white-sand forests of Peru (Loreto region: loreto.wsf.PE),
14 Colombia (Guainía region: guainía.wsf.CO, and Caquetá region: caquetá.wsf.CO), and the
15 western Brazil (Acre region: acre.wsf.BR). The checklist of the white-sand forests of northern
16 Peru were extracted from García-Villacorta *et al.* (2003), supplemented with a more regional
17 study of its woody flora (Fine *et al.* 2010) as well as collections made by other botanists and
18 projects in the same region as recorded in the Missouri Botanical Garden’s Tropicos database
19 (Tropicos-Peru 2013). The following studies were used to compile the checklists of vascular
20 white-sand floras of Colombia (guainía.wsf.CO, and caquetá.wsf.CO), and Brazil (acre.wsf.BR):
21 Cortés & Franco (1997), Arbeláez (2003), Silveira (2003), Cárdenas-López (2007), and Ferreira
22 (2009).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

TAXONOMIC INCLUSION AND STANDARDIZATION.—To have a standardized database, all checklists and flora treatments were checked for synonyms and illegitimate names using the Taxonomic Name Resolution Service v3.0 (Boyle *et al.* 2013, TNRS 2013), which is an online tool that matches a plant checklist against agreed plant taxonomies. Only native vascular plants (gymnosperms, angiosperms, and ferns) were included in the database, and all cultivated, naturalized and hybrid species were excluded. The Missouri Botanical Garden’s Tropicos database was the chosen source for taxonomic matching. In very few cases, especially for recently described species that are still in the process of inclusion in taxonomic databases, resolving species names was achieved by consulting The Plant List website (The Plant List 2013). Intraspecific names (sub-species, varieties, forms) were maintained as much as possible in the database because they may represent taxonomic variation confined to white-sand habitats (e.g. white-sand specialists, cryptic undescribed species), and because taxonomic revisions tend to find new species when revising taxa occurring in these habitats (e.g. Cuatrecasas 1961, Struwe & Albert 2004, Daly & Fine 2011). Therefore, including sub-specific taxa may be useful for a better understanding of floristic patterns in relation to white-sand forests.

The taxonomy at the family level for angiosperms follows the Angiosperm Phylogeny Group III system (The Angiosperm Phylogeny Group 2009). Prior to the analysis, plant families or genera with strictly aquatic habit, or not well represented in forested habitats, were excluded from the database, including Elatinaceae, Nymphaeaceae, Pontederiaceae, Alismataceae, Salviniaceae, Onagraceae, Poaceae, Ceratophyllaceae, Cyperaceae, Hydrocharitaceae,

1 Hydroleaceae, Mayacaceae, Potamogetonaceae, Typhaceae, Lentibularaceae, Cabombaceae,
2 *Pista*, *Montrichardia*, and *Lemna* (Araceae).

3

4 DISTRIBUTIONAL PATTERNS IN WESTERN AMAZON WHITE-SAND FORESTS.—To study the
5 distributional patterns of species from these four floristic checklists, we searched for specimens
6 of each taxon in the Missouri Botanical Garden herbarium online database Tropicos
7 (<http://www.tropicos.org>) and determined the ecoregions in which they occur following Olson *et*
8 *al.* (2001). We did not count a species as being present in an ecoregion if it was only represented
9 by one specimen or if the identification was dubious, based on visual verification of the
10 specimens at MO. In addition, we classified each species in the white-sand dataset into one of
11 three categories: white-sand specialist, poor-soil specialist, or habitat generalist. For this study,
12 white-sand specialists are defined as species occurring exclusively on white-sand soils; poor-soil
13 specialists are species that can be found in both white-sand soils as well as other oligotrophic
14 habitats (e.g. igapó forests, sandy-clay soils); and generalist species are those occurring on white-
15 sand soils, other nutrient-poor soils and any other habitats of the Neotropical region (e.g. clay-
16 rich upland forests, flooded forests, swamp forests, montane forests, savanna, dry forests). The
17 assignment of species to each category was based on field knowledge of species habitat
18 preferences supplemented by review of herbarium label descriptions citing the habitat where
19 specimens were collected (e.g. white-sand forest, varillal, campina forest, campinarana, suelo
20 arenoso, Amazon caatinga, suelo de arenisca). Species with dubious taxonomic identification
21 were not included in the analysis.

22

1 Distributional analyses were conducted for the three groups of species separately as well
2 as together as one group to explore which species are shared amongst ecoregions occurring
3 within the limits of Amazonia (Fig. 1). White-sand forests from Colombia (guainía.wsf.CO, and
4 caquetá.wsf.CO) occur at the margin of the western Guiana Shield (Fig. 1), and previous studies
5 have shown that they have strong phylogeographic connections (i.e. they share a large number of
6 species) with the Guiana Shield region (Cortés & Franco 1997, Cortés *et al.* 1998, Arbeláez
7 2003). Thus, in order to further evaluate the floristic relationship of western Amazon white-sand
8 forests outside of the Guiana Shield, we conducted distributional analyses both including and
9 excluding the Colombian white-sand datasets.

10

11 FLORISTIC RELATIONSHIPS OF WESTERN AMAZON WHITE-SAND FORESTS.—To carry out an
12 analysis of the floristic relationships of western Amazon white-sand forests, province-level plant
13 checklists were compiled for each of the countries with territories in Amazonia as defined here:
14 Bolivia, Brazil, Colombia, Ecuador, Peru, Venezuela, Guyana, French Guiana, and Surinam (Fig.
15 1; Appendix 1). Political unit definitions vary depending on the country (e.g. state in Brazil,
16 department in Peru and Colombia), and we use the name “provinces” throughout this article to
17 refer to all such political units. Because of their relatively small extent, the four provinces
18 located in the Ecuadorian Amazon were treated in the analysis as one unit. In total, data were
19 collated for 26 provinces. For developing this database the following floristic treatments were
20 used: checklist of Peru (Brako & Zarucchi 1993, Tropicós-Peru 2013), checklist of Ecuador
21 (Jorgensen & León-Yáñez 1999, Tropicós-Ecuador 2013), checklist of Bolivia (Tropicós-Bolivia
22 2013) checklist of Brazil (Forzza *et al.* 2010a, b), checklist of the Colombian Amazon (SINCHI
23 2013), and checklist of the Guiana Shield region (Funk *et al.* 2007). These checklists and floras

1 were used to create a presence-absence matrix of species with which we conducted analyses of
2 floristic composition. White-sand specialists found in provinces in which the four white-sand
3 checklists were embedded (CO.GN, CO.CQ, PE.LO, and BR.AC) were excluded from the
4 province-level lists. Including white-sand species in these lists would not have been appropriate
5 with the floristic dissimilarity metric that we used to conduct clustering analyses (see below),
6 because we would have obtained a floristic dissimilarity of zero between the white-sand list and
7 that of the floristic province in which it was embedded.

8

9 CLUSTER ANALYSIS.—In order to evaluate the floristic distance between study units we created a
10 dissimilarity matrix by employing the one-complement of the Simpson similarity index
11 (Simpson 1943) , which measures the proportion of the more species-poor site that is not nested
12 within the more species-rich site (Tuomisto 2010):

13

$$14 \quad \text{Simpson}_{dist} = 1 - \frac{a}{\min(b, c) + a}$$

15

16 Where a is the number of species present in both sites; b is the number of species
17 restricted to one site; and c is the number of species restricted to the other site (Koleff *et al.*
18 2003). We then performed a hierarchical clustering of the floristic checklists based on this
19 dissimilarity matrix using the function *hclust* in the R statistical environment (R Core Team
20 2015).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15

There are multitudes of clustering algorithms for different kinds of data and applications. Yet, evaluations of the benefits of different algorithms are seldom carried out in ecology or biogeography. Assessment of dendrograms is possible by correlating the original dissimilarity matrix with a cophenetic matrix obtained from a dendrogram (Sokal & Rohlf 1962). The algorithm with the highest Pearson cophenetic correlation value will be the one that best represents the original dissimilarity matrix in the topology of the resulting dendrogram (Sokal & Sneath 1963). In order to choose which of four clustering algorithms (Ward, Average linkage (UPGMA), Single linkage, and Complete linkage) gave the best representation of floristic relationships, we calculated the correlation of distances between sites in the resulting hierarchical clusters with the dissimilarity matrix obtained with the Simpson distance index. The analysis of cophenetic correlation among the four algorithms ranged from 0.28 to 0.63 (Appendix 2), with the best performing being Average Linkage. We checked and ensured that ties in dissimilarity values among clusters were not affecting our results using algorithms in the *recluster* package (Dapporto *et al.* 2013).

16
17
18
19
20
21
22

We assessed the statistical support for clusters using multi-scale bootstrapping, implemented with the *pvclust* function in the *pvclust* package (Suzuki & Shimodaira 2006). We used 1000 bootstrap replicates with 10 different sampling levels and focused on approximately unbiased (AU) p-values as a measure of statistical support (Suzuki & Shimodaira 2006). The *pvclust* algorithm also facilitates the estimation of the standard error of each cluster, which provides a measure to diagnose outliers not strongly supported by the data.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

ORDINATION.—As a complement to the floristic patterns revealed by the clustering analysis we performed Non-metric Multidimensional Scaling (NMDS), which extracts a reduced number of axes from the multidimensional space where the positions of the sites are defined (Minchin 1987, Borcard *et al.* 2011).

We used the R package *vegan* (Oksanen *et al.* 2015) to implement NMDS in the manner recommended by Minchin (1987), using the function *metaMDS* with different random start configurations and a final scaling of the results, with the function *postMDS*, along the first dimension for a better interpretation. To avoid reaching an unstable solution where only a local optimum of stress is found, we ran the analysis from different random spatial configurations up to 500 times. Convergence to the same stress value from these random spatial configurations indicated that a global optimum had been reached. The dissimilarity matrix for the NMDS analysis was constructed using the Simpson distance index.

RESULTS

PATTERNS IN HABITAT SPECIALIZATION.—Overall, a total of 1,180 vascular plant species comprising 133 families, and 491 genera were found to occur in the four western Amazon white-sand forest sites. The species distributional dataset resulted in 69,986 unique plant records representing these species' occurrences across the Neotropics. At the species level, 43% of the total vascular flora occurring on white-sand forests was found to be habitat generalist (i.e.

1 occurring in white-sands forests as well as diverse other habitat types), 34% poor-soil specialist
2 (i.e. occurring in white-sand forests and other oligotrophic habitats), and 23% white-sand
3 specialist (i.e. restricted to white-sand forests). Appendix 3 summarizes the richness of families,
4 genera, and species found in the three designated habitat preference categories. Of the total plant
5 records, 74% (51,790 records) corresponded to “habitat generalists,” 21% to “poor-soil
6 specialists” (14,723 records), and 5% to “white-sand specialists” (3,473 records).

7

8 DISTRIBUTION PATTERNS ACROSS ECOREGIONS.—A large proportion of the species of white-sand
9 forests of the western Amazon occurred within ecoregions of the Guiana Shield when this region
10 was analysed as a unit. When excluding the Colombian white-sand checklists from the analysis,
11 we found that 65% of white-sand specialist species (56 out of 85 species) in Loreto, Peru and/or
12 Acre, Brazil were distributed in the Guiana Shield while 35% are endemic to these white-sand
13 areas (29 species) (Appendix 5). When considering white-sand specialists of the full dataset (i.e.
14 including Colombian white-sand checklists), 88.8% of the total (248 white-sand specialist
15 species out of 279 species, Appendix 4) occurred also within the limits of the Guiana Shield
16 region, with the remaining white-sand specialists being endemic to the western Amazon. When
17 analysing what were the proportions of white-sand specialist from Peru (loreto.wsf.PE), and
18 western Brazil (acre.wsf.BR) found in the Guiana Shield region, the three ecoregions with the
19 highest percentage of western Amazon white-sand species were the Caquetá Moist Forests
20 (30%), Guayanan Highlands Moist Forests (33%), and Negro-Branco Moist Forests (43%)
21 (Appendix 5). These proportions changed slightly when analysed using the full white-sand
22 dataset: Caquetá Moist Forests (69%), Guayanan Highlands Moist Forests (51%), and Negro-
23 Branco Moist Forests (49%). Similar patterns were found when looking at the Guiana Shield

1 ecoregions with the lowest proportion of white-sand specialists using both datasets: Rio Negro
2 campinarana (7-8%), and Guyanan Savanna (9-17%). These ecoregions specific locations, and
3 their shared proportions of white-sand specialists using both datasets, are shown in Figure 2.

4

5 FLORISTIC SIMILARITY ANALYSIS.—A total of 26 floristic datasets at the level of provinces were
6 compiled and compared with four western Amazon white-sand forests. There were a total of
7 26,887 vascular plant species in the floristic dataset from 2865 genera and 268 APG III families.
8 Appendix 1 gives a summary of the area and number of vascular plant species found at each
9 province. The number of species in the white-sand forest checklists ranged from 363
10 (acre.wsf.BR) to 955 species (guainia.wsf.CO). The number of species in the province checklists
11 ranged from 607 (Vichada, Colombia) to 8,355 (Amazonas, Brazil).

12

13 The hierarchical cluster analysis indicates that the four western Amazon white-sand
14 floras cluster first with the non-white-sand flora in which they are geographically embedded,
15 rather than with each other (Appendix 6). The four white-sand forests are then found in a large
16 cluster, with surrounding and neighbouring provinces (Fig. 3), rather than falling with provinces
17 of the core Guiana Shield area (VE.DA, GF, SU, GY, VE.BO, BR.RR).

18

19 Relatively high levels of unbiased bootstrap support values (AU) were found by the
20 *pvclust* randomization procedure for all clusters containing white-sand forests sites (Fig. 3). In
21 particular, the cluster grouping the four western Amazonian white-sand floras along with

1 neighbouring non-white-sand floras had an AU pv-value of 84 (Appendix 6). It is clear from the
2 *pvclust* analysis, however, that guainía.wsf.CO is nested within two floras more representative of
3 the western lowlands of the Guiana Shield area (CO.GN, and VE.AM (AU p-value = 76)) than to
4 the other three western Amazonian white-sand sites. Overall, cluster topology support values
5 estimated by *pvclust* (AU p-values) ranged from 59 to 100% (Appendix 6).

6

7 The standard error of the majority AU bootstrap values in *pvclust* was close to 0 for the
8 majority of clusters (Appendix 7), which gives confidence that the existence of these
9 phytogeographic clusters is supported by data. Only cluster 6, which does not include white-sand
10 forests, had a relative high standard error in the bootstrap procedure (Appendix 7).

11

12 ORDINATION.—An NMDS ordination with two axes recovered similar phytogeographic
13 patterns to cluster analysis results as interpreted by the relative distances between floras and
14 white-sand forests portrayed in the ordination space (Fig. 3). The Shepard plot indicates that
15 there is a good fit of the ordination distance among sites against the original dissimilarity
16 distance (Appendix 8). Adding additional axes to the analysis did not result in any substantial
17 reductions in the stress value. White-sand forests of Acre (acre.wsf.BR), Loreto (loreto.wsf.PE),
18 and Caquetá (caquetá.wsf.CO) are closer to each other floristically than they are to the white-
19 sand forests of Guainía (guainía.wsf.CO).

20

21

1 DISCUSSION

2

3 PATTERNS OF HABITAT SPECIALIZATION.—Viewed from the air and explored on the ground,
4 Amazonian white-sand forests are no doubt distinctive in physiognomy and structure compared
5 to neighbouring upland forests on clay or sandy-clay soils (Anderson 1981, Prance 1996,
6 Coomes & Grubb 1996, Bongers *et al.* 1985, Duivenvoorden & Lips 1995, García-Villacorta *et*
7 *al.* 2003, Fine *et al.* 2010, Silveira 2003, Vicentini 2004, Ferreira 2009). In terms of species
8 composition, however, our results show that only about a quarter (23%) of the total 1183
9 vascular plant species inhabiting western Amazonian white-sand forests are specialized to these
10 forests while the vast majority (77%) also occur in non-white-sand habitats (Table 2). This
11 finding shows that a large majority of the species diversity found in white-sand forest can be
12 attributed to plants from other habitats. Previous work on white-sand floristics and
13 phytogeography has often examined taxonomic sub-sets of floras or taken a restricted (sub-
14 regional) geographic approach, which despite showing the influence of plants also found in other
15 habitats, may have overestimated the proportion of true white-sand soil specialists. For instance,
16 a floristic study of the white-sand forests of the Brazilian Amazon found that the majority of the
17 species (54.5%) occurring in the vegetation type were restricted to it, whereas 23.6% also
18 occurred in non-white-sand *terra firme* forests, 20% in *igapó* forests, and 2.6% in *varzea* forests
19 (Anderson 1978, cited in Anderson (1981)). Similarly, an analysis of tree plots in white-sand and
20 neighbouring forests in Peru found that 52% of tree species in white sand forests were
21 specialists, while the rest were facultative specialists (9%), or habitat generalists (39%) (Fine *et*
22 *al.* 2010).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

In contrast, we have used a relatively strict criterion to determine white-sand specialists, *i.e.* that they are found only in white-sand forests, and we therefore may have underestimated specialization in white-sand forests compared to a classification based on species relative abundance. For example, a previous regional study from Madre de Dios, Peru found that only 15-26% of all studied species were restricted to a single forest type or habitat (Pitman *et al.* 1999), while a subsequent study in the same region that used relative abundance information found that 76.5% of dominant tree species were habitat specialists, *i.e.* significantly more abundant in one habitat type (Phillips *et al.* 2003). In accordance with this pattern, a previous study of white-sand forests in Loreto, Peru, found that floristic differentiation between white-sand and *terra firme* forests was greater when one considered relative abundances of species rather than just presence/absence information (Fine *et al.* 2010). Thus, in our study, while some species may not have been classified as white-sand specialists using our strict presence/absence criterion, they may be much more abundant in white-sand forest than other forest types, thus potentially qualifying as specialists using abundance-based criteria. In other words, while there are many species present in white-sand forests that are also found in other forest types, the forest does tend to be dominated by white-sand specialists (Fine *et al.* 2010) or by poor-soil specialists as defined here, *i.e.* species with preference for waterlogged or oligotrophic soils (Freitas 1996, García-Villacorta *et al.* 2003, Vicentini 2004). Similarly, a recent study of swamp forests in the Ecuadorian Amazon found that just 8.6% of tree species in swamp forest are specialists on swamp forest, but that these specialists represent 43.6% of stems in the swamp forest (Pitman *et al.* 2014).

1 Another potential caveat of our results is that the number of white-sand specialist species
2 may increase in the future via the discovery of cryptic species. For example, based upon detailed
3 molecular and morphological studies, populations confined to white-sand habitats and previously
4 assigned to *Protium subserratum* (Burseraceae) were found to be distinct from populations of
5 that occurred on non-white sand soils, and it was suggested that they should be recognised at
6 species level (Daly & Fine 2011, Fine *et al.* 2013). Similarly, a study on reproductive biology,
7 phenotypic differences, and ecological preferences within the *Pagamea coriacea* complex
8 (Rubiaceae) identified two sympatric species exploiting different gradient combinations of light
9 and drainage within white-sand forests (Esteves & Vicentini 2013). Elucidating the number of
10 cryptic undescribed species, and potentially incipient species undergoing ecological adaptation
11 into white-sands forests, will remain uncertain without more taxonomic, field, and molecular-
12 based studies.

13
14 In any case, our result that just 23% of species in western Amazon white-sand forest are
15 restricted to that habitat type is much lower than previous estimates. We suggest that the
16 ecological conditions of other oligotrophic habitats are similar enough to white sand forests such
17 that they function like meta-ecosystems (Loreau *et al.* 2003, Gravel *et al.* 2010), harbouring
18 species that are able to disperse into white sand habitats, and survive there, albeit at low
19 densities. Likewise, the dispersal of propagules via mass-effect from surrounding habitats
20 (Shmida & Wilson 1985, Holt 1993) may be an additional factor influencing the similarity
21 values obtained when using presence/absence data rather than abundance data (cf. Vormisto,
22 Svenning, *et al.* 2004).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19

DISTRIBUTIONAL PATTERNS AND PHYTOGEOGRAPHICAL CONNECTIONS.—Of the total western Amazonian white-sand specialist species, 88% of them occurred in floras within the Guiana Shield region, whereas 12% are endemic to the western Amazon (i.e. restricted to the four white-sand areas studied here). This pattern of phytogeographic connection was still high when only non-Guiana Shield white-sand forests (loreto.wsf.PE, and acre.wsf.BR) were considered (65% of the white-sand specialists were shared with floras within the Guiana Shield region). Among all ecoregions, Caquetá Moist Forests, Guayanan Highlands Moist Forests, and Negro-Branco Moist Forests shared the highest proportions of western Amazon white-sand specialists (Appendix 5). These ecoregions are located at the central and western part of the Guiana Shield region (Fig. 2b) which supports results from studies of Colombian white-sand forests showing strong phytogeographic links with the Guiana Shield flora (Cortés *et al.* 1998, Giraldo-Cañas 2001, Arbeláez & Duivenvoorden 2004). On the other hand, the ecoregions with the highest percentage of shared western Amazon white-sand specialists, when using only species from loreto.wsf.PE (Peru), and acre.wsf.BT (Brazil), are located in the western Amazon (Iquitos varzea, Napo moist forests, and Southwest Amazon moist forests) (Fig. 2a; Appendix 5). It remains unexplained to what degree the white-sand forests contained in these ecoregions is influencing the compositional similarity of the white-sand forests we studied here.

20 DISPERSAL AND FLORISTIC RELATIONSHIPS.—All white-sand forests clustered with the flora of the
21 province where they belong geographically (Fig. 3). Given the overall large number of non-
22 white-sand specialist species in western Amazon white-sand floras, this result suggests that the

1 floristic patterns are driven by the overwhelming number of non-white-sand specialist species;
2 hinting at an important influence of regional dispersal processes (Ricklefs 1987, Cornell &
3 Lawton 1992, Holt 1993, Latham & Ricklefs 1993, Cottenie 2005, Ricklefs 2008). Collectively,
4 our results support the hypothesis of a flora constructed via both long-distance dispersal of
5 white-sand specialists from distant white-sand habitats and more local dispersal of poor-soil
6 specialists and generalist species from ecologically similar oligotrophic habitats occurring in the
7 same areas in which white-sand forests are embedded.

8
9 Certain combinations of edaphic factors that are present in white-sand habitats promote
10 colonization by species from physiognomically distinct habitat types, such as nutrient-poor *terra*
11 *firme* clayey or sandy-clay soils and waterlogged habitats (Freitas 1996, García-Villacorta *et al.*
12 2003, Vicentini 2004). In this regard, peat-accumulating palm swamps (Tuomisto *et al.* 1994,
13 Lähteenoja *et al.* 2009), and old terraces of *terra firme* forests on hilly areas (García-Villacorta *et*
14 *al.* 2010) may represent habitats with similar high stress and/or low resource availability (i.e.
15 poor drainage conditions or nutrient-poor soils, respectively) that may be used by poor-soil plant
16 specialists as dispersal corridors to reach isolated white-sand forests (García-Villacorta *et al.*
17 2010). Ancient hilly terraces on poor oligotrophic clay soils are found to be a common feature at
18 certain drainage divides of the western Amazon (e.g. Stallard 2011). These hilly terraces can
19 have high erosion rates (Stallard 1988), which accelerates the leaching of soil nutrients
20 (Laurance *et al.* 2010, Vormisto, Tuomisto, *et al.* 2004), and they can be extensive and
21 interconnected in some areas (García-Villacorta *et al.* 2010). Similarly, some non-specialist
22 white-sand species have been found in swamp habitats of the Amazon floodplain (Tuomisto *et*
23 *al.* 1994, Lähteenoja *et al.* 2009, Lähteenoja & Page 2011, García-Villacorta *et al.* 2011, Dávila

1 *et al.* 2013, Draper *et al.* 2014), which suggest that edaphic conditions in these habitats may
2 facilitate dispersal of white-sand species.

3

4 THE EFFECT OF GEOGRAPHICAL DISTANCE BETWEEN FLORAS—Geographical proximity may help
5 to explain some part of the observed phytogeographical affinities of white-sand floras. The
6 Amazonas provinces in Brazil and Venezuela (BR.AM, and VE.AM) as well as Colombian
7 provinces are geographically adjacent and were found to have close links to western Amazonian
8 white-sand forests. Geographical distance may also help to explain why *acre.wsf.BR* (Brazil) and
9 *loreto.wsf.PE* (Peru) are more similar to each other and to *caquetá.wsf.CO* (Colombia) than to
10 *guainía.wsf.CO* (Colombia), the latter being closer in geographic distance to the Guiana Shield
11 region (Fig. 2). A correlation analysis of the geographic distance and floristic matrices (Mantel
12 test) in the dataset found a significant association (Appendix 9), implying that spatially adjacent
13 sites are floristically more similar, a pattern often driven by dispersal limitation (Nekola & White
14 1999). In this context, several studies of plant dispersal syndromes in Amazonian white-sand
15 forests have suggested that long-distance dispersal or stepping-stone dispersal may be an
16 important ecological process in the maintenance of Amazonian white-sand forests (Macedo &
17 Prance 1978, Prance & Schubart 1978).

18

19 SYNTHESIS AND IMPLICATIONS FOR CONSERVATION.—Floristically, we found that western
20 Amazon white-sand forests include a large number of plant species that are not restricted to
21 white-sand habitats. Regardless, at the local and regional level white-sand forests possess a
22 significant number of endemics that add to both regional and beta diversity in the Amazon and

1 should remain a conservation priority. It is possible that this finding may be influenced by
2 cryptic, undescribed species –i.e., some morphologically indistinguishable populations found on
3 white sand may actually represent different species. The high proportion of species not
4 specialized to white-sand habitats may result from both immigration of species from other
5 nutrient-poor/water-logged habitats and mass effect dispersal of species from neighbouring
6 habitats with richer soils. Cluster analysis and NMDS ordination concurred that the white-sand
7 forests of the western Amazon are floristically most similar to the non-white sand floras of the
8 geographic regions to which they belong. More broadly, the composition of white-sand forests of
9 the western Amazon is more similar to floras of the western and central Guiana Shield region
10 than to other floras in Amazonia, which implies that long-distance dispersal processes may be
11 important in shaping its species composition. There was significant distance decay in similarity
12 of overall floristic composition, which implies that dispersal processes are playing an important
13 role in driving current floristic assemblage patterns. Dispersal processes may have more
14 importance than local species interactions in structuring Amazonian white-sand plant
15 communities, perhaps via stepping-stone dispersal processes across ecologically similar habitat
16 types like Amazon peatlands, and hilly terraces on oligotrophic soils. The long term conservation
17 of Amazonian white-sand forests will require the maintenance of regional dispersal processes,
18 necessitating corridors connecting these archipelagos of habitat islands across the Amazon and
19 the Guiana Shield.

20

21

22

1 **ACKNOWLEDGEMENTS**

2 The Missouri Botanical Garden's A. H. Gentry Fellowship provided financial support to RG for
3 visiting its herbarium to study Amazonian white-sand specimen collections. RG thanks Dr. Kalle
4 Ruokolainen for encouragement and advice in an early stage of this study. The Sibbald Trust
5 Fund, of the Royal Botanic Garden of Edinburgh, provided generous support to RG during the
6 preparation of this manuscript. A Principal's Career Scholarship supported the lead author as part
7 of his doctoral studies in the University of Edinburgh, Scotland, UK.

8

9 **LITERATURE CITED**

10 AB'SABER, A. N. 1982. The Palaeoclimate and Palaeoecology of Brazilian Amazonia. *In* G. T.
11 Prance (Ed.) *Biological Diversification in the Tropics*. pp. 41–59, Columbia University
12 Press.

13 ANDERSON, A. B. 1981. White-sand vegetation of Brazilian Amazonia. *Biotropica* 199–210.

14 ARBELÁEZ, M. V. 2003. Diversity and dynamics of the vegetation on the sandstone plateaus of
15 the Colombian Amazonia. PhD Dissertation. University of Amsterdam. Available at:
16 <http://dare.uva.nl/document/68172> [Accessed May 18, 2015].

17 ARBELÁEZ, M. V., and J. F. DUIVENVOORDEN. 2004. Patterns of plant species composition on
18 Amazonian sandstone outcrops in Colombia. *J. Veg. Sci.* 15: 181–188.

19 BERRY, P. E., O. HUBER, and B. K. HOLST. 1995. Floristic analysis and phytogeography. *Flora*
20 *Venezuelan Guayana* 1: 161–191.

21 BONGERS, F., D. ENGELEN, and H. KLINGE. 1985. Phytomass structure of natural plant
22 communities on spodosols in southern Venezuela: the Bana woodland. *Vegetatio* 63:
23 13–34.

24 BORCARD, D., F. GILLET, and P. LEGENDRE. 2011. *Numerical Ecology with R*. Springer Science &
25 Business Media.

26 BOYLE, B., N. HOPKINS, Z. LU, J. A. R. GARAY, D. MOZZHERIN, T. REES, N. MATASCI, M. L. NARRO,
27 W. H. PIEL, S. J. MCKAY, S. LOWRY, C. FREELAND, R. K. PEET, and B. J. ENQUIST. 2013.
28 The taxonomic name resolution service: an online tool for automated standardization of
29 plant names. *BMC Bioinformatics* 14: 16.

- 1 BRAKO, L., and J. L. ZARUCCHI. 1993. Catalogue of the Flowering Plants And Gymnosperms of
2 Peru / Catalogo De Las Angiospermas Y Gimnospermas Del Peru. Missouri Botanical
3 Garden, St. Louis, Mo.
- 4 CÁRDENAS-LÓPEZ, D. ed. 2007. Flora del escudo Guayanés en Inírida (Guainía, Colombia).
5 SINCHI, Bogotá, Colombia.
- 6 CLAPPERTON, C. M. 1993. Quaternary Geology and Geomorphology of South America. Elsevier.
- 7 COOMES, D. A., and P. J. GRUBB. 1996. Amazonian caatinga and related communities at La
8 Esmeralda, Venezuela: forest structure, physiognomy and floristics, and control by soil
9 factors. *Vegetatio* 122: 167–191.
- 10 COOMES, D. A., and P. J. GRUBB. 1998. Responses of juvenile trees to above- and belowground
11 competition in nutrient-starved amazonian rain forest. *Ecology* 79: 768–782.
- 12 COOPER, A. 1979. Muri and white sand savannah in Guyana, Surinam and French Guiana.
13 *Ecosyst. World*.
- 14 CORNELL, H. V., and J. H. LAWTON. 1992. Species Interactions, Local and Regional Processes,
15 and Limits to the Richness of Ecological Communities: A Theoretical Perspective. *J.*
16 *Anim. Ecol.* 61: 1–12.
- 17 CORTÉS, R., and P. FRANCO. 1997. Análisis panbiogeográfico de la flora de Chiribiquete,
18 Colombia. *Caldasia* 19: 465–478.
- 19 CORTÉS, R., P. FRANCO, and J. O. RANGEL-C. 1998. La flora vascular de la Sierra de
20 Chiribiquete, Colombia. *Caldasia* 20: 103–141.
- 21 COTTENIE, K. 2005. Integrating environmental and spatial processes in ecological community
22 dynamics. *Ecol. Lett.* 8: 1175–1182.
- 23 CUATRECASAS, J. 1961. A taxonomic revision of the Humiriaceae. Smithsonian Institution.
- 24 DALY, D. C., and P. V. A. FINE. 2011. A New Amazonian Section of *Protium* (Burseraceae)
25 including Both Edaphic Specialist and Generalist Taxa. *Studies in Neotropical*
26 *Burseraceae* XVI. *Syst. Bot.* 36: 939–949.
- 27 DAPPORTO, L., M. RAMAZZOTTI, S. FATTORINI, G. TALAVERA, R. VILA, and R. L. H. DENNIS. 2013.
28 recluster: an unbiased clustering procedure for beta-diversity turnover. *Ecography* 36:
29 1070–1075.
- 30 DÁVILA, N., I. HUAMANTUPA, M. RIOS, W. TRUJILLO, and C. VRISENDORP. 2013. Vegetation and
31 Flora. *In* N. C. A. Pitman, E. Ruelas-Inzunza, C. Vriesendorp, D. F. Stotz, T. Wachter, A.
32 del Campo, D. Alvira, B. Rodríguez-Grández, R. S. Chase, A. R. Sáenz, and P. Soria
33 (Eds.) *Perú: Eye-Campuya-Algodón. Rapid Biological Inventories Report* 25. pp. 231–
34 243, The Field Museum of Natural History, Chicago.
- 35 DRAPER, F. C., K. H. ROUCOUX, I. T. LAWSON, E. T. A. MITCHARD, E. N. H. CORONADO, O.
36 LÄHTEENOJA, L. T. MONTENEGRO, E. V. SANDOVAL, R. ZARÁTE, and T. R. BAKER. 2014.

- 1 The distribution and amount of carbon in the largest peatland complex in Amazonia.
2 Environ. Res. Lett. 9: 124017.
- 3 DUBROEUCQ, D., and B. VOLKOFF. 1998. From Oxisols to Spodosols and Histosols: evolution of
4 the soil mantles in the Rio Negro basin (Amazonia). CATENA 32: 245–280.
- 5 DUIVENVOORDEN, J. F., and J. M. LIPS. 1995. A land-ecological study of soils, vegetation, and
6 plant diversity in Colombian Amazonia.
- 7 DUTILLEUL, P., J. D. STOCKWELL, D. FRIGON, and P. LEGENDRE. 2000. The Mantel Test versus
8 Pearson's Correlation Analysis: Assessment of the Differences for Biological and
9 Environmental Studies. J. Agric. Biol. Environ. Stat. 5: 131–150.
- 10 ESTEVES, S. DE M., and A. VICENTINI. 2013. Cryptic species in *Pagamea coriacea* sensu lato
11 (*Rubiaceae*): evidence from morphology, ecology and reproductive behavior in a
12 sympatric context. Acta Amaz. 43: 415–428.
- 13 FERREIRA, C. A. C. 2009. Análise comparativa do ecossistema campina na Amazônia brasileira.
14 Tese de Doutorado. UFAM, Manaus.
- 15 FINE, P. V. A., F. ZAPATA, D. C. DALY, I. MESONES, T. M. MISIEWICZ, H. F. COOPER, and C. E. A.
16 BARBOSA. 2013. The importance of environmental heterogeneity and spatial distance in
17 generating phylogeographic structure in edaphic specialist and generalist tree species of
18 *Protium* (*Burseraceae*) across the Amazon Basin. J. Biogeogr. 40: 646–661.
- 19 FINE, P. V., R. GARCÍA-VILLACORTA, N. C. PITMAN, I. MESONES, and S. W. KEMBEL. 2010. A
20 floristic study of the white-sand forests of Peru. Ann. Mo. Bot. Gard. 283–305.
- 21 FLORA DO BRASIL. 2013. Brazilian Flora Checklist. Available at: <http://floradobrasil.jbrj.gov.br>
22 [Accessed January 1, 2013].
- 23 FORZZA, R. C. ET AL. 2010a. Catálogo de plantas e fungos do Brasil - Vol. 1. SciELO - JBRJ, Rio
24 de Janeiro.
- 25 FORZZA, R. C. ET AL. 2010b. Catálogo de plantas e fungos do Brasil - Vol. 2. SciELO - JBRJ, Rio
26 de Janeiro.
- 27 FREITAS, L. A. 1996. Caracterización florística y estructural de cuatro comunidades boscosas de
28 terraza baja en la zona de Jenaro Herrera, Amazonía Peruana. IIAP, Iquitos, Peru.
- 29 FUNK, V. A., T. HOLLOWELL, P. BERRY, C. KELLOFF, S. N. ALEXANDER, and OTHERS. 2007.
30 Checklist of the plants of the Guiana Shield (Venezuela: Amazonas, Bolivar, Delta
31 Amacuro; Guyana, Surinam, French Guiana). Department of Botany, National Museum
32 of Natural History, Washington, D. C.
- 33 GARCÍA-VILLACORTA, R. 2015. Integrating molecular biogeography and community ecology to
34 understand the evolution of habitat specialization in Amazonian forests. PhD
35 Dissertation. University of Edinburgh, Edinburgh, Scotland, UK.
- 36 GARCÍA-VILLACORTA, R., M. AHUITE R., and M. OLÓRTEGUI Z. 2003. Clasificación de bosques
37 sobre arena blanca de la Zona Reservada Allpahuayo-Mishana. Folia Amaz. 14: 11–28.

- 1 GARCÍA-VILLACORTA, R., N. DÁVILA, R. FOSTER, I. HUAMANTUPA, and C. VRIESENDORP. 2010.
 2 Flora y Vegetación/Flora and Vegetation. *In* M. P. Gilmore, C. Vriesendorp, W. S.
 3 Alverson, A. del Campo, R. von May, C. W. López, and S. O. Ríos (Eds.) Perú: Maijuna.
 4 Rapid Biological Inventories Report 22. pp. 176–182, The Field Museum of Natural
 5 History, Chicago.
- 6 GARCÍA-VILLACORTA, R., and B. E. HAMMEL. 2004. A noteworthy new species of *Tovomita*
 7 (*Clusiaceae*) from Amazonian white sand forests of Peru and Colombia. *Brittonia* 56:
 8 132–135.
- 9 GARCÍA-VILLACORTA, R., I. HUAMANTUPA, Z. CORDERO, N. PITMAN, and C. VRIESENDORP. 2011.
 10 Flora y Vegetación/Flora and Vegetation. *In* N. C. A. Pitman, C. Vriesendorp, D. K.
 11 Moskovitz, R. von May, D. Alvira, T. Wachter, D. F. Stotz, and A. del Campo (Eds.)
 12 Perú: Yaguas-Cotuhé. Rapid Biological Inventories Report 23. pp. 86–97, The Field
 13 Museum of Natural History, Chicago.
- 14 GENTRY, A. H. 1986. Endemism in tropical versus temperate plant communities. *In* M. E. Soulé
 15 (Ed.) *Conservation Biology: the science of scarcity and diversity*. pp. 153–181, Sinauer
 16 Associates, Sunderland, Massachusetts.
- 17 GENTRY, A. H., and R. ORTIZ. 1993. Patrones de composición florística en la Amazonía
 18 peruana. *In* R. Kalliola, M. Puhakka, and W. Danjoy (Eds.) R. Kalliola, M. Puhakka, and
 19 W. Danjoy. *Amazonía peruana: vegetación húmeda en el llano subandino*. Proyecto
 20 Amazonia, Universidad de Turku (PAUT) y Oficina Nacional de Evaluación de Recursos
 21 Naturales (ONERN), Jyväskylä. pp. 155–166, PAUT-ONERN.
- 22 GIRALDO-CAÑAS, D. 2001. Relaciones fitogeográficas de las sierras y afloramientos rocosos de
 23 la Guayana colombiana: un estudio preliminar. *Rev. Chil. Hist. Nat.* 74: 353–364.
- 24 GRAVEL, D., N. MOUQUET, M. LOREAU, and F. GUICHARD. 2010. Patch Dynamics, Persistence,
 25 and Species Coexistence in Metaecosystems. 176: 289–302.
- 26 HAMMOND, D. S. 2005a. Biophysical features of the Guiana Shield. *In* D. S. Hammond (Ed.)
 27 *Tropical forests of the Guiana Shield*. pp. 15–194, CABI Publishing.
- 28 HAMMOND, D. S. 2005b. *Tropical forests of the Guiana shield: ancient forests in a modern world*.
 29 CABI.
- 30 HERMOZA, W., S. BRUSSET, P. BABY, W. GIL, M. RODDAZ, N. GUERRERO, and R. BOLAÑOS. 2005.
 31 The Huallaga foreland basin evolution: Thrust propagation in a deltaic environment,
 32 northern Peruvian Andes. *J. South Am. Earth Sci.* 19: 21–34.
- 33 HOLT, R. D. 1993. Ecology at the mesoscale: the influence of regional processes on local
 34 communities. *In* R. E. Ricklefs and D. Schluter (Eds.) *Species diversity in ecological*
 35 *communities*. pp. 77–88, University of Chicago Press, Chicago.
- 36 HOORN, C. 1994. Fluvial palaeoenvironments in the intracratonic Amazonas Basin (Early
 37 Miocene-early Middle Miocene, Colombia). *Palaeogeogr. Palaeoclimatol. Palaeoecol.*
 38 109: 1–54.

- 1 HORBE, A. M. C., M. A. HORBE, and K. SUGUIO. 2004. Tropical Spodosols in northeastern
2 Amazonas State, Brazil. *Geoderma* 119: 55–68.
- 3 JORGENSEN, P. M., and S. LEÓN-YÁNEZ eds. 1999. Catalogue of the vascular plants of Ecuador.
4 Botanical Garden Press, Saint Louis, MO.
- 5 KLINGE, H. 1965. Podzol Soils in the Amazon Basin. *J. Soil Sci.* 16: 95–103.
- 6 KOLEFF, P., K. J. GASTON, and J. J. LENNON. 2003. Measuring beta diversity for presence–
7 absence data. *J. Anim. Ecol.* 72: 367–382.
- 8 KUBITZKI, K. 1989. Amazon lowland and Guayana highland- historical and ecological aspects of
9 the development of their floras. *Amaz. Kiel* 11: 1–12.
- 10 LÄHTEENOJA, O., and S. PAGE. 2011. High diversity of tropical peatland ecosystem types in the
11 Pastaza-Marañón basin, Peruvian Amazonia. *J. Geophys. Res. Biogeosciences* 116:
12 G02025.
- 13 LÄHTEENOJA, O., K. RUOKOLAINEN, L. SCHULMAN, and M. OINONEN. 2009. Amazonian peatlands:
14 an ignored C sink and potential source. *Glob. Change Biol.* 15: 2311–2320.
- 15 LATHAM, R. E., and R. E. RICKLEFS. 1993. Global Patterns of Tree Species Richness in Moist
16 Forests: Energy-Diversity Theory Does Not Account for Variation in Species Richness.
17 *Oikos* 67: 325–333.
- 18 LAURANCE, S. G. W., W. F. LAURANCE, A. ANDRADE, P. M. FEARNESIDE, K. E. HARMS, A. VICENTINI,
19 and R. C. C. LUIZÃO. 2010. Influence of soils and topography on Amazonian tree
20 diversity: a landscape-scale study. *J. Veg. Sci.* 21: 96–106.
- 21 LEGENDRE, P., and L. F. LEGENDRE. 2012. Numerical ecology 3rd edition. Elsevier, Amsterdam.
- 22 LOREAU, M., N. MOUQUET, and R. D. HOLT. 2003. Meta-ecosystems: a theoretical framework for
23 a spatial ecosystem ecology. *Ecol. Lett.* 6: 673–679.
- 24 LUCAS, Y., A. CHAUVEL, R. BOULET, G. RANZANI, and F. SCATOLINI. 1984. Transição latossolos-
25 podzóis sobre a Formação Barreiras na região de Manaus, Amazônia. *Rev. Bras. Ciênc.*
26 *Solo* 8: 325–335.
- 27 LUCAS, Y., C. R. MONTES, S. MOUNIER, M. LOUSTAU CAZALET, D. ISHIDA, R. ACHARD, C. GARNIER,
28 B. COULOMB, and A. J. MELFI. 2012. Biogeochemistry of an Amazonian podzol-ferralsol
29 soil system with white kaolin. *Biogeosciences* 9: 3705–3720.
- 30 MACEDO, M., and G. T. PRANCE. 1978. Notes on the vegetation of Amazonia II. The dispersal of
31 plants in Amazonian white sand campinas: the campinas as functional islands. *Brittonia*
32 30: 203–215.
- 33 MANTEL, N. 1967. The Detection of Disease Clustering and a Generalized Regression
34 Approach. *Cancer Res.* 27: 209–220.
- 35 MENDONÇA, B. A. F. DE, F. N. B. SIMAS, C. E. G. R. SCHAEFER, E. I. FERNANDES FILHO, J. F. DO
36 VALE JÚNIOR, and J. G. F. DE MENDONÇA. 2014. Podzolized soils and paleoenvironmental

- 1 implications of white-sand vegetation (Campinarana) in the Viruá National Park, Brazil.
2 *Geoderma Reg.* 2–3: 9–20.
- 3 MINCHIN, P. R. 1987. An evaluation of the relative robustness of techniques for ecological
4 ordination. *In* I. C. Prentice and E. van der Maarel (Eds.) *Theory and models in*
5 *vegetation science. Advances in vegetation science.* pp. 89–107, Springer Netherlands.
- 6 NEKOLA, J. C., and P. S. WHITE. 1999. The Distance Decay of Similarity in Biogeography and
7 *Ecology. J. Biogeogr.* 26: 867–878.
- 8 OKSANEN, J., F. G. BLANCHET, R. KINDT, P. LEGENDRE, P. R. MINCHIN, R. B. O'HARA, G. L.
9 SIMPSON, P. SOLYMOS, M. H. H. STEVENS, and H. WAGNER. 2015. *vegan: Community*
10 *Ecology Package Available at: <http://cran.r-project.org/web/packages/vegan/index.html>*
11 *[Accessed May 20, 2015].*
- 12 OLSON, D. M., E. DINERSTEIN, E. D. WIKRAMANAYAKE, N. D. BURGESS, G. V. N. POWELL, E. C.
13 UNDERWOOD, J. A. D'AMICO, I. ITOUA, H. E. STRAND, J. C. MORRISON, C. J. LOUCKS, T. F.
14 ALLNUTT, T. H. RICKETTS, Y. KURA, J. F. LAMOREUX, W. W. WETTENGEL, P. HEDAO, and K.
15 R. KASSEM. 2001. Terrestrial Ecoregions of the World: A New Map of Life on Earth A
16 new global map of terrestrial ecoregions provides an innovative tool for conserving
17 biodiversity. *BioScience* 51: 933–938.
- 18 PHILLIPS, O. L., P. N. VARGAS, A. L. MONTEAGUDO, A. P. CRUZ, M.-E. CHUSPEZANS, W. G.
19 SÁNCHEZ, M. YLI-HALLA, and S. ROSE. 2003. Habitat Association among Amazonian
20 Tree Species: A Landscape-Scale Approach. *J. Ecol.* 91: 757–775.
- 21 PITMAN, N. C. A., J. E. G. ANDINO, M. AULESTIA, C. E. CERÓN, D. A. NEILL, W. PALACIOS, G.
22 RIVAS-TORRES, M. R. SILMAN, and J. W. TERBORGH. 2014. Distribution and abundance of
23 tree species in swamp forests of Amazonian Ecuador. *Ecography* 37: 902–915.
- 24 POTTER, P. E. 1994. Modern sands of South America: composition, provenance and global
25 significance. *Geol. Rundsch.* 83: 212–232.
- 26 PRANCE, G. T. 1996. Islands in Amazonia. *Philos. Trans. R. Soc. B Biol. Sci.* 351: 823–833.
- 27 PRANCE, G. T., and H. O. SCHUBART. 1978. Notes on the vegetation of Amazonia I. A preliminary
28 note on the origin of the open white sand campinas of the lower Rio Negro. *Brittonia* 30:
29 60–63.
- 30 RÄSÄNEN, M., and A. LINNA. 1998. Geología y geofomas del area de Iquitos, Peru. *In* R. Kalliola
31 and Paredes (Eds.) *Geoecología y desarrollo integrado de la zona de Iquitos, Perú.* pp.
32 60–136.
- 33 R CORE TEAM. 2015. R: A language and environment for statistical computing. R Foundation for
34 Statistical Computing, Vienna, Austria Available at: <http://www.R-project.org/>.
- 35 REVILLA, J. 1974. Descripción de los tipos de vegetación en Mishana, Río Nanay. Loreto, Perú.
36 PAHO. Project AMRO-0719 Report. Pan Am. Health Organ. Wash. DC.
- 37 RICHARDS, P. W. 1941. Lowland tropical podzols and their vegetation. *Nature* 148: 129–131.

- 1 RICKLEFS, R. E. 1987. Community Diversity: Relative Roles of Local and Regional Processes.
2 Science 235: 167–171.
- 3 RICKLEFS, R. E. 2008. Disintegration of the Ecological Community. *Am. Nat.* 172: 741–750.
- 4 ROSSETTI, D. F., T. C. BERTANI, H. ZANI, E. H. CREMON, and E. H. HAYAKAWA. 2012. Late
5 Quaternary sedimentary dynamics in Western Amazonia: Implications for the origin of
6 open vegetation/forest contrasts. *Geomorphology* 177–178: 74–92.
- 7 ROSSI, J. P. 1996. Statistical tool for soil biology : 11. Autocorrelogram and Mantel test. *Eur. J.*
8 *Soil Biol.* 32: 195–203.
- 9 SEYLER, F., F. MULLER, G. COCHONNEAU, L. GUIMARÃES, and J. L. GUYOT. 2009. Watershed
10 delineation for the Amazon sub-basin system using GTOPO30 DEM and a drainage
11 network extracted from JERS SAR images. *Hydrol. Process.* 23: 3173–3185.
- 12 SHMIDA, A., and M. V. WILSON. 1985. Biological Determinants of Species Diversity. *J. Biogeogr.*
13 12: 1–20.
- 14 SILVEIRA, M. 2003. *Vegetação e flora das campinaranas do Sudoeste Amazônico*. Universidade
15 Federal do Acre, Departamento de Ciências da natureza.
- 16 SIMPSON, G. G. 1943. Mammals and the nature of continents. *Am. J. Sci.* 241: 1–31.
- 17 SINCHI. 2013. Herbario virtual. Herbario Amazónico Colombiano (COAH). Available at:
18 http://www.sinchi.org.co/coleccionesbiologicas/index.php?option=com_herbariov_oc&Itemid=29 [Accessed January 1, 2013].
19
- 20 SOKAL, R. R., and F. J. ROHLF. 1962. The Comparison of Dendrograms by Objective Methods.
21 *Taxon* 11: 33–40.
- 22 SOKAL, R. R., and P. H. A. SNEATH. 1963. *Principles of Numerical Taxonomy*. W. H. Freeman.
- 23 SPRUCE, R. 1908. *Notes of a Botanist on the Amazon and Andes*. Macmillan, London.
- 24 STALLARD, R. F. 1988. Weathering and Erosion in the Humid Tropics. *In* A. Lerman and M.
25 Meybeck (Eds.) *Physical and Chemical Weathering in Geochemical Cycles*. NATO ASI
26 Series. pp. 225–246, Springer Netherlands.
- 27 STALLARD, R. F. 2011. Landscape processes: geology, hidrology, and soils. *In* N. C. A. Pitman,
28 C. Vriesendorp, D. K. Moskovitz, R. von May, D. Alvira, T. Wachter, D. F. Stotz, and A.
29 del Campo (Eds.) *Perú: Yaguas-Cotuhé. Rapid Biological Inventories Report 23*. pp.
30 199–210, The Field Museum of Natural History, Chicago.
- 31 STRUWE, L., and V. A. ALBERT. 2004. A Monograph of Neotropical *Potalia Aublet* (Gentianaceae:
32 *Potalieae*). *Syst. Bot.* 29: 670–701.
- 33 SUZUKI, R., and H. SHIMODAIRA. 2006. Pvcust: an R package for assessing the uncertainty in
34 hierarchical clustering. *Bioinformatics* 22: 1540–1542.

- 1 THE ANGIOSPERM PHYLOGENY GROUP. 2009. An update of the Angiosperm Phylogeny Group
2 classification for the orders and families of flowering plants: APG III. *Bot. J. Linn. Soc.*
3 161: 105–121.
- 4 THE PLANT LIST. 2013. Version 1.1. Published on the Internet. <http://www.theplantlist.org/>
5 (accessed 1st March). Available at: <http://www.theplantlist.org/> [Accessed March 18,
6 2013].
- 7 TNRS. 2013. The Taxonomic Name Resolution Service.
- 8 TROPICOS-BOLIVIA. 2013. Catalogue of the vascular plant of Bolivia. Available at:
9 <http://www.tropicos.org/Project/BO> [Accessed January 1, 2013].
- 10 TROPICOS-ECUADOR. 2013. Catalogue of the vascular plant of Ecuador. Available at:
11 <http://www.tropicos.org/Project/CE> [Accessed January 1, 2013].
- 12 TROPICOS-PERU. 2013. Catalogue of the vascular plant of Peru. Available at:
13 <http://www.tropicos.org/Project/PE> [Accessed January 1, 2013].
- 14 TUOMISTO, H. 2010. A diversity of beta diversities: straightening up a concept gone awry. Part 2.
15 Quantifying beta diversity and related phenomena. *Ecography* 33: 23–45.
- 16 TUOMISTO, H., A. LINNA, and R. KALLIOLA. 1994. Use of digitally processed satellite images in
17 studies of tropical rain forest vegetation. *Int. J. Remote Sens.* 15: 1595–1610.
- 18 VICENTINI, A. 2004. A vegetação ao longo de um gradiente edáfico no Parque Nacional do Jaú.
19 *In* H. Borges, S. Iwanaga, C. C. Duriganand, and M. R. Pinheiro (Eds.) *Janelas para a*
20 *biodiversidade no Parque Nacional do Jaú: uma estratégia par ao estudo da*
21 *biodiversidade na Amazônia.* pp. 117–143, Fundação Vitória Amazônica/WWF/IBAMA,
22 Manaus.
- 23 VORMISTO, J., J.-C. SVENNING, P. HALL, and H. BALSLEV. 2004. Diversity and Dominance in Palm
24 (Arecaceae) Communities in Terra firme Forests in the Western Amazon Basin. *J. Ecol.*
25 92: 577–588.
- 26 VORMISTO, J., H. TUOMISTO, and J. OKSANEN. 2004. Palm distribution patterns in Amazonian
27 rainforests: What is the role of topographic variation? *J. Veg. Sci.* 15: 485–494.
- 28

FIGURE LEGENDS

FIGURE 1. The Amazon and the Guiana Shield region (dashed area) with political division acronyms used in the floristic analysis overlaid on an elevation map (darker areas indicate higher elevations). Approximate locations of studied white-sand forests: 1 = acre.wsf.BR (Acre region, Brazil), 2 = loreto.wsf.PE (Loreto region, Peru), 3 = caquetá.wsf.CO (Caquetá region, Colombia), 4 = guainía.wsf.CO (Guainía region, Colombia). Province acronyms in Appendix 1.

FIGURE 2. Proportion of western Amazon white-sand specialists shared among different ecoregions within Amazonia constructed using (a) only loreto.wsf.PE and acre.wsf.BR, and (b) full white-sand dataset. non-GS WA ws = non-Guiana Shield western Amazon white-sand specialists. WS specialists = full list of white-sand specialists. Sites 1 and 2 = non-Guiana Shield western Amazon white-sand areas; 3 and 4 = Guiana Shield western Amazon white-sand sites. GS limit = Guiana Shield limit. Note change in the proportion and ecoregions of shared white-sand specialists according to the dataset used. Site names as in Fig. 1.

FIGURE 3. Relationships of western Amazon white-sand forests with provinces/states in the Amazon and Guiana regions as represented by non-metric Multidimensional Scaling (NMDS) ordination. Lines connecting the sites represent cluster analysis result. WSF = western Amazon white-sand forests. Province acronyms in Appendix 1.

FIGURES

Figure 1





Figure. 2.

a)



b)





Figure 3



Appendix 1. Number of vascular plant species and source used to build the database ordered by ascending number of species. Studied western Amazon white-sand forests in bold.

Site code	Province	Area (km ²)	Country	No. spp.	Source
acre.wsf.BR	Acre	-	Brazil	363	(Silveira 2003, Ferreira 2009)
CO.VD	Vichada	38,734.93	Colombia	607	(SINCHI 2013)
caquetá.wsf.CO	Caquetá	-	Colombia	657	(Cortés <i>et al.</i> 1998, Arbeláez & Duivenvoorden 2004)
loreto.wsf.PE	Loreto	-	Peru	731	(García-Villacorta <i>et al.</i> 2003, Fine <i>et al.</i> 2010, Tropicós-Peru 2013)
guainía.wsf.CO	Guainía	-	Colombia	955	(Cárdenas-

López 2007)





PE.UC	Ucayali	105,078.38	Peru	(Brako & Zarucchi 1993, Tropicós-Peru 2013)
CO.GN	Guainía	68,819.14	Colombia	(SINCHI 2013)
CO.VP	Vaupés	53,242.63	Colombia	(SINCHI 2013)
VE.DA	Delta Amacuro	38,230.4	Venezuela	(Funk <i>et al.</i> 2007)
CO.GV	Guaviare	55,570.16	Colombia	(SINCHI 2013)
PE.MD	Madre de Dios	84,420.81	Peru	(Brako & Zarucchi 1993, Tropicós-Peru 2013)
BR.AP	Amapá	141,105.57	Brazil	(Forzza <i>et al.</i> 2010a, b, Flora

				do Brasil 2013)
BR.RR	Roraima	224,384.08	Brazil	(Forzza <i>et al.</i> 2010a, b, Flora do Brasil 2013)
BR.RO	Rondonia	236,376.67	Brazil	(Forzza <i>et al.</i> 2010a, b, Flora do Brasil 2013)
CO.CQ	Caquetá	90,029.75	Colombia	(SINCHI 2013)
BR.MT	Mato Grosso	599,681.47	Brazil	(Forzza <i>et al.</i> 2010a, b, Flora do Brasil 2013)
CO.AM	Amazonas	107,462.42	Colombia	(SINCHI 2013)
BR.AC	Acre	152,729.51	Brazil	(Forzza <i>et al.</i>

				2010a, b, Flora do Brasil 2013)
SR	Surinam	146,011.49	Surinam	(Funk <i>et al.</i> 4886 2007)
GF	French Guiana	83,014.98	French Guiana	(Funk <i>et al.</i> 5166 2007)
PE.LO	Loreto	375,550.2	Perú	(Brako & Zarucchi 1993, Tropicos-Peru 5271 2013)
BO.AM	Amazon region	681,909.86	Bolivia	(Tropicos- Bolivia 2013) 5378
BR.PA	Pará	892,481.47	Brazil	(Forzza <i>et al.</i> 2010a, b, Flora do Brasil 6098 2013)
GY	Guyana	209,549.47	Guyana	(Funk <i>et al.</i> 6938 2007)

VE.BO	Bolivar	193,997.81	Venezuela	(Funk <i>et al.</i> 2007)
VE.AM	Amazonas	179,579.86	Venezuela	(Funk <i>et al.</i> 2007)
ECU	Ecuadorian Amazon	105653.6	Ecuador	(Jorgensen & León-Yáñez 1999, Tropicos- Ecuador 2013)
BR.AM	Amazonas	1'570,659.01	Brazil	(Forzza <i>et al.</i> 2010a, b, Flora do Brasil 2013)

Appendix 2. Cophenetic correlation results of the five assessed clustering algorithms. The highest cophenetic correlation (Average linkage, $r = 0.629$) is the one that best represent the dissimilarity matrix in its resulting dendrogram followed by Complete clustering ($r = 0.579$).



Appendix 3. Summary statistics and ecological preferences of 1,180 vascular plant species found in four western Amazonian white-sand forests.

Ecological preference	No. families with species in these categories	No. genera with species in these categories	No. species (%)	Records
habitat generalist	103	295	509 (43)	51,790
poor-soil specialist	89	208	406 (34)	14,723
white-sand specialist	65	160	277 (23)	3473
Total	133	491	1180 (100)	69,986

Appendix 4. Species in the western Amazon white-sand specialist dataset with records found within the Guiana Shield region.

Family/Species	No. records
Acanthaceae	1
<i>Aphelandra lamprantha</i>	1
Annonaceae	47
<i>Anaxagorea brachycarpa</i>	14
<i>Anaxagorea manausensis</i>	1
<i>Annona paludosa</i>	5
<i>Bocageopsis canescens</i>	8
<i>Cymbopetalum alkekengi</i>	0
<i>Duguetia arenicola</i>	2
<i>Guatteriaopsis sessiliflora</i>	0
<i>Oxandra asbeckii</i>	4
<i>Oxandra leucodermis</i>	3
<i>Pseudoxandra revoluta</i>	0
<i>Tetrameranthus duckei</i>	10
<i>Tetrameranthus pachycarpus</i>	0
<i>Trigynaea lanceipetala</i>	0
Apocynaceae	54
<i>Mandevilla annulariifolia</i>	6
<i>Mandevilla caurensis</i>	4
<i>Mandevilla nerioides</i>	8
<i>Molongum lucidum</i>	4
<i>Odontadenia killipii</i>	6
<i>Odontadenia kochii</i>	7
<i>Parahancornia peruviana</i>	2
<i>Parahancornia surrogata</i>	17
Aquifoliaceae	30
<i>Ilex divaricata</i>	28
<i>Ilex spruceana</i>	2
Araceae	5
<i>Philodendron auyantepuiense</i>	5
Araliaceae	11
<i>Schefflera japurensis</i>	11
Arecaceae	11
<i>Euterpe catinga</i>	5
<i>Mauritia carana</i>	6

Asteraceae	26
<i>Calea abelioides</i>	16
<i>Chromolaena tyleri</i>	2
<i>Gongylolepis martiana</i>	7
<i>Stenopadus colombianus</i>	1
Begoniaceae	1
<i>Begonia lutea</i>	1
Bignoniaceae	54
<i>Anemopaegma oligoneuron</i>	9
<i>Digomphia densicoma</i>	34
<i>Distictella obovata</i>	11
Bonnetiaceae	64
<i>Bonnetia paniculata</i>	11
<i>Bonnetia sessilis</i>	53
Bromeliaceae	67
<i>Ananas paraguayensis</i>	23
<i>Brocchinia hechtoides</i>	23
<i>Brocchinia paniculata</i>	8
<i>Pitcairnia patentiflora</i>	1
<i>Vriesea socialis</i>	12
Burmanniaceae	6
<i>Burmannia dasyantha</i>	5
<i>Burmannia vaupesana</i>	1
Burseraceae	44
<i>Protium calanense</i>	5
<i>Protium carolense</i>	16
<i>Protium heptaphyllum</i> subsp. <i>ulei</i>	18
<i>Protium leptostachyum</i>	5
Calophyllaceae	34
<i>Caraipa longipedicellata</i>	8
<i>Caraipa savannarum</i>	19
<i>Caraipa tereticaulis</i>	6
<i>Caraipa utilis</i>	0
<i>Haploclathra cordata</i>	1
<i>Haploclathra paniculata</i> var. <i>paniculata</i>	0
Chrysobalanaceae	92
<i>Couepia amaraliae</i>	6
<i>Couepia bracteosa</i>	11
<i>Couepia canomensis</i>	7
<i>Couepia williamsii</i>	0
<i>Exellodendron coriaceum</i>	9
<i>Hirtella punctillata</i>	12
<i>Hirtella revillae</i>	0
<i>Licania gracilipes</i>	7

<i>Licania hebantha</i>	4
<i>Licania intrapetiolaris</i>	28
<i>Licania stewardii</i>	5
<i>Parinari montana</i>	3
Clusiaceae	77
<i>Clusia cerroana</i>	23
<i>Clusia chiribiquetensis</i>	15
<i>Clusia fockeana</i>	9
<i>Clusia huberi</i>	11
<i>Clusia opaca</i>	8
<i>Clusia renggerioides</i>	11
<i>Tovomita calophyllophylla</i>	0
Combretaceae	7
<i>Terminalia yapacana</i>	7
Connaraceae	1
<i>Connarus guggenheimii</i>	1
Droseraceae	5
<i>Drosera esmeraldae</i>	5
Ebenaceae	6
<i>Diospyros myrmecocarpa</i>	3
<i>Diospyros tessmannii</i>	0
<i>Lissocarpa kating</i>	3
Emmotaceae	0
<i>Emmotum floribundum</i>	0
Ericaceae	14
<i>Vaccinium euryanthum</i>	14
Eriocaulaceae	127
<i>Paepalanthus fasciculatus</i>	34
<i>Paepalanthus formosus</i>	4
<i>Rondonanthus capillaceus</i>	27
<i>Syngonanthus allenii</i>	1
<i>Syngonanthus reflexus</i>	29
<i>Syngonanthus simplex</i>	7
<i>Syngonanthus tenuis</i> var. <i>bulbifer</i>	7
<i>Syngonanthus trichophyllus</i>	11
<i>Syngonanthus umbellatus</i>	7
Erythroxylaceae	13
<i>Erythroxylum kapplerianum</i>	13
Euphorbiaceae	117
<i>Conceveiba terminalis</i>	23
<i>Croton araracuarae</i>	3
<i>Croton chiribiquetensis</i>	1
<i>Croton mollis</i>	10
<i>Croton scutatus</i>	8

<i>Croton spiraeifolius</i>	3
<i>Hevea guianensis</i> var. <i>lutea</i>	4
<i>Hevea nitida</i>	9
<i>Hevea nitida</i> var. <i>toxicodendroides</i>	1
<i>Mabea subsessilis</i>	18
<i>Micrandra elata</i>	3
<i>Micrandra spruceana</i>	18
<i>Senefelderopsis chiribiquetensis</i>	16
Euphroniaceae	19
<i>Euphronia hirtelloides</i>	19
Fabaceae	145
<i>Abarema barbouriana</i> var. <i>arenaria</i>	6
<i>Abarema microcalyx</i>	9
<i>Aldina heterophylla</i>	3
<i>Calliandra vaupesiana</i>	7
<i>Dicymbe uaiparuensis</i>	2
<i>Dimorphandra macrostachya</i> subsp. <i>glabrifolia</i>	2
<i>Dimorphandra vernicosa</i>	17
<i>Heterostemon mimosoides</i>	21
<i>Inga gereauana</i>	0
<i>Inga neblinensis</i>	8
<i>Jacqueshuberia loretensis</i>	0
<i>Macrolobium arenarium</i>	1
<i>Macrolobium discolor</i> var. <i>egranulosum</i>	3
<i>Macrolobium limbatum</i> var. <i>limbatum</i>	3
<i>Macrolobium limbatum</i> var. <i>propinquum</i>	0
<i>Macrolobium molle</i>	8
<i>Macrolobium suaveolens</i>	7
<i>Senna kuhlmannii</i>	0
<i>Swartzia benthamiana</i> var. <i>benthamiana</i>	13
<i>Tachigali ptychophysca</i>	4
<i>Taralea oppositifolia</i>	31
Gentianaceae	30
<i>Curtia conferta</i>	3
<i>Irlbachia nemorosa</i>	12
<i>Tachia guianensis</i>	15
<i>Tachia loretensis</i>	0
Humiriaceae	43
<i>Humiria balsamifera</i>	13
<i>Humiria balsamifera</i> var. <i>balsamifera</i>	9
<i>Humiria balsamifera</i> var. <i>floribunda</i>	1
<i>Humiria crassifolia</i>	11
<i>Humiriastrum piraparanense</i>	9
<i>Vantanea spichigeri</i>	0

Hymenophyllaceae	17
<i>Trichomanes arbuscula</i>	8
<i>Trichomanes sprucei</i>	1
<i>Trichomanes superbum</i>	2
<i>Trichomanes vandenboschii</i>	1
<i>Trichomanes vittaria</i>	5
Icacinaceae	28
<i>Emmotum acuminatum</i>	6
<i>Emmotum floribundum</i>	18
<i>Pleurisanthes flava</i>	4
Lauraceae	13
<i>Aniba heterotepala</i>	0
<i>Endlicheria arunciflora</i>	5
<i>Endlicheria chrysovelutina</i>	0
<i>Endlicheria citriodora</i>	1
<i>Mezilaurus opaca</i>	0
<i>Mezilaurus triunca</i>	0
<i>Ocotea alata</i>	0
<i>Ocotea atrata</i>	2
<i>Ocotea esmeraldana</i>	4
<i>Ocotea immersa</i>	1
<i>Sextonia pubescens</i>	0
Lindsaeaceae	27
<i>Lindsaea meifolia</i>	6
<i>Lindsaea pendula</i>	11
<i>Lindsaea rigidiuscula</i>	10
Loganiaceae	20
<i>Bonyunia minor</i>	20
<i>Potalia coronata</i>	0
Lythraceae	15
<i>Cuphea annulata</i>	13
<i>Cuphea kubeorum</i>	2
Malpighiaceae	29
<i>Blepharandra heteropetala</i>	5
<i>Byrsonima amoena</i>	8
<i>Diacidia galphimoides</i>	16
Malvaceae	18
<i>Pachira amazonica</i>	5
<i>Pachira brevipes</i>	5
<i>Pachira coriacea</i>	5
<i>Pachira fuscolepidota</i>	3
Melastomataceae	97
<i>Acanthella sprucei</i>	23
<i>Clidemia epibaterium</i>	12

<i>Clidemia heteroneura</i>	12
<i>Clidemia pycnaster</i>	3
<i>Comolia microphylla</i>	15
<i>Ernestia tenella</i>	4
<i>Graffenrieda fantastica</i>	3
<i>Macairea theresiae</i>	4
<i>Miconia crassinervia</i>	4
<i>Pachyloma coriaceum</i>	5
<i>Siphanthera fasciculata</i>	1
<i>Tococa rotundifolia</i>	11
Metaxyaceae	0
<i>Metaxya lanosa</i>	0
Myrtaceae	40
<i>Myrcia clusiifolia</i>	20
<i>Myrcia revolutifolia</i>	20
Nyctaginaceae	14
<i>Neea obovata</i>	14
Ochnaceae	62
<i>Ouratea roraimae</i>	12
<i>Ouratea spruceana</i>	21
<i>Sauvagesia erioclada</i>	1
<i>Sauvagesia fruticosa</i>	10
<i>Sauvagesia guianensis subsp. araracuarensis</i>	1
<i>Wallacea insignis</i>	17
Olacaceae	8
<i>Chaunochiton angustifolium</i>	7
<i>Curupira tefeensis</i>	1
Orchidaceae	41
<i>Duckeella pauciflora</i>	2
<i>Encyclia leucantha</i>	5
<i>Epistephium hernandii</i>	9
<i>Epistephium lucidum</i>	4
<i>Epistephium parviflorum</i>	16
<i>Octomeria erosilabia</i>	2
<i>Octomeria taracuana</i>	1
<i>Stenocoryne longicornis</i>	1
<i>Trichosalpinx orbicularis</i>	1
Orobanchaceae	4
<i>Buchnera rubriflora</i>	4
Pentaphragmaceae	27
<i>Ternstroemia campincola</i>	3
<i>Ternstroemia klugiana</i>	0
<i>Ternstroemia punctata</i>	5
<i>Ternstroemia pungens</i>	19

Phyllanthaceae	4
<i>Phyllanthus myrsinites</i>	4
Piperaceae	2
<i>Piper fonteboanum</i>	1
<i>Piper froesii</i>	1
<i>Piper mituense</i>	0
Polypodiaceae	16
<i>Cochlidium tepuiense</i>	16
Primulaceae	7
<i>Cybianthus reticulatus</i>	7
Proteaceae	1
<i>Euplassa saxicola</i>	1
Rapateaceae	42
<i>Duckea junciformis</i>	7
<i>Duckea squarrosa</i>	9
<i>Monotrema arthrophyllum</i>	5
<i>Monotrema xyridoides</i>	15
<i>Rapatea elongata</i>	4
<i>Schoenocephalium martianum</i>	2
Rhabdodendraceae	17
<i>Rhabdodendron amazonicum</i>	17
Rubiaceae	537
<i>Ferdinandusa guainiae</i>	6
<i>Ferdinandusa hirsuta</i>	3
<i>Ferdinandusa sprucei</i>	5
<i>Ixora intensa</i>	9
<i>Ladenbergia lambertiana</i>	57
<i>Pagamea acrensis</i>	16
<i>Pagamea aracaensis</i>	5
<i>Pagamea coriacea</i>	49
<i>Pagamea guianensis</i>	54
<i>Pagamea macrophylla</i>	4
<i>Pagamea montana</i>	21
<i>Pagamea plicata</i>	17
<i>Pagamea thyrsoiflora</i>	40
<i>Palicourea nitidella</i>	64
<i>Platycarpum orinocense</i>	1
<i>Platycarpum orinocense</i> var. <i>orinocense</i>	5
<i>Remijia macrocnemia</i>	9
<i>Remijia roraimae</i>	13
<i>Retiniphyllum concolor</i>	22
<i>Retiniphyllum scabrum</i>	40
<i>Retiniphyllum schomburgkii</i>	72
<i>Retiniphyllum truncatum</i>	18

<i>Rudgea krukovii</i>	1
<i>Sipaneopsis huberi</i>	4
<i>Stachyococcus adinanthus</i>	2
Rutaceae	17
<i>Adiscanthus fusciflorus</i>	5
<i>Decagonocarpus cornutus</i>	5
<i>Decagonocarpus oppositifolius</i>	6
<i>Hortia vandelliana</i>	1
<i>Leptothyrsa sprucei</i>	0
<i>Ravenia biramosa</i> var. <i>peruviana</i>	0
Salicaceae	62
<i>Euceraea nitida</i>	30
<i>Laetia coriacea</i>	20
<i>Laetia cupulata</i>	12
Sapindaceae	1
<i>Paullinia splendida</i> fo. <i>chrysocarpa</i>	1
Sapotaceae	67
<i>Chrysophyllum bombycinum</i>	0
<i>Chrysophyllum sanguinolentum</i> subsp. <i>spurium</i>	7
<i>Elaeoluma schomburgkiana</i>	8
<i>Pradosia schomburgkiana</i>	39
<i>Pradosia schomburgkiana</i> subsp. <i>schomburgkiana</i>	13
Schlegeliaceae	0
<i>Schlegelia cauliflora</i>	0
Selaginellaceae	6
<i>Selaginella coarctata</i>	6
Siparunaceae	3
<i>Siparuna micrantha</i>	3
Thymelaeaceae	17
<i>Tepuianthus colombianus</i>	2
<i>Tepuianthus savannensis</i>	15
Vochysiaceae	13
<i>Ruizterania retusa</i>	13
Xyridaceae	80
<i>Abolboda acicularis</i> var. <i>granularis</i>	2
<i>Abolboda americana</i>	9
<i>Abolboda grandis</i>	8
<i>Abolboda grandis</i> var. <i>grandis</i>	8
<i>Abolboda macrostachya</i> var. <i>macrostachya</i>	8
<i>Abolboda pulchella</i>	6
<i>Xyris araracuare</i>	2
<i>Xyris esmeraldae</i>	17
<i>Xyris lomatophylla</i>	5
<i>Xyris spruceana</i>	10

<i>Xyris subglabrata</i>	4
<i>Xyris wurdackii</i> subsp. <i>caquetensis</i>	1
Total No. Records	2503
Total No. Species	279

Appendix 5. Ecoregions with at least 20% of white-sand specialists using the (a) whole white-sand specialist dataset compared to using only the (b) non-Guiana Shield white-sand specialist dataset. Note changes in the proportions by ecoregions according to the dataset used.


a) Full white-sand dataset:
caquetá.wsf.CO (Colombia),
guainía.wsf.CO (Colombia),
acre.wsf.BR (Brazil), lore.wsf.PE
(Peru)

b) Non-Guiana Shield white-sand dataset: acre.wsf.BR (Brazil), loreto.wsf.PE (Peru)


	No.	%		No.	%
Ecoregion	species	species	Ecoregion	species	species
<hr/>					
Caquetá moist forests	191	69%	Iquitos varzea	86	97%
<hr/>					
Guayanan					
Highlands moist forests	142	51%	Napo moist forests	56	63%
<hr/>					
Negro-Branco moist forests	137	49%	Southwest Amazon moist forests	43	48%
<hr/>					
Iquitos varzea	95	34%	Negro-Branco moist forests	38	43%

			Guayanan Highlands		
Tepuis	93	34%	moist forests	29	33%
<hr/>					
Japurá-Solimoes- Negro moist forests	64	23%	Caquetá moist forests	27	30%
<hr/>					
Guianan moist forests	61	22%	Japurá- Solimoes- Negro moist forests	25	28%
<hr/>					
Napo moist forests	59	21%	Uatuma- Trombetas moist forests	25	28%
<hr/>					
Uatuma- Trombetas moist forests	56	20%	Solimoes- Japurá moist forest	22	25%
<hr/>					
Solimoes-Japurá moist forest	55	20%	Tepuis	21	24%


Appendix 6. Cluster analysis of Amazon and Guiana Shield sites with bootstrap support values using Average linkage as clustering algorithm and 1000 bootstrap runs. AU bootstrap p-values at each node (left), standard bootstrap values (right), and cluster number (below). Western Amazon white-sand forests are marked with asterisks: acre.wsf.BR (Brazil), caquetá.wsf.CO (Colombia), guainía.wsf.CO (Colombia), and loreto.wsf.PE (Peru).



Appendix 7. Standard error of AU bootstrap p-values of each identified cluster by pvelust. Cluster 6 showed relatively high standard error (appendix 5).



Appendix 8. Shepard plot of NMDS ordination plot of Amazon-Guianan floras including four western Amazonian white-sand forests. Stress = 0.26. Iterations = 500. Dimensions = 2.



Appendix 9. Floristic similarity as a function of geographic distance between pairs of floras in the Andes-Guiana Shield region. Statistical significance from Mantel test. To test whether geographical distance between the studied floras had an influence on the floristic patterns recovered by the ordination and cluster analysis, we performed a Mantel test (Mantel 1967, Rossi 1996, Dutilleul *et al.* 2000, Legendre & Legendre 2012). To create the distance matrix we extracted the geographical coordinates of each province by using its polygon centroid in a Geographical Information System (GIS) that were then used to calculate geographical (Euclidean) distances between floras. We ran 999 permutations on the floristic dissimilarity matrix at a significance p-value of 0.05. We found a significant positive Mantel correlation between the floristic distance and geographical distance matrices (Mantel's $r = 0.4866$, $p = 0.0001$). Based on this result we reject the null hypothesis that these two matrices are not related and hence the geographical separation of sites may explain a portion of the observed floristic dissimilarities. A Mantel test including only the four white-sand sites against their geographical distances also found a marginally significant result ($r = 0.5818$, $p = 0.083$).

