



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Noise-processing by signaling networks

Citation for published version:

Kontogeorgaki, S, Sánchez-García, RJ, Ewing, RM, Zygalakis, KC & MacArthur, BD 2017, 'Noise-processing by signaling networks' *Scientific Reports*, vol. 7, no. 1, 532, pp. 532. DOI: 10.1038/s41598-017-00659-x

Digital Object Identifier (DOI):

[10.1038/s41598-017-00659-x](https://doi.org/10.1038/s41598-017-00659-x)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Scientific Reports

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Noise-processing by signaling networks

Styliani Kontogeorgaki¹, Rubén J. Sánchez-García^{1,5}, Rob M. Ewing^{2,5}, Konstantinos C. Zygalkis³, and Ben D. MacArthur^{1,4,5,*}

¹Mathematical Sciences, University of Southampton, SO17 1BJ, UK

²Biological Sciences, University of Southampton, SO17 1BJ, UK

³School of Mathematics, University of Edinburgh, EH9 3FD, UK

⁴Centre for Human Development, Stem Cells and Regeneration, Faculty of Medicine, University of Southampton, SO17 1BJ, UK

⁵Institute for Life Sciences, University of Southampton, SO17 1BJ, UK

*Correspondence to bdm@soton.ac.uk

ABSTRACT

Signaling networks mediate environmental information to the cell nucleus. To perform this task effectively they must be able to integrate multiple stimuli and distinguish persistent signals from transient environmental fluctuations. However, the ways in which signaling networks process environmental noise are not well understood. Here we outline a mathematical framework that relates a network's structure to its capacity to process noise, and use this framework to dissect the noise-processing ability of signaling networks. We find that complex networks that are dense in directed paths are poor noise processors, while those that are sparse and strongly directional process noise well. These results suggest that while cross-talk between signaling pathways may increase the ability of signaling networks to integrate multiple stimuli, too much cross-talk may compromise the ability of the network to distinguish signal from noise. To illustrate these general results we consider the structure of the signaling network that maintains pluripotency in mouse embryonic stem cells, and find an incoherent feedforward loop structure involving Stat3, Tfc2l1, Esrrb, Klf2 and Klf4 is particularly important for noise-processing. Taken together these results suggest that noise-processing is an important function of signaling networks and they may be structured in part to optimize this task.

Introduction

Cellular identities are regulated by a variety of complex, interconnected, molecular regulatory networks, including signaling networks, metabolic networks and core transcriptional regulatory networks¹⁻⁹. Signaling networks are of particular importance in maintaining robust cellular identities since they mediate noisy environmental information from the local cellular micro-environment to the cell nucleus^{7,10-14}. In order to perform this task effectively they must be able to transmit complex environmental information robustly, and failure to do this has been linked to cancer initiation and progression, as well as defects in embryonic development¹⁵⁻¹⁷. Much of what is known about signaling networks comes from the detailed reductionist analysis of their constituent signaling pathways. Several of these have been studied in great detail, and the core components and biochemical mechanisms of signal transduction in pathways such as Wnt, TGF- β and MAP Kinase signaling are now well-defined. These signaling pathways function in a wide diversity of different biological processes and systems, and they are known to have a central role in maintaining pluripotency and specifying cell identities, for example^{18,19}. A long-standing question of interest is why, despite the myriad of biological processes and systems that involve signaling, there are only a few distinct, but widely conserved and re-used pathways²⁰. An emerging feature that may in part explain this observation is that signaling pathway 'modules' are interconnected in many different ways and cross-talk between pathways has been shown for most signaling pathways, with mechanisms ranging from transcriptional activation of pathway components through to direct interactions between proteins in different pathways²⁰. In addition to cross-talk between pathways, specific feedback mechanisms allow for the homeostatic control of signaling activity. A well-defined example in the Wnt signaling pathway is the transcriptional activation of the Dickkopf proteins which negatively regulate Wnt signaling by binding to Wnt receptors in response to Wnt activation²¹. However, while much is now known about the function of specific signaling pathways, very little known about how cross-talk between pathways affects information-processing. In the context of signal processing it has been suggested that promiscuity in the protein-protein interactions is a major source of intrinsic noise, and that signaling pathways have evolved features (for example receptor clustering) to better distinguish signal from noise²². Although some studies have considered the ways in which noise propagates through regulatory networks²³⁻²⁷, the general mechanisms by which signaling networks distinguish persistent environmental signals from the noise that is inherent to the cellular micro-environment are still not well understood. To address this problem, here we outline a general mathematical framework which relates a network's structure to its capacity to process noise, and use this framework to dissect the noise-processing ability of signaling

networks. As an illustrative example we examine the noise-processing ability of the network that maintains pluripotency in mouse embryonic stem cells.

Embryonic stem (ES) cells are found naturally in the pre-implantation embryo and are able to give rise to all embryonic lineages, a property known as pluripotency. The molecular basis of pluripotency has been extensively studied, and it is now known that activation of a small number of core transcription factors – including Oct3/4, Sox2, and Nanog along with other secondary factors such as Myc, Klf4 and Lin28 – is sufficient to maintain the pluripotent state^{5,6,28–35}. Indeed, forced expression of combinations of these factors in somatic cells is sufficient to induce pluripotency *de novo*^{28,30,36}. Although this central transcriptional circuit is self-sustaining when shielded from external stimulation³⁷, it is known that a network of signaling pathways which process extra-cellular environmental information are also essential both to maintenance of, and exit from, the pluripotent state³⁸. Importantly, while the core transcriptional circuitry is broadly similar in mouse and human pluripotent cells³³, their dependency on external signaling is markedly different: mouse ES cells are dependent on Lif/Stat signaling^{39,40}, Bmp⁴¹ and canonical Wnt³⁷ to promote self-renewal, while Fgf/Erk signaling disrupts pluripotency^{37,42–44}; by contrast human ES cell self-renewal is independent of Lif⁴⁵, yet requires Activin and Fgf^{46,47} signaling and human ES cells undergo differentiation when exposed to Bmp⁴⁷.

The remainder of the paper is organized as follows: we begin by outlining our general mathematical theory, as well as setting our assumptions, before establishing a mathematical formula that makes the connection between network structure and noise-processing explicit. To illustrate these results we then use this expression to investigate the structure of the regulatory network for pluripotency in mouse ES cells. This network is chosen since it is particularly well characterized³⁴ and so constitutes a good test model. We find that certain elements in this network, particularly incoherent feedforward structures, are particularly important for its noise-processing ability. Interestingly, these elements are distinct from the core feedback structures that are known to maintain the pluripotent ground state⁴⁸, suggesting that different portions of this network perform different regulatory tasks.

Results

Our concern is with how a network G processes noise from an external source. In the context of signaling networks the nodes in the network are molecules in the signaling cascades, and edges are regulatory interactions (e.g. phosphorylation etc.) between molecules. Since signaling networks pass information from the cell exterior to the nucleus, we assume that the network G is inherently directed: the presence of an edge (i, j) indicates that node i exerts a regulatory effect on node j but not necessarily vice versa. Since regulatory interactions may be activatory or inhibitory we also allow each edge to have positive or negative weight representing the strength of activation or inhibition respectively. We denote the weight of edge (i, j) by A_{ij} . Assuming that there are n nodes in G the $n \times n$ adjacency matrix \mathbf{A} then describes the strength of all interactions in the system.

In general the regulatory interactions between nodes may be highly nonlinear. However, to better understand the relationship between network structure and function we will assume here that the dynamics are linear. By doing so we are effectively considering the linearization near to a fixed point in the nonlinear dynamics; this rationale for studying the linear case has been taken elsewhere⁴⁹. In the absence of external fluctuations, the dynamics of the system are described by the following system of ordinary differential equations (ODEs):

$$\frac{d\mathbf{x}}{dt} = -\mathbf{M}\mathbf{x}, \quad \mathbf{M} = d\mathbf{I} - \mathbf{A}, \quad (1)$$

where \mathbf{x} is the vector of node states (for example, protein concentrations), \mathbf{I} is the $n \times n$ identity matrix, and we have assumed that all nodes decay at the same rate d , which then sets a timescale for the dynamics. Without loss of generality we may take $d = 1$ since this may always be achieved by suitable re-scaling. Given the linearity of this system there are only two possible types of long-term behavior: convergence toward a stable fixed point or divergence to infinity. We will assume that only the first behavior can happen, *i.e.* convergence to a stable fixed point is the only physically realistic scenario. This occurs whenever the real parts of the eigenvalues of \mathbf{M} are all strictly positive. Properties of the network G for which Eq. (1) admits a stable solution have been discussed at length, and it is known that sparse modular networks confer stability, for example^{49,50}. However, our concern here is not with stability *per se* but rather with the effect that external noise has on the magnitude of fluctuations around a stable equilibrium. To investigate this we will consider the following stochastic differential equation associated with Eq. (1)

$$\frac{d\mathbf{x}}{dt} = -\mathbf{M}\mathbf{x} + \boldsymbol{\Sigma} \frac{d\mathbf{W}}{dt}, \quad (2)$$

where $\mathbf{W}(t)$ is a standard n -dimensional Brownian motion (and $d\mathbf{W}/dt$ is therefore n -dimensional Gaussian white noise). Eq. (2) describes a multivariate Ornstein-Uhlenbeck process. Whenever Eq. (1) admits a stable solution, Eq. (2) is ergodic and therefore admits a unique invariant measure⁵¹. Furthermore, by the linearity of Eq. (2) we know that $\mathbf{x}(t)$ is distributed

according to a multivariate normal with mean $e^{-\mathbf{M}t} \mathbf{x}(0)$ and covariance matrix $\mathbf{K}(t)$ given by,

$$\mathbf{K}(t) = \langle \mathbf{x}(t) - \langle \mathbf{x}(t) \rangle, (\mathbf{x}(t) - \langle \mathbf{x}(t) \rangle)^T \rangle = \int_0^t e^{-\mathbf{M}(t-s)} \boldsymbol{\Sigma} \boldsymbol{\Sigma}^T e^{-\mathbf{M}(t-s)^T} ds. \quad (3)$$

If Eq. (2) describes an ergodic process then

$$\lim_{t \rightarrow \infty} \mathbf{K}(t) = \mathbf{K}, \quad (4)$$

where \mathbf{K} satisfies the Liapunov equation⁵¹

$$\mathbf{M}\mathbf{K} + \mathbf{K}\mathbf{M}^T = \boldsymbol{\Sigma}\boldsymbol{\Sigma}^T. \quad (5)$$

Although this is the standard formulation⁵¹, instead of working with Eq. (5) we will work directly with the Eq. (3) as it ultimately allows a more transparent assessment the effects of network structure on the stationary covariance of the system.

Since our purpose is to determine the way in which input noise is processed by the network G it is natural to consider a single noisy input to the system, which represents the fluctuating extra-cellular environment, and a single output, representing the computational core of the network. To do so we may, without loss of generality, chose a labeling of the nodes such that the first node is the noisy input and the n -th node is the output. Thus, we set $\boldsymbol{\Sigma} = (\sigma, 0, \dots, 0)^T$ and we are interested in calculating the variance of the n -th node in the network, which is given by K_{nn} , relative to the magnitude of the input noise. If the input fluctuations carry no information, then this is a measure of the extent to which the network suppresses or amplifies random environmental fluctuations. If the fluctuations contain important environmental information, then this is a measure of the extent to which the target node can ‘sense’ this extra-cellular information. From Eq. (3) the limiting covariance, in index notation and using Einstein summation notation, is

$$K_{ij} = \int_0^\infty \left(e^{-\mathbf{M}(t-s)} \right)_{il} (\boldsymbol{\Sigma}\boldsymbol{\Sigma}^T)_{lm} \left(e^{-\mathbf{M}^T(t-s)} \right)_{mj} ds. \quad (6)$$

Now, since $(\boldsymbol{\Sigma}\boldsymbol{\Sigma}^T)_{lm} = \sigma^2$ when $l = m = 1$ and zero otherwise, we have

$$K_{ij} = \sigma^2 \int_0^\infty e^{-2(t-s)} \left(e^{\mathbf{A}(t-s)} \right)_{il} \left(e^{\mathbf{A}^T(t-s)} \right)_{1j} ds. \quad (7)$$

Although it is not immediately transparent, this expression connects strongly to the structure of the network via the fact that (i, j) -th entry of the k -th power of the adjacency matrix is the total weight of all walks from node i to node j with length k (the weight $w(P)$ of a walk P from node i to node j is the product of its edge weights, $\prod_{(i,j) \in P} A_{ij}$). Thus, the exponential of the adjacency matrix of a network is a weighted sum of all walks between nodes i and j , and so is a simple measure of network ‘communicability’⁵². Using the notation $\beta_{ijk} = A_{ij}^k = \sum_{P_k} w(P_k)$, we may re-write the exponential terms in Eq. (7) as

$$\left(e^{\mathbf{A}(t-s)} \right)_{ij} = \sum_{k=0}^\infty \frac{\beta_{ijk}}{k!} (t-s)^k. \quad (8)$$

Making use of this connection and using the shorthand $\beta_{ik} = \beta_{i1k}$ we then obtain,

$$K_{ij} = \sigma^2 \sum_{k=0}^\infty \sum_{l=0}^\infty \int_0^\infty \frac{\beta_{ik}\beta_{jl}}{k!l!} (t-s)^{k+l} e^{-2(t-s)} ds, \quad (9)$$

where $\beta_{ij0} = I_{ij}$ (the $n \times n$ identity matrix). Since

$$\lim_{t \rightarrow \infty} \int_0^t (t-s)^n e^{-2(t-s)} ds = \frac{n!}{2^{n+1}}, \quad (10)$$

we may furthermore simplify Eq. (9) to

$$K_{ij} = \frac{\sigma^2}{2} \sum_{k=0}^\infty \sum_{l=0}^\infty \frac{1}{2^{k+l}} \binom{k+l}{l} \beta_{ik}\beta_{jl}. \quad (11)$$

Since the variance of the noisy input is given by $K_{11} = \sigma^2/2$, we may investigate the noise-processing ability of the network by considering the ratio

$$R = \frac{K_{nn}}{K_{11}} = \sum_{k=0}^\infty \sum_{l=0}^\infty \frac{1}{2^{k+l}} \binom{k+l}{l} \beta_k \beta_l, \quad (12)$$

where we have further simplified notation by setting $\beta_{nk} = \beta_k$. When $R > 1$ noise is amplified by the network; when $R < 1$ noise is suppressed by the network. Importantly, if the process described by Eq. (2) is ergodic then R is finite and depends only on the structure of the network G . This formula therefore provides an explicit connection between network architecture and noise-processing; our interest is to determine how R is affected by different network architectures. To do so we note that Eq. (12) has a natural interpretation in terms of random walks on G , as follows.

Since each walk P from the input to the target has an associated weight $w(P)$, a pair of (possibly intersecting) walks P, Q from the input to the target also has an associated weight $w(P, Q) = w(P)w(Q)$, the product of the edge weights involved. If we write P_k and P_l for arbitrary walks from the input to the target of length k , and l respectively, then the product $\beta_k \beta_l$ can be written as

$$\beta_k \beta_l = \sum_{P_k} w(P_k) \sum_{P_l} w(P_l) = \sum_{P_k, P_l} w(P_k, P_l). \quad (13)$$

Substituting this into Eq. (12) and rewriting the second sum in terms of $m = k + l$ gives,

$$R = \sum_{m=0}^{\infty} \sum_{k=0}^m \sum_{P_k, P_{m-k}} \frac{1}{2^m} \binom{m}{k} w(P_k, P_{m-k}), \quad (14)$$

from which it can be seen that R is a weighted sum of all pairs of walks through G from the source node to the target, with the relative importance of each walk-pair determined by a coefficient drawn from a binomial distribution $B(m, 1/2)$, where m is the length of the walk-pair. The appearance of binomial probabilities arises as the natural probability measure for pairs of random walkers on the network G . To see this consider two independent random walkers starting at the same time at the input node. At each time step, one walker is chosen with probability $1/2$, and that walker moves through G choosing available edges with equal probability (*i.e.* if the walker is at node i then each outgoing edge from node i is chosen with probability $1/d_i^{\text{out}}$ where d_i^{out} is the out-degree of node i). The probability that after precisely $m = k + l$ steps both walkers are at the target node is $\binom{m}{k} 2^{-m}$. Thus, the inner sum in Eq. (14) is the expected weight of a pair of walks from input to target, with respect to the probability measure generated by two independent random walkers [the presence of two random walkers rather than one, as might be expected, arises from the fact that \mathbf{K} depends upon two exponential terms, $e^{A(t-s)}$ and $e^{A^T(t-s)}$, in Eq. (7)]. If G is a directed acyclic graph (in which there are no feedback loops), then all random walks have finite length and the first sum in Eq. (14) has finitely many terms. However, if cycles are present in the network then random walks may be arbitrarily long and the first sum will correspondingly have infinitely many terms. Positive feedback loops add infinitely many positive terms to the sum, and therefore always serve to amplify noise with respect to similarly structured acyclic networks; negative feedback loops add both positive and negative terms to the sum and may amplify or reduce noise with respect to similarly structured acyclic networks, depending on the particular arrangement of inhibitory edges in the network. This observation is consistent with previous studies on the effect of positive and negative feedback loops in noise propagation, particularly in biological regulatory networks^{25,26,53,54}. As the random walks are independent of the edge weights, the importance of each edge to the noise-processing ability of the network has 2 contributions: 1) the probability that either of the random walkers traverses that edge (which depends solely on its position within the network relative to the source and target); and 2) the extent to which it contributes to any walk it participates in (which depends solely on its weight).

It is worth noting here that although random walkers are a natural way to explore directed networks^{55,56}, if the matrix \mathbf{A} is normal (that is, if it commutes with its transpose \mathbf{A}^T ; a strong condition that is not typically satisfied by directed networks but does occur if the network G is undirected, for example), then a much simpler related result for the trace of the covariance matrix $\text{Tr}(\mathbf{K})$ may be obtained. In the rest of this section we will study the particular case that \mathbf{A} is normal. Although not directly relevant to signaling networks, this case does nevertheless provide insight into Eq. (12) by relating noise-processing to well-known network-theoretic notions.

First, let us take the sum of the ratios K_{jj}/K_{11} in Eq. (12) for all output nodes j ,

$$\bar{R} = \sum_{j=1}^n \frac{K_{jj}}{K_{11}}, \quad (15)$$

$$= \sum_{j=1}^n \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \frac{1}{2^{k+l}} \binom{k+l}{l} \beta_{jk} \beta_{jl}, \quad (16)$$

$$= \sum_{m=0}^{\infty} \sum_{k=0}^m \sum_{j=1}^n \frac{A_{1j}^k A_{1j}^{m-k}}{2^k 2^{m-k}} \binom{m}{k}. \quad (17)$$

In general this sum cannot be simplified and we resort to interpreting in terms of random walkers, as above. However, if the




Figure 1. (Left) The regulatory network for pluripotency derived by Dunn and co-workers in³⁴. (Right) The reduced network that we study here in which Lif is taken as a noisy input and Oct4 is taken as the target. Since CH and PD cannot be reached from Lif via a walk on this network, we can exclude these nodes, along with Tcf3 and ERK, from our analysis. Edges that participate in feedback loops in the network are shown in red.

matrices \mathbf{A} and \mathbf{A}^T commute then we can use the binomial formula to expand the m -th power of the symmetric part of \mathbf{A} as,

$$\mathbf{A}_s^m = \left(\frac{\mathbf{A} + \mathbf{A}^T}{2} \right)^m = \sum_{k=0}^m \mathbf{A}^k (\mathbf{A}^T)^{m-k} \frac{1}{2^k} \frac{1}{2^{m-k}} \binom{m}{k}. \quad (18)$$

The $(1, 1)$ -entry of the matrix above is precisely the double inner sum in Eq. (17), so we may write

$$\bar{R} = \left[\sum_{m=0}^{\infty} \left(\frac{\mathbf{A} + \mathbf{A}^T}{2} \right)^m \right]_{11}. \quad (19)$$

If $\|\mathbf{A}_s\| < 1$ [a condition which implies stability in Eq. (1)], then the geometric series in Eq. (19) converges to

$$\sum_{m=0}^{\infty} \left(\frac{\mathbf{A} + \mathbf{A}^T}{2} \right)^m = (\mathbf{I} - \mathbf{A}_s)^{-1} = \mathbf{M}_s^{-1}, \quad (20)$$

where $\mathbf{M}_s = (\mathbf{M} + \mathbf{M}^T)/2$ is the symmetric part of \mathbf{M} . By taking into account all possible walks through the network, this is a simple variation on the exponential of the adjacency matrix as a measure of network communicability, although in this case the communicability of G is taken with respect to the analytic function $(1-x)^{-1}$, rather than $\exp(x)$ ⁵⁷. Using this result we finally obtain,

$$\bar{R} = [\mathbf{M}_s^{-1}]_{11}. \quad (21)$$

Thus, in order to determine the noise-processing ability of the network using this measure, we need only calculate the matrix inverse of the symmetric part of \mathbf{M} , and take the $(1, 1)$ -entry (where without loss of generality the first node is the input node). This result is strongly related to the Laplacian matrix of the network G : if \mathbf{A} is symmetric and normalized so that each row (or column) sums to 1 then \mathbf{M}_s is precisely the normalized Laplacian of G , which is well-known to be closely related to network connectivity^{58,59}.

To illustrate how Eq. (12) works in practice, we now consider a couple of examples.

Signaling cascade

The simplest example network is a signaling cascade, consisting of a chain of $m = n - 1$ interactions between n nodes. To facilitate a transparent illustration we shall assume that all the weights in the network are the same. In this case, we may index the nodes such that $A_{ij} = a$ for edges $(i, i + 1)$, where $i = 1, 2, \dots, n - 1$, and is zero otherwise. Eq. (12) then gives

$$R = \frac{(2m)!}{m!^2} \left(\frac{a}{2} \right)^{2m}. \quad (22)$$

This result coincides with the expression derived in⁶⁰. For large m we may use Stirling's approximation to obtain

$$R \sim \frac{a^{2m}}{\sqrt{\pi m}}, \quad (23)$$

from which it can be seen that the signaling pathway suppresses noise if

$$|a| < (\pi m)^{1/4m}. \quad (24)$$

Three conclusions are apparent from this result: (1) since the variance of the target node depends upon the magnitude of the interaction strength squared, the sign of the interactions (*i.e.* whether they are activating or inhibiting) does not affect the ability of the cascade to process noise; (2) since R is monotonic decreasing with m , the effect of input noise diminishes with longer cascades (for fixed a); (3) if $|a| < 1$ then $R < 1$ for any m and noise is diminished by the cascade; while if $|a| > 1$ noise may be amplified by the cascade depending on the magnitude of a relative to m . In general since R decreases with a and m , this analysis suggests that long signaling cascades with weak interactions process noise better than short cascades with stronger interactions. This result is in accordance with previous studies on noise-processing by transcriptional cascades⁶¹.

Feedforward loop

In reality signaling cascades do not operate in isolation; rather cross-talk between pathways means that many different paths from the input to the output may exist, and each may process different aspects of the extra-cellular signal. The simplest example of such a network is the feedforward loop motif, a commonly occurring structure in biological regulatory networks which is known to be involved with in a range of biological functions, including distinguishing persistent signals from noise⁶². The simplest feedforward loop consists of three nodes with two paths from the source (node 1) to the target (node 3): one direct and one indirect, via an intermediary (node 2). In this case, assuming that all interactions are of equal weight we obtain

$$R = \beta_1^2 \frac{1}{2} + \beta_1 \beta_2 \frac{3}{2^2} + \beta_2^2 \frac{3}{2^3}. \quad (25)$$

By contrast to the simple signaling cascade, the sign of the edges in the feedforward loop do affect its noise-processing ability. If all edges are positive then $\beta_1, \beta_2 > 0$ (all paths in the network are positive) and the target receives a consistent signal from the source. In this case the feedforward loop is said to be coherent⁶². However, if $\beta_1 < 0$ or $\beta_2 < 0$ (which occurs if either one or three of the edges is negative) then the target receives a inconsistent signal from the source. In this case, the feedforward loop is said to be incoherent⁶². Denoting the noise-processing ratios in the coherent and incoherent cases by R^+ and R^- respectively, it follows from Eq. (25) that $R^+ > R^-$, and therefore that the incoherent feedforward loop is better at processing noise. This general conclusion also holds when the paths from the source to the target are of arbitrary length: incoherence of pairs of paths through the network tends to lead to better noise-processing. These results are consistent with previous studies on the effect of positive and negative paths in feedforward loops on noise propagation^{25, 63–66}

Noise-processing by stem cells

In order to apply these general results we now consider signal transduction in the regulatory network for pluripotency. The skeleton of this network has recently been inferred from analysis of correlations between expression patterns of important regulatory factors³⁴ and is illustrated in Fig. 1. The behavior of this network is determined by input from three extra-cellular factors commonly added to ES cell culture media preparations: the cytokine leukemia inhibitory factor (Lif), and selective inhibitors of glycogen synthase kinase 3 (TGF- β) (Chiron99021, denoted CH) and mitogen-activated protein kinase kinase (Mek) (PD0325901, denoted PD). Since it is known that stimulation of Lif signaling is sufficient to maintain pluripotency *in vitro*, we will choose Lif as the noisy source in our analysis. When Lif is present in the extra-cellular environment it binds to the Stat3 receptor^{67,68}, and activates signaling pathways that stimulate the core transcriptional regulatory network for pluripotency in mouse ES cells^{35,69}. At the heart of this core network are the trio of transcription factors, Oct4, Sox2 and Nanog^{29,70,71}. Since Oct4 is well-established as the most central factor in this core, we will consider the propagation of a noisy signal from (the input) Lif through the network to (the output) Oct4. Although the sign of the interactions (*i.e.* whether they are activatory or inhibitory) is known³⁴ their strength is unknown. In the absence of this information we assume that all interactions are of equal unit strength since this represents the most economic model. To investigate how network structure affects noise-processing, we sought to determine how the ratio R changes upon targeted removal of different interactions from this network by comparing the R -values of perturbed networks with that of the unperturbed network (denoted R_{full}). To uncover some of the structural determinants of noise-processing we also calculated two simple network measures based upon our interpretation of Eq. (12): (1) $c = p^+ / (p^+ + p^-)$, where p^+ and p^- are the number of positive and negative paths from Lif to Oct4 respectively. This is a simple measure of structural coherence; and (2) f , the total number of feedback loops in the network, as a measure of network complexity. To determine how network properties varied with removal of specific edges, we also determined how these measures changed upon targeted removal of different interactions from the network by comparing values with those of the unperturbed network (denoted c_{full} and f_{full} respectively). The results of this analysis are summarized in Table 1.

The shortest paths from Lif to Oct4 in this network have length 4; there are two such paths: (1) Lif \rightarrow Stat3 \rightarrow Tfcp2l1 \rightarrow Esrrb \rightarrow Oct4; and (2) Lif \rightarrow Stat3 \rightarrow Klf4 \rightarrow Klf2 \rightarrow Oct4. The first of these paths is negative (due to the inhibitory interaction

Interaction removed	Noise-processing (R/R_{full})	Coherence (c/c_{full})	Feedback (f/f_{full})
Tfcp211 \rightarrow Sall4	0.02	0.68	0.64
Sox2 \rightarrow Oct4	0.12	0.60	0.37
Sall4 \rightarrow Klf2	0.18	0.98	0.45
Sall4 \rightarrow Sox2	0.18	0.86	0.18
Stat3 \rightarrow Gbx2	0.32	0.98	0
Gbx2 \rightarrow Klf4	0.32	0.98	0
Nanog \rightarrow Sox2	0.83	0.94	0.18
Klf2 \rightarrow Nanog	1	1.10	0.27
Esrrb \rightarrow Tfcp211	1.12	0.95	0.36
Nanog \rightarrow Esrrb	1.19	1.25	0.45
Klf4 \rightarrow Tfcp211	1.99	1.06	0
Stat3 \rightarrow Klf4	4.70	0.98	0
Klf4 \rightarrow Klf2	8.68	0.92	0
Klf2 \rightarrow Oct4	12.10	0.86	0.18
Stat3 \rightarrow Tfcp211	12.18	1.03	0
Tfcp211 \rightarrow Esrrb	24.62	1.23	0.18
Esrrb \rightarrow Oct4	25.68	1.53	0.27

Table 1. The effect that targeted removal of interactions on network noise-processing. The first column identifies the edge removed from the network; the second column shows the effect of targeted removal of the given edge on the ratio R by comparison with that of the unperturbed network; the third column shows the effect of targeted removal of the given edge has on network coherence; the fourth column shows the effect of targeted removal of the given edge has on network feedback. Edges that emanate from Oct4 do not contribute to the noise processing capacity of the network and their removal does not affect R so they are excluded from this table. Since all paths from Lif to Oct4 pass through the edge Lif \rightarrow Stat3 its removal disconnects the network; this edge is also accordingly excluded from the table. Interactions are ordered by column 1.

Esrrb \rightarrow Oct4) while the second is positive. Thus, when taken together this pair of paths forms an incoherent feedforward loop; since these are the shortest paths in the network, we anticipate from Eq. (12) that this incoherent feedforward loop will have an important role in noise-processing in this network. Indeed, this is what is observed: if any of the elements of this structure are removed, then the noise-processing capacity of this network is severely inhibited and the ratio R increases substantially (see Table 1). By contrast, we also anticipate that since feedback loops introduce arbitrarily long walks in the network [and therefore contribute infinitely many terms to the sum in Eq. (12)] removal of edges which participate strongly in the feedback structure of the network will result in a substantial reduction in its noise-processing capacity. Again, this is what is observed: when edges which participate in large numbers of feedback loops are removed, the ratio R decreases substantially (see Table 1). Although we have focused on the effect of edge removal on two particular properties (coherence and feedback) each edge in the network is likely to have multiple functional roles, and an individual edge may be important for both coherence and feedback (for example, since Esrrb has an important role both in mediating environmental signals to the computational core and maintaining the feedback structure of the core, edges such as Tfcp211 \rightarrow Esrrb and Esrrb \rightarrow Oct4 naturally have a dual role). For this reason, while we observe general trends between noise-processing and coherence/feedback, we do not, as expected, see strong correlations (see Fig. 1). However, when taken together, these results indicate that those interactions in the feedback rich core transcriptional circuitry (shown in red in Fig.2) that are needed to maintain a self-perpetuating pluripotent identity^{5,48} have a tendency to amplify extrinsic noise. To compensate, a distinct set of interactions between auxiliary factors structured into a set of incoherent feedforward loops tend to suppress environmental noise, and thereby ensure that environmental signals are robustly mediated to this core circuit.

Discussion

In this paper we have investigated the noise-processing by networks. To do so we have combined a stochastic approach (similar to that taken by Anderson and co-workers in Ref.⁶⁰) with graph-theoretic notions to derive a simple expression that relates a network's structure to its noise-processing ability. This expression is easily calculated even for large networks [particularly if the network is undirected, see Eq. (21)], so provides an economic measure that may be used to examine the structure of naturally occurring networks, and guide the design of man-made networks. In practical applications it is often desirable to maintain




Figure 2. Plots of the data from Table 1. Removal of edges that result in an increase of coherence in the network tend to diminish the system’s noise-processing ability, while removal of edges which reduce the overall feedback structure of the network tend to improve the system’s noise-processing ability. Red lines show linear regression.

signal intensity while noise is simultaneously controlled. In such cases some noise-minimizing strategies, such as introducing long or incoherent paths through the network, may not be desirable or feasible. It would be interesting to investigate further how the interplay between noise-processing and signal maintenance shapes the structure of biological regulatory networks, for example using recent methods from the theory of adaptive networks⁷². Similarly, it would also be interesting to examine how network structures affect noise-processing more generally, taking into account more complex network dynamics such as limit cycles. To illustrate our results we have considered the structure of the network that maintains pluripotency in mouse ES cells, and found that important network structures, distinct from those that maintain the core pluripotent state, are responsible for noise processing in this system, suggesting that different features of this network are responsible for different regulatory tasks. Accordingly, we anticipate that the structure of many natural networks may be determined, in part, to optimally process noise. It will be interesting to elucidate the extent to which cross-talk between pathways in natural networks, which typically have to process multiple complex signals, is shaped by the trade-off between signal integration and noise-processing.

References

1. Barabási, A. L. & Oltvai, Z. N. Network biology: Understanding the cell’s functional organization. *Nat. Rev. Genet.* **5**, 101–113 (2004).
2. Tkačik, G. & Bialek, W. Cell biology: Networks, regulation, pathways. In *Encyclopedia of complexity and systems science*, 719–741 (Berlin: Springer, 2009).
3. Newman, M. E. J. *Networks. An introduction* (OUP, 2010).
4. Estrada, E. *The structure of complex networks* (OUP, 2011).
5. Macarthur, B. D., Ma’ayan, A. & Lemischka, I. R. Systems biology of stem cell fate and cellular reprogramming. *Nat. Rev. Mol. Cell Biol.* **10**, 672–681 (2009).
6. Martello, G. & Smith, A. The nature of embryonic stem cells. *Annu. Rev. Cell Dev. Biol.* **30**, 647–675 (2014).
7. Jeong, H., Mason, S. P., Barabási, A. L. & Oltvai, Z. N. Lethality and centrality in protein networks. *Nature* **411**, 41–42 (2001).
8. Blais, A. & Dynlacht, B. D. Constructing transcriptional regulatory networks. *Genes Dev.* **19**, 1499–1511 (2005).
9. Alberts, B. *et al. Molecular biology of the cell* (Garland science, Taylor and Francis Group, LLC, 2002).
10. Adler, E. M., Gough, N. R. & Ray, L. B. 2015: Signaling breakthroughs of the year. *Sci. Signal.* **9**, eg1 (2016).
11. Irish, J. M. *et al.* Single cell profiling of potentiated phospho-protein networks in cancer cells. *Cell* **118**, 217–228 (2015).
12. Verkaar, F., Cadigan, K. M. & Amerongen, V. R. Celebrating 30 years of wnt signaling. *Sci. Signal.* **5**, mr2 (2012).
13. Levine, E. & Hwa, T. Stochastic fluctuations in metabolic pathways. *Proc. Natl. Acad. Sci. USA.* **104**, 9224–9229 (2007).
14. Schreiber, S. L. & Bernstein, B. E. Signaling network model of chromatin. *Cell* **111**, 771–778 (2015).

15. Song, J. *et al.* A protein interaction between β -catenin and dnmt1 regulates wnt signaling and dna methylation in colorectal cancer cells. *Mol. Cancer Res.* **13**, 969–981 (2015).
16. Valenta, T., Hausmann, G. & Basler, K. The many faces and functions of β -catenin. *EMBO J.* **31**, 2714–36 (2012).
17. Clevers, H., Loh, K. M. & Nusse, R. An integral program for tissue renewal and regeneration: Wnt signaling and stem cell control. *Science* **346**, 1248012 (2014).
18. Niwa, H. Wnt: What's needed to maintain pluripotency? *Nat. Cell Biol.* **13**, 1024–1026 (2011).
19. Kim, H. *et al.* Modulation of β -catenin function maintains mouse epiblast stem cell and human embryonic stem cell self-renewal. *Nat. Commun.* **4**, 2403 (2013).
20. Attisano, L. & Wrana, J. L. Signal integration in $\text{tgf-}\beta$, wnt, and hippo pathways. *FI000Prime Rep.* **5** (2013).
21. Niida, A. *et al.* Dkk1, a negative regulator of wnt signaling, is a target of the beta-catenin/tcf pathway. *Oncogene* **23**, 8520–8526 (2004).
22. Ladbury, J. E. & Arold, S. T. Noise in cellular signaling pathways: causes and effects. *Trends Biochem. Sci.* **37**, 173–178 (2012).
23. Pedraza, J. M. & van Oudenaarden, A. Noise propagation in gene networks. *Science* **307**, 1965–1969 (2005).
24. Paulsson, J. Summing up the noise in gene networks. *Nature* **427**, 415–418 (2004).
25. Hornung, G. & Barkai, N. Noise propagation and signaling sensitivity in biological networks: A role for positive feedback. *PLoS Comput. Biol.* **4**, e8 (2008).
26. Hooshangi, S. & Weiss, R. The effect of negative feedback on noise propagation in transcriptional gene networks. *Chaos* **16**, 026108 (2006).
27. Rué, P., Domedel-Puig, N., Garcia-Ojalvo, J. & Pons, A. Integration of cellular signals in chattering environments. *Prog. Biophys. Mol. Biol.* **110**, 106–112 (2012).
28. Takahashi, K. & Yamanaka, S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* **126**, 663–676 (2006).
29. Boyer, L. A. *et al.* Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* **122**, 947–956 (2005).
30. Yu, J. *et al.* Induced pluripotent stem cell lines derived from human somatic cells. *Science* **318**, 1917–1920 (2007).
31. Dahéron, L. *et al.* LIF/STAT3 signaling fails to maintain self-renewal of human embryonic stem cells. *Stem Cells* **22**, 770–778 (2004).
32. Ye, S., Li, P., Tong, C. & Ying, Q. Embryonic stem cell self-renewal pathways converge on the transcription factor Tfcp2l1. *EMBO J.* **32**, 2548–60 (2013).
33. Kim, J., Chu, J., Shen, X., Wang, J. & Orkin, S. H. An extended transcriptional network for pluripotency of embryonic stem cells. *Cell* **132**, 1049–1061 (2008).
34. Dunn, S. J., Martello, G., Yordanov, B., Emmott, S. & Smith, A. G. Defining an essential transcription factor program for naïve pluripotency. *Science* **344**, 1156–60 (2014).
35. Huang, G., Ye, S., Zhou, X., Liu, D. & Ying, Q. L. Molecular basis of embryonic stem cell self-renewal: From signaling pathways to pluripotency network. *Cell. Mol. Life Sci.* **72**, 1741–1757 (2015).
36. Takahashi, K. *et al.* Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* **131**, 861–872 (2007).
37. Ying, Q.-L. *et al.* The ground state of embryonic stem cell self-renewal. *Nature* **453**, 519–523 (2008).
38. Stumpf, P. S., Ewing, R. & MacArthur, B. D. Single-cell pluripotency regulatory networks. *Proteomics* **16**, 2303–2312 (2016).
39. Williams, R. L., Hilton, D. J. & Nicolai, N. A. Myeloid leukaemia inhibitory factor maintains the developmental potential of embryonic stem cells. *Nature* **336**, 15 (1988).
40. Smith, A. G. *et al.* Inhibition of pluripotential embryonic stem cell differentiation by purified polypeptides. *Nature* **336**, 688–690 (1988).
41. Ying, Q.-L., Nichols, J., Chambers, I. & Smith, A. Bmp induction of id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with stat3. *Cell* **115**, 281–292 (2003).

42. Burdon, T., Stracey, C., Chambers, I., Nichols, J. & Smith, A. Suppression of shp-2 and erk signalling promotes self-renewal of mouse embryonic stem cells. *Dev. Biol.* **210**, 30–43 (1999).
43. Stavridis, M. P., Lunn, J. S., Collins, B. J. & Storey, K. G. A discrete period of fgf-induced erk1/2 signalling is required for vertebrate neural specification. *Development* **134**, 2889–2894 (2007).
44. Kunath, T. *et al.* Fgf stimulation of the erk1/2 signalling cascade triggers transition of pluripotent embryonic stem cells from self-renewal to lineage commitment. *Development* **134**, 2895–2902 (2007).
45. Sato, N., Meijer, L., Skaltsounis, L., Greengard, P. & Brivanlou, A. H. Maintenance of pluripotency in human and mouse embryonic stem cells through activation of wnt signaling by a pharmacological gsk-3-specific inhibitor. *Nat. Med.* **10**, 55–63 (2004).
46. Amit, M. *et al.* Clonally derived human embryonic stem cell lines maintain pluripotency and proliferative potential for prolonged periods of culture. *Dev. Biol.* **227**, 271–278 (2000).
47. Xu, R.-H. *et al.* Basic FGF and suppression of BMP signaling sustain undifferentiated proliferation of human es cells. *Nat. Methods* **2**, 185–190 (2005).
48. MacArthur, B. D. *et al.* Nanog-dependent feedback loops regulate murine embryonic stem cell heterogeneity. *Nat. Cell Biol.* **14**, 1139–1147 (2012).
49. May, R. M. Will a large complex system be stable? *Nature* **238**, 413–414 (1972).
50. Allesina, S. & Tang, S. Stability criteria for complex ecosystems. *Nature* **483**, 205–208 (2012).
51. Gardiner, C. W. *Stochastic methods: a handbook for the natural and social sciences* (Berlin: Springer, 2009).
52. Estrada, E. & Higham, D. J. Network properties revealed through matrix functions. *SIAM Rev.* **52**, 696–714 (2010).
53. Pedraza, J. M. Noise propagation in gene networks. *Science* **307**, 1965–1969 (2005).
54. Zhang, H., Chen, Y. & Chen, Y. Noise propagation in gene regulation networks involving interlinked positive and negative feedback loops. *PLOS ONE* **7**, 1–8 (2012).
55. Rosvall, M. & Bergstrom, C. T. Maps of random walks on complex networks reveal community structure. *Proc. Natl. Acad. Sci. USA.* **105**, 1118–1123 (2008).
56. Chung, F. Laplacians and the Cheeger inequality for directed graphs. *Ann. Comb.* **9**, 1–19 (2005).
57. Estrada, E. & Higham, D. J. Network properties revealed through matrix functions. *SIAM Rev.* **52**, 696–714 (2010).
58. Mohar, B. The Laplacian spectrum of graphs. In *Graph theory, combinatorics, and applications.*, Wiley-Intersci. Publ., 871–898 (Wiley: New York, 1991).
59. Luxburg, U. V. A tutorial on spectral clustering. *Stat. Comput.* **17**, 395–416 (2006).
60. Anderson, D. F., Mattingly, J. C., Nijhout, H. F. & Reed, M. C. Propagation of fluctuations in biochemical systems, i: Linear ssc networks. *Bull. Math. Biol.* **69**, 1791–1813 (2007).
61. Pilkiewicz, K. R. & Mayo, M. L. Fluctuation sensitivity of a transcriptional signaling cascade. *Phys. Rev. E* **94**, 032412 (2016).
62. Alon, U. *An introduction to systems biology: design principles of biological circuits* (Chapman Hall, 2007).
63. Shahrezaei, V., Ollivier, J. F. & Swain, P. S. Colored extrinsic fluctuations and stochastic gene expression. *Mol. Syst. Biol.* **1**–9 (2008).
64. Osella, M., Bosia, C., CorÃ¡j, D. & Caselle, M. The role of incoherent microRNA-mediated feedforward loops in noise buffering. *PLOS Comput. Biol.* **7** (2011).
65. Wang, P., LÃĀe, J. & Ogorzalek, M. J. Global relative parameter sensitivities of the feed-forward loops in genetic networks. *Neurocomputing* **78**, 155–165 (2012). Selected papers from the 8th International Symposium on Neural Networks (ISNN 2011).
66. Ghosh, B., Karmakar, R. & Bose, I. Noise characteristics of feed forward loops. *Physical biology* **2**, 36 (2005).
67. Matsuda, T. *et al.* STAT3 activation is sufficient to maintain an undifferentiated state of mouse embryonic stem cells. *EMBO J.* **18**, 4261–4269 (1999).
68. Niwa, H., Burdon, T., Chambers, I. & Smith, A. Self-renewal of pluripotent embryonic stem cells is mediated via activation of STAT3. *Genes Dev.* **12**, 2048–2060 (1998).
69. Yu, J. & Thomson, J. Pluripotent stem cell lines. *Genes Dev.* **22**, 1987–1997 (2008).

70. Niwa, H., Miyazaki, J. & Smith, A. G. Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat. Genet.* **24**, 372–376 (2000).
71. Nichols, J. *et al.* Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. *Cell* **95**, 379–391 (1998).
72. Gross, T. & Blasius, B. Adaptive coevolutionary networks: a review. *Journal of the Royal Society Interface* **5**, 259–271 (2008).

Additional Information

The authors declare no competing financial interests.

Acknowledgments

This work was funded by BBSRC Grant No. BB/L000512/1 and by PhD funding from the Institute for Life Sciences, University of Southampton. The authors would like to thank Enrico Mossotto, Rosanna Smith and Patrick Stumpf for useful feedback on the manuscript.

Author contributions statement

SK, RSG, KCZ and BDM conducted the mathematical modeling and network analysis. SK, RSG, RE, KCZ and BDM interpreted the results. All authors wrote the paper.