



Vegetable oil as fat replacer inhibits formation of heterocyclic amines and 2 polycyclic aromatic hydrocarbons in reduced fat pork patties

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Lu, F., Kuhnle, G. K. and Cheng, Q. (2017) Vegetable oil as fat replacer inhibits formation of heterocyclic amines and 2 polycyclic aromatic hydrocarbons in reduced fat pork patties. *Food Control*, 81. pp. 113-125. ISSN 0956-7135 doi: <https://doi.org/10.1016/j.foodcont.2017.05.043> Available at <http://centaur.reading.ac.uk/70536/>

It is advisable to refer to the publisher's version if you intend to cite from the work.

Published version at: http://ac.els-cdn.com/S0956713517302852/1-s2.0-S0956713517302852-main.pdf?_tid=f5d9a0cc-42f6-11e7-aa61-00000aab0f6c&acdnat=1495901591_36523bfad4dcbb6a29079af52fcc6489

To link to this article DOI: <http://dx.doi.org/10.1016/j.foodcont.2017.05.043>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Accepted Manuscript

Vegetable oil as fat replacer inhibits formation of heterocyclic amines and polycyclic aromatic hydrocarbons in reduced fat pork patties

Fei Lu, Gunter K. Kuhnle, Qiaofen Cheng



PII: S0956-7135(17)30285-2
DOI: 10.1016/j.foodcont.2017.05.043
Reference: JFCO 5649
To appear in: *Food Control*
Received Date: 05 March 2017
Revised Date: 25 May 2017
Accepted Date: 26 May 2017

Please cite this article as: Fei Lu, Gunter K. Kuhnle, Qiaofen Cheng, Vegetable oil as fat replacer inhibits formation of heterocyclic amines and polycyclic aromatic hydrocarbons in reduced fat pork patties, *Food Control* (2017), doi: 10.1016/j.foodcont.2017.05.043

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Vegetable oil as fat replacer inhibits formation of heterocyclic amines and**
2 **polycyclic aromatic hydrocarbons in reduced fat pork patties**

3 Fei Lu, Gunter K. Kuhnle and Qiaofen Cheng*

4 Department of Food and Nutritional Sciences, University of Reading, Whiteknights, PO Box
5 226, Reading RG6 6AP, UK

6 *Corresponding author – Email: q.cheng@reading.ac.uk

7 **Abstract**

8 Formation of heterocyclic amines (HCAs) and polycyclic aromatic hydrocarbons
9 (PAHs) was examined to evaluate the impact of using vegetable oil as fat
10 replacement on carcinogen formation in meat product. Pork patties were formulated
11 with 40% fat replacement by olive oil, sunflower oil or grape seed oil, respectively
12 and cooked at 180°C or 220°C. Control patties contained the highest amount of
13 HCAs compared with all other patties at both temperatures. Olive oil and sunflower
14 oil replacement completely inhibited formation of MeIQ (2-amino-3, 4-
15 methylimidazo[4,5-f]quinoline), while grape seed oil completely inhibited MeIQx (2-
16 amino-3,8-dimethylimidazo[4,5-f]quinoxaline), 4,8-DiMeIQx (2-amino-3,4,8-
17 trimethylimidazo[4,5-f]quinoxaline) and PhIP (2-amino-1-methyl-6-phenylimidazo[4,5-
18 b]pyridine) in patties. Grape seed oil achieved the highest inhibition capacity
19 compared with sunflower oil and olive oil. HCAs increased significantly with cooking
20 temperature ($p < 0.05$), but no difference was observed in total PAHs for patties
21 cooked at different temperature ($p > 0.05$). In conclusion, fat replacement with
22 sunflower oil, olive oil or grape seed oil in pork patties could reduce the formation of
23 HCAs without compromising eating quality.

24 **Key words:** Antioxidants; Fat modification; Maillard reactions; Oxidation.

25 **Chemical compounds studied in this article:**

26 2-amino-3-methylimidazo[4,5-f]quinoline (PubChem CID: 53462); 2-Amino-3,4-
27 dimethylimidazo[4,5-f]quinoline (PubChem CID: 62274); 2-Amino-3,8-
28 dimethylimidazo[4,5-f]quinoxaline (PubChem CID: 62275); 2-amino-3,4,8-
29 trimethylimidazo[4,5-f]quinoxaline (PubChem CID: 104739); 2-Amino-1-methyl-6-
30 phenylimidazo[4,5-b]pyridine (PubChem CID: 1530); Benzo[a]pyrene (PubChem CID:
31 2336); Benz[a]anthracene (PubChem CID: 5954)

32 **1. Introduction**

33 Fat plays an important role in the human diet. It not only creates a unique sensation
34 of food, but also helps maintain health. The consumption of pork in the world has
35 dramatically increased from 18 to 110 million tons per year (1950-2010) (Brown,
36 2013). Research found that increased saturated fatty acid intake could elevate the
37 risk of cardiovascular disease, but monounsaturated fatty acids (MUFA) and
38 polyunsaturated fatty acids (PUFA) could reduce the risk and maintain
39 cardiovascular health (McAfee et al., 2010; Sadler, 2014). Therefore, changing fatty
40 acids profile of meat products by replacing saturated fatty acids with unsaturated
41 fatty acids has attracted lots of attention in both academic research and meat
42 processors. Adding olive oil could dramatically increase the percentage of MUFA in
43 final products, whereas sunflower oil and grape seed oil could greatly raise the level
44 of PUFA in fat replaced meat products (Gunstone, 2002; Matthäus, 2008).
45 Rodríguez-Carpena et al. (2012) successfully replaced 50% fat with avocado,
46 sunflower and olive oil in cooked pork patties and reported that avocado and olive oil
47 could even offer better aroma to the final products than control ones. Vural and

48 Javidipour (2002) successfully substituted beef fat in Frankfurters with the mixture of
49 interesterified palm, cottonseed, and olive oil without changing physical parameters
50 and total sensory scores. Choi et al. (2010) used pre-emulsified grape seed oil and 2%
51 rice bran fibre to develop pork batters with 50% fat replacement and reported that
52 the fat-reduced pork batters could achieve the comparable eating quality with control
53 samples. Domínguez, Agregán, Gonçalves, and Lorenzo (2016) replaced 100% pork
54 back fat with olive oil in pork pâté, which significantly increase the content of
55 tocopherol and MUFA in cooked products without altering physio-chemical properties.
56 Domínguez, Pateiro, Agregán, and Lorenzo (2017) and Lorenzo, Munekata, Pateiro,
57 Campagnol, and Domínguez (2016) replaced 25%-75% backfat with olive oil,
58 microencapsulated fish oil and the mixture of fish oil and olive oil, which significantly
59 increased the percentage of PUFA in frankfurter type sausage and Spanish
60 *salchichón*. These results indicate that vegetable oil could be used successfully to
61 replace fat partially or completely to offer products comparable eating quality with
62 healthier fatty acids profile, i.e. high level of MUFA and PUFA.

63 However, unsaturated fatty acids in vegetable oils may pose risk in domestic cooking
64 due to their oxidation and decomposition at high temperature. For example, linoleic
65 acid was found associated with the formation of potentially toxic compounds, such as
66 free radicals, aldehydes and ketones (Guillén & Uriarte, 2012a; Katragadda et al.,
67 2010). These reactive oxygen species (ROS) initiated by unsaturated fatty acids
68 peroxidation could induce the decomposition of Amadori compounds and generate
69 1- and 3- deoxysone that are intermediates for Strecker aldehydes, pyrazines and
70 pyridines in Maillard reaction. Consequently, it might promote the formation of
71 heterocyclic amines (HCAs) (Morello, Shahidi & Ho, 2002; Turesky, 2010; Zamora &
72 Hidalgo, 2007). Effect of fatty acids/oils on the formation of HCAs has been

73 documented in previous research. Johansson et al. (1995) reported that the higher
74 level of MeIQx and DiMeIQx were found in burgers fried in rapeseed oil containing
75 high level of oleic acid with high peroxides values, compared with butter, margarine
76 and sunflower oil. Zamora et al. (2012) stated that both primary and secondary lipid
77 oxidation products, hydroperoxides, such as methyl 13-hydroperoxyoctadeca-9,11-
78 dienoate and alkenals could accelerate the formation of PhIP in chemical model
79 system. Some hydroperoxides generated from the decomposition of the unsaturated
80 hydrocarbons during heating, such as linolenate acid can also undertake
81 aromatization and de-hydrocyclization, further cleave into benzaldehydes and other
82 benzene ring-containing compounds, which are precursors of polycyclic aromatic
83 hydrocarbons (PAHs) (Chen & Chen, 2001; Lorenzo et al., 2011; Lorenzo, Purriños,
84 Fontán, & Franco, 2010; Singh, Varshney & Agarwal, 2016).

85 HCAs, PAHs and N-nitrous compounds are well-known carcinogens which were
86 detected in processed meat products (Hasnol, Jinap & Sanny, 2014; Jinap et al.,
87 2013; Liao et al., 2010; Oz & Kaya, 2010; Salmon, Knize & Felton, 1997). HCAs are
88 mainly formed with the presence of free amino acids, carbohydrates and creatine
89 under high cooking temperature (Rahman et al., 2014). IARC (1993) classified the
90 following 5 aminoimidazoarenes (AIAs) compounds as human carcinogens, including
91 2-amino-3-methylimidazo[4,5-f]quinoline (IQ), 2-amino-3,4-methylimidazo[4,5-
92 f]quinoline (MeIQ), 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline (MeIQx), 2-amino-
93 3,4,8-trimethylimidazo[4,5-f]quinoxaline (4, 8-DiMeIQx) and 2-amino-1-methyl-6-
94 phenylimidazo[4,5-b]pyridine (PhIP). PAHs are hydrocarbons that contain two or
95 more benzene rings, such as pyrene, anthracene and naphthalene. They can be
96 formed through incomplete combustion or pyrolysis of organic components, including
97 fat, protein and carbohydrates at the temperature over 200 °C. Grilling, roasting and

98 smoking meat products likely contain high level of PAHs (Alomirah et al., 2011).
99 Benz[a]anthracene (BaA) and benzo[a]pyrene (BaP) are the most potent
100 carcinogenic PAHs in processed meat products (PHE, 2008). The metabolite of BaP,
101 BaP-7,8-diol-9,10-epoxide, has been reported with the highest tumour-inducing
102 activity due to causing DNA adducts (Purcaro, Moret & Conte, 2013).

103 Vegetable oils contain various antioxidants such as vitamin E, β -carotenes and
104 phenolic compounds (Ramírez-Anaya et al., 2015). These antioxidants have been
105 characterized as free radical scavengers during cooking, which might inhibit the
106 formation of carcinogens (Janoszka, 2011; Wong, Cheng & Wang, 2012). Cheng,
107 Chen and Wang (2007) reported that marinating beef patties with phenolic
108 compounds such as epicatechin gallate, rosmarinic acids and carnosic acid could
109 significantly reduce HCAs by 24%-70% in final cooked products. Balogh et al. (2000)
110 found that HCAs (IQ, MeIQ, MeIQx, DiMeIQx and PhIP) were inhibited by 45%-75%
111 when sprayed 1% vitamin E (w/w) on the surface of beef patties before frying.
112 Therefore, in the concern of the carcinogen level in processed meat products,
113 replacing saturated fat with vegetable oils rich in unsaturated fatty acids needs to be
114 justified. Thus, the objectives of this study were to (1) explore the effect of partially
115 replacing pork back fat with sunflower oil, olive oil and grape seed oil on the
116 formation of HCAs and PAHs; (2) examine the effect of different cooking
117 temperatures on the formation of carcinogens in fat reduced pork patties.

118 **2. Material and methods**

119 **2.1 Materials**

120 Three batches of lean pork leg and pork back fat with 40.3% SFA, 43.4% MUFA and
121 10.0% PUFA (McCance & Widdowson, 2002) were purchased from Jennings
122 Caversham (Reading, UK) at different time point to consider the batch effect.

123 Excess visible fat on pork legs was trimmed, then minced by a Kenwood Food
124 processor (Chef Titanium KM010, 4.6, Kenwood Limited) and vacuum packed
125 separately. Raw materials were stored at -18 °C and defrosted 24h at 4 °C before
126 use. Commercial grape seed oil (Waitrose[®], produced in Italy) with 12.4% SFA, 20.2%
127 MUFA, 68.2% PUFA, 10-15mg tocopherols and 5.9-11.5mg/100g polyphenols (Bail,
128 Stuebiger, Krist, Unterweger, & Buchbauer, 2008), sunflower oil (Morrisons[®],
129 produced in UK) with 14.3% SFA, 20.5% MUFA, 63.3% PUFA and 50mg
130 tocopherols (McCance & Widdowson, 2002) and refined olive oil (Filippo[®], phenols
131 were removed by industrial process, produced in Italy) with 14.3% SFA, 73.0%
132 MUFA, 8.2% PUFA and 100-300mg tocopherols (McCance & Widdowson, 2002)
133 were purchased from local supermarket (Reading, UK). Oils were kept in refrigerator
134 (4°C) before making patties and further analysis.

135 The standards IQ (2-amino-3-methyl-imidazo [4,5-f] quinoline), MeIQ (2-amino-3,4-
136 dimethyl-imidazo [4,5-f] quinoline), MeIQx (2-amino-3,8-dimthylimidazo [4,5-
137 f]quinoxaline), 4,8-DiMeIQx (2-amino-3,4,8-trimethyl-imidazo [4,5-f] quinoxaline),
138 PhIP (2-amino-1-methyl-6-phenylimidazo [4,5-b] pyridine, BaA (Benz[a]anthracene)
139 and BaP (benzo[a]pyrene) were purchased from *Toronto Research Chemicals*
140 (Toronto, Canada). Ammonium acetate, triethylamine, acetonitrile (HPLC grade),
141 bovine serum albumin (BSA), dinitrophenylhydrazine (DNPH), ethyl acetate 99.5%
142 0.9000g/ml, 6M guanidine HCl (pH 6.5), hydrochloric acid solution 0.1M, methanol
143 (HPLC grade), HPLC grade water, sodium hydroxide 1M, perchloric acid (99.8%),
144 sodium phosphate buffer (pH 6.5), thiobarbituric acid (TBA), and trichloroacetic acid
145 (TCA) were purchased from *Fisher Scientific* (Loughborough, UK). 2,2-Azobis(2-
146 methylpropionamide) dihydrochloride granular 97% (ABAP), 2,2-Azino-bis(3-
147 ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS), (±)-6-Hydroxy-

148 2,5,7,8-tetramethylchromane-2-carboxylic acid (*Trolox*), phosphate buffer solution
149 0.1 M and phosphoric acid were purchased from *Sigma-Aldrich* (Gillingham, UK).
150 The solid-phase extraction *Extrelut NT 20* columns and diatomaceous earth refill
151 material were purchased from *Merck* (Darmstadt, Germany). Bond Elut propyl-
152 sulfonic acid (PRS) cartridges (100 mg, 10 ml), C-18 cartridges (7 ml) were
153 purchased from *VWR Inc* (Lutterworth, UK).

154 **2.2 Procedures for preparing and cooking pork patties**

155 The formulation of control patties (as shown in Table 2) included 700g lean pork
156 mince, 180g distill water, 100g pork back fat and 20g salt per kilogram. For the fat
157 partially replaced patties, 40% of pork back fat was replaced with sunflower oil, olive
158 oil or grape seed oil respectively. Overall, there were 4 types of pork patties
159 prepared in this study, i.e. control (100% back fat, C patties), sunflower oil patties (S
160 patties), olive oil patties (O patties) and grape seed oil patties (G patties). All
161 ingredients were homogenized at 5000rpm for 5min in the Kenwood Food processor
162 to reach a uniform emulsion. Each patty was weighed 100g, shaped in a foil cup
163 (9.0cm diameter * 2.5cm height) for standardization.

164 Patties were cooked in an air-forced oven at either 180 °C or 220 °C. Cooking was
165 terminated when core temperature of patties reached to 73 °C. After cooking, pork
166 patties were covered by foil and chilled in cold room at 4 °C for 24 hours. Physical
167 properties including texture and colour were measured on the following day of
168 cooling. While part of samples were homogenized and stored in -18 °C for further
169 chemical analysis. Cooking loss was determined according to the equation: Cooking
170 loss (%) = $(W_r - W_c)/W_r$, where W_r was the weight of raw pork patties, and W_c was

171 weight of cooked pork patties. All the treatments were replicated three times. For
172 each replicate, 8 patties were made for each treatment.

173 **2.3 Composition analysis: pH, moisture, fat and protein content**

174 pH was measured by inserting a pH meter (68X243601, Oakton Instruments, USA)
175 into the mixture by blending 5g meat sample with 45ml distill water (Puangsombat et
176 al., 2011). The moisture content was determined by drying 3g meat samples in an
177 oven at 100°C for 24 hours according to AOAC methods (Horwitz & Latimer, 2005).
178 The fat content was determined by drying sample in an oven for 4 hours firstly then
179 using Soxhlet extraction system. The protein content was determined using the
180 Kjeldahl method (Horwitz & Latimer, 2005).

181 **2.4 Lipid/protein oxidation and antioxidant capacity**

182 **2.4.1 Lipid oxidation--Thiobarbituric acid-reactive substances (TBARS) value**

183 The degree of lipid oxidation in samples was expressed by TBARS values, which
184 were determined by the method reported by Rodríguez-Carpena et al. (2012). 5 g
185 well blended pork patty was homogenized with 15 ml perchloric acid (3.86%) and
186 0.5ml BHT (4.2% in ethanol) in a beaker, which was immersed in an ice bath to
187 minimize oxidative reactions in samples during extraction. The mixture was then
188 filtered and centrifuged at 3000 rpm for 4 min, 2 ml supernatant was mixed with 2 ml
189 thiobarbituric acid (0.02 M) in test tube. The test tubes were then placed in a boiling
190 water bath (100 °C) for 45 min. After cooling, the absorbance was measured at
191 532nm using a spectrophotometer (6315, Bibby Scientific Ltd, UK). The standard
192 curve was prepared using 1,1,3,3-tetraethoxypropane (TEP) in 3.86% perchloric acid
193 with the concentration of 0, 0.5, 1.0, 2.5, 5.0 and 10.0µM.

194 **2.4.2 Total protein carbonyl value (Protein oxidation)**

195 The degree of protein oxidation can be evaluated by calculating the total carbonyl
196 value according to the method described by Rodríguez-Carpena et al. (2012). 1g of
197 well blended pork patties was homogenized at 1:10 (w/v) in 20 mM sodium
198 phosphate buffer containing 0.6 M NaCl (pH 6.5) for 30 s. Two equal aliquots of
199 0.2ml mixture were then dispensed in 2 ml eppendorf tubes, respectively. 1ml cold
200 trichloroacetic acid (TCA) (10%, w/w) was added into tubes and centrifuged for 5 min
201 at 5000 rpm. One pellet was mixed with 1 ml 2 M HCl in order to measure protein
202 concentration, while the other pellet was mixed with 1 ml of 0.2% (w/v)
203 dinitrophenylhydrazine (DNPH) in 2 M HCl in order to measure carbonyl
204 concentration. Both tubes were incubated for 1 h at room temperature. Subsequently,
205 1 ml 10% TCA was added into tubes and pellets were washed twice with 1 ml
206 ethanol: ethyl acetate (1:1, v/v) to remove excess of DNPH. The pellets were then
207 mixed with 1 ml of 20 mM sodium phosphate buffer containing 6 M guanidine HCl
208 (pH 6.5), stirred and centrifuged for 2 min at 5000 rpm to remove insoluble fragments.
209 Protein concentration was calculated from absorption at 280 nm using BSA as
210 standard. The amount of carbonyls was expressed as nmol of carbonyl per mg of
211 protein using an absorption coefficient of $21.0 \text{ nM}^{-1} \text{ cm}^{-1}$ for absorbance at 370 nm
212 for DNPH.

213 **2.4.3 Trolox equivalent antioxidant capacity (TEAC) of vegetable oil/ back fat**

214 TEAC was used to evaluate the total antioxidant capacity of vegetable oils and pork
215 back fat. The measuring procedures were based on the method reported by van den
216 Berg et al. (1999). An ABTS radical solution was prepared by mixing 2.5 mM ABAP
217 with 20 mM ABTS²⁻ stock solution in 100 mM phosphate buffer (pH 7.4), which

218 contained 150 mM NaCl. The solution was covered with foil and heated at 60°C for
219 12 min, then cooled down to room temperature. 40 µl of the sample solution was
220 mixed with 1960 µl of the freshly prepared ABTS/ABAP solution. Difference of
221 absorbance at 734 nm in 6 min was recorded. A calibration curve was made by
222 measuring the difference of absorbance in 6min for *Trolox* at the concentration of 0,
223 0.5, 1.0, 2.5, 5.0, 7.5 and 10.0 µM. The TEAC of vegetable oil or back fat was
224 presented on a molar basis to *Trolox* (µmol Trolox/100g).

225 **2.5 Analysis of HCAs**

226 HCAs extraction and purification were following the procedures reported by
227 Puangsombat et al. (2011). To minimize the variation and bias due to the unevenly
228 distribution of sauce on the surface of meat, all samples were blended well with
229 sauce before measuring. 3g ground meat sample was mixed with 12ml 1M sodium
230 hydroxide firstly, then the mixture was transferred into an *Extrelut 20 column* with
231 17g diatomaceous earth. The HCAs were eluted by 60ml ethyl acetate in *Extrelut*
232 *column*, and transferred into PRS cartridge which was pre-conditioned with 7ml ethyl
233 acetate. A PRS cartridge was then washed with 6ml 0.1M HCl, 15ml methanol/0.1M
234 HCl (45/55, v/v) and 2ml pure water to remove interferences from the PRS cartridge.
235 The HCAs were then eluted by 20ml 0.5M ammonium acetate (pH 8.5) from the PRS
236 cartridge and transferred into a C-18 cartridge that was conditioned with 5ml
237 methanol and 5ml pure water. Finally, HCAs were eluted with 1ml
238 methanol/ammonium hydroxide (9/1, v/v) from C-18 cartridge into 2ml vial, followed
239 by drying the mixture under nitrogen stream for 1.5h at room temperature. The
240 residue in the vial was dissolved with 1ml methanol and submitted for HPLC analysis.
241 1ml of mixtures containing 5 standard compounds (IQ, MeIQ, MeIQx, 4, 8-DiMeIQx

242 and PhIP) (5 ng/ml) was spiked into samples before extraction for measuring the
243 recovery rate. Three replicates were carried out for each sample.

244 IQ, MeIQ, MeIQx, 4, 8-DiMeIQx and PhIP were analysed using HPLC (HP1635
245 Series, Agilent ChemStation, Agilent Technologies, Kidlington, UK) connected with a
246 diode array UV detector (RF 2000). The HCAs were separated gradually by a
247 reversed-phase Luna 5u C18 column (250 × 4.60 mm, 5 µm, 100A, Product No:
248 00G-4252-E0, Phenomenex, UK). Mobile phase A was 0.01 M triethylamine
249 (adjusted pH 3.6 with phosphoric acid) and phase B was acetonitrile (>99%, HPLC
250 grade). The solvent contained 95% A and 5% B at beginning, then linearly changed
251 to 75% A and 25% B within 30 min at flow rate 1.0 ml/min. The temperature of
252 column was 40°C. The UV detector was set at 252 nm (Puangsombat et al., 2011).

253 **2.6 Analysis of PAHs**

254 5g meat sample was homogenised with 15ml 1M NaOH for 1 h. The homogenized
255 sample was then mixed with 17g diatomaceous earth and loaded in an *Extrelut 20*
256 *column*. Elution of PAHs started from *Extrelut column*, and was followed by propyl
257 sulphonic acid (PRS) cartridges with 60 ml of CH₂Cl₂ containing 5% toluene. The
258 dichloromethane solution was then evaporated to small amount (0.5-1ml) and the
259 rest of the solvent was dried under a nitrogen stream. The residue was re-dissolved
260 in 1ml n-hexane and transferred to the top of a glass column packed with silica-gel
261 (10 g). PAHs were then eluted by 25 ml of n-hexane and 60 ml of 60:40 (v/v) n-
262 hexane-CH₂Cl₂ mixtures. After evaporation to dryness the residues were dissolved
263 in acetonitrile (spiked and unspiked samples) before the HPLC analysis (Janoszka,
264 2011). 50ng of 2 standard mixtures (BaA and BaP) was spiked for measuring the
265 recovery rate.

266 BaA and BaP were analysed using HPLC (HP1635 Series, Agilent ChemStation,
267 Agilent Technologies, Kidlington, UK) connected with a fluorescence detector.
268 Mixture of 84% acetonitrile (>99%, HPLC grade) and 16% water (HPLC grade) were
269 used as a mobile phase under isocratic conditions. The separations were performed
270 at 40°C under isocratic conditions with flow rate 1.0 ml/min. The fluorescence
271 detection was performed by applying the following excitation (Ex)/emission (Em)
272 wavelength program: 280/410 nm from 0 to 8.50 min (BaA), 376/410 nm from 8.50 to
273 15 min (BaP) (Janoszka, 2011).

274 **2.7 Recovery rate of HCAs and PAHs**

275 The 5 standard HCA (IQ, MeIQ, MeIQx, 4, 8-DiMeIQx and PhIP) and 2 PAHs (BaA
276 and BaP) compounds were identified through the retention time of the peaks, and
277 the quantity of each individual compound was determined according to the standard
278 calibration curves, which was established by the standard solution at 0.5ng/ml,
279 5ng/ml and 50ng/ml. Limit of detection (LOD) for IQ, MeIQ, MeIQx, 4, 8-DiMeIQx,
280 PhIP, BaA and BaP were for 0.02 ng/g, 0.01 ng/g, 0.02 ng/g, 0.05 ng/g, 0.03 ng/g,
281 0.07 ng/g and 0.06 ng/g. The average recoveries of these 5 HCAs based on
282 triplicates were 60.01% for IQ, 61.76% for MeIQ, 53.64% for MeIQx, 60.57% for 4,8-
283 DiMeIQx and 55.98% for PhIP. Results were comparable with several published data
284 (Gibis, Kruwinnus & Weiss, 2015; Messner & Murkovic, 2004; Oz & Cakmak, 2016;
285 Yao et al., 2013). The recovery rate for BaA and BaP was 54.37 and 49.54%
286 respectively, which was comparable with published results of 50% - 115%
287 (Farhadian, Jinap, Abas, & Sakar, 2010; Ishizaki, Saito, Hanioka, Narimatsu, &
288 Kataoka, 2010; Iwasaki et al., 2010; Janoszka, 2011). Recovery rate could be
289 affected by sorbing materials, flow rate through cartridges, organic modifier quality
290 and/or content, interfering effects of eluting solvents (Buseti, Heitz, Cuomo, Badoer,

291 & Traverso, 2006). Stevens, Hamstra, Hegeman & Scharrer (2006) reported that
292 overlapping peaks on chromatogram caused by insufficient separation procedures
293 could result in recovery rate over 110%, and AOAC (2016) recommends that 40-120%
294 recovery rate is acceptable for compounds at 1ng/g concentration.

295 **2.8 Inhibitory rate of HCAs and PAHs**

296 Inhibitory rate was determined according to the equation:

$$297 \text{ Inhibitory rate (\%)} = (A_c - A_t) / A_c \times 100$$

298 where A_c was the total amount of HCAs/PAHs in control samples (ng/g), and A_t was
299 the total amount of HCAs/PAHs in fat partially replaced patties (S/ O/ G patties)
300 (ng/g).

301 **2.9 Physical parameters**

302 **2.9.1 Colour**

303 Colour feature including L^* , a^* and b^* was measured using Hunter Lab Colour
304 instrument (Hunter Associates Laboratory, Virginia, USA, 2003). Each sample was
305 measured at 3 different locations, while the average was recorded as the colour
306 feature of the sample.

307 **2.9.2 Texture**

308 Texture profile analysis (TPA) was performed at room temperature with a *Stable*
309 *system* Texture Analyzer (Middleboro, USA). Cylinder samples (D=18mm, H=22mm)
310 were prepared using a stainless cork borer. Before the analysis, samples were
311 tempered at room temperature (20°C) for 30min. The settings for texture analysis
312 were: load cell 5 kg, head speed 1.6 mm/s, and compression depth 10.0 mm. The

313 calculation of TPA values was based on the compression curve with force (y-axis)
314 and time (x-axis). Values for hardness (N) was defined as the absolute peak force in
315 the 1st compression cycle, cohesiveness as the area of work in the 2nd compression
316 divided by the area of work in the 1st compression, chewiness as the product of
317 hardness*cohesiveness*springiness, and springiness as the force that sample return
318 to its initial pattern after compression (Sánchez del Pulgar, Gázquez & Ruiz-
319 Carrascal, 2012). Each sample was measured at least 5 times, and the average was
320 recorded as the value of the sample.

321 **2.10 Statistical analysis**

322 Each treatment had three replicates. Statistical significance test was carried out by
323 using SPSS Statistics 21 (IBM, 2014). The significant difference in chemical
324 composition, physical property, levels of HCAs and PAHs for the 8 treatments were
325 carried out by one-way analysis of variance (ANOVA) at the significant level 0.05,
326 and *Duncan* test was selected for multiple comparison if equal variances assumed,
327 otherwise *Tamhane's T2* test was used. To analyse the effect of factors and the
328 interaction between factors (cooking temperature and replacing oil type), two-way
329 ANOVA was employed at the significant level 0.05. Multivariate linear regression
330 model was employed to explore the effect of multi independent factors, including lipid
331 oxidation, protein oxidation and antioxidant capacity of oils on the formation of HCAs
332 and PAHs at the significant level 0.05. Pearson correlation was employed for the
333 measurements as a prerequisite for the regression.

334 **3. Results and discussion**

335 **3.1 Effect of replacing oil and cooking temperature on proximate composition** 336 **and physical parameters of reduced fat pork patties**

337 3.1.1 Proximate composition

338 Table 2 shows that the effect of replacing oil and cooking temperature on proximate
339 composition and pH values in patties. Type of replacing oil did not pose any effect on
340 all proximate composition and pH, but cooking temperature had significant influence
341 on moisture and cooking loss ($p < 0.01$). Interaction between type of oil and cooking
342 temperature was only observed in cooking loss ($p < 0.01$).

343 Moisture content varies from $63.36 \pm 0.37\%$ to $67.53 \pm 0.26\%$ in fat partially
344 replaced patties, which are consistent with the results reported by Rodríguez-
345 Carpena et al. (2012). They found that pork patties with 50% fat replaced with
346 sunflower oil, olive oil and avocado oil had moisture level at 61.48-63.39% when
347 patties were cooking at 170 °C for 18 min in a forced-air oven. There was no
348 significant difference in moisture content for all the patties (with or without fat
349 replacement) ($p = 0.206$), which indicated that replacing back fat with vegetable oils
350 did not affect moisture level in the final products. However, cooking temperature
351 significantly affected the moisture content in the final products ($p < 0.05$). Higher
352 cooking temperature at 220 °C led to low moisture level in cooked patties, compared
353 with low cooking temperature at 180 °C ($p < 0.01$), which were 63.58% vs 69.15% for
354 control, 63.36% vs 65.91% for olive oil treatment, 63.49% vs 66.34% for sunflower
355 oil treatment and 65.90% vs 67.53% for grape seed oil treatment. Low moisture level
356 in the final products are directly associated with high cooking loss, as cooking loss is
357 mainly composed of water and water soluble nutrients such as proteins (Sánchez del
358 Pulgar et al., 2012). Pork patties with or without fat replacement had cooking loss
359 ranging from 20.30%- 24.75%, which was consistent with the results reported by
360 Rodríguez-Carpena, Morcuende and Estévez (2011). They found that cooked pork

361 patties at 170 °C for 18 min had cooking loss at 20.69%- 22.20%. Fat and protein
362 content ranged from 9.49%-10.01% and 15.03%-15.34% respectively, as expected,
363 fat and protein content in fat modified patties were comparable with these in C
364 patties. pH ranged from 5.85 to 5.99. Cooking temperature and type of oil did not
365 affect pH of patties, there was no interaction between temperature and type of oil as
366 well ($p>0.05$).

367 **3.1.2 Physical parameters**

368 **3.1.2.1 Texture**

369 Texture property of cooked pork patties was examined including hardness,
370 cohesiveness, springiness and chewiness through a typical texture profile analysis.
371 Results were listed in Table 3. Cooking temperature affected hardness,
372 cohesiveness and chewiness ($p \leq 0.05$), but had no effect on springiness ($p>0.05$).
373 Fat replacement with vegetable oils did not affect any of the texture attributes
374 ($p>0.05$). There was no interaction observed between oil replacement and
375 temperature for hardness, chewiness and springiness except cohesiveness ($p>0.05$).
376 High cooking temperature led to high hardness. Control patties cooked at 220°C had
377 significantly higher hardness ($26.65 \pm 3.15\text{N}$) than that cooked at 180°C ($20.14 \pm 2.81\text{N}$)
378 ($p<0.05$). Roldán et al. (2013) observed that the elevating cooking temperature
379 resulted in higher hardness, while the increased hardness in pork patties might be
380 associated with high cooking loss. There was no difference observed in hardness for
381 fat modified patties regardless of temperature ($p>0.05$), which agreed with the results
382 reported by Rodríguez-Carpena et al. (2011). They stated that patties that partially
383 (50%) replaced with sunflower oil, olive oil and avocado oil had same hardness with
384 control patties. However, Hur, Jin and Kim (2008) reported that olive oil replacement

385 in pork patties resulted in low hardness compared with control patties. The cooking
386 temperature could help to explain the disagreement. In their study, patties were
387 cooked at 100 °C in water bath, while samples were cooked in convection oven at
388 180°C or 220°C. The hardening effect of high cooking temperature could be
389 neutralized by the softening effect of to the replacing vegetable oil. As a result, there
390 was no difference observed in hardness for fat modified patties and control sample
391 ($p>0.05$).

392 Temperature affected cohesiveness greatly ($p=0.001$), because the texture
393 parameters are mainly determined by denaturation of the structural protein system,
394 i.e. actomyosin complex and collagen (Palka & Daun, 1999). Type of oil did not have
395 any effect on the cohesiveness of pork patties ($p>0.05$). Rodríguez-Carpena et al.
396 (2012) also reported that there was no difference in cohesiveness between control
397 and patties with 50% fat substitution using avocado, sunflower or olive oil. The
398 interaction between temperature and type of oil on cohesiveness was observed
399 ($p<0.05$). For control, olive and sunflower oil treatment, patties cooked at 220°C had
400 higher cohesiveness than these cooked at 180°C ($p<0.05$), but there was no
401 difference in cohesiveness for grape seed oil samples cooked at different
402 temperatures ($p>0.05$). This interaction between cooking temperature and type of oil
403 could be explained by the emulsion stability of oil/fat emulsion. Youssef and Barbut
404 (2009) and Rodríguez-Carpena et al. (2011) reported that vegetable oil with high
405 polyunsaturated fatty acids (PUFA) had small fat globules in meat emulsion, which
406 could offer a stronger fat-protein interaction. Grape seed oil contained high level of
407 PUFA compared with back fat and olive oil, which led to a stable emulsion in G
408 patties. In addition, polyphenol compounds in grape seed oil emulsion could help
409 maintain the protein functionality through inhibition of protein oxidation during

410 processing (Ganhão, Morcuende & Estévez, 2010). Thus, grape seed oil/meat
411 emulsion was less sensitive to temperature changes in relation to cohesiveness of
412 the final products.

413 For chewiness, temperature significantly affected it ($p < 0.05$). High cooking
414 temperature resulted in high chewiness. Chewiness remained similar for all patties
415 cooked at 180 °C (from 3.94 to 4.11 N.s), but increased to 5.92 ± 0.77 N.s in C patties,
416 and 5.35 ± 0.41 N.s in S patties that cooked at 220 °C. Greater hardness,
417 cohesiveness and chewiness at higher elevating cooking temperature could be due to
418 the more severe denaturation of myosin (40-60 °C) and actin (66-73 °C) (Sánchez del
419 Pulgar et al., 2012). In addition, chewiness is also associated with the water retention
420 in meat products (Roldán et al., 2014). Patties cooked at 180°C had significantly
421 higher moisture content than those cooked at 220°C (as shown in Table 2).
422 Therefore, high chewiness would be expected in samples cooked at 220°C due to
423 high moisture loss.

424 3.1.2.2 Colour

425 Effects of vegetable oil and cooking temperature on colour characteristics of cooked
426 pork patties including lightness (L^*), redness (a^*) and yellowness (b^*) were
427 summarized in Table 3. Temperature significantly affected all three parameters ($p \leq$
428 0.05), especially a^* with $p = 0.005$. Different vegetable oils did not have any impact
429 on the colour parameters, while the interaction between oil and temperature was
430 observed in yellowness (b^*). Patties cooked at 220 °C had lower L^* than those
431 cooked at 180 °C ($p < 0.01$), which agreed with the results of Sánchez del Pulgar et al.
432 (2012). The decrease in surface lightness could be attributed to the brown pigments
433 formed from caramelization of sugars and Maillard reaction when samples were

434 cooked at temperature over 90 °C (Girard, 1992). In addition, the lightness was also
435 associated with the moisture content in meat products. Qiao et al. (2001) reported
436 that there was positive correlation between lightness and moisture content in broiler
437 breast fillet. Presence of heme pigments, containing 90-95% myoglobin in muscles
438 gives meat red colour (a^*). At 180 °C, a higher a^* value was found in all pork patties
439 with oil replacement than control patties ($p < 0.05$). The antioxidants in these
440 vegetable oils, such as vitamin E could prohibit the oxidation of oxymyoglobin and
441 lead to a high redness in the final products (Hui, 2001; Sánchez del Pulgar et al.,
442 2012). Cooking temperature could significantly affect a^* as well. All patties that
443 cooked at 220 °C had significantly lower a^* than those cooked at 180 °C. The
444 reduction of a^* caused by increased temperature could be associated with the
445 denaturation of myoglobin (Nollet, 2012). Liao, Xu and Zhou (2009) found that a^* of
446 stir fried pork floss decreased significantly by 30% when cooking temperature
447 increased from 125 °C to 150 °C.

448 Yellowness b^* ranged from 15.41-18.73 in all cooked patties. Both type of oil and
449 cooking temperature had no effect on b^* values, but the interaction between type of
450 oil and cooking temperature was observed. Jamali et al. (2016) also found that b^*
451 value in beef patties was not affected by cooking temperature (160 °C and 220 °C).
452 The results of b^* in control samples (16.98-18.73) were comparable with Vittadini et
453 al. (2005).

454 **3.2 Effects of vegetable oils and cooking temperature on the formation of** 455 **HCAs**

456 Concentration of HCAs (IQ, MeIQ, MeIQx, 4, 8-DiMeIQx and PhIP) in control patties
457 and fat modified patties cooked at 180°C and 220°C were listed in Table 4. Type of

oil affected all individual HCAs compound except IQ, cooking temperature significantly affected the total amount of HCAs and all individual HCAs compounds except MelQx. Interaction between oil and cooking temperature was observed in total HCAs, IQ, MeIQ, 4, 8-DiMeIQx and PhIP, but not MelQx. At both temperatures, all fat modified patties had significantly lower amount of MeIQ, 4, 8-DiMeIQx and total HCAs than control patties ($p < 0.05$). MelQx, 4, 8-DiMeIQx and PhIP were not detectable in G patties. Tocopherols (average 50mg/100g in sunflower oil, 100-300mg/g in refined olive oil and 10-15mg/g in grape seed oil) and polyphenols (common profile in grape seed oil: catechin, epicatechin and epicatechin gallate) in these oils could play roles in reducing the final HCAs in patties (Bail et al., 2008; McCance & Widdowson, 2002; Rombaut et al., 2014). Tocopherols have been found to block dialkyl-pyrazine radicals for further reaction with creatine to form HCAs, or react with precursors of 4, 8-DiMeIQx to inhibit the formation of HCAs (Pearson et al., 1992; Vitaglione & Fogliano, 2004). Phenolic compounds could also prevent the formation of imidazoquinoxaline-type HCAs through trapping pyrazine cation radicals and some other carbon-centred radicals generated either from pyrazine cation radicals or different pathway during Maillard reaction (Kato et al., 1996). In addition, phenolic compounds have the ability to directly trap phenylacetaldehyde, which is a major intermediate during the formation of PhIP (Cheng et al., 2007).

Total HCAs ranged from not detected (Nd) to 140.57 ± 22.03 ng/g. Control patties cooked at both cooking temperatures contained significantly higher amount of total HCAs (67.56 ± 17.29 ng/g and 140.57 ± 22.03 ng/g), followed by S patties (5.98 ± 1.10 ng/g and 23.88 ± 2.44 ng/g) and O patties (4.11 ± 0.87 ng/g and 20.03 ± 2.25 ng/g), while G patties achieved the lowest total HCAs in both temperatures (Nd and 1.90 ± 0.04 ng/g). Control samples cooked at 220°C contained all types of HCAs,

483 whereas none of HCA compounds were detected in G patties cooked at 180°C. The
484 dominating compounds of HCAs were MeIQ (59.70 ± 0.98ng/g) and 4, 8-DiMeIQx
485 (43.37 ± 15.67ng/g) in C patties, while PhIP in S (14.78 ± 1.49ng/g) and O patties
486 (22.70 ± 1.95ng/g). The total HCAs in C patties were higher than some published
487 results. Vangnai et al. (2014) reported MeIQx (7.59 ± 0.43ng/g), PhIP (13.12 ± 0.72
488 ng/g) and total HCAs (22.35 ± 1.17 ng/g) in fried pork loins cooked at 204 °C for 16
489 minutes. The total level of HCAs in pan-fried well-done pork was 49.7ng/g with
490 cooking ended at 80 °C core temperature (Iwasaki et al., 2010). The sampling
491 procedure for measuring HCAs could help explain the difference. HCAs were
492 extracted from the 2mm outer layer surface of samples in this study, while lots of
493 researchers extracted HCAs from entirely ground samples. The precursors of HCAs,
494 such as creatine, glucose and free amino acids would migrate to the surface of meat
495 and enhance Maillard reactions during cooking (Gibis & Weiss, 2015). As a result,
496 surface could accumulate much higher level of HCAs compared with internal part of
497 the sample. Therefore, a higher concentration of HCAs would be expected than that
498 exacted from entirely ground samples.

499 IQ was detected up to 3.88 ng/g in cooked patties, which was in the range of 0.7-5.3
500 ng/g in fried ground beef patties (Balogh et al., 2000). At 180 °C, IQ was not detected
501 apart from O patties, but cooking at 220°C generated high level of IQ in all patties
502 (p<0.05). Different vegetable oils did not affect the formation of IQ (p>0.05), but
503 interaction between vegetable oil and temperature was observed in formation of IQ.
504 IQ is generally formed through reactions between creatinine, pyridine radicals and
505 formaldehydes (Vitaglione & Fogliano, 2004). Vegetable oils could decompose into
506 hydroperoxides, and then aldehydes and ketones at high cooking temperature,
507 which further react with amino acids in Maillard reactions (Johansson, Skog, &

508 Jagerstad, 1993; Zamora & Hidalgo, 2007). Olive oils with high level of oleic acid
509 could decompose into aldehydes (-CHO) much faster (3-15 times) than sunflower oil
510 and grape seed oil containing high amount of linoleic and linolenic acid at 190°C
511 (Guillén & Uriarte, 2012b). This might explain why IQ was only detected in olive oil
512 patties at 180°C. But the pathway of formation of HCAs in the real meat system was
513 more complicated than chemical model system, and antioxidants in oils should also
514 be considered (Johansson, Skog, & Jagerstad, 1993). Therefore, further
515 investigation is needed to understand promoting mechanism and role of antioxidants
516 in oils. The highest MeIQ was found in control patties cooked at 220 °C (59.70±0.98
517 ng/g), followed by in control patties cooked at 180 °C (18.26±14.46 ng/g). Janoszka
518 (2010) reported that 6.28 ng/g MeIQ was detected in pan-fried pork patties cooked at
519 170°C for 12 minutes, which was similar with this study. Formation of MeIQ was
520 completely inhibited by olive oil and sunflower oil at both cooking temperatures.
521 Grape seed oil could only inhibit formation of MeIQ at low cooking temperature, while
522 1.31ng/g was detected in G patties cooked at 220°C. The inhibitory effect on MeIQ
523 could be attributed to the antioxidants such as vitamin E and polyphenols in
524 vegetable oils. Rounds et al. (2012) and Liao et al. (2009) also reported that vitamin
525 E and polyphenols could reduce the formation of MeIQ. Balogh et al. (2000) found
526 that vitamin E had stronger inhibitory effect on MeIQ with reduction rate 64.3% than
527 phenolic compound in Oleoresin rosemary extract with reduction rate 47.9% in fried
528 beef patties. Since olive oil and sunflower oil contain higher level of vitamin E than
529 grape seed oil, therefore, stronger inhibition of MeIQ would be expected in O patties
530 and S patties. Cooking temperature did not affect MeIQx level in patties ($p>0.05$), but
531 formation of MeIQx was significantly affected by different vegetable oils. S patties
532 and O patties had similar MeIQx with control sample. For G patties cooked at both

533 temperatures, there was no MeIQx detected. However, sunflower oil and olive oil did
534 not affect MeIQx in patties, although both oils were rich of vitamin E. No MeIQx was
535 detected at all which was in agreement with Rounds et al. (2012), who also reported
536 grape seed extract could completely inhibit the formation of MeIQx in cooked beef
537 patties. Temperature significantly increased formation of 4, 8-DiMeIQx as evidenced
538 in C patties cooked at 180 °C and 220 °C (25.66 ± 1.51 ng/g and 43.37 ± 15.67 ng/g)
539 ($p < 0.05$). All vegetable oils effectively reduced 4, 8-DiMeIQx in patties. Grape seed
540 oil was the most effective one among the three vegetable oil as 4, 8-DiMeIQx was
541 not detected in G patties cooked at both temperatures.

542 PhIP ranged from Nd to 24.07 ± 1.99 ng/g. A similar level of PhIP (18.4 ± 11.5 ng/g) in
543 the fried pork patties was reported by Zhang et al. (2013), when patties were cooked
544 at 180°C for 5 min. At 180 °C, PhIP (11.43 ± 6.33 ng/g) was totally inhibited by all 3
545 vegetable oils, but only grape seed oil could completely inhibit the formation of PhIP
546 at both cooking temperatures. The stronger inhibitory effect on PhIP in G patties
547 could be attributed to the phenolic compounds in grape seed oil. Gibis and Weiss
548 (2012), Jamali et al. (2016) and Oguri et al. (1998) found that catechin, epicatechin
549 and epicatechin-3-O-gallate in grape seed extract may be responsible for 50%-90%
550 reduction of PhIP in both oven cooked beef patties and chemical model system.
551 Zamora and Hidalgo (2015) suggested phenolic compounds could effectively
552 scavenge the carbonyl compounds in the Strecker degradation of phenylalanine to
553 produce phenylacetaldehyde (major intermediate in the development of PhIP).
554 Temperature significantly affected PhIP level in pork patties ($p < 0.05$). PhIP
555 increased significantly in C patties from 11.43 ± 6.33 ng/g to 24.07 ± 1.99 ng/g, O
556 patties from Nd to 14.78 ± 1.49 ng/g and S patties from Nd to 22.70 ± 1.95 ng/g when
557 cooking temperature increased from 180 °C to 220 °C ($p < 0.05$). The results agreed

558 with Gibis and Weiss (2012) and Wong et al. (2012) that PhIP level was directly
559 related to the cooking temperature. At 175-200 °C, only very low level of PhIP (0-
560 6.91 ng/g) could be formed even at varied cooking time, but it could increase
561 dramatically to 31.80ng/g with prolonged cooking time if temperature went above
562 200 °C.

563 Type of vegetable oil significantly affected the level of total HCAs in cooked patties
564 ($p<0.05$), but temperature did not have any effect ($p>0.05$). Interaction between
565 types of vegetable oil and cooking temperature was observed as well ($p<0.05$). In the
566 current study, reduction of total HCAs by 85.75%-93.90% was found in O patties,
567 83.01%-91.15% in S patties, while G patties achieved the highest reduction rate at
568 98.64%-100% (Table 4). Antioxidants in the vegetable oils could be responsible for
569 the reduction of total HCAs as a strong negative correlation ($r=-0.618$, $p<0.01$) was
570 disclosed between total HCAs level and antioxidant capacity (TEAC) of oils (Table 6).
571 Grape seed oil had the highest TEAC value with 0.71 ± 0.01 $\mu\text{mol Trolox}/100\text{g}$,
572 followed by olive oil (0.52 ± 0.05 $\mu\text{mol Trolox}/100\text{g}$) and sunflower oil (0.18 ± 0.04 μmol
573 $\text{Trolox}/100\text{g}$), while pork back fat had the lowest TEAC value (0.09 ± 0.02 μmol
574 $\text{Trolox}/100\text{g}$) (Figure 1). Therefore, reduction of HCAs in vegetable oils, especially
575 grape seeds oil was expected compared with control sample. Results were
576 comparable with findings of Matthäus (2008) and Castelo-Branco and Torres (2012).
577 Balogh et al. (2000) found that 1% vitamin E spray on the surface of beef patties
578 could reduce the concentrations of IQ, MeIQ, MeIQx, DiMeIQx and PhIP significantly
579 by 45% to 75%, because vitamin E could remove free radicals in Maillard reactions.
580 Similar result was also reported by Lan, Kao and Che (2004). They found that 70%
581 of total HCAs (IQ, MeIQ, MeIQx, 4, 8-DiMeIQx and PhIP) were prohibited when 0.2%
582 α -tocopherol was added into ground pork 1h before cooking. Phenolic compounds,

583 such as catechin, epicatechin-3-O-gallate, oligomer procyanidins and tocopherols in
584 grape seed oil contributed to its antioxidant capacity (Agostini et al., 2012; Crews et
585 al., 2006; Matthäus, 2008). Vitaglione and Fogliano (2004) suggested that mixture of
586 antioxidant compounds could perform better than single antioxidant as they could
587 inhibit various pathways in different steps of reactions. Therefore, phenolic
588 compounds might work synergistically with tocopherols to enhance each other to
589 inhibit the formation of HCAs. However, the synergistic effect between different
590 antioxidants needs to be further examined.

591 Cooking temperature significantly affected total HCAs in pork patties ($p < 0.01$) (Table
592 4). Patties cooked at 220°C had significantly higher level of total HCAs than these at
593 180°C ($p < 0.01$). Effect of temperature on the formation of HCAs was well examined
594 in previous research (Knize et al., 1994; Liao et al., 2009; Oz & Kaya, 2011).
595 Thermal processing has vital influence on the formation of polar HCAs (IQ, MeIQ,
596 MeIQx, DiMeIQx and PhIP), which are formed in meat products when samples are
597 cooked at 160-250 °C, typical domestically cooking temperature. High cooking
598 temperature generated more diverse types of HCAs, but also stimulate the
599 accumulation of the amount of HCAs on the surface of meat products (Olsson &
600 Pickova, 2005; Skog, Johansson & Jaegerstad, 1998).

601 **3.3 Effects of vegetable oils and cooking temperature on the formation of** 602 **PAHs**

603 Concentration of PAHs (BaA and BaP) in cooked pork patties with different cooking
604 temperature was listed in Table 5. The range of total PAHs was from 1.59 ± 0.26 ng/g
605 to 3.84 ± 0.21 ng/g. The dominating compound of PAHs was BaP in all samples. BaA
606 ranged from 0.14-0.31 ng/g in cooked patties, while BaP ranged from 1.44 to

607 3.53ng/g. Temperature did not affect the formation of both BaA and BaP, but type of
608 vegetable oil had significant effect on the formation ($p<0.05$). Interaction between
609 type of oil and cooking temperature was also observed in both compounds (Table 5).
610 BaP level in this study (1.44-3.53 ng/g) are consistent with results reported by Nisha
611 et al. (2015) and Janoszka (2011), i.e. 1.52 ng/g of BaP in the oven broiled pork and
612 1.61 ng/g BaP in oven grilled pork chop (17min at 170°C). Olive oil and grape seed
613 oil showed inhibitory effect on BaP when patties cooked at 220 °C, but no effect or
614 even promoting effect was observed at 180 °C. On the contrast, sunflower oil offered
615 inhibition on BaP at 180 °C, but promotion at 220 °C. As BaP is one of the highest
616 toxic potency compounds during meat cooking, EU Commission has regulated that
617 the updated limit of BaP occurring in processed meat and seafood products is 2 ng/g
618 (Purcaro et al., 2013; Wretling et al., 2010). Among all the patties, only O patties
619 cooked at 220 °C and S patties cooked at 180 °C met the safety regulation of BaP.
620 Therefore, it is necessary to develop any procedures or alternative methods that
621 reduce the amount of BaP to safety limit.

622 The effect of oil and interaction between type of oil and cooking temperature on the
623 formation of PAHs were significant ($p<0.05$). Cooking temperature did not affect the
624 formation of PAHs ($p>0.05$). S patties cooked at 220 °C had the highest total PAHs
625 (3.84 ± 0.21 ng/g), followed by G patties (3.46 ± 0.16 ng/g) cooked at 180 °C and C
626 patties (3.28 ± 0.07 ng/g) cooked at 220 °C. O patties cooked at 220°C obtained the
627 lowest PAHs. PAHs were mainly associated with the pyrolysis of fat undertaken at
628 high temperature (Viegas et al., 2012). Therefore, smoking point of vegetable oils
629 may help explain the difference in PAHs. Sunflower oil and grape seed oil contain
630 high content of PUFAs, especially linolenic acid and linolenic acid have lower smoke
631 points (grape seed oil 216°C, sunflower oil 227°C) which makes them easy to

632 decompose, compared with olive oil (smoke point 242°C). The decomposition of oil
633 could generate more reactive free radicals to accelerate the production of PAHs
634 (Chen & Lin, 1997; Elmore et al., 2002). They also concluded that hydroperoxides
635 from lipid oxidation, could subsequently generate cyclic compounds through
636 intramolecular reaction, and PUFA could undergo further polymerization. In addition,
637 vegetable oils themselves contained BaP, which might increase the total amount of
638 PAHs in cooked meat. Fromberg, Højgard and Duedahl-olesen (2007) reported that
639 olive oil, sunflower oil, and grape seed oil approximately contained 0.12 ng/g, 0.4
640 ng/g and 1.0 ng/g BaP respectively. As a result, high level of PAHs was expected in
641 sunflower oil and grape seed oil samples. Although vegetable oils contain
642 antioxidants, the inhibitory effect on PAHs formation was not observed consistently.
643 In S patties, the inhibitory efficiency at both temperature were 7.75% and 51.52%,
644 but olive oil and grape seed oil increased the formation of PAHs by 17.07% and
645 34.11%, respectively. It shows that antioxidants in vegetable oils were not involved in
646 the formation of PAHs to a great extent, which is further confirmed by the correlation
647 analysis. As indicated in Table 6, there is no correlation relationship observed
648 between antioxidant capacity of oil (TEAC) and total PAHs. The impact of
649 tocopherols and phenolic compounds on the formation of PAHs in processed meat
650 were not well documented. In vitro study, Zhu et al. (2014) found that vitamin E
651 intake could significantly prohibit free radicle induced by BaP and protect cellular
652 damage in human lung, but the effect antioxidants on formation of PAHs in food
653 products has been scarce.

654 **3.4 Correlation between lipid oxidation, protein oxidation and the formation of** 655 **HCAs and PAHs**

656 In Figure 2, S patties, O patties and G patties had significantly lower TBARS values
657 than control samples ($p < 0.05$), while G patties had the lowest TBARS value
658 compared with O and S patties ($p = 0.001$). This inhibitory effect against lipid oxidation
659 could be attributed to the antioxidants (tocopherols and phenolic compounds) within
660 the oils, since a significantly negative relationship was found between TBARS values
661 and antioxidant capacity of oils ($r = -0.764$, $p < 0.01$). Wong et al. (2015) reported that
662 0.1-0.4mmol vitamin E could inhibit 45% of lipid oxidation in beef patties, by
663 obstructing the formation of some key aldehydes and ketones during lipid oxidation.
664 Similar results were achieved by Ahn, Grün and Fernando (2002) as well. Frankel
665 (1998) proposed that α -tocopherol could prevent the chain propagating and remove
666 free radicals through reacting with either singlet oxygen or peroxy radicals.
667 Consequently lipid oxidation was reduced. Meanwhile, polyphenols, such as
668 epicatechin (EC) oligomer procyanidins were also sufficient to inhibit lipid oxidation
669 by reducing free radicals and preventing chain propagation in cooked pork and beef
670 (Rojas & Brewer, 2007). They could also chelate metals (iron and copper in meat) or
671 react with ROS, and then turn into non-radical species. As a result, reactions were
672 terminated (Roman et al., 2013). Moreover, Ahn et al. (2002) and Tang et al. (2001)
673 reported that polyphenols such as catechin, epicatechin were more efficient in
674 inhibiting lipid oxidation than α -tocopherol at the same concentration in cooked meat.
675 This could explain why G patties had the lowest TBARS values than S and O patties.
676 Although Gunstone (2002) stated that a higher degree of unsaturation of fatty acids
677 could be easier to trigger the lipid oxidation and interacted with Maillard reaction, the
678 presence of antioxidants should be also considered in the formation of HCAs.

679 Protein carbonyls are produced from protein oxidative degradation in meat products,
680 which were used to analyse degree of protein oxidation (Figure 3). Significant effect

681 of vegetable oils on the protein oxidation was observed ($p=0.001$). C patties had a
682 significantly higher level of protein carbonyls (12.11 mmol/kg) than other 3 fat
683 modified patties ($p<0.05$). The protein oxidation could be inhibited by the presence of
684 antioxidants in oil, as negative correlation between antioxidant capacity of oils (TEAC)
685 and protein carbonyl level was found with $r=-0.606$, $p<0.01$, as indicated in Table 6.
686 Botsoglou et al. (2014) found that protein carbonyl value could be reduced from 3.25
687 mmol/kg to 2.25 mmol/kg in cooked pork patties when 50mg/kg α -tocopherol was
688 added. Vuorela et al. (2005) reported that phenolic compounds, including
689 vinylsyringol and sinapic acid in rapeseed oil had good antioxidant capacity against
690 protein oxidation in cooked pork patties. Ganhão et al. (2010) also found that
691 arbutus-berries extract containing catechins significantly reduced protein oxidation
692 by chelating heme iron in cooked patties. However, there was no difference in the
693 protein carbonyl level among O, S and G patties ($p>0.05$).

694 In the development of HCAs under high temperature cooking, both lipid oxidation
695 and protein oxidation are involved with Maillard reactions in the meat system
696 (Johansson, Skog & Jagerstad, 1993; Zamora & Hidalgo, 2007). Lot of researchers
697 believed that formation of HCAs could be primarily related to interactions between
698 free radicals generated from lipid oxidation and free radicals produced in Maillard
699 reactions (Hwang & Ngadi, 2002; Skog et al., 1998). Therefore, it is useful to explore
700 the relationship between lipid/protein oxidation and the formation of HCAs. In this
701 study, correlation analysis was conducted between TBARS/protein carbonyl values
702 and concentration of total HCAs in fat modified cooked patties (Table 6). Significant
703 positive correlation was disclosed between total HCAs and TBARS ($r=0.826$, $p<0.01$)
704 and between HCAs and protein carbonyl ($r=0.778$, $p<0.01$), which further confirmed
705 that both lipid oxidation and protein oxidation participated the formation of HCAs

706 during cooking process. In order to further examine the relationship between lipid
707 oxidation/protein oxidation/antioxidant capacity of lipids and total HCAs in cooked
708 patties, multivariate linear regression model was displayed below,

709 $\text{Total HCAs} = -42.37 + 108.26 * \text{TBARS} + 5.647 * \text{Protein Carbonyl}$.

710 It can be seen from the equation that TBARS (lipid oxidation) played a predominant
711 role on the formation of HCAs, compared with protein carbonyl (protein oxidation).
712 The factor 'TEAC (antioxidant capacity of lipids)' has been removed from the model,
713 because the strong correlation between TEAC and TBARS/protein carbonyl
714 indicates that variance accounted for TEAC could be well accounted by
715 TBARS/protein carbonyl. In cross-validation, adding/removing 'TEAC' caused little
716 change in adjusted R square of the predicted models, which indicated variance
717 caused by TEAC could be well explained by other independent factors in the model
718 (Field, 2009).

719 Free radicals, such as aldehydes and ketones generated from lipids oxidation could
720 interact with Maillard reactions by reacting with the polar head of an amino group to
721 produce more HCAs (Jägerstad et al., 1998; Zamora & Hidalgo, 2007). On the other
722 hand, active protein carbonyl residues, such as alkyl, peroxy radicals that formed by
723 muscle protein oxidation can be initialized by lipid oxidation, metal ions and other
724 peroxidized compounds (Cai et al., 2002). Subsequently, these carbonyls could
725 interact with Maillard reaction via Schiff base and then generate Strecker aldehydes,
726 which are intermediates of imidazoquinolines and imidazoquinoxalines (Estévez,
727 2011; Soladoye et al., 2015). Researchers also suggested that lipid oxidation could
728 trigger protein oxidation by reacting with heme iron that released from myoglobin
729 (Ganhão et al., 2010; Vuorela et al., 2005). In this work, there was no correlation

730 observed between TBARS /protein carbonyl and PAHs ($p>0.05$), which indicated that
731 the involvement of lipid and protein oxidation in the formation of PAHs was only at
732 null level. Thus, no linear regression model was fitted. The antioxidants in vegetable
733 oils could not inhibit the formation of PAHs, which was evidenced by null correlation
734 between TEAC and PAHs ($p>0.05$).

735 **4. Conclusions**

736 Control patties contained the highest amount of HCAs and relatively higher PAHs at
737 180 °C and 220 °C. All 3 fat modified patties had significantly lower HCAs, which
738 could be attributed to antioxidants, such as tocopherols and polyphenol compounds
739 existing in the oils. The negative correlation ($r= -0.618$, $p<0.01$) between the
740 antioxidant capacity of lipids and the total amount of HCAs could be useful evidence
741 to support this claim. Both lipid and protein oxidation contributed to the formation of
742 HCAs, which were supported by the positive relationship between TBARS/ protein
743 carbonyl values and total HCAs with $r= 0.826$ and 0.788 ($p<0.01$), respectively. Olive
744 oil and sunflower oil completely prohibited MeIQ, whereas grape seed oil could
745 inhibit MeIQx, 4, 8-DiMeIQx and PhIP. Grape seed oil could achieve the highest
746 inhibitive effect on the formation of HCAs. However, effect of vegetable oils on the
747 formation of PAHs was not consistent, which could be attributed to complexity of oil
748 decomposition and antioxidants in the oils. The involvement of lipid oxidation and
749 protein oxidation in formation of PAHs was limited or at a minimum level.
750 Antioxidants in oils could not reduce the total amount of PAHs effectively. Therefore,
751 it is necessary to explore other methods to reduce PAHs in processed meat. Due to
752 the synergistic effect of antioxidants, it would be interesting to explore replacing pork
753 back fat with a mixture of several types of oils to reach an optimum fatty acids profile,

754 instead of replacing pork back fat with one single type of oil. Overall, replacing pork
755 back fat with vegetable oils in processed meat products could offer healthier meat
756 products with reduced HCAs without compromising eating quality.

757 **Acknowledgement**

758 We thank C. Humphrey, C. Busey, M. Puranik and other technicians at University of
759 Reading for the technical support.

760 **References**

- 761 Agostini, F., Bertussi, R. A., Agostini, F., Atti dos Santos, A. C., Rossato, M., &
762 Vanderlinde, R. (2012). Supercritical Extraction from Vinification Residues:
763 Fatty Acids, α -Tocopherol, and Phenolic Compounds in the Oil Seeds from
764 Different Varieties of Grape. *The ScientificWorld Journal*, 2012, 1-9.
765 doi:10.1100/2012/790486
- 766 Ahn, J., Grün, I. U., & Fernando, L. N. (2002). Antioxidant Properties of Natural Plant
767 Extracts Containing Polyphenolic Compounds in Cooked Ground Beef. *Food*
768 *Chemistry and Toxicology*, 67, 1364-1369.
- 769 Alomirah, H., S., A.-Z., S., A.-H., Zaghloul, S., Sawaya, W., Ahmed, N., & Kannan, K.
770 (2011). Concentration and dietary exposure to polycyclic aromatic
771 hydrocarbons (PAHs) from grilled and smoked foods. *Food Control*, 22, 2028-
772 2035.
- 773 AOAC. (2016). *Official methods of analysis of AOAC International*. Guidelines for
774 Standard Method Performance Requirements: Appendix F.
775 http://www.eoma.aocac.org/app_f.pdf Retrieved on 24//05/2017.
- 776 Bail, S., Stuebiger, G., Krist, S., Unterweger, H., & Buchbauer, G. (2008).
777 Characterisation of various grape seed oils by volatile compounds,
778 triacylglycerol composition, total phenols and antioxidant capacity. *Food*
779 *Chemistry*, 108(3), 1122-1132.
- 780 Balogh, Z., Gray, J. I., Gomaa, E. A., & Booren, A. M. (2000). Formation and
781 inhibition of heterocyclic aromatic amines in fried ground beef patties. *Food*
782 *and Chemical Toxicology*, 38(5), 395-401.
783 doi:http://dx.doi.org/10.1016/S0278-6915(00)00010-7
- 784 Botsoglou, E., Govaris, A., Ambrosiadis, I., Fletourisc, D., & Papageorgioud, G.
785 (2014). Effect of olive leaf (*Olea europea* L.) extracts on protein and lipid
786 oxidation in cooked pork meat patties enriched with n-3 fatty acids. *Journal of*
787 *the Science of Food and Agriculture*, 94, 227-234.
- 788 Brown, L. R. (2013). Moving up the food Chain. Retrieved from
789 <http://permaculturenews.org/2013/11/27/moving-food-chain/>
- 790 Cai, W., Gao, Q.-d., Zhu, L., Peppia, M., He, C., & Vlassara, H. (2002). Oxidative
791 Stress-Inducing Carbonyl Compounds From Common Foods: Novel
792 Mediators of Cellular Dysfunction. *Molecular Medicine*, 8(7), 337-346.

- 793 Castelo-Branco, V. N., & Torres, A. G. (2012). Generalized linear model describes
794 determinants of total antioxidant capacity of refined vegetable oils. *European*
795 *Journal of Lipid Science and Technology*, 114(3), 332-342.
- 796 Chen, B. H., & Chen, Y. C. (2001). Formation of Polycyclic Aromatic Hydrocarbons
797 in the Smoke from Heated Model Lipids and Food Lipids. *Journal of*
798 *agricultural and food chemistry*, 49, 5238-5243.
- 799 Chen, B. H., & Lin, Y. S. (1997). Formation of Polycyclic Aromatic Hydrocarbons
800 during Processing of Duck Meat. *Journal of agricultural and food chemistry*,
801 45, 1394-1403.
- 802 Cheng, K. W., Chen, F., & Wang, M. (2007). Inhibitory activities of dietary phenolic
803 compounds on heterocyclic amine formation in both chemical model system
804 and beef patties. *Molecular Nutrition & Food Research*, 51(8), 969-976.
- 805 Choi, Y.-S., Choi, J.-H., Han, D.-J., Kim, H.-Y., Lee, M.-A., Kim, H.-W., . . . Kim, C.-J.
806 (2010). Optimization of replacing pork back fat with grape seed oil and rice
807 bran fiber for reduced-fat meat emulsion systems. *Meat science*, 84(1), 212-
808 218. doi:http://dx.doi.org/10.1016/j.meatsci.2009.08.048
- 809 Crews, C., Hough, P., Godward, J., Brereton, P., Lees, M., Guiet, S., & Winkelmann,
810 W. (2006). Quantitation of the Main Constituents of Some Authentic Grape-
811 Seed Oils of Different Origin. *Journal of agricultural and food chemistry*, 54,
812 6261-6265.
- 813 Domínguez, R., Agregán, R., Gonçalves, A., & Lorenzo, J. M. (2016). Effect of fat
814 replacement by olive oil on the physico-chemical properties, fatty acids,
815 cholesterol and tocopherol content of pâté. *Grasas y Aceites*, 67(2), 133.
- 816 Domínguez, R., Pateiro, M., Agregán, R., & Lorenzo, J. M. (2017). Effect of the
817 partial replacement of pork backfat by microencapsulated fish oil or mixed fish
818 and olive oil on the quality of frankfurter type sausage. *Journal of Food*
819 *Science and Technology*, 54(1), 26-37.
- 820 Elmore, J. S., Campo, M. M., Enser, M., & Mottram, D. S. (2002). Effect of Lipid
821 Composition on Meat-like Model Systems Containing Cysteine, Ribose, and
822 Polyunsaturated Fatty Acids. *Journal of agricultural and food chemistry*, 50,
823 1126-1132.
- 824 Estévez, M. (2011). Protein carbonyls in meat systems: A review. *Meat science*,
825 89(3), 259-279. doi:http://dx.doi.org/10.1016/j.meatsci.2011.04.025
- 826 Farhadian, A., Jinap, S., Abas, F., & Sakar, Z. I. (2010). Determination of polycyclic
827 aromatic hydrocarbons in grilled meat. *Food Control*, 21, 606-610.
- 828 Field, A. (2009). *Discovering statistics using SPSS : (and sex and drugs and rock 'n'*
829 *roll)*. 3rd ed. London: Sage.
- 830 Frankel, E. N. (1998). *Lipid oxidation: Chapter 8 Antioxidatns*. Dundee: Oily Press.
- 831 Fromberg, A., Højgaard, A., & Duedahl-olesen, L. (2007). Analysis of polycyclic
832 aromatic hydrocarbons in vegetable oils combining gel permeation
833 chromatography with solid-phase extraction clean-up. *Food Additives and*
834 *Contaminants*, 24(7), 758-767.
- 835 Ganhão, R., Morcuende, D., & Estévez, M. (2010). Protein oxidation in emulsified
836 cooked burger patties with added fruit extracts: Influence on colour and
837 texture deterioration during chill storage. *Meat science*, 85(3), 402-409.
838 doi:http://dx.doi.org/10.1016/j.meatsci.2010.02.008
- 839 Gibis, M., Kruwinnus, M., & Weiss, J. (2015). Impact of different pan-frying
840 conditions on the formation of heterocyclic aromatic amines and sensory
841 quality in fried bacon. *Food Chemistry*, 168, 383-389.
842 doi:http://dx.doi.org/10.1016/j.foodchem.2014.07.074

- 843 Gibis, M., & Weiss, J. (2012). Antioxidant capacity and inhibitory effect of grape seed
844 and rosemary extract in marinades on the formation of heterocyclic amines in
845 fried beef patties. *Food Chemistry*, *134*, 766-774.
- 846 Gibis, M., & Weiss, J. (2015). Impact of Precursors Creatine, Creatinine, and
847 Glucose on the Formation of Heterocyclic Aromatic Amines in Grilled Patties
848 of Various Animal Species. *Journal of Food Science*, *80*(11), C2430-C2439.
- 849 Girard, J. P. (1992). Chapter 2: Cooking. In M.Laroche (Ed.), *Technology of Meat*
850 *and Meat Products*. Chichester, England: Ellis Horwood Limited.
- 851 Guillén, M., & Uriarte, P. (2012a). Aldehydes contained in edible oils of a very
852 different nature after prolonged heating at frying temperature: Presence of
853 toxic oxygenated α,β unsaturated aldehydes. *Food Chemistry*, *131*(3), 915-
854 926. doi:http://dx.doi.org/10.1016/j.foodchem.2011.09.079
- 855 Guillén, M. D., & Uriarte, P. S. (2012b). Study by ¹H NMR spectroscopy of the
856 evolution of extra virgin olive oil composition submitted to frying temperature
857 in an industrial fryer for a prolonged period of time. *Food Chemistry*, *134*(1),
858 162-172. doi: https://doi.org/10.1016/j.foodchem.2012.02.083
- 859 Gunstone, F. D. (2002). *Vegetable oils in food technology : composition, properties*
860 *and uses*. Oxford Blackwell.
- 861 Hasnol, N. D. S., Jinap, S., & Sanny, M. (2014). Effect of different types of sugars in
862 a marinating formulation on the formation of heterocyclic amines in grilled
863 chicken. *Food Chemistry*, *145*, 514-521.
- 864 Horwitz, W., & Latimer, G. W. (2005). *Official methods of analysis of AOAC*
865 *International. 18th Edition*. Gaithersburg AOAC International.
- 866 Hui, Y. H. (2001). Chapter 3: Meat colour. In O. Y. a. J. West (Ed.), *Meat science*
867 *and applications*. New York: Marcel Dekker.
- 868 Hur, S. J., Jin, S. K., & Kim, I. S. (2008). Effect of extra virgin olive oil substitution for
869 fat on quality of pork patty. *Journal of the Science of Food and Agriculture* *88*,
870 1231-1237.
- 871 Hwang, D. K., & Ngadi, M. (2002). Kinetics of Heterocyclic Amines Formation in
872 Meat Emulsion at Different Fat Contents. *LWT - Food Science and*
873 *Technology*, *35*(7), 600-606. doi:http://dx.doi.org/10.1006/fstl.2002.0913
- 874 IARC. (1993). Some naturally occurring substances: food items and constituents,
875 heterocyclic aromatic amines and mycotoxins. *IARC Monographs on the*
876 *Evaluation of Carcinogenic Risks to Humans*, *56*, 165-195.
- 877 Ishizaki, A., Saito, K., Hanioka, N., Narimatsu, S., & Kataoka, H. (2010).
878 Determination of polycyclic aromatic hydrocarbons in food samples by
879 automated on-line in-tube solid-phase micro extraction coupled with high-
880 performance liquid chromatography-fluorescence detection. *Journal of*
881 *Chromatography A*, *1217*, 5555-5563.
- 882 Iwasaki, M., Kataoka, H., Ishihara, J., Takachi, R., Hamada, G. S., Sharma, S., . . .
883 Tsugane, S. (2010). Heterocyclic amines content of meat and fish cooked by
884 Brazilian methods. *Journal of Food Composition and Analysis*, *23*, 61-69.
- 885 Jägerstad, M., Skog, K., Arvidsson, P., & Solyakov, A. (1998). Chemistry, formation
886 and occurrence of genotoxic heterocyclic amines identified in model systems
887 and cooked foods. *Z Lebensm Unters Forsch A*, *207*, 419-427.
- 888 Jamali, M. A., Zhang, Y., Teng, H., Li, S., Wang, F., & Peng, Z. (2016). Inhibitory
889 Effect of *Rosa rugosa* Tea Extract on the Formation of Heterocyclic Amines in
890 Meat Patties at Different Temperatures. *Molecules*, *21*(2), 173.
- 891 Janoszka, B. (2010). Heterocyclic amines and azaarenes in pan-fried meat and its
892 gravy fried without additives and in the presence of onion and garlic. *Food*

- 893 *Chemistry*, 120(2), 463-473.
894 doi:<http://dx.doi.org/10.1016/j.foodchem.2009.10.039>
- 895 Janoszka, B. (2011). HPLC-fluorescence analysis of polycyclic aromatic
896 hydrocarbons (PAHs) in pork meat and its gravy fried without additives and in
897 the presence of onion and garlic. *Food Chemistry*, 126(3), 1344-1353.
898 doi:<http://dx.doi.org/10.1016/j.foodchem.2010.11.097>
- 899 Jinap, S., Mohd-Mokhtar, M. S., Farhadian, A., Hasnol, N. D. S., Jaafar, S. N., &
900 Hajeb, P. (2013). Effects of varying degrees of doneness on the formation of
901 Heterocyclic Aromatic Amines in chicken and beef satay. *Meat science*, 94,
902 202-207.
- 903 Johansson, M., Skog, K., & Jagerstad, M. (1993). Effects of edible oils and fatty
904 acids on the formation of mutagenic heterocyclic amines in a model system.
905 *Cardnogenesis*, 14(1), 89-94.
- 906 Johansson, M.A.E., Fredholm, L., Bjerne, I. & Jägerstad, M. (1995). Influence of
907 frying fat on the formation of heterocyclic amines in fried beefburgers and pan
908 residues. *Food and Chemical Toxicology*, 33(12), pp.993-1004.
- 909 Kato, T., Harashima, T., Moriya, N., Kikugawa, K., & Hiramoto, K. (1996). Formation
910 of the mutagenic/carcinogenic imidazoquinoxaline-type heterocyclic amines
911 through the unstable free radical Maillard intermediates and its inhibition by
912 phenolic antioxidants. *Carcinogenesis*, 17(11), 2469-2476.
- 913 Katragadda, H. R., Fullana, A., Sidhu, S., & Carbonell-Barrachina, Á. A. (2010).
914 Emissions of volatile aldehydes from heated cooking oils. *Food Chemistry*,
915 120(1), 59-65. doi:<http://dx.doi.org/10.1016/j.foodchem.2009.09.070>
- 916 Knize, M. G., Dolbeare, F. A., Carroll, K. I., Moore, D. H., & Felton, J. S. (1994).
917 Effect of cooking time and temperature on the heterocyclic amine content of
918 fried beef patties. *Food and Chemical Toxicology*, 32(7), 595-603.
- 919 Lan, C. M., Kao, T. H., & Che, B. H. (2004). Effects of heating time and antioxidants
920 on the formation of heterocyclic amines in marinated foods. *Journal of*
921 *Chromatography B*, 802, 27-37.
- 922 Liao, G. Z., Wang, G. Y., Xu, X. L., & Zhou, G. H. (2010). Effect of cooking methods
923 on the formation of heterocyclic aromatic amines in chicken and duck breast.
924 *Meat science*, 85(1), 149-154.
- 925 Liao, G. Z., Xu, X. L., & Zhou, G. H. (2009). Effects of cooked temperatures and
926 addition of antioxidants on formation of heterocyclic aromatic amines in pork
927 floss. *Journal of Food Processing and Preservation*, 33(2), 159-175.
- 928 Lorenzo, J. M., Munekata, P. E. S., Pateiro, M., Campagnol, P. C. B., & Domínguez,
929 R. (2016). Healthy Spanish salchichón enriched with encapsulated n- 3 long
930 chain fatty acids in konjac glucomannan matrix. *Food Research International*,
931 89, 289-295.
- 932 Lorenzo, J. M., Purriños, L., Bermudez, R., Cobas, N., Figueiredo, M., & García
933 Fontán, M. C. (2011). Polycyclic aromatic hydrocarbons (PAHs) in two
934 Spanish traditional smoked sausage varieties: "Chorizo gallego" and "Chorizo
935 de cebolla". *Meat science*, 89(1), 105-109. doi:
936 <http://dx.doi.org/10.1016/j.meatsci.2011.03.017>
- 937 Lorenzo, J. M., Purriños, L., Fontán, M. C. G., & Franco, D. (2010). Polycyclic
938 aromatic hydrocarbons (PAHs) in two Spanish traditional smoked sausage
939 varieties: "Androlla" and "Botillo". *Meat science*, 86(3), 660-664. doi:
940 <http://dx.doi.org/10.1016/j.meatsci.2010.05.032>
- 941 Matthäus, B. (2008). Virgin grape seed oil: Is it really a nutritional highlight?
942 *European Journal of Lipid Science and Technology*, 110, 645-650.

- 943 McAfee, A. J., McSorley, E. M., Cuskelly, G. J., Moss, B. W., Wallace, J. M. W.,
944 Bonham, M. P., & Fearon, A. M. (2010). Red meat consumption: An overview
945 of the risks and benefits. *Meat science*, *84*(1), 1-13.
946 doi:<http://dx.doi.org/10.1016/j.meatsci.2009.08.029>
- 947 McCance, R. A., & Widdowson, E. M. (2002). McCance and Widdowson's The
948 composition of foods. Section 2.4: Fats and oils (6th ed.). Cambridge, London:
949 The Royal Society of Chemistry, Food Standards Agency.
- 950 Messner, C., & Murkovic, M. (2004). Evaluation of a new model system for studying
951 the formation of heterocyclic amines. *Journal of Chromatography B*, *802*(1),
952 19-26. doi:<http://dx.doi.org/10.1016/j.jchromb.2003.11.015>
- 953 Morello, M. J., Shahidi, F., & Ho, C. (2002). *Free radicals in food : chemistry,*
954 *nutrition, and health effects. Chapter 5: Radical Induced Formation of D-*
955 *Glucosone from Amadori Compounds*. Washington DC: American Chemical
956 Society.
- 957 Nisha, A. R., Dinesh, K. V., Arivudainambi, S., Umer, M., & Khan, M. S. (2015).
958 Polycyclic aromatic hydrocarbons in processed meats: A toxicological
959 perspective. *Research Journal of Chemistry and Environment*, *19*(6), 72-76.
- 960 Nollet, L. M. L. (2012). Chapter 3: Attributes of Muscle Food: Color, Texture, Flavor.
961 In L. M. L. Nollet (Ed.), *Handbook of Meat, Poultry and Seafood Quality*.
962 Oxford, UK: Blackwell Publishing Ltd.
- 963 Oguri, A., Suda, M., Totsuka, Y., Sugimura, T., & Wakabayashi, K. (1998). Inhibitory
964 effects of antioxidants on formation of heterocyclic amines. *Mutation*
965 *Research*, *42*, 237-245.
- 966 Olsson, V., & Pickova, J. (2005). The Influence of Production Systems on Meat
967 Quality, with Emphasis on Pork. *AMBIO: A Journal of the Human Environment*,
968 *34*(4), 338-343.
- 969 Oz, F., & Cakmak, I. H. (2016). The effects of conjugated linoleic acid usage in
970 meatball production on the formation of heterocyclic aromatic amines. *LWT -*
971 *Food Science and Technology*, *65*, 1031-1037.
- 972 Oz, F., & Kaya, M. (2010). Effect of Cooking Techniques and Levels on the
973 Formation of Heterocyclic Aromatic Amines in Chicken and Fish. *Journal of*
974 *Animal and Veterinary Advances*, *9*(8), 1259-1264.
- 975 Oz, F., & Kaya, M. (2011). The inhibitory effect of black pepper on formation of
976 heterocyclic aromatic amines in high-fat meatball. *Food Control*, *22*, 596-600.
- 977 Palka, K., & Daun, H. (1999). Changes in texture, cooking losses, and myofibrillar
978 structure of bovine M. semitendinosus during heating. *Meat science*, *51*(3),
979 237-243. doi:[http://dx.doi.org/10.1016/S0309-1740\(98\)00119-3](http://dx.doi.org/10.1016/S0309-1740(98)00119-3)
- 980 Pearson, A. M., Chen, C., Gray, J. I., & Aust, S. D. (1992). Mechanism (s) involved in
981 meat mutagen formation and inhibition. *Free Radical Biology and Medicine*,
982 *13*(2), 161-167.
- 983 PHE. (2008). *Public Health England: Polycyclic Aromatic Hydrocarbons*. England.
- 984 Puangsombat, K., Gadgil, P., Houser, T. A., Hunt, M. C., & Smith, J. S. (2011).
985 Heterocyclic amine content in commercial ready to eat meat products. *Meat*
986 *Science*, *88*, 227-233.
- 987 Purcaro, G., Moret, S., & Conte, L. S. (2013). Overview on polycyclic aromatic
988 hydrocarbons: Occurrence, legislation and innovative determination in foods.
989 *Talanta*, *105*, 292-305. doi:<http://dx.doi.org/10.1016/j.talanta.2012.10.041>
- 990 Qiao, M., Fletcher, D., Smith, D., & Northcutt, J. (2001). The effect of broiler breast
991 meat color on pH, moisture, water-holding capacity, and emulsification
992 capacity. *Poultry Science*, *80*(5), 676-680.

- 993 Rahman, U., Sahar, A., Khan, M. I., & Nadeem, M. (2014). Production of heterocyclic
994 aromatic amines in meat: Chemistry, health risks and inhibition. A review.
995 *LWT - Food Science and Technology*, *59*, 229-233.
- 996 Ramírez-Anaya, J. d. P., Samaniego-Sánchez, C., Castañeda-Saucedo, M. C.,
997 Villalón-Mir, M., & de la Serrana, H. L.-G. (2015). Phenols and the antioxidant
998 capacity of Mediterranean vegetables prepared with extra virgin olive oil using
999 different domestic cooking techniques. *Food Chemistry*, *188*, 430-438.
1000 doi:<http://dx.doi.org/10.1016/j.foodchem.2015.04.124>
- 1001 Rodríguez-Carpena, J. G., Morcuende, D., & Estévez, M. (2011). Partial
1002 Replacement of Pork Back-Fat by Vegetable Oils in Burger Patties: Effect on
1003 Oxidative Stability and Texture and Color Changes during Cooking and
1004 Chilled Storage. *Journal of Food Science*, *76*(7), 1025-1031.
- 1005 Rodríguez-Carpena, J. G., Morcuende, D., & Estévez, M. (2012). Avocado,
1006 sunflower and olive oils as replacers of pork back-fat in burger patties: Effect
1007 on lipid composition, oxidative stability and quality traits. *Meat science*, *90*(1),
1008 106-115. doi:<http://dx.doi.org/10.1016/j.meatsci.2011.06.007>
- 1009 Rojas, M. C., & Brewer, M. S. (2007). Effect of Natural Antioxidants on Oxidative
1010 Stability of Cooked, Refrigerated Beef and Pork. *Journal of Food Science*, *72*,
1011 S282-S288.
- 1012 Roldán, M., Antequera, T., Martín, A., Mayoral, A. I., & Ruiz, J. (2013). Effect of
1013 different temperature–time combinations on physicochemical, microbiological,
1014 textural and structural features of sous-vide cooked lamb loins. *Meat science*,
1015 *93*(3), 572-578. doi:<http://dx.doi.org/10.1016/j.meatsci.2012.11.014>
- 1016 Roldán, M., Antequera, T., Pérez-Palacios, T., & Ruiz, J. (2014). Effect of added
1017 phosphate and type of cooking method on physico-chemical and sensory
1018 features of cooked lamb loins. *Meat science*, *97*(1), 69-75.
- 1019 Roman, O., Heyd, B., Broyart, B., Castillo, R., & Maillard, M. (2013). Oxidative
1020 reactivity of unsaturated fatty acids from sunflower, high oleic sunflower and
1021 rapeseed oils subjected to heat treatment, under controlled conditions. *LWT -*
1022 *Food Science and Technology*, *52*, 49-59.
- 1023 Rombaut, N., Savoie, R., Thomasset, B., Bélliard, T., Castello, J., Van Hecke, É., &
1024 Lanoisellé, J.-L. (2014). Grape seed oil extraction: Interest of supercritical fluid
1025 extraction and gas-assisted mechanical extraction for enhancing polyphenol
1026 co-extraction in oil. *Comptes Rendus Chimie*, *17*(3), 284-292.
1027 doi:<http://dx.doi.org/10.1016/j.crci.2013.11.014>
- 1028 Rounds, L., Havens, C. M., Feinstein, Y., Friedman, M., & Ravishankar, S. (2012).
1029 Plant extracts, spices, and essential oils inactivate *Escherichia coli* O157: H7
1030 and reduce formation of potentially carcinogenic heterocyclic amines in
1031 cooked beef patties. *Journal of agricultural and food chemistry*, *60*(14), 3792-
1032 3799.
- 1033 Sadler, M. J. (2014). Food, Nutrients and Food Ingredients with Authorities EU Health
1034 Claims (Vol. 1). Cambridge, UK: Woodhead Publishing.
- 1035 Salmon, C. P., Knize, M. G., & Felton, J. S. (1997). Effects of Marinating on
1036 Heterocyclic Amine Carcinogen Formation in Grilled Chicken. *Food and*
1037 *Chemical Toxicology*, *35*, 433-441.
- 1038 Sánchez del Pulgar, J., Gázquez, A., & Ruiz-Carrascal, J. (2012). Physico-chemical,
1039 textural and structural characteristics of sous-vide cooked pork cheeks as
1040 affected by vacuum, cooking temperature, and cooking time. *Meat science*,
1041 *90*(3), 828-835. doi:<http://dx.doi.org/10.1016/j.meatsci.2011.11.024>

- 1042 Singh, L., Varshney, J. G., & Agarwal, T. (2016). Polycyclic aromatic hydrocarbons'
1043 formation and occurrence in processed food. *Food Chemistry*, 199, 768-781.
1044 doi:<http://dx.doi.org/10.1016/j.foodchem.2015.12.074>
- 1045 Skog, K. I., Johansson, M. A. E., & Jaègerstad, M. I. (1998). Carcinogenic
1046 Heterocyclic Amines in Model Systems and Cooked Foods: A Review on
1047 Formation, Occurrence and Intake. *Food and Chemical Toxicology*, 36, 879-
1048 896.
- 1049 Stevens, J. M., Hamstra, A., Hegeman, T., & Scharrer, G. (2006). Factors and
1050 Parameters that Directly Affect Recovery of Collected Fractions in Preparative
1051 HPLC. 24.
1052 [http://www.gilson.com/Resources/Factors_and_Parameters_that_Directly_Aff](http://www.gilson.com/Resources/Factors_and_Parameters_that_Directly_Affect_Recovery_of_Collected_Fractions_in_Preparative_HPLC.pdf)
1053 [ect_Recovery_of_Collected_Fractions_in_Preparative_HPLC.pdf](http://www.gilson.com/Resources/Factors_and_Parameters_that_Directly_Affect_Recovery_of_Collected_Fractions_in_Preparative_HPLC.pdf)
- 1054 Soladoye, O. P., Juárez, M. L., Aalhus, J. L., Shand, P., & Estévez, M. (2015).
1055 Protein Oxidation in Processed Meat: Mechanisms and Potential Implications
1056 on Human Health. *Comprehensive Reviews in Food Science and Food Safety*,
1057 14(2), 106–122.
- 1058 Tang, S., Sheehan, D., Buckley, D. J., Morrissey, P. A., & Kerry, J. P. (2001). Anti-
1059 oxidant activity of added tea catechins on lipid oxidation of raw minced red
1060 meat, poultry and fish muscle. *International Journal of Food Science &*
1061 *Technology*, 36(6), 685-692.
- 1062 Turesky, R. J. (2010). Heterocyclic Aromatic Amines, Potential human carcinogens. .
1063 *Advances in molecular toxicology*, 4, 37-74.
- 1064 van den Berg, R., Haenen, G. R. M. M., van den Berg, H., & Bast, A. (1999).
1065 Applicability of an improved Trolox equivalent antioxidant capacity (TEAC)
1066 assay for evaluation of antioxidant capacity measurements of mixtures. *Food*
1067 *Chemistry*, 66(4), 511-517. doi:[http://dx.doi.org/10.1016/S0308-](http://dx.doi.org/10.1016/S0308-8146(99)00089-8)
1068 [8146\(99\)00089-8](http://dx.doi.org/10.1016/S0308-8146(99)00089-8)
- 1069 Vangnai, K., Houser, T. A., Hunt, M. C., & Smith, J. S. (2014). Effect of enhancement
1070 on the formation of heterocyclic amines in cooked pork loins: Preliminary
1071 studies. *Meat science*, 98(2), 88-93.
1072 doi:<http://dx.doi.org/10.1016/j.meatsci.2014.05.007>
- 1073 Viegas, O., Amaro, L. F., Ferreira, I. M. P. L. V. O., & Pinho, O. (2012). Inhibitory
1074 Effect of Antioxidant-Rich Marinades on the Formation of Heterocyclic
1075 Aromatic Amines in Pan-Fried Beef. *Journal of Agriculture and Food*
1076 *Chemistry*, 60, 6235-6240.
- 1077 Vitaglione, P., & Fogliano, V. (2004). Use of antioxidants to minimize the human
1078 health risk associated to mutagenic/carcinogenic heterocyclic amines in food.
1079 *Journal of Chromatography B*, 802, 189-199.
- 1080 Vittadini, E., Rinaldi, M., Chiavaro, E., Barbanti, D., & Massini, R. (2005). The effect
1081 of different convection cooking methods on the instrumental quality and yield
1082 of pork Longissimus dorsi. *Meat science*, 69(4), 749-756.
1083 doi:<http://dx.doi.org/10.1016/j.meatsci.2004.11.005>
- 1084 Vuorela, S., Salminen, H., Makela, M., Kivikari, R., Karonen, M., & M.Heinonen.
1085 (2005). Effect of Plant Phenolics on Protein and Lipid Oxidation in Cooked
1086 Pork Meat Patties. *Journal of agricultural and food chemistry*, 53, 8492-8497.
- 1087 Vural, H., & Javidipour, I. (2002). Replacement of beef fat in Frankfurters by
1088 interesterified palm, cottonseed and olive oils. *European Food Research and*
1089 *Technology*, 214(6), 465-468.

- 1090 Wong, D., Cheng, K., & Wang, M. (2012). Inhibition of heterocyclic amine formation
1091 by water-soluble vitamins in Maillard reaction model systems and beef patties.
1092 *Food Chemistry*, 133, 760-766.
- 1093 Wong, D., Hu, X., Tao, N., Wang, X., & Wang, M. (2015). Effect and mechanism of
1094 pyridoxamine on the lipid peroxidation and stability of polyunsaturated fatty
1095 acids in beef patties. *Journal of the Science of Food and Agriculture*.
1096 doi:10.1002/jsfa.7522
- 1097 Wretling, S., Eriksson, A., Eskhult, G. A., & Larsson, B. (2010). Polycyclic aromatic
1098 hydrocarbons (PAHs) in Swedish smoked meat and fish. *Journal of Food*
1099 *Composition and Analysis*, 23, 264-272.
- 1100 Yao, Y., Peng, Z., Shao, B., Wan, K., Wang, F., Zhang, Y., . . . Hui, T. (2013). Effects
1101 of frying and boiling on the formation of heterocyclic amines in braised
1102 chicken. *Poultry Science*, 92(11), 3017-3025.
- 1103 Youssef, M. K., & Barbut, S. (2009). Effects of protein level and fat/oil on emulsion
1104 stability, texture, microstructure and color of meat batters. *Meat science*, 82(2),
1105 228-233. doi:http://dx.doi.org/10.1016/j.meatsci.2009.01.015
- 1106 Zamora, R., Alcón, E. and Hidalgo, F.J. (2012). Effect of lipid oxidation products on
1107 the formation of 2-amino-1-methyl-6-phenylimidazo [4, 5-b] pyridine (PhIP) in
1108 model systems. *Food chemistry*, 135(4), 2569-2574.
- 1109 Zamora, R., & Hidalgo, F. J. (2007). Coordinate Contribution of Lipid Oxidation and
1110 Maillard Reaction to the Nonenzymatic Food Browning. *Critical Reviews in*
1111 *Food Science and Nutrition*, 45(1), 49-59.
- 1112 Zamora, R., & Hidalgo, F. J. (2015). 2-Amino-1-methyl-6-phenylimidazo[4,5-b]-
1113 pyridine (PhIP) formation and fate: an example of the coordinate contribution
1114 of lipid oxidation and Maillard reaction to the production and elimination of
1115 processing-related food toxicants. *Royal Society of Chemistry*, 5, 9709-9721.
- 1116 Zhang, Y., Yu, C., Mei, J., & Wang, S. (2013). Formation and mitigation of
1117 heterocyclic aromatic amines in fried pork. *Food Additives & Contaminants:*
1118 *Part A*, 30(9), 1501-1507.
- 1119 Zhu, W., Cromie, M. M., Cai, Q., Lv, T., Singh, K., & Gao, W. (2014). Curcumin and
1120 Vitamin E Protect against Adverse Effects of Benzo[a]pyrene in Lung
1121 Epithelial Cells. *PLoS ONE* 9(3), e92992. .
1122 doi:doi:10.1371/journal.pone.0092992

1123

1124

1125

1126

1127 Table 1: Formulation of control and fat reduced pork patties

Ingredients in recipe represent as g/1kg					
Treatment	Lean pork leg	Oil	Pork back fat	Salt	Water
Control, pork back fat (C patties)	700	0	100	20	180
Fat replaced with Sunflower oil (S patties)	700	40	60	20	180
Fat replaced with Olive oil (O patties)	700	40	60	20	180
Fat replaced with Grape seed oil (G patties)	700	40	60	20	180

1128

1129

1130

1131

1132

1133

1134

1135 Table 2: Proximate composition and pH values of 4 types of patties cooked at 180 °C and 220 °C ^{a,b}.

Treatment	Cooking temperature(°C)	Moisture (%)	Cooking loss	Fat (%)	Protein (%)	pH
Control	180	69.15±0.30 ^d	20.51±1.59 ^a	10.01±0.85 ^a	15.19±1.56 ^a	5.99±0.01 ^a
	220	63.58±0.60 ^a	24.75±1.24 ^d	9.98±1.05 ^a	15.03±1.69 ^a	5.85±0.02 ^b
Olive oil	180	65.91±0.16 ^b	20.30±0.60 ^a	9.62±1.21 ^a	15.21±1.98 ^a	5.85±0.02 ^b
	220	63.36±0.37 ^a	23.21±0.84 ^{bcd}	9.49±1.25 ^a	15.09±1.37 ^a	5.86±0.01 ^b
Sunflower oil	180	66.34±0.63 ^b	22.54±0.29 ^{bc}	9.68±1.17 ^a	15.28±1.59 ^a	5.88±0.01 ^b
	220	63.49±0.15 ^a	24.23±0.76 ^{cd}	9.70±1.05 ^a	15.17±1.94 ^a	5.87±0.02 ^b
Grapeseed oil	180	67.53±0.26 ^c	21.88±0.31 ^{ab}	9.76±1.14 ^a	15.34±1.32 ^a	5.87±0.02 ^b
	220	65.90±0.09 ^b	24.72±1.09 ^d	9.68±1.32 ^a	15.17±1.46 ^a	5.87±0.01 ^b
p-value (Type of oil)		0.21	0.43	0.74	0.54	0.065
p-value (Temperature)		<0.01	<0.01	0.33	0.28	0.16
p-value (Interaction between oil* temperature)		0.18	<0.01	0.49	0.15	0.16

1136 ^a Results with different letters in the same column are significantly different at the level p<0.05.

1137 ^b Values represented as the Mean ± standard deviation (SD), n=3.

1138 Table 3: Texture parameters (hardness, cohesiveness and chewiness) and colour parameters (lightness L*, redness a* and
 1139 yellowness b*) in 4 types of patties cooked at 180 °C and 220 °C ^{a,b}

Treatment	Cooking temperature (°C)	Hardness (N)	Cohesiveness	Chewiness (N.s)	Springiness	L*	a*	b*
Control	180	20.14±2.81 ^a	0.34±0.02 ^{ab}	4.06±0.77 ^{ab}	0.71±0.03 ^a	70.71±0.08 ^{bc}	2.30±0.11 ^c	16.98±0.80 ^{abc}
	220	26.65±3.15 ^b	0.37±0.01 ^d	5.92±0.77 ^c	0.77±0.04 ^a	65.80±1.82 ^a	1.46±0.22 ^{ab}	18.73±1.15 ^c
Olive oil	180	18.02±3.58 ^a	0.34±0.01 ^a	3.94±0.41 ^a	0.72±0.04 ^a	69.80±1.76 ^{bc}	2.82±0.08 ^d	16.23±0.71 ^{ab}
	220	21.75±0.37 ^a	0.38±0.01 ^d	5.35±0.41 ^{bc}	0.75±0.02 ^a	65.33±3.69 ^a	1.18±0.02 ^a	17.44±1.90 ^{bc}
Sunflower oil	180	17.98±2.45 ^a	0.35±0.01 ^{abc}	3.96±0.35 ^a	0.73±0.05 ^a	70.10±0.57 ^{bc}	2.86±0.13 ^d	15.41±0.73 ^a
	220	19.40±2.81 ^a	0.38±0.02 ^d	4.34±0.99 ^{ab}	0.74±0.03 ^a	65.97±2.05 ^a	1.43±0.22 ^{ab}	17.59±0.45 ^{bc}
Grape seed oil	180	17.86±1.12 ^a	0.37±0.01 ^{cd}	4.11±0.30 ^{ab}	0.77±0.01 ^a	71.41±1.74 ^c	3.24±0.37 ^d	17.03±0.38 ^{abc}
	220	21.55±2.37 ^a	0.36±0.01 ^{bc}	5.22±1.02 ^{abc}	0.77±0.04 ^a	68.15±0.47 ^{ab}	1.66±0.43 ^b	18.47±0.15 ^c
p-value (Type of oil)		0.10	0.87	0.50	0.341	0.062	0.54	0.65
p-value (Temperature)		0.05	0.001	0.001	0.085	0.05	0.005	0.05
p-value (Interaction between oil*temperature)		0.41	0.02	0.33	0.36	0.083	0.84	0.04

1140 ^a Results with different letters in the same column are significantly different at the level p<0.05.

1141 ^b Values represented as the Mean ± SD, n=3.

1142 Table 4: Heterocyclic amines in cooked pork patties with partial replacement of fat by vegetable oils at 180 °C and 220 °C ^{a,b,c}

Treatment	Cooking temperature	IQ (ng/g)	MeIQ (ng/g)	MeIQx (ng/g)	4,8-DiMeIQx (ng/g)	PhIP (ng/g)	Total (ng/g)	Inhibitory efficiency
Control	180	Nd	18.26±14.46 ^a	8.34±1.78 ^{ab}	25.66±1.51 ^b	11.43±6.33 ^a	67.56±17.29 ^c	N/a
	220	3.88±3.50 ^a	59.70±0.98 ^b	13.45±7.43 ^b	43.37±15.67 ^c	24.07±1.99 ^b	140.57±22.03 ^d	N/a
Olive oil	180	0.58±0.01 ^b	Nd	3.50±0.68 ^a	Nd	Nd	4.11±0.87 ^a	93.90%
	220	1.30±0.42 ^b	Nd	2.52±0.36 ^a	1.31±0.22 ^a	14.78±1.49 ^a	20.03±2.25 ^b	85.75%
Sunflower oil	180	Nd	Nd	4.32±0.50 ^a	1.02±0.50 ^a	Nd	5.98±1.10 ^a	91.15%
	220	0.64±0.16 ^b	Nd	4.31±0.55 ^a	5.12±0.35 ^a	22.70±1.95 ^b	23.88±2.44 ^b	83.01%
Grape seed oil	180	Nd	Nd	Nd	Nd	Nd	Nd	100%
	220	0.59±0.04 ^b	1.31±0.06 ^c	Nd	Nd	Nd	1.90±0.04 ^a	98.64%
p-value (Type of oil)		0.12	<0.01	<0.01	<0.01	<0.01	<0.01	-
p-value (Temperature)		0.037	<0.01	0.37	0.039	<0.01	<0.01	-
p-value (Interaction between oil*temperature)		0.040	<0.01	0.24	0.035	<0.01	<0.01	-

1143 ^a Results with different letters in the same column are significantly different at the level $p < 0.05$.

1144 ^b Values represented as the Mean \pm SD, $n=3$.

1145 ^c Nd: Not Detected.

1146

1147 Table 5: Polycyclic aromatic hydrocarbons in in cooked pork patties with partial replacement of fat by vegetable oils at 180 °C and
 1148 220 °C ^{a,b}

Treatment	Cooking temperature (°C)	BaA (ng/g)	BaP(ng/g)	Total PAHs (ng/g)	Inhibitory efficiency
Control	180	0.15±0.01 ^a	2.44±0.37 ^c	2.58±0.36 ^c	N/a
	220	0.21±0.03 ^b	3.08±0.06 ^d	3.28±0.07 ^d	N/a
Olive oil	180	0.15±0.02 ^a	2.24±0.40 ^{bc}	2.38±0.40 ^{bc}	7.75%
	220	0.15±0.01 ^a	1.44±0.27 ^a	1.59±0.26 ^a	51.52%
Sunflower oil	180	0.14±0.01 ^a	1.88±0.17 ^{ab}	2.02±0.16 ^{ab}	21.71%
	220	0.31±0.02 ^c	3.53±0.20 ^e	3.84±0.21 ^e	-17.07%
Grape seed oil	180	0.18±0.01 ^{ab}	3.29±0.15 ^d	3.46±0.16 ^d	-34.11%
	220	0.18±0.05 ^{ab}	2.51±0.07 ^c	2.71±0.07 ^c	17.38%
p-value (Type of oil)		<0.01	<0.01	0.031	-
p-value (Temperature)		0.1	0.076	0.43	-
p-value (Interaction between oil* temperature)		<0.01	<0.01	<0.01	-

1149 ^a Results with different letters in the same column are significantly different at the level p<0.05.

1150 ^b Values represented as the Mean ± SD, n=3.

1151


1152 Table 6: Pearson correlation coefficient (p) between the level of total HCAs/PAHs (ng/g) and TBARS, protein carbonyl and TEAC

	TEAC	HCAs	PAHS
TBARS	-0.764**	0.826**	-0.154
Protein carbonyl	-0.606**	0.778**	0.019
TEAC	-	-0.618**	0.301

1153 ** Significant level 0.01


1154

Figure 1: Trolox Equivalent Antioxidant capacity (umol Trolox/100g) of pork backfat and 3 vegetable oils.




Results with different letters are significantly different at the level $p < 0.05$.

Figure 2: TBARS values in pork patties cooked at 180 °C and 220 °C.



Results with different letters are significantly different at the level $p < 0.05$.

Figure 3: Protein carbonyl values in pork patties cooked at 180 °C and 220 °C.



Results with different letters are significantly different at the level $p < 0.05$

- Antioxidants in these oils could inhibit the formation of HCAs
- Grape seed oil achieved the highest inhibition capacity compared with sunflower oil and olive oil
- Lipid oxidation plays key role in the formation of HCAs
- Cooking temperature did not affect the total PAHs in pork patties

ACCEPTED MANUSCRIPT