
Contributed Paper

Use of a seagrass residency index to apportion commercial fishery landing values and recreation fisheries expenditure to seagrass habitat service

Emma L. Jackson,^{*†} Siân E. Rees,^{*} Catherine Wilding,^{*‡} and Martin J. Attrill^{*}

^{*}Marine Institute, Plymouth University, Drake Circus, Plymouth, PL4 8AA, United Kingdom

[†]School of Medical and Applied Sciences, Central Queensland University, Bryan Jordan Drive, P.O. Box 1319, Gladstone, Queensland 4680, Australia

[‡]Cornwall Wildlife Trust, Five Acres, Allet, Truro, Cornwall, TR4 9DJ, United Kingdom

Abstract: *Where they dominate coastlines, seagrass beds are thought to have a fundamental role in maintaining populations of exploited species. Thus, Mediterranean seagrass beds are afforded protection, yet no attempt to determine the contribution of these areas to both commercial fisheries landings and recreational fisheries expenditure has been made. There is evidence that seagrass extent continues to decline, but there is little understanding of the potential impacts of this decline. We used a seagrass residency index, that was trait and evidence based, to estimate the proportion of Mediterranean commercial fishery landings values and recreation fisheries total expenditure that can be attributed to seagrass during different life stages. The index was calculated as a weighted sum of the averages of the estimated residence time in seagrass (compared with other habitats) at each life stage of the fishery species found in seagrass. Seagrass-associated species were estimated to contribute 30%–40% to the value of commercial fisheries landings and approximately 29% to recreational fisheries expenditure. These species predominantly rely on seagrass to survive juvenile stages. Seagrass beds had an estimated direct annual contribution during residency of €58–91 million (4% of commercial landing values) and €112 million (6% of recreation expenditure) to commercial and recreational fisheries, respectively, despite covering <2% of the area. These results suggest there is a clear cost of seagrass degradation associated with ineffective management of seagrass beds and that policy to manage both fisheries and seagrass beds should take into account the socioeconomic implications of seagrass loss to recreational and commercial fisheries.*

Keywords: commercial fisheries, ecosystem services, habitat valuation, seagrass, recreational fisheries

El Uso de un Índice de Residencia de Pastos Marinos para Distribuir Valores de Desembarco de Pesquerías Comerciales y Gastos de Pesquerías Recreativas para Servicios de Hábitat de Pastos Marinos

Resumen: *Donde los pastos marinos dominan las costas, estos tienen un papel fundamental en el mantenimiento de poblaciones de especies explotadas. Por esto, los pastos marinos en el Mediterráneo merecen protección, pero no se ha hecho intento alguno por determinar la contribución de estas áreas a los atraques de las pesquerías comerciales y los gastos de las pesquerías recreativas. Existe evidencia de que la distribución de los pastos marinos continúa declinando, pero hay poco entendimiento de los impactos potenciales de esta disminución. Usamos un índice de residencia de pastos marinos, con base en la evidencia del ciclo de vida, para estimar la proporción de valores de desembarco de pesquerías comerciales y de gastos totales de pesquerías recreativas que se le puede atribuir a los pastos marinos durante diferentes etapas de vida. El índice se calculó como la suma ponderada de los promedios de tiempos de residencia estimada en los pastos marinos (en comparación con otros hábitats) durante cada etapa de vida de las especies objeto de pesquerías que se encuentran en los pastos. Se estimó que las especies asociadas a los pastos contribuyen entre el 30*

§Address for correspondence: School of Medical and Applied Sciences, Building 604, Central Queensland University - Marina Campus, Bryan Jordan Drive, Gladstone, Queensland 4680, Australia, email emma.jackson@cqu.edu.au
Paper submitted May 5, 2014; revised manuscript accepted September 7, 2014.

y el 40% al valor de desembarco de las pesquerías comerciales y aproximadamente al 29% del gasto de las pesquerías recreativas. Estas especies dependen predominantemente de los pastos marinos para sobrevivir a las etapas juveniles. Durante la residencia, se estimó que los pastos marinos tuvieron una contribución anual directa de €58-91 millones (4% de los valores de desembarco comercial) y de €112 millones (6% del gasto recreativo) de las pesquerías comerciales y recreativas, respectivamente, a pesar de tener una cobertura menor al 2% del área. Estos resultados sugieren que existe un costo claro de la degradación de pastos marinos asociado con el manejo poco efectivo de los mismos y que la política de manejo tanto de las pesquerías como de los pastos marinos debe considerar las implicaciones socio-económicas de la pérdida de los pastos para las pesquerías comerciales y recreativas.

Palabras Clave: pastos marinos, pesquerías comerciales, pesquerías recreativas, servicios ambientales, valoración de hábitat

Introduction

Fisheries provide food security and socioeconomic and recreation benefits to humans (Pauly 2009). In the Mediterranean, commercial fisheries target a wide diversity of species (Rätz et al. 2010), and over 80% of fishing vessels registered in 2008 were artisanal fishing boats operating mostly inshore and targeting local markets for sale of their landings (Sacchi 2011). Total commercial first sale landings of species in the Mediterranean is approximately €1.6–2.2 billion/year (Hutniczac & Roth 2012). Recreational fishing activities also contribute to the economy (Toivonen et al. 2004; Pawson et al. 2007). In Europe alone, €25 billion is spent per year by anglers on equipment, transportation, and lodgings (Dillon 2004). Both recreational and commercial fishing activities depend on the physical resource and functioning of marine ecosystems (Bell 1997; Rees et al. 2010).

Seagrass meadows are considered to have a fundamental role in maintaining populations of commercially and recreationally exploited fisheries species by indirectly supporting coastal food webs (Vizzini et al. 2002) and directly providing one or more of the following: permanent habitat, allowing full life cycle completion; temporary nursery area for juvenile development; feeding area for various life-history stages; and refuge from predation (Jackson et al. 2001). These provisions have been called habitat services (TEEB 2010), and they improve fisheries and shellfisheries' quality and quantity and secure supply.

A seagrass fringe is found on almost all coasts of the Mediterranean (Green & Short 2003); estimated coverage is 2.5–5.5 million ha (Buia et al. 2000). The primary species of seagrass is the Mediterranean endemic *Posidonia oceanica*. These meadows are protected under the European Council (EC) Habitats Directive within designated protected areas (European Marine Sites), and EC Regulation 1626/94 excludes mobile fishing gear use within 1.5 nautical miles of the coast, where the seagrass occurs. Despite this protection, seagrass is in decline in the Mediterranean (Langmead et al. 2007) and many areas lack enforcement of protection measures. Principle 4 of the ecosystem approach (as defined by the CBD

2000) states that in order to recognize “potential gains from management, there is a need to understand and manage the ecosystem in an economic context.” This requires an understanding of the economic associations of ecosystem services and markets through delineation of links between ecosystem state changes and impacts of these changes on human welfare. Identifying the services and underlying process and examining ways of valuing ecosystem services helps provide an economic argument for conservation and can inform decisions on the cost and benefits of development options.

Methods to provide evidence of the economic importance of seagrass ecosystem services in relation to commercial and recreational fisheries have developed over the last 15 years (Scott et al. 2000; Rönnbäck et al. 2007; Unsworth et al. 2010), but they remain largely biased by a lack of consideration of temporary seagrass use within a species' lifecycle, a consideration of only commercial fishery landings values, or an over reliance on expert opinion (but see McArthur et al. 2003; Blandon & zu Ermgassen 2014). A specific seagrass meadow's importance for habitat provision depends on pre- and postsettlement processes (Jackson et al. 2001), the availability and suitability of other areas, and the preferences, ontogenetic habitat shifts, and survival rates of different species. Often no data on survival and recruitment to adult stocks exist or are difficult to measure in the field, while age at maturity and maximum age information is more easily accessed and habitat use for different species is available for many areas.

To address the effect of ontogenetic shifts in habitat association, Scott et al. (2000) formulated a seagrass residency index (SRI). They used residency as a surrogate for dependency and a weighted sum of the averages of the estimated residence time in seagrass (compared with other habitats) at each life stage. Scott et al. (2000) collated species life history information and habitat use from expert knowledge, which not only limited the number of species which could be incorporated in their analysis, but also limited the study to a small geographical region.

We sought to objectively assess the economic importance of seagrass as fish habitat. We used biological life

history trait databases and published reports of habitat preference to identify potential residency in seagrass and to apportion seagrass contribution to the fisheries stock. In addition to apportioning the direct commercial fishery landings values and recreational fishery expenditure associated with seagrass, we considered the implication of seagrass loss for regional fisheries policy.

Methods

Identifying Commercial and Recreationally Important Species

Species were identified as being commercially and recreationally targeted if they were listed on the General Fisheries Commission for the Mediterranean (GFCM) fisheries capture statistics (1950–2010) (Food and Agriculture Organisation 2012). All species names were checked, updated and, where necessary, amalgamated, based on the World Register of Marine Species (WoRMS) (www.marinespecies.org).

Calculating Seagrass Residency Indices

The extent of seagrass association varies among species. For example, black seabream (*Spondyllosoma cantharus*) is abundant on seagrass meadows, but it also occurs in a variety of other habitat types, whereas blackspot seabream (*Pagellus bogaraveo*) appears to be relatively ubiquitous. To differentiate habitat use, we calculated SRI (Scott et al. 2000) values for the relative amount of time a species spends in seagrass during different life stages relative to other habitats. If juvenile (highest mortality rate) survival is low, the species' contribution to fished stock will be low. Therefore, the amount of time spent in seagrass as a juvenile was also weighted relative to the adult stage. The final SRI was calculated as

$$\text{SRI}_i = 1 - \exp\{-[\exp(-m_i(t_{ji} - t_{ai}))x_i + y_i]\}, \quad (1)$$

where \exp is the exponential; for each species i , m is natural mortality; t_{ji} is the time (years) spent as a juvenile; t_{ai} is the time (years) spent as an adult; x_i is the proportion of time spent in seagrass as a juvenile; and y_i is the proportion of time spent in seagrass as an adult (Scott et al. 2000). The proportions of time spent in seagrass as an adult and as a juvenile were calculated as

$$x_i = \frac{(t_{ji}/t_i)}{(1/H_{ji})} \quad (2)$$

and

$$y_i = \frac{(t_{ai}/t_i)}{(1/H_{ai})}, \quad (3)$$

where t is the average maximum age, H_a is the number of habitats the species has been recorded as using (association) as an adult and H_j is the number of habitats

the species has been recorded as using (association) as a juvenile.

A systematic literature review was performed on habitat use (presence or absence) of commercially and recreationally important Mediterranean species. Where information was available, dominant or key habitats were noted, both for the species overall and for each life cycle stage. We used peer-reviewed literature, online databases, and factsheets (Fishbase, SeaLifebase, Larvalbase, Marine Species Information Portal, MarLIN, BIOTIC, Arkive, FAO factsheets, and FAO Adrimed). Key habitats were primarily derived from these widely acknowledged sources, unless specific mention was made of reliance on a particular habitat in the literature. Habitat association at each life stage for each species was ranked as 0 (no record found), 1 (rarely recorded or unusual), or 2 (consistently present in some abundance or many instances in the literature). This ranking was to illustrate confidence and was not used in the weighting calculations (Supporting Information).

For each species, we recorded the minimum, maximum, and mean duration of each life stage (juvenile and adult; years). These data were sourced from BIOTIC, Fishbase, and peer-reviewed literature. The juvenile stage included egg development time, larval phase duration, and juvenile stages up to sexual maturity or adulthood. For maximum adult age, the average maximum age was used where available, rather than the oldest on record, to prevent unrealistic skewing of results in fished populations. For sequential hermaphrodites the age of first becoming reproductively active was used; for gonochoristic species, with different male and female maturation ages, an average was taken (accounting for sex ratio, usually 1:1). We used Mediterranean data when possible and data from areas with similar water temperature otherwise. Where species groups (usually genus) were used, data were obtained to represent the most common species in the Mediterranean.

To calculate the weighting coefficient for the juvenile life stage, an estimate of natural mortality (m) was required. Some values for m are available in the literature (e.g., Pauly 1980); however, because temperature affects m , all scores were re-estimated using the life history tool in Fishbase (Froese et al. 2005). This tool applies the maximum length and water temperature as follows:

$$m = 10^{(0.566 - 0.718 * \log(L_{inf}))} + 0.02 * T, \quad (4)$$

where T is the mean annual water temperature; L_{inf} is the length that the fish within a population would reach if they were to grow indefinitely (also known as asymptotic length [Froese & Pauly 2000]). The mean Mediterranean Sea temperature (surface to 150 m) of 16.3 °C was taken from the annual average for 1960–2000 (Beuvier et al. 2010). Because many commercially valuable fish (particularly pelagic egg and larval stages) are found at shallower depths, where warmer temperatures allow for

faster growth, this is likely to be a conservative estimate. Asymptotic length (L_{inf}) was stated using accepted protocol for maximum length (i.e., fork length for scombroids [tuna and tuna-like fishes] and total length for all other fishes). Because we used length as a proxy for weight, natural mortality will be underestimated in eel-like fish and overestimated in sphere-shaped fish. Finally, age at first maturity (average age at which fish of a given population mature for the first time) was collated using the Fishbase life history tool or taken from peer reviewed publications (Supporting Information). Life history information and calculated SRI are shown in Table 1.

Commercial Fisheries Landings Values

To determine Mediterranean commercial fisheries landings value (CFV), data were extracted from the Commercial Harvest Database (Hutniczac & Roth 2012); this database evaluates European fishing in monetary terms. To generate commercial fishing landings values in the Mediterranean, time-series data for landings in Mediterranean fishing zones (2006–2008) were derived from the GFCM database (Food and Agriculture Organisation 2012). Landings data are recorded as live weight equivalent. Data for the price per ton (ungutted fish) for landings were derived from the European Union statistics portal, as estimated by the national authorities when reporting to the European Commission (European Commission 2012). Mediterranean landings of commercially important fish from 2006 to 2008 achieved an annual average first sale estimated value of €1.9 billion (SD €294.8 million) (Hutniczac & Roth 2012).

Recreational Fisheries Expenditure

Recreational fishing is defined here as “fishing activities exploiting marine living aquatic resources from which it is prohibited to sell or trade the catches obtained” (General Fisheries Commission for the Mediterranean 2010). For management purposes in the GFCM region, recreational fishing is divided into the following categories: leisure fishing, fishing practiced for pleasure; sport fishing, fishing contest practiced within an established institutional framework which sets rules, collects data on catches, and reveals to the public the outcomes of the event; charter fishing, recreational fishing practiced from a rented boat with a captain or fishing guide on board for leisure or sport purposes; and underwater fishing, fishing practiced as a sport or for leisure by snorkeling without the help of mechanical devices (e.g., a scooter).

Pesca turismo, or tourism fishing, is not included as a definition of recreational fishing because it is not solely geared toward recreational fishing (Gaudin & Young 2007; General Fisheries Commission for the Mediterranean 2012).

A variety of sources were used to determine Mediterranean recreational fishing expenditure. A literature search was conducted to identify data related to the recreational fishing activity of a Mediterranean country. Where primary studies were conducted, the study at the largest spatial scale (e.g., national level) was included for analysis. If a study included primary data on the number of recreational fishers (e.g., fishing licenses), then the broad expenditure per year for the number of fishers was derived from the regional economic expenditure for recreational fishers (demonstrated in Cisneros-Montemayor & Sumaila 2010). All recreational expenditure is expressed as expenditure per year by recreational fishers in a particular country. Types of expenditure included in each study varied, so expenditure could not be compared among countries unless reference was made to the original study. Annual expenditure of Mediterranean recreational fishers was therefore estimated at €2.5 billion. A full valuation table for recreation fishing in the Mediterranean is in the Supporting Information.

Application of SRI to Landings and Expenditure

To apportion the commercial CFVs and recreational fisheries expenditure to seagrass habitat provision service in the Mediterranean (CFV_{seagrass}), the Commercial Harvest Database Mediterranean section (Hutniczac & Roth 2012) was filtered to include those commercial fish species (icom in Eq. 5 associated with seagrass meadows [$n = 56$]). The average, minimum, and maximum annual landings values for the years 2006–2008 were calculated for each species. Commercial landings values related to pooled species (e.g., *Epinephelus* spp.) are included because the Commercial Harvest Database provides catch data on both individual species and broad species groups depending on the reporting requirements of the country where landings were made. We did not consider this double counting of commercial landings values because the fish is landed and categorized for reporting only once. If scores existed for broad species groups only, then we applied the SRI for the most abundant species in that group:

$$\text{CFV}_{\text{seagrass}} = \sum \text{CFV}_{\text{Mediterranean}}^{(\text{icom})} \times \text{SRI}^{(\text{icom})}. \quad (5)$$

To apply the SRI to recreation expenditure (RFV_{seagrass}), it was necessary to determine those species favored by recreational fishers. To provide a measure of the proportion of recreation expenditure to assign to fish species targeted by recreational fisheries, the mean catch composition of recreational fishers in Spain, France, and Italy was used as a proxy for the wider Mediterranean recreational fisheries (Gordoa et al. 2004). This demonstrated that 27.2% of the recreational catch was species associated with seagrass ($n = 7$, irec in Eq. 6).

Table 1. Organism life history traits and proportion of time spent in seagrass at different life stages used to calculate the seagrass residency index (SRI) (for details see Supporting Information).^a

Common name	Scientific name	tMat (years)	tMax (years)	m	x	y	tj	ta	SRI value
<i>Fish</i>									
axillary seabream	<i>Pagellus acarne</i>	2	9	1.35	0.042	0.506	-1.54	-9.00	0.397
black seabream	<i>Spondylosoma canthbarus</i>	4	15	1.46	0.052	0.153	-1.96	-10.00	0.142
blotched picarel	<i>Spicara maena</i>	2.5	9.1	0.98	0.055	0.041	-1.88	-9.10	0.040
bluefish	<i>Pomatomus saltatrix</i>	2	9	1.66	0	0.086	-0.65	-6.00	0.083
bogue	<i>Boops boops</i>	1	6	1.33	0	0.083	-1.37	-11.00	0.080
common dentex	<i>Dentex dentex</i>	1	28	1.63	0.008	0.121	-4.60	-26.00	0.114
common pandora	<i>Pagellus erythrinus</i>	1	20	1.33	0.079	0.407	-0.65	-20.00	0.334
dusky grouper	<i>Epinephelus marginatus</i>	5	26	1.72	0	0.114	-2.51	-15.00	0.108
european conger	<i>Conger conger</i>	5	15	1.94	0.037	0.074	-2.05	-6.00	0.071
european eel	<i>Anguilla anguilla</i>	8.9	41	0.14	0.035	0.041	-2.16	-10.10	0.041
European flounder	<i>Platichthys flesus</i>	3	15	1.40	0.034	0.110	-1.59	-14.00	0.104
European pilchard(= Sardine)	<i>Sardina pilchardus</i>	2.5	10	1.19	0.030	0.151	-0.15	-4.00	0.140
European seabass	<i>Dicentrarchus labrax</i>	2	14	1.58	0.023	0.105	-1.55	-10.00	0.099
garfish	<i>Belone belone</i>	2	8.4	1.50	0.059	0.190	-2.57	-12.00	0.173
gilthead seabream	<i>Sparus aurata</i>	1	12	1.50	0.019	0.063	-5.75	-41.04	0.061
grass goby	<i>Zosterisessor ophiocephalus</i>	4.3	16.6	0.38	0.122	0.297	-2.95	-16.61	0.257
pouting(= bib)	<i>Trisopterus luscus</i>	1	5	1.44	0	0.200	-0.67	-5.00	0.181
red mullet	<i>Mullus barbatus</i>	2.7	10.1	1.23	0	0.037	-2.51	-6.01	0.036
red porgy	<i>Pagrus pagrus</i>	3	12	1.60	0.061	0.188	-3.53	-15.00	0.171
saddled seabream	<i>Oblada melanura</i>	2	11	1.37	0	0.091	-0.35	-5.00	0.087
salema porgie	<i>Sarpa salpa</i>	1.8	11	1.44	0.031	0.075	-4.64	-15.00	0.073
sand streenbras	<i>Litobognathus mormyrus</i>	2.5	12	1.39	0	0.096	-1.54	-11.00	0.092
surmullet	<i>Mullus surmuletus</i>	1	8	1.29	0.013	0.252	-0.65	-8.00	0.238
white seabream	<i>Diplodus sargus</i>	2	10	1.42	0.010	0.100	-2.02	-12.00	0.095
whiting	<i>Merlangius merlangus</i>	1.5	10	1.41	0.092	0.121	-0.68	-28.00	0.114
<i>Arthropoda</i>									
caranote shrimp	<i>Melicerius kerathurus</i>	0.29	2.5	0.94	0.0232	0.221	-0.15	-2.56	0.200
common prawn	<i>Palaemon serratus</i>	0.58	5	1.50	0	0.089	-1.60	-9.00	0.085
spinous spider crab	<i>Maja squinado</i>	2.5	6	1.50	0	0.073	-0.67	-2.07	0.070
common shrimp	<i>Crangon crangon</i>	1	2	1.50	0	0.071	-0.67	-12.00	0.069
Mediterranean Green crab	<i>Carcinus aestuarii</i>	0.5	7	1.50	0	0.028	-0.30	-7.00	0.027
<i>Molluscs</i>									
common cuttlefish	<i>Sepia officinalis</i>	1	2	1.30	0.125	0.250	-1.57	-8.40	0.221
common octopus	<i>Octopus vulgaris</i>	0.28	4	1.31	0	0.145	-1.09	-10.00	0.135
Mediterranean mussel	<i>Mytilus galloprovincialis</i>	1	50	1.50	0	0	-2.56	-9.00	0.001

^aAbbreviations: tMat, age at maturity; tMax maximum age; m, mortality; x, proportion of time (years) spent in seagrass as a juvenile; y, the proportion of time (years) spent in seagrass as an adult.

Therefore, 27.2% of RFVMediterranean was carried forward for analysis. Where recreation expenditure is associated with broad species groups only, then the application of the SRI for the most abundant species on that group was applied:

$$\text{RFV seagrass} = \sum \text{RFV Mediterranean}^{(\text{irec})} \times \text{SRI}^{(\text{irec})}. \quad (6)$$

Results

The total annual landings value of commercial species that were, at some stage in their life history, associated with seagrass was approximately €681.2 million/year (35% of the total landings value of commercial fisheries in the Mediterranean [Table 2]). When the SRI scores were applied to determine the proportion of landings value directly attributable to the presence of seagrass, approximately €77.7 million (€58.3 million to €91.4 million; approximately 4% of the Mediterranean CFV) was directly associated with seagrass presence (Table 2). No Mediterranean fisheries' species was identified as being totally dependent on seagrass. Even the most strongly associated species, the axillary seabream (*Pagellus acarne*), showed some association with other habitats. From an ecological perspective, the commercial species that would be most affected by the loss of seagrass (highest SRI scores) are axillary seabream (*Pagellus acarne*), scorpionfishes and rockfishes (Scorpaenidae and Scorpaena), common pandora (*Pagellus erythrinus*), poor cod (*Trisopterus minutus*), caramote shrimp (*Melicertus kerathurus*), and great Atlantic scallop (*Pecten maximus*) (Table 1). From an economic perspective, the largest losses in landings if seagrass were to decline would be in commercial fisheries that target cuttlefish (Sepiidae, Sepiolidae), scorpionfishes (Scorpaenidae), octopuses (Octopodidae), European anchovy (*Engraulis encrasicolus*), and European pilchard (*Sardina pilchardus*). Despite having a minimal association with seagrass, the latter 2 species are so widely landed in the Mediterranean that the economic impact on the fisheries could be substantial.

The total annual expenditure of recreational fishers pursuing catches of species that, at some stage in their life history, depend on the presence of seagrass was estimated at €723.3 million (approximately 28% of all Mediterranean recreational fishing activity; Table 2). When the SRI scores were applied to determine the proportion of that expenditure that could be directly attributed to the presence of seagrass, €112.6 million of that expenditure (approximately 6% of RFVMediterranean) could be attributed the presence of seagrass (Table 2). In terms of impacts on the recreation industry, from an ecological perspective the species that would be most affected by the loss of seagrass are scorpionfishes and rockfish (Scorpaenidae, *Scorpaena* spp.), porgies, seabreams (Sparidae), cuttlefish (Sepiidae), garfish (*Belone belone*), and pouting (*Trisopterus luscus*)

(Table 1). From an economic perspective, the largest losses in associated expenditure if seagrass were to decline would be from the recreational fisheries that target seabass (*Dicentrarchus* spp.), cuttlefish (Sepiidae, Sepiolidae), and scorpionfishes (Scorpaenidae).

Discussion

By providing an essential habitat, we found that seagrass meadows had an estimated direct contribution of 4% to the total value of landings of commercial fisheries and 6% to the total expenditure of recreational fisheries, equating to a total of approximately €78 million and €112 million per year, respectively. The total area of *Posidonia oceanica* was estimated at about 5.5 million ha (Buia et al. 2000), or 2% of the surface area of the Mediterranean Sea; thus, this habitat disproportionately contributed to these Mediterranean fisheries in terms of "habitat service" (TEEB 2010). We found that seagrass had higher direct market value (in terms of landings values and expenditure) to recreational fisheries than to commercial fisheries, due to the different target species of the 2 fishing sectors and the different markets under which each sector operates. This highlights that studies which consider only the value of commercial fisheries landings will greatly underestimate the contribution of seagrass (or other habitats) to wider regional economic activity (e.g., sport fishing). Additionally, we demonstrated that any loss or degradation of seagrass is likely to have a disproportionate negative economic impact on fishery sectors with smaller inshore fisheries.

Previous studies assessing the economic importance of seagrass in relation to fishery species habitats focused on generic assessments which do not consider potential specificity or nonlinearity in service provision. For example, Rönnbäck et al. (2007) carried out a semiquantitative valuation of seagrass habitat secondary production by noting habitat association, market values, and landings of commercially fished species (see Rönnbäck et al. 2007). Although useful for assessing the potential implications of ecosystem state change on local markets, such an assessment does not consider the reliance of different life stages on different habitats, the effect of seagrass quality and density on fish production, or the indirect importance of species which, while not themselves commercially harvested, may be significant prey consumed by commercial species. The incorporation of seagrass quality and food web dynamics would greatly improve the accuracy of economic estimates, but this relies on the collation of extensive primary data. Such data on the reliance of different life stages on seagrass meadows are available in the literature, however, and provide a more realistic view of the contribution of seagrasses to commercial and recreationally important stocks. Without the application of the SRI (i.e., examining only habitat association of species [sensu Rönnbäck et al. 2007]), our data suggest

Table 2. Commercial and recreational fisheries value for species associated with seagrass habitat in the Mediterranean, their calculated seagrass residency index (SRI), and the proportion of value directly attributable to the presence of seagrass.^a

Common name	Scientific name	CFV ^b Med ^c €	SD	RFV Med €	SRI	CFV Seagrass €	RFV Seagrass €	CFV Seagrass low €	CFV Seagrass high €	
<i>Chordata</i>										
axillary seabream	<i>Pagellus acarne</i>	147,570	137,200		0.397	58,643		4,121	113,164	
black seabream	<i>Spondylitosa cantharus</i>	1,767,301	1,477,695		0.142	250,433		41,038	459,827	
blue whiting	<i>Micromesistius poutassou</i>	9,841,546	2,780,560	7,396,395	0.40×10^{-6}	4	3	3	5	
Bluefish	<i>Pomatomus saltatrix</i>	1,787,761	668,534	11,834,231	0.083	148,095	980,326	92,715	203,475	
Bogue	<i>Boops boops</i>	14,385,930	877,107		0.080	1,150,235		1,080,106	1,220,365	
Brill	<i>Scophthalmus rhombus</i>	388,178	46,240		0.49×10^{-5}	19		17	21	
chub mackerel	<i>Scomber japonicus</i>	4,865,828	4,359,487		0.000	193		20	367	
common dentex	<i>Dentex dentex</i>	5,416,204	4,708,978		0.114	615,035		80,309	1,149,761	
common pandora	<i>Pagellus erythrinus</i>	5,265,870	2,516,492		0.334	1,759,713		918,769	2,600,657	
conger eel	<i>Conger conger</i>	0	0	23,668,463	0.105	0	2,485,499	0	0	
dentex nei	<i>Dentex spp</i>	360,844	123,360		0.121	43,656		28,731	58,580	
dusky grouper	<i>Epinephelus marginatus</i>	2,912,992	1,168,600		0.108	314,594		188,389	440,799	
European anchovy	<i>Engraulis encrasicolus</i>	157,778,239	30,900,822		0.058	9,174,719		7,377,853	10,971,585	
European conger	<i>Conger conger</i>	2,652,773	1,423,742		0.071	189,400		87,749	291,051	
European eel	<i>Anguilla anguilla</i>	615,515	343,020		0.041	24,990		11,063	38,916	
European flounder	<i>Platichthys flesus</i>	597,160	558,014		0.104	62,024		4,066	119,982	
European pilchard	<i>Sardina pilchardus</i>	62,134,439	2,634,731		0.140	8,724,513		8,354,561	9,094,465	
European plaice	<i>Pleuronectes platessa</i>	4,123	3,007		Negligible					
European seabass	<i>Dicentrarchus labrax</i>	6,531,419	4,064,888	112,425,198	0.099	649,622	11,181,934	245,324	1,053,920	
Garfish	<i>Belone belone</i>	769,285	80,782	14,792,789	0.173	133,421	2,565,581	119,410	147,431	
gilthead seabream	<i>Sparus aurata</i>	12,177,915	6,505,716	124,259,429	0.061	742,647	7,577,725	345,908	1,139,386	
gobies nei	<i>Gobiidae</i>	2,254,728	1,226,544		0.257	580,293		264,621	895,965	
gray gurnard	<i>Eutrigla gurnardus</i>	6,429	6,114		0	0		0	0	
groupers, seabasses, nei	<i>Serranidae</i>	1,088,132	675,840	5,917,116	0.099	108,227	588,523	41,007	175,447	
gurnards, scarabins nei	<i>Triglidae</i>	11,694,972	1,128,523		0	0		0	0	
Mediterranean shore crab	<i>Carcinus aestuarii</i>	39,089	5,602		0.027	1,068	0	915	1,221	
picarels nei	<i>Spicara spp</i>	14,543,792	2,332,605	1,479,279	0.040	587,550	59,741	493,148	681,552	
poor cod	<i>Trisopterus minutus</i>	4,015,989	324,080		0	0		0	0	
porgies, seabreams	<i>Sparidae</i>	18,452	13,048	144,969,334	0.234	4,315	33,904,852	1,264	7,367	
Pouting	<i>Trisopterus luscus</i>	121,999	70,174	1,479,279	0.181	22,115	268,148	9,394	34,835	
red mullet	<i>Mullus barbatus</i>	61,239,560	6,182,432		0.036	2,203,010		1,980,606	2,425,415	
red porgy	<i>Pagrus pagrus</i>	4,266,635	2,160,089		0.171	729,470		360,158	1,098,783	
round sardinella	<i>Sardinella aurita</i>	1,757,707	1,522,274		0	0		0	0	
saddled seabream	<i>Oblada melanura</i>	3,648,798	3,244,191		0.087	317,078		35,160	598,996	
Salema	<i>Sarpa salpa</i>	1,734,583	1,616,959		0.073	126,050		8,548	243,553	
sand steenbras	<i>Litobognathus mormyrus</i>	4,212,103	3,066,262		0.092	387,042		105,289	668,795	
Sargo breams nei	<i>Diplodus spp</i>	7,332,731	6,398,666		0.095	697,802		88,888	1,306,716	
scorpionfishes nei	<i>Scorpaenidae</i>	14,250,854	9,686,110	41,419,810	0.337	4,799,688	13,950,192	1,537,406	8,061,970	

Continued

Table 2. Continued

Common name	Scientific name	CFV ^b Med ^c €	SD	RFV Med €	SRI	CFV Seagrass €	RFV Seagrass €	CFV Seagrass low €	CFV Seagrass high €
<i>Chordata</i>									
scorpionfishes rockfishes, nei	<i>Scorpaena spp</i>	187,896	5,327	17,751,347	0.337	63,284	5,978,654	61,489	65,078
seabasses nei	<i>Dicentrarchus spp</i>	3,677,300	1,186,949	112,425,198	0.099	365,748	11,181,933	247,693	483,803
<i>Anthropoda</i>									
caramote prawn	<i>Melicertus kerathurus</i>	32,103,164	10,760,179		0.20	642,063		426,860	857,267
common spiny lobster	<i>Palinurus elephas</i>	11,081,150	8,070,822		Negligible				
common prawn	<i>Palaemon serratus</i>	524,877	303,831		0.085	44,607		18,786	70,429
common shrimp	<i>Crangon crangon</i>	7,325,478	5,890,433		0.069	504,998		98,928	911,068
spinous spider crab	<i>Maja squinado</i>	40,785	15,972		0.070	2,868		1,745	3,991
surmullet	<i>Mullus surmuletus</i>	56,871,034	7,046,634		0.238	13,509,937		11,835,982	15,183,893
surmullets nei	<i>Mullus spp</i>	6,970,261	787,195	32,544,136	0.238	1,655,813	7,730,987	1,468,812	1,842,814
white seabream	<i>Diplodus sargus</i>	1,596,195	638,931		0.095	151,898		91,096	212,700
whiting	<i>Merlangius merlangus</i>	4,084,420	624,845		0.114	464,883		393,764	536,002
wrasses, hogfishes nei	<i>Labridae</i>	19,241	2,275	5,917,116	0.111	2,134	656,415	1,882	2,387
<i>Mollusca</i>									
common cuttlefish	<i>Sepia officinalis</i>	4,653,712	3,932,323		0.221	1,029,397		159,571	1,899,224
common octopus	<i>Octopus vulgaris</i>	20,831,324	18,341,194		0.135	2,811,764			
cuttlefish nei	<i>Sepiidae, Sepiolidae</i>	85,116,576	8,449,894	54,733,320	0.221	18,827,720	12,106,967	16,958,610	20,696,830
great Atlantic scallop	<i>Pecten maximus</i>	5,938	0	0	Negligible				
octopuses nei	<i>Octopodidae</i>	22,504,114	2,664,100	10,354,952	0.135	3,037,554	1,397,688	2,677,960	3,397,148
scallops nei	<i>Pectinidae</i>	982,259	974,137		0	0		0	0
Total		681,203,166		723,367,390		77,720,132	112,615,167	58,349,734	91,467,006

^aAbbreviations: CFV, commercial fisheries value; RFV, recreational fisheries value; Med, Mediterranean; nei, not elsewhere identified in Food and Agriculture Organization landings.

^bAveraged over 3 years (2006–2008) and the ranges of these and the corresponding CFV seagrass are presented.

^cThe 2006–2008 average value of landings.

that the economic importance of seagrasses would be overestimated 10-fold. Targeted studies such as ours are not reductionist and do not detract from the broader socioecological importance of seagrass protection. Rather, they provide realistic parameters and a strong socioeconomic argument for fisheries managers to influence seagrass protection strategies.

In addition to functioning as important nursery and foraging habitat for fish, shellfish (Jackson et al. 2001), and wildfowl (Ganter 2000), seagrasses are also thought to oxygenate and stabilize sediments, providing shoreline stabilization and erosion protection (Koch et al. 2009) and increasing potential economic value in terms of shoreline and property protection. Seagrasses are also considered a foundation species because they provide habitat and enhance ecosystem biodiversity and are carbon sequestration and nutrient cycling hubs (Kennedy et al. 2010). Habitat use represents only part of the secondary production importance of seagrasses; there is strong evidence that seagrass meadows have an important role in coastal primary production, for example. Rates of productivity for seagrass alone are large: *Posidonia oceanica* can fix 550–1000 g C m⁻²·yr⁻¹ (Mazzella et al. 1992). Cebrián et al. (1997) examined the fate of leaf and blade production of 4 Mediterranean seagrass species and found that >80% of seagrass production was consumed by herbivores or decomposed by detritivores, which may enhance food production for fishes.

The data we present here should be applied conservatively in terms of how commercial and recreational fisheries may be affected by habitat loss, most crucially because the study does not account for other nonlinearities in the provision of habitat for fishery species (i.e., species of seagrass, location, and the implications of degradation [Duarte 2000]). Variability in how commercial CFV and economic expenditure are spatially apportioned to seagrass meadows would not be surprising because biological resources and ecological services provided by seagrasses are based on plant physical structure and the large meadows they form (Boström et al. 2006).

The role of seagrass meadows as a nursery ground for many marine species, including those of commercial and recreational value, is well documented (Jackson et al. 2001; Heck et al. 2003), but evidence of this nursery function is often confined to studies which identify high juvenile density (Valle et al. 1999; Guidetti 2000). The question is whether seagrass meadows merely concentrate juveniles or whether the residents actually gain a selective advantage over individuals inhabiting other habitats (Beck et al. 2001). If the juveniles in seagrass meadows (or other habitats) fail to reach maturity, then these areas do not function as productive nurseries. Lacking habitat-specific survival data, we weighted habitats equally; a species found in many different habitats will have a lower SRI. Seagrasses may improve survival by providing shelter and food, however. They may also

promote planktonic larval settlement and, for those species that do not have a pelagic larval phase, may act directly as spawning areas (e.g., cuttlefish [*Sepia officinalis*] [Ezzedine-Najai et al. 1997]). It is therefore likely that any decline in seagrass quality or extent will affect the entire CFV Mediterranean (€681million) and RFV Mediterranean (€723 million). Consequently, due to the nature of fisheries in the Mediterranean, where inshore areas support a large population of artisanal and subsistence fishers, any socioeconomic impacts resulting from a decline in seagrass could have a disproportionate effect on small to medium size fishing operations that are the cultural backbone of many Mediterranean coastal communities. It is imperative that artisanal and subsistence fishers be included and properly accounted for in seagrass protection strategies (e.g., Marine Protected Areas, MPAs) (Rodriguez-Rodriguez 2014).

While limits on fishing with mobile gears exist in the Mediterranean (Vlachopoulou et al. 2013), in some areas certain species of fish and shellfish are still directly harvested from seagrass (Gonzalez-Correa et al. 2005; Cardona et al. 2007), and we incorporated these species in our study. These extractions can have negative impacts on the seagrass itself (Ardizzone et al. 2000) and nontarget species, including juvenile fishery species for which seagrass is nursery habitat, and therefore compromise future value to fisheries (Rees et al. 2010). In some areas where conservation of the seagrass meadows has resulted in shifts in fishing practice, fishing the edges of the meadows with static gear has become increasingly productive (Savage et al. 2012). Knowledge of the links between seagrass meadows and commercial CFVs and the expenditure of recreational fishers may, however, not affect wider policies (e.g., on climate change), but this type of information may be useful for local fisheries managers where trawling in seagrasses remains a direct impact (Vlachopoulou et al. 2013). While initial ecosystem service valuations have been carried out for Mediterranean *Posidonia oceanica* meadows in terms of carbon sequestration (Luisetti et al. 2013), attempts to quantify other ecosystem services are still needed in order to estimate the full economic value of seagrass meadows. Such information has significant implications for management and policy implementation, particularly in regards to the EU Marine Strategy Framework Directive, which does not link descriptors of good environmental status that are relevant to seagrass and seagrass ecosystem services. For example, descriptor 1 (biological diversity) will examine the state of seagrass meadows, but seagrass meadows are not considered in descriptor 3, population of commercial fish and shellfish, or descriptor 4, elements of marine food webs.

The MSFD explicitly requires an economic and sociological analysis of the use of coastal and marine waters to determine the “cost of degradation” of the marine environment. We have demonstrated that there is a clear

cost of degradation associated with ineffective management of seagrass meadows and that there is a potentially disproportionate impact of such management on small-scale artisanal fisheries that operate in inshore waters. European Union policy to manage both fisheries and seagrass meadows should align to take into account the socioeconomic implications of seagrass degradation on both recreational and commercial fisheries.

Acknowledgments

The research leading to these results received funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement number 226675, under the guidance and support of Professor Laurence Mee (1951-2014). We thank A. Ghermandi for information contributing to the economic valuation and M. Graziano for assisting with the collation of life history data.

Supporting Information

Life history data collated for the calculation of the SRI, including source information (Appendix S1), and a full valuation table for recreational fishing in the Mediterranean, including source information (Appendix S2), are available online. The authors are solely responsible for their content and functionality of these materials. Queries (other than the absence of material) should be directed to the corresponding author.

Literature Cited

Ardizzone GD, Tucci P, Somaschini A, Belluscio A. 2000. Is bottom trawling partly responsible for the regression of *Posidonia oceanica* meadows in the Mediterranean Sea? Pages 37-46 in Kaiser MJ, de Groot SJ, editors. The effect of fishing on non-target species and habitats: biological conservation and socio-economic issues. Blackwell, Oxford.

Beck MW, et al. 2001. The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates. *Bioscience* **51**:633-641.

Bell FW. 1997. The economic valuation of saltwater marsh supporting marine recreational fishing in the southeastern United States. *Ecological Economics* **21**:243-254.

Beuvier J, Sevault F, Herrmann M, Kontoyiannis H, Ludwig W, Rixen M, Stanev E, Béranger K, Somot S. 2010. Modeling the Mediterranean Sea interannual variability during 1961-2000: focus on the Eastern Mediterranean transient. *Journal of Geophysical Research: Oceans* (1978-2012) **115**:17.

Blandon A, zu Ermgassen PS. 2014. Quantitative estimate of commercial fish enhancement by seagrass habitat in southern Australia. *Estuarine, Coastal and Shelf Science* **141**:1-8.

Boström C, Jackson EL, Simenstad CA. 2006. Seagrass landscapes and their effects on associated fauna: a review. *Estuarine, Coastal and Shelf Science* **68**:383-403.

Buia MC, Gambi MC, Zupo V. 2000. Structure and functioning of Mediterranean seagrass ecosystems: an overview. *Biologia Marina Mediterranea* **7**:167-190.

Cardona L, López D, Sales M, de Caralt S, Díez I. 2007. Effects of recreational fishing on three fish species from the *Posidonia oceanica* meadows off Minorca (Balearic archipelago, western Mediterranean). *Scientia Marina* **71**:811-820.

CBD. 2000. Convention on Biological Diversity. Available from <http://69.90.183.227/doc/legal/cbd-un-en.pdf> (accessed March 12, 2011).

Cebrián J, Duarte C, Marbà N, Enriquez S. 1997. Magnitude and fate of the production of four co-occurring western Mediterranean seagrass species. *Marine Ecology Progress Series* **155**:29-44.

Cisneros-Montemayor A, Sumaila U. 2010. A global estimate of benefits from ecosystem-based marine recreation: potential impacts and implications for management. *Journal of Bioeconomics* **12**:245-268.

Dillon B. 2004. A bio-economic review of recreational angling for Bass (*Dicentrarchus labrax*). Scarborough Centre for Coastal Studies, University of Hull.

Duarte CM. 2000. Marine biodiversity and ecosystem services: an elusive link. *Journal of Experimental Marine Biology and Ecology* **250**:117-131.

European Commission. 2012. Eurostat; Available from http://epp.eurostat.ec.europa.eu/portal/page/portal/statistics/search_database (accessed February 6, 2012).

Ezzedine-Najai S, Ros J, Guerra A. 1997. Tagging of the cuttlefish, *Sepia officinalis* L. (Cephalopoda: Decapoda), in the Gulf of Tunis. *Scientia Marina* **61**:59-65.

Food and Agriculture Organisation. 2012. FishStat Plus - Universal software for fishery statistical time series. Fisheries and Aquaculture Department, Food and Agriculture Organisation of the United Nations, Rome

Froese R, Lourdes M, Palomares D, Pauly D. 2005. Estimation of life-history key facts. Available from <http://www.fishbase.org/manual/key%20facts.htm> (accessed February 22, 2012).

Froese R, Pauly D. 2000. FishBase 2000 Concepts, design and data sources: estimation of life-history key facts. Available from <http://www.fishbase.org/manual/key%20facts.htm> (accessed December 20, 2012).

Ganter B. 2000. Seagrass (*Zostera* spp.) as food for brent geese (*Branta bernicla*): an overview. *Helgoland Marine Research* **54**:63-70.

Gaudin C, Young CD. 2007. Recreational fisheries in the Mediterranean countries: a review of existing legal frameworks. Page 104 in General fisheries commission for the Mediterranean, editor. Studies in reviews. Food and Agriculture Organisation of the United Nations, Rome.

General Fisheries Commission for the Mediterranean. 2010. Report of the transversal workshop on monitoring of recreational fisheries in the GFCM area. Palma de Mallorca, Spain, 20-22nd October 2010. Page 30. General Fisheries Commission for the Mediterranean.

General Fisheries Commission for the Mediterranean. 2012. Report of the 12th session of the sub-committee on economic and social sciences (SCESS). Page 18. Food and Agriculture Organisation, Rome.

Gonzalez-Correa JM, Bayle JT, Sanchez-Lizasa JL, Valle C, Sanchez-Jerez P, Ruiz JM. 2005. Recovery of deep *Posidonia oceanica* meadows degraded by trawling. *Journal of Experimental Marine Biology and Ecology* **320**:65-76.

Gordoa A, Franquesa R, Calliart B, de la Serna J, Di Natale A, Ordan M. 2004. Sport fishing: an informative and economic alternative for tuna fishing in the Mediterranean (SFITUM). EC project 02/C132/11/41. Final Report. Green, E. P., F. T. Short 2003. World Atlas of Seagrass. UNEP WCMC, London.

Green EP, Short FT, Eds. 2003. World atlas of seagrasses. University of California Press, Berkeley.

Guidetti P. 2000. Differences among fish assemblages associated with nearshore *Posidonia oceanica* seagrass beds, rocky-algal reefs and unvegetated sand habitats in the Adriatic Sea. **50**:515-529.

- Heck KL, Hays G, Orth RJ. 2003. Critical evaluation of the nursery role hypothesis for seagrass meadows. *Marine Ecology-Progress Series* **253**:123–136.
- Hutniczac B, Roth E. 2012. Welfare benefits from fisheries in the ecosystem: commercial harvest database. Pages 1–6. Knowledge-based Sustainable Management for Europe's Seas (KnowSeas) Deliverable 4.2 Aggregate Assessment of Benefits and Costs. European Union Seventh Framework Programme.
- Jackson EL, Rowden AA, Attrill MJ, Bossey SJ, Jones MB. 2001. The importance of seagrass beds as a habitat for fishery species. *Oceanography and Marine Biology: An Annual Review* **39**:269–303.
- Kennedy H, Beggins J, Duarte CM, Fourquean JW, Holmer M, Marbá N, Middleburg JJ. 2010. Seagrass sediments as a global carbon sink: isotopic constraints. *Global Biogeochemical Cycles* **24**:1–8.
- Koch EW, et al. 2009. Non-linearity in ecosystem services: temporal and spatial variability in coastal protection. *Frontiers in Ecology and the Environment* **7**:29–37.
- Langmead O, McQuatters-Gollop A, Mee LD, editors. 2007. European lifestyles and marine ecosystems: exploring challenges for mapping Europe's seas. University of Plymouth Marine Institute, Plymouth.
- Luisetti T, Jackson EL, Turner RK. 2013. Valuing the European 'coastal blue carbon' storage benefit. *Marine Pollution Bulletin* **71**:101–106.
- Mazzella L, Buia MC, Gambi MC, Lorenti M, Russo GF, Scipione MB, Zupo V. 1992. Plant-animal trophic relationships in the *Posidonia oceanica* ecosystem of Mediterranean Sea: a review. Pages 165–187 in John DM, Hawkins SJ, Price JH, editors. Plant-animal interactions in the marine benthos. Clarendon Press, Oxford.
- McArthur LC, Boland JW, Brown B, Jones G. 2003. Investigation of the influence of seagrass on fisheries catch level using isotonic regression. *Environmental Modeling & Assessment* **8**:285–290.
- Pauly D. 1980. On the interrelationships between natural mortality, growth parameters, mean environmental temperature in 175 fish stocks. *Journal du Conseil* **39**:175–192.
- Pauly D. 2009. Beyond duplicity and ignorance in global fisheries. *Scientia Marina* **73**:215–224.
- Pawson MG, Tingley DPG, Glenn H. 2007. EU contract FISH/2004/011 on Sport Fisheries (or Marine Recreational Fisheries) in the EU. Prepared for for The European Commission Directorate-General for Fisheries. Page 242. The Centre for Environment, Fisheries and Aquaculture Science (Cefas).
- Rätz H-J, Dörner H, Scott R, Barbas T. 2010. Complementary roles of European and national institutions under the Common Fisheries Policy and the Marine Strategy Framework Directive. *Marine Policy* **34**:1028–1035.
- Rees SE, Rodwell LD, Attrill MJ, Austen MC, Mangi SC. 2010. The value of marine biodiversity to the leisure and recreation industry and its application to marine spatial planning. *Marine Policy* **34**:868–875.
- Rodriguez-Rodriguez D. 2014. Artisanal fisheries in the Mediterranean. Med PAN, Marseille, France.
- Rönnbäck P, Kautsky N, Pihl L, Troell M, Söderqvist T, Wennhage H. 2007. Ecosystem goods and services from Swedish coastal habitats: identification, valuation, implications of ecosystem shifts. *AMBIO: A Journal of the Human Environment* **36**:534–544.
- Sacchi J. 2011. Analysis of economic activities in the Mediterranean: fishery and aquaculture sectors. Page 84. Plan Bleu UNEP/MAP Regional Activity Centre, Valbonne.
- Savage C, Thrush SF, Lohrer AM, Hewitt JE. 2012. Ecosystem services transcend boundaries: estuaries provide resource subsidies and influence functional diversity in coastal Benthic communities. *PLoS One* **7** DOI: 10.1371/journal.pone.0042708.
- Scott LC, Boland JW, Edyvane KS, Jones G. 2000. Development of a seagrass-fish habitat model—I: a seagrass residency index for economically important species. *Environmetrics* **11**:541–552.
- TEEB. 2010. The economics of ecosystems and biodiversity: mainstreaming the economics of nature: a synthesis of the approach, conclusions and recommendations of TEEB. United Nations Environment Programme and the European Commission, Progress Press, Malta, pp. 39.
- Toivonen AL, Roth E, Navrud S, Gudbergsson G, Appelblad H, Bengtsson B, Tuunainen P. 2004. The economic value of recreational fisheries in Nordic countries. *Fisheries Management and Ecology* **11**: 1–14.
- Unsworth RK, Cullen LC, Pretty JN, Smith DJ, Bell JJ. 2010. Economic and subsistence values of the standing stocks of seagrass fisheries: potential benefits of no-fishing marine protected area management. *Ocean & Coastal Management* **53**:218–224.
- Valle CF, O'Brien JW, Wiese KB. 1999. Differential habitat use by *California halibut*, *Paralichthys californicus*, barred sand bass, *Paralabrax nebulifer*, and other juvenile fishes in Alamos Bay, California. *Fishery Bulletin* **97**:646–660.
- Vizzini S, Sarà G, Michener RH, Mazzola A. 2002. The role and contribution of the seagrass *Posidonia oceanica* (L.) Delile organic matter for secondary consumers as revealed by carbon and nitrogen stable isotope analysis. *Acta Oecologica* **23**:277–285.
- Vlachopoulou EI, Wilson A, Miliou A. 2013. Disconnects in EU and Greek fishery policies and practices in the eastern Aegean Sea and impacts on *Posidonia oceanica* meadows. *Ocean & Coastal Management* **76**:105–113.

