The Open University

Open Research Online

The Open University's repository of research publications and other research outputs

The effect of a cold atmospheric pressure plasma jet on DNA

Conference or Workshop Item

How to cite:

Ptasińska, Sylwia; Bahnev, Blagovest; Stypczynska, Agnieszka; Bowden, Mark; Braithwaite, Nicholas St.J. and Mason, Nigel J. (2009). The effect of a cold atmospheric pressure plasma jet on DNA. In: XXVI International Conference on Photonic, Electronic and Atomic Collisions, Journal of Physics: Conference Series, 22-28 Jul 2009, Kalamazoo, Michigan, USA.

For guidance on citations see \underline{FAQs} .

 \odot 2009 IOP Publishing Ltd

Version: Version of Record

Link(s) to article on publisher's website: http://dx.doi.org/doi:10.1088/1742-6596/194/15/152029

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data <u>policy</u> on reuse of materials please consult the policies page.

oro.open.ac.uk



The effect of a cold atmospheric pressure plasma jet on DNA

This article has been downloaded from IOPscience. Please scroll down to see the full text article. 2009 J. Phys.: Conf. Ser. 194 152029 (http://iopscience.iop.org/1742-6596/194/15/152029) View the table of contents for this issue, or go to the journal homepage for more

Download details: IP Address: 94.12.107.211 The article was downloaded on 20/10/2010 at 12:07

Please note that terms and conditions apply.

Journal of Physics: Conference Series 194 (2009) 152029

IOP Publishing

doi:10.1088/1742-6596/194/15/152029

The effect of a cold atmospheric pressure plasma jet on DNA

Sylwia Ptasińska¹, Blagovest Bahnev, Agnieszka Stypczyńska, Mark Bowden, Nicholas St. J. Braithwaite and Nigel J. Mason

Department of Physics & Astronomy, The Open University, Milton Keynes, MK7 6AA, United Kingdom

Synopsis In this study, we have investigated the interaction of a 'cold' Atmospheric Pressure Plasma Jet (APPJ) with dry plasmid DNA. The plasmid DNA is observed to be very sensitive to short-term exposures to low energy electrons, ions and excited neutral species as well as to UV light, which are all components of the APPJ fed by helium. In order to determine the mechanisms that lead to strand breaks in the DNA, the APPJ exposure was carried out over a range of operating conditions.

Recently there has been increased interest in the use of non-thermal atmospheric pressure plasmas (see Fig.1) as a tool in medical applications [1] e.g. in surgery and dentistry. However, most of these applications are still in an early stage of their development mainly due to the lack of fundamental knowledge of plasma interactions with living tissue at both a macroand a microscopic level.




Figure. 1. The experimental set-up.

In order to obtain a better understanding of the effect of 'cold' plasma on bio-molecules, we have exposed plasmid DNA to a helium plasma generated using a non-thermal atmospheric pressure plasma jet. Damage to the DNA induced by plasma exposure was determined by gel electrophoresis for different irradiation times. The amount of undamaged DNA molecules decreased by 30-60% after 10 sec of irradiation (see Fig.2).

The distance between the tip of the visible plasma jet and the DNA was varied (see Fig.2). It was found that initially damage fell off rapidly with distance but damage was still observed more than 10 cm from the jet. We have also explored damage as a function of helium flow rate and the amount of UV light.




Figure. 2. The percentage amount of undamaged plasmid DNA as a function of irradiation time (closed circles) and distance of a sample from the visible plasma (open circles).

Recently we have started studies on effect of water and proteins on DNA damage using plasma exposure. We expect that the presence of water will increase the damage to DNA due to the formation of highly reactive radicals during the APPJ interaction. In contrast the presence of proteins around DNA molecules may shield them from the plasma-generated species and so reduce the amount of damage. Results will be reported at the conference.

References

[1] Fridman G, Friedman G, Gustol A, Shekhter A B, Vasilets V N, Fridman A 2008 *Plasma Process*. *Polim.* **5** 503-533

¹E-mail: s.ptasinska@open.ac.uk