



Open Research Online

The Open University's repository of research publications and other research outputs

Well-Quasi-Order for Permutation Graphs Omitting a Path and a Clique

Journal Item

How to cite:

Atminas, Aistis; Brignall, Robert; Korpelainen, Nicholas; Lozin, Vadim and Vatter, Vincent (2015). Well-Quasi-Order for Permutation Graphs Omitting a Path and a Clique. *Electronic Journal of Combinatorics*, 22(2), article no. P2.20.

For guidance on citations see [FAQs](#).

© 2015 The Authors

Version: Version of Record

Link(s) to article on publisher's website:

<http://www.combinatorics.org/ojs/index.php/eljc/article/view/v22i2p20>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Well-Quasi-Order for Permutation Graphs Omitting a Path and a Clique

Aistis Atminas¹

DIMAP and Mathematics Institute
University of Warwick, Coventry, UK

a.atminas@warwick.ac.uk

Robert Brignall²

Department of Mathematics and Statistics
The Open University, Milton Keynes, UK

r.brignall@open.ac.uk

Nicholas Korpelainen²,

Mathematics Department
University of Derby, Derby, UK

n.korpelainen@derby.ac.uk

Vadim Lozin^{1,3}

DIMAP and Mathematics Institute
University of Warwick, Coventry, UK

v.lozin@warwick.ac.uk

Vincent Vatter^{2,4}

Department of Mathematics
University of Florida, Gainesville, Florida, USA

vatter@ufl.edu

Submitted: Jan 31, 2014; Accepted: Apr 2, 2015; Published: XX

Mathematics Subject Classifications: 05A05, 05A15

Abstract

We consider well-quasi-order for classes of permutation graphs which omit both a path and a clique. Our principle result is that the class of permutation graphs omitting P_5 and a clique of any size is well-quasi-ordered. This is proved by giving a structural decomposition of the corresponding permutations. We also exhibit three infinite antichains to show that the classes of permutation graphs omitting $\{P_6, K_6\}$, $\{P_7, K_5\}$, and $\{P_8, K_4\}$ are not well-quasi-ordered.

¹Atminas and Lozin gratefully acknowledge support from DIMAP – the Center for Discrete Mathematics and its Applications at the University of Warwick.

²Brignall, Korpelainen, and Vatter were partially supported by EPSRC Grant EP/J006130/1.

³Lozin was partially supported by EPSRC Grants EP/I01795X/1 and EP/L020408/1.

⁴Vatter was partially supported by the National Security Agency under Grant Number H98230-12-1-0207 and the National Science Foundation under Grant Number DMS-1301692. The United States Government is authorized to reproduce and distribute reprints not-withstanding any copyright notation herein.

1 Introduction

While the Minor Theorem of Robertson and Seymour [17] shows that the set of all graphs is well-quasi-ordered under the minor relation, it is well known that this set is not well-quasi-ordered under the induced subgraph order. Consequently, there has been considerable interest in determining which classes of graphs are well-quasi-ordered under this order. Here we consider finite graphs and permutation graphs which omit both a path P_k and a clique K_ℓ . Our main result, proved in Section 4, is that permutation graphs which avoid both P_5 and a clique K_ℓ are well-quasi-ordered under the induced subgraph order for every finite ℓ . We also prove, in Section 5, that the three classes of permutation graphs defined by forbidding $\{P_6, K_6\}$, $\{P_7, K_5\}$ and $\{P_8, K_4\}$ respectively are not well-quasi-ordered, by exhibiting an infinite antichain in each case.

We begin with elementary definitions. We say that a *class* of graphs is a set of graphs closed under isomorphism and taking induced subgraphs. A class of graphs is *well-quasi-ordered* (*wqo*) if it contains neither an infinite strictly decreasing sequence nor an infinite antichain (a set of pairwise incomparable graphs) under the induced subgraph ordering. Note that as we are interested only in classes of finite graphs, wqo is synonymous with a lack of infinite antichains.

Given a permutation $\pi = \pi(1) \cdots \pi(n)$, its corresponding *permutation graph* is the graph G_π on the vertices $\{1, \dots, n\}$ in which i is adjacent to j if both $i \leq j$ and $\pi(i) \geq \pi(j)$. This mapping is many-to-one, because, for example, $G_{231} \cong G_{312} \cong P_3$. Given permutations $\sigma = \sigma(1) \cdots \sigma(k)$ and $\pi = \pi(1) \cdots \pi(n)$, we say that σ is *contained* in π and write $\sigma \leq \pi$ if there are indices $1 \leq i_1 < \cdots < i_k \leq n$ such that the sequence $\pi(i_1) \cdots \pi(i_k)$ is in the same relative order as σ . For us, a *class* of permutations is a set of permutations closed downward under this containment order, and a class is wqo if it does not contain an infinite antichain.

The mapping $\pi \mapsto G_\pi$ is easily seen to be *order-preserving*, i.e., if $\sigma \leq \pi$ then G_σ is an induced subgraph of G_π . Therefore if a class \mathcal{C} of permutations is wqo then the associated class of permutation graphs $\{G_\pi : \pi \in \mathcal{C}\}$ must also be wqo. However, it is possible to exploit the many-to-one nature of this mapping to construct infinite antichains of permutations which do not correspond to antichains of permutation graphs (though currently there are no examples of this in the literature). For this reason, when showing that classes of permutation graphs are wqo we instead prove the stronger result that the associated permutation classes are wqo, but when constructing infinite antichains, we must construct antichains of permutation graphs.

A summary of our results is shown in Figure 1, with our new contributions in the upper-right highlighted. The rest of this paper is organised as follows: in Section 2 we briefly summarise the status of the analogous question for non-permutation graphs. Section 3 sets up the necessary notions from the study of permutation classes, before the proof of the well-quasi-orderability of P_5 , K_ℓ -free permutation graphs in Section 4. Section 5 contains three non-wqo results, Section 6 briefly presents some enumerative consequences of our results, and the final section contains a few concluding remarks about the three remaining open cases.




Figure 1: This figure shows the known wqo results for classes of graphs and permutation graphs avoiding paths and cliques, including the results of this paper. Filled circles indicate that all graphs avoiding the specified path and clique are wqo. Half-filled circles indicate that the corresponding class of permutation graphs are wqo, but that the corresponding class of all graphs are not wqo. Empty circles indicate that neither class is wqo. Note that for the three unknown cases (indicated by question marks), it is known that the corresponding class of graphs contains an infinite antichain.

2 Non-Permutation Graphs

When the graphs needn't be permutation graphs, well-quasi-ordering is of course harder to attain. On the side of wqo, graphs avoiding K_2 are trivially wqo and graphs avoiding P_4 (co-graphs) are well-known to be wqo, so there are only two nontrivial results, namely graphs avoiding K_3 and P_5 or P_6 . Of course, it suffices to show that K_3, P_6 -free graphs are wqo, and this was recently proved by Atminas and Lozin [6].

On the side of non-wqo, two infinite antichains are required: one (from [13]) in the class of graphs with neither P_5 (or even $2K_2$) nor K_4 , and one (from [14]) in the class omitting both P_7 and K_3 . For completeness, we outline both constructions here.

For graphs which omit P_5 and K_4 , Korpelainen and Lozin [13] construct an infinite antichain by adapting a correspondence between permutations and graphs due to Ding [8]. The correspondence we require for our antichain can be described as follows, and is accompanied by Figure 2. For a permutation π of length n , first note that π can be thought of as a structure with n points, equipped with two linear orderings.

With this in mind, form a graph B_π which consists of three independent sets, U , V and W , each containing n vertices. Let $U = \{u_1, u_2, \dots, u_n\}$. Between U and V , there is a *chain graph*: vertex u_i in U is adjacent to i vertices of V , and for $i > 1$ the neighbourhood of u_{i-1} in V is contained in the neighbourhood of u_i in V . Note that this containment of neighbourhoods defines a linear ordering on the vertices of U : $u_1 < u_2 < \dots < u_n$.

Next, between U and W , we build another chain graph. This time, vertex $u_{\pi(i)}$




Figure 2: The construction of B_π from a permutation π .

has i neighbours in W , and for $i > 1$ the neighbourhood of $u_{\pi(i)}$ in W contains the neighbourhood of $u_{\pi(i-1)}$ in W . This defines a second linear ordering on U , namely $u_{\pi(1)} \prec u_{\pi(2)} \prec \dots \prec u_{\pi(n)}$, and hence π has been encoded in B_π . Finally, to complete the construction, between V and W there is a complete bipartite graph, i.e., every edge is present.

Now it is routine to verify that B_π does not contain $2K_2 \leq P_5$ or K_4 for any π . Moreover for permutations σ and π , we have $\sigma \leq \pi$ if and only if $B_\sigma \leq B_\pi$ as induced subgraphs. Thus, one may take any infinite antichain of permutations (for example, the “increasing oscillating” antichain), and encode each element of the antichain as a graph, yielding an infinite antichain in the class of $2K_2, K_4$ -free graphs.

For graphs which omit P_7 and K_3 , a modification of this construction was used by [14] to answer a question of Ding [8], who had asked whether P_7 -free bipartite graphs are wqo. Starting with B_π , the modification “splits” every vertex u of U into two, $u^{(1)}$ and $u^{(2)}$, with an edge between them. The new vertex $u^{(1)}$ takes the neighbourhood of u with V , while $u^{(2)}$ takes the neighbourhood of u with W . The result is a graph with four independent sets $U^{(1)}, U^{(2)}, V$ and W each of size n , with a perfect matching between $U^{(1)}$ and $U^{(2)}$, a complete bipartite graph between V and W , and chain graphs between $U^{(1)}$ and V , and between $U^{(2)}$ and W .

This construction yields a $2P_3$ -free bipartite graph. Moreover, the permutation π is still encoded in such a way as to ensure an infinite antichain of permutations maps to an infinite antichain of graphs.

3 Structural Tools

Instead of working directly with permutation graphs, we establish our wqo results for the corresponding permutation classes (which, by our observations in the introduction, is a stronger result). The permutations 24153 and 31524 are the only permutations which correspond to the permutation graph P_5 , and thus to establish our main result we must determine the structure of the permutation class

$$\text{Av}(24153, 31524, \ell \cdots 21).$$

Considering the wqo problem from the viewpoint of permutations has the added benefit of allowing us to make use of the recently developed tools in this field. In particular, we

utilise grid classes and the substitution decomposition. Thus before establishing our main result we must first introduce these concepts.

We frequently identify a permutation π of length n with its *plot*, the set $\{(i, \pi(i)) : 1 \leq i \leq n\}$ of points in the plane. We say that a rectangle in the plane is *axis-parallel* if its top and bottom sides are parallel with the x -axis while its left and right sides are parallel with the y -axis. Given natural numbers i and j we denote by $[i, j]$ the closed interval $\{i, i + 1, \dots, j\}$ and by $[i, j)$ the half-open interval $\{i, i + 1, \dots, j - 1\}$. Thus the axis-parallel rectangles we are interested in may be described by $[x, x'] \times [y, y']$ for natural numbers x, x', y , and y' .

Monotone grid classes are a way of partitioning the entries of a permutation (or rather, its plot) into monotone axis-parallel rectangles in a manner specified by a $0/\pm 1$ matrix. In order for these matrices to align with plots of permutations, we index them with Cartesian coordinates. Suppose that M is a $t \times u$ matrix (thus M has t columns and u rows). An M -gridding of the permutation π of length n consists of a pair of sequences $1 = c_1 \leq \dots \leq c_{t+1} = n + 1$ and $1 = r_1 \leq \dots \leq r_{u+1} = n + 1$ such that for all k and ℓ , the entries of (the plot of) π that lie in the axis-parallel rectangle $[c_k, c_{k+1}) \times [r_\ell, r_{\ell+1})$ are increasing if $M_{k,\ell} = 1$, decreasing if $M_{k,\ell} = -1$, or empty if $M_{k,\ell} = 0$.

We say that the permutation π is M -griddable if it possesses an M -gridding, and the *grid class* of M , denoted by $\text{Grid}(M)$, consists of the set of M -griddable permutations. We further say that the permutation class \mathcal{C} is M -griddable if $\mathcal{C} \subseteq \text{Grid}(M)$, and that this class is *monotone griddable* if there is a finite matrix M for which it is M -griddable.

Grid classes were first described in this generality (albeit under a different name) by Murphy and Vatter [16], who studied their wqo properties. To describe their result we need the notion of the *cell graph* of a matrix M . This graph has vertex set $\{(i, j) : M_{i,j} \neq 0\}$ and (i, j) is adjacent to (k, ℓ) if they lie in the same row or column and there are no nonzero entries lying between them in this row or column. We typically attribute properties of the cell graph of M to M itself; thus we say that M is a forest if its cell graph is a forest.

Theorem 3.1 (Murphy and Vatter [16]). *The grid class $\text{Grid}(M)$ is wqo if and only if M is a forest.*

(This is a slightly different form of the result than is stated in [16], but the two forms are equivalent as shown by Vatter and Waton [19], who also gave a much simpler proof of Theorem 3.1.)

The monotone griddable classes were characterised by Huczynska and Vatter [12]. In order to present this result we also need some notation. Given permutations σ and τ of respective lengths m and n , their *sum* is the permutation $\pi \oplus \tau$ whose plot consists of the plot of τ above and to the right of the plot of σ . More formally, this permutation is defined by

$$\sigma \oplus \tau(i) = \begin{cases} \sigma(i) & \text{for } 1 \leq i \leq m, \\ \tau(i - m) + m & \text{for } m + 1 \leq i \leq m + n. \end{cases}$$

The obvious symmetry of this operation (in which the plot of τ lies below and to the right of the plot of σ) is called the *skew sum* of σ and τ and is denoted $\sigma \ominus \tau$. We can now state the characterisation of monotone griddable permutation classes.

Theorem 3.2 (Huczynska and Vatter [12]). *The permutation class \mathcal{C} is monotone griddable if and only if it does not contain arbitrarily long sums of 21 or skew sums of 12.*

The reader might note that the classes we are interested in are not monotone griddable, let alone M -griddable for a forest M . However, our proof will show that these classes can be built from a monotone griddable class via the “substitution decomposition”, which we define shortly. Before this, though, we must introduce a few more concepts concerning monotone grid classes, the first two of which are alternative characterisations of monotone griddable classes.

Given a permutation π , we say that the axis-parallel rectangle R is *monotone* if the entries of π which lie in R are monotone increasing or decreasing (otherwise R is *non-monotone*). We say that the permutation π can be *covered by s monotone rectangles* if there is a collection \mathfrak{R} of s monotone axis-parallel rectangles such that every point in the plot of π lies in at least one rectangle in \mathfrak{R} . Clearly if $\mathcal{C} \subseteq \text{Grid}(M)$ for a $t \times u$ matrix M then every permutation in \mathcal{C} can be covered by tu monotone rectangles. To see the other direction, note that every permutation which can be covered by s monotone rectangles is M -griddable for some matrix M of size at most $(2s - 1) \times (2s - 1)$. There are only finitely many such matrices, say $M^{(1)}, \dots, M^{(m)}$, so their *direct sum*,

$$\left(\begin{array}{cccc} & & & M^{(m)} \\ & & & \\ & & \ddots & \\ M^{(1)} & & & \end{array} \right)$$

is a finite matrix whose grid class contains all such permutations.¹

This characterisation of monotone griddability is recorded in Proposition 3.3 below, which also includes a third characterisation. We say that the line L *slices* the rectangle R if $L \cap R \neq \emptyset$. If $\mathcal{C} \subseteq \text{Grid}(M)$ for a $t \times u$ matrix then for every permutation $\pi \in \mathcal{C}$ there is a collection of $t + u$ horizontal and vertical lines (the grid lines) which slice every non-monotone axis-parallel rectangle of π . Conversely, every such collection of lines defines a gridding of π , completing the sketch of the proof of the following result.

Proposition 3.3 (specialising Vatter [18, Proposition 2.3]). *For a permutation class \mathcal{C} , the following are equivalent:*

- (1) \mathcal{C} is monotone griddable,
- (2) there is a constant ℓ such that for every permutation $\pi \in \mathcal{C}$ the set of non-monotone axis-parallel rectangles of π can be sliced by a collection of ℓ horizontal and vertical lines, and
- (3) there is a constant s such that every permutation in \mathcal{C} can be covered by s monotone rectangles.

¹By adapting this argument it follows that if every permutation in the class \mathcal{C} lies in the grid class of a forest of size at most $t \times u$, then \mathcal{C} itself lies in the grid class of a (possibly much larger) forest.

We now move on to the substitution decomposition, which will allow us to build the classes we are interested in from grid classes of forests. An *interval* in the permutation π is a set of contiguous indices $I = [a, b]$ such that the set of values $\pi(I) = \{\pi(i) : i \in I\}$ is also contiguous. Given a permutation σ of length m and nonempty permutations $\alpha_1, \dots, \alpha_m$, the *inflation* of σ by $\alpha_1, \dots, \alpha_m$ — denoted $\sigma[\alpha_1, \dots, \alpha_m]$ — is the permutation of length $|\alpha_1| + \dots + |\alpha_m|$ obtained by replacing each entry $\sigma(i)$ by an interval that is order isomorphic to α_i in such a way that the intervals themselves are order isomorphic to σ . Thus the sum and skew sum operations are particular cases of inflations: $\sigma \oplus \tau = 12[\sigma, \tau]$ and $\sigma \ominus \tau = 21[\sigma, \tau]$. Given two classes \mathcal{C} and \mathcal{U} , the *inflation* of \mathcal{C} by \mathcal{U} is defined as

$$\mathcal{C}[\mathcal{U}] = \{\sigma[\alpha_1, \dots, \alpha_m] : \sigma \in \mathcal{C}_m \text{ and } \alpha_1, \dots, \alpha_m \in \mathcal{U}\}.$$

The class \mathcal{C} is said to be *substitution closed* if $\mathcal{C}[\mathcal{C}] = \mathcal{C}$. The *substitution closure*, $\langle \mathcal{C} \rangle$, of a class \mathcal{C} is defined as the smallest substitution closed class containing \mathcal{C} . A standard argument shows that $\langle \mathcal{C} \rangle$ exists, and by specialising a result of [4] we obtain the following.

Theorem 3.4 (Albert, Ruškuc, and Vatter [4, specialisation of Theorem 4.4]). *If the matrix M is a forest then the class $\langle \text{Grid}(M) \rangle$ is wqo.*

The permutation π is said to be *simple* if the only ways to write it as an inflation are trivial — that is, it can only be written either as the inflation of a permutation by singletons, or as the inflation of a singleton by a permutation. Thus if π has length n , it is simple if and only if its only intervals have length 0, 1, and n . Thus every permutation can be expressed as the inflation of a simple permutation. Moreover, in most cases, this decomposition is unique. A permutation is said to be *sum (resp., skew) decomposable* if it can be expressed as the sum (resp., skew sum) of two shorter permutations. Otherwise it is said to be *sum (resp., skew) indecomposable*.

Proposition 3.5 (Albert and Atkinson [1]). *Every permutation π except 1 is the inflation of a unique simple permutation σ . Moreover, if $\pi = \sigma[\alpha_1, \dots, \alpha_m]$ for a simple permutation σ of length $m \geq 4$, then each interval α_i is unique. If π is an inflation of 12 (i.e., is sum decomposable), then there is a unique sum indecomposable α_1 such that $\pi = \alpha_1 \oplus \alpha_2$. The same holds, mutatis mutandis, with 12 replaced by 21 and sum replaced by skew.*

We close this section by noting how easily this machinery can show that permutation graphs omitting both P_k and K_3 are wqo for all k , a result originally due to Korpelainen and Lozin [14]. The corresponding permutations all avoid 321 and thus lie in the grid class of an *infinite* matrix, known as the infinite staircase (see Albert and Vatter [5]). Moreover, the sum indecomposable permutations which avoid 321 and the two permutations corresponding to P_k can be shown to lie in the grid class of a finite staircase,

$$\begin{pmatrix} & & & & 1 & 1 \\ & & & & & \\ & & \ddots & \ddots & & \\ & 1 & 1 & & & \\ 1 & 1 & & & & \end{pmatrix}.$$




Figure 3: The impossible configuration for a simple permutation in Proposition 4.1.

Finally, it follows by an easy application of Higman’s Theorem [11] that if the sum indecomposable permutations in a class are wqo, then the class itself is wqo. (In this case, the wqo conclusion also follows by Theorem 3.4.)

4 Permutation Graphs Omitting P_5 and K_ℓ

In this section we prove that the class of permutations corresponding to permutation graphs omitting P_5 and K_ℓ ,

$$\text{Av}(24153, 31524, \ell \cdots 21),$$

is wqo. Our proof basically consists of two steps. First, we show that the simple permutations in these classes are monotone griddable, and then we show that these griddings can be refined to forests. The conclusion then follows from Theorem 3.4.

Given a set of points in the plane, their *rectangular hull* is defined to be the smallest axis-parallel rectangle containing all of them. We begin with a very simple observation about these simple permutations.

Proposition 4.1. *For every simple permutation $\pi \in \text{Av}(24153, 31524)$, either its greatest entry lies to the left of its least entry, or its leftmost entry lies above its rightmost entry.*

Proof. Suppose, for a contradiction, that π is a simple permutation in $\text{Av}(24153, 31524)$ such that its greatest entry lies to the right of its least, and its leftmost entry lies below its rightmost entry. Thus, these four extremal entries form the pattern 2143, and the situation is as depicted in Figure 3(i). Since π is simple, regions A and B cannot both be empty, so, without loss of generality, suppose that A is non-empty and label the greatest entry in this region as point x .

Since π is simple, the rectangular hull of the leftmost entry, the least entry, and the point x cannot be an interval in π . Therefore, there must be a point either in B , or in that part of C lying below x . Take the rightmost such point, and label it y . If y is in region C , we immediately encounter the contradiction illustrated in Figure 3(ii): our choices of x and y , and the forbidden permutation 24153 causes the permutation to be sum decomposable. Therefore, y is placed in region B , and we have the picture depicted in Figure 3(iii). Since π is simple, there must now be a point in the region labelled

D . However, in order for this permutation to not be sum decomposable, we must insert another point in the region to the right of region D , and below the greatest entry in D , but this would force an occurrence of 31524. \square

The class $\text{Av}(24153, 31524, \ell \cdots 21)$ is closed under group-theoretic inversion (because $24153^{-1} = 31524$ and $\ell \cdots 21^{-1} = \ell \cdots 21$), so we may always assume that the latter option in Proposition 4.1 holds.

The rest of our proof adapts several ideas from Vatter [18]. Two rectangles in the plane are said to be *dependent* if their projections onto either the x - or y -axis have nontrivial intersection, and otherwise they are said to be *independent*. A set of rectangles is called independent if its members are pairwise independent. Thus an independent set of rectangles may be viewed as a permutation, and it satisfies the Erdős-Szekeres Theorem (every permutation of length at least $(a-1)(b-1)+1$ contains either $12 \cdots a$ or $b \cdots 21$). We construct independent sets of rectangles in the proofs of both Propositions 4.3 and 4.5. In these settings, the rectangles are used to capture “bad” areas in the plot of a permutation, and our desired result is obtained by slicing the rectangles with horizontal and vertical lines in the sense of Proposition 3.3. The following result shows that we may slice a collection of rectangles with only a few lines, so long as we can bound its independence number.

Theorem 4.2 (Gyárfás and Lehel [10]). *There is a function $f(m)$ such that for any collection \mathfrak{R} of axis-parallel rectangles in the plane which has no independent set of size greater than m , there exists a set of $f(m)$ horizontal and vertical lines which slice every rectangle in \mathfrak{R} .*

Our next two results both rest upon Theorem 4.2.

Proposition 4.3. *For every ℓ , the simple permutations in $\text{Av}(24153, 31524, \ell \cdots 21)$ are contained in $\text{Grid}(M)$ for a finite 0/1 matrix M .*

Proof. By Proposition 3.3 it suffices to show that there is a function $g(\ell)$ such that every permutation in $\text{Av}(24153, 31524, \ell \cdots 21)$ can be covered by $g(\ell)$ increasing axis-parallel rectangles (i.e., rectangles which only cover increasing sets of points). We prove this statement by induction on ℓ . For the base case, we can take $g(2) = 1$. Now take a simple permutation $\pi \in \text{Av}(24153, 31524, \ell \cdots 21)$ for $\ell \geq 3$ and suppose that the claim holds for $\ell - 1$.

By Proposition 4.1, we may assume that the leftmost entry of π lies above its rightmost entry. Let π_t be the permutation formed by all entries of π lying above its rightmost entry and π_b the permutation formed by all entries of π lying below its leftmost entry (as shown in Figure 4). Thus every entry of π corresponds to an entry in π_t , to an entry in π_b , or to entries in both permutations. Moreover, both π_t and π_b avoid $(\ell - 1) \cdots 21$.

We would like to apply induction to find monotone rectangle coverings of both π_t and π_b but of course these permutations needn't be simple. Nevertheless, if π_t is the inflation of the simple permutation σ_t and π_b of σ_b then both σ_t and σ_b can be covered by $g(\ell - 1)$ increasing axis-parallel rectangles by induction. Now we stretch these rectangles so that




Figure 4: On the left, the division of π used in Proposition 4.3. In the centre and on the right, the final contradictions in the proof.

they cover the corresponding regions of π . By adapting the proof of Proposition 3.3, we may then extend this rectangle covering to a gridding of size at most $4g(\ell - 1) \times 4g(\ell - 1)$. While this gridding needn't be monotone, inside each of its cells we see points which correspond either to inflations of increasing sequences of σ_t or of σ_b . Let \mathfrak{L} denote the grid lines of this gridding.

We now say that the axis-parallel rectangle R is *bad* if it is fully contained in a cell of the above gridding and the points it covers contain a decreasing interval in either π_t or π_b . Further let \mathfrak{R} denote the collection of all bad rectangles. We aim to show that there is a collection of $f(2\ell(\ell - 1))$ lines which slice every bad rectangle, where f is the function defined in Theorem 4.2. This, together with Proposition 3.3 and the comments before it, will complete the proof because these lines together with \mathfrak{L} will give a gridding of π of bounded size.

Theorem 4.2 will give us the desired lines if we can show that \mathfrak{R} has no independent set of size greater than $2\ell(\ell - 1) + 1$. Suppose to the contrary that \mathfrak{R} does contain an independent set of this size. Thus at least $\ell(\ell - 1) + 1$ of these bad rectangles lie in one of π_t or π_b ; suppose first that these $\ell(\ell - 1) + 1$ bad rectangles lie in π_t . Because π_t avoids $(\ell - 1) \cdots 21$, the Erdős-Szekeres Theorem implies that at least $\ell + 1$ of its bad rectangles occur in increasing order (when read from left to right). Because π itself is simple, each such rectangle must be separated, and this separating point must lie in $\pi_b \setminus \pi_t$ since (by definition) the points inside this bad rectangle form a decreasing interval in π_t . Appealing once more to Erdős-Szekeres we see that two such separating points must themselves lie in increasing order, as shown in the centre of Figure 4. However, this is a contradiction to our assumption that π avoids 31524 (given by the solid points). As shown on the rightmost pane of this figure, if the $\ell(\ell - 1) + 1$ bad rectangles lie in π_b we instead find a copy of 24153. \square

A *submatrix* of a matrix is obtained by deleting any collection of rows and columns from the matrix. Our next result shows that in $\text{Av}(24153, 31524, \ell \cdots 21)$ the simple permutations can be gridded in a matrix which does not contain $\begin{pmatrix} 1 & 1 \\ 1 & * \end{pmatrix}$ as a submatrix, i.e., in this matrix, there is no non-zero cell with both a non-zero cell below it in the same column, and a non-zero cell to its right in the same row. (The $*$ indicates an entry that




Figure 5: Propagating a side of a hull in \mathfrak{H} , and its corresponding propagation tree.

can be either 0 or 1.)

The following result is in some sense the technical underpinning of our entire argument. We advise the reader to note during the proof that if the hulls in \mathfrak{H} are assumed to be increasing, then the resulting gridding matrix M will be 0/1, not 0/ \pm 1.

Proposition 4.4. *Suppose that π is a permutation and \mathfrak{H} is a collection of m monotone rectangular hulls which cover the entries of π satisfying*

(H1) *the hulls in \mathfrak{H} are pairwise nonintersecting,*

(H2) *no single vertical or horizontal line slices through more than k hulls in \mathfrak{H} , and*

(H3) *no hull in \mathfrak{H} is dependent both with a hull to its right and a hull beneath it.*

*Then there exists a function $f(m, k)$ such that π is M -griddable for a 0/ \pm 1 matrix M of size at most $f(m, k) \times f(m, k)$ which does not contain $\begin{pmatrix} \pm 1 & \pm 1 \\ \pm 1 & * \end{pmatrix}$ as a submatrix.*

Proof. We define a gridding of π using the sides of the hulls in \mathfrak{H} . For a given side from a given hull, form a gridline by extending it to the edges of the permutation. By our hypothesis (H2), this line can slice at most k other hulls.

Whenever a hull is sliced in this way, a second gridline, perpendicular to the first, is induced so that all of the entries within the hull are contained in the bottom left and top right quadrants (for a hull containing increasing entries), or top left and bottom right quadrants (for a hull containing decreasing entries) defined by the two lines. This second gridline may itself slice through the interior of at most k further hulls in \mathfrak{H} , and each such slice will induce another gridline, and so on. We call this process the *propagation* of a line. See Figure 5 for an illustration. For a given propagation sequence, the *propagation tree* has gridlines for vertices, is rooted at the original gridline for the side, and has an edge between two gridlines if one induces the other in this propagation.

Before moving on, we note that it is clear that the propagation tree is connected, but less obvious that it is in fact a tree. This is not strictly required in our argument (we will only need to bound the number of vertices it contains), but if the tree were to contain a cycle it would have to correspond to a cyclic sequence of hulls, and this is impossible without contradicting hypothesis (H3).

In order to bound the size of a propagation tree, we first show that it has height at most $2m - 1$. In the propagation tree of a side from some hull $H_0 \in \mathfrak{H}$, take a sequence H_1, H_2, \dots, H_p of hulls from \mathfrak{H} corresponding to a longest path in the propagation tree, starting from the root. Thus, H_1 is sliced by the initial gridline formed from the side of H_0 , and H_i is sliced by the gridline induced from H_{i-1} for $i = 2, \dots, p$.

We now define a word $w = w_1 w_2 \cdots w_p$ from this sequence, based on the position of hull H_i relative to hull H_{i-1} . For $i = 1, 2, \dots, p$, let

$$w_i = \begin{cases} \mathbf{u} & \text{if } H_i \text{ lies above } H_{i-1} \\ \mathbf{d} & \text{if } H_i \text{ lies below } H_{i-1} \\ \mathbf{l} & \text{if } H_i \text{ lies to the left of } H_{i-1} \\ \mathbf{r} & \text{if } H_i \text{ lies to the right of } H_{i-1} \end{cases}$$

Note that by the process of inducing perpendicular gridlines, successive letters in w must alternate between $\{\mathbf{u}, \mathbf{d}\}$ and $\{\mathbf{l}, \mathbf{r}\}$. Moreover, since no hull interacts with hulls both below it and to its right, w cannot contain a \mathbf{ur} or \mathbf{ld} factor. In other words, after the first instance of \mathbf{u} or \mathbf{l} , there are no more instances of \mathbf{r} or \mathbf{d} . This means that w consists of a (possibly empty) alternating sequence $\mathbf{drdr} \cdots$ or $\mathbf{rdrd} \cdots$ followed by a (possibly empty) alternating sequence $\mathbf{ulul} \cdots$ or $\mathbf{lulu} \cdots$.

Any alternating sequence of the form $\mathbf{drdr} \cdots$ or $\mathbf{rdrd} \cdots$ can have at most $m - 1$ letters, as each hull in \mathfrak{H} (other than H_0) can be sliced at most once in such a sequence. Similarly, any alternating sequence of the form $\mathbf{ulul} \cdots$ or $\mathbf{lulu} \cdots$ can have at most $m - 1$ letters. Consequently, we have $p \leq 2m - 2$, and thus every propagation sequence has length at most $2m - 1$, as required.

Since each gridline in the propagation tree has at most k children, this means that the propagation tree for any given side has at most

$$1 + k + k^2 + \cdots + k^{2m-1} < k^{2m}$$

vertices, yielding a gridding of π with fewer than $4mk^{2m}$ gridlines, and we may take this number to be $f(m, k)$. The gridding matrix M is then naturally formed from the cells of this gridding of π : each empty cell corresponds to a 0 in M , each cell containing points in decreasing order corresponds to a -1 in M , and each cell containing points in increasing order corresponds to a 1 in M .

Finally, we verify that M satisfies the conditions in the proposition. The process of propagating gridlines ensures that each rectangular hull in \mathfrak{H} is divided into cells no two of which occupy the same row or column of the M -gridding. This means that there can be no $\begin{pmatrix} \pm 1 & \pm 1 \\ \pm 1 & * \end{pmatrix}$ submatrix of M with two cells originating from the same hull in \mathfrak{H} .

Thus, the cells of a submatrix of the form $\begin{pmatrix} \pm 1 & \pm 1 \\ \pm 1 & * \end{pmatrix}$ must be made up from points in distinct hulls in \mathfrak{H} , but this is impossible since \mathfrak{H} contains no hulls which are dependent with hulls both below it and to its right. \square

We now apply this proposition to refine the gridding provided to us by Proposition 4.3.




Figure 6: Two independent bad rectangles. The filled points form a copy of 24153, irrespective of the relative orders of the splitting points.

Proposition 4.5. *For every ℓ , the simple permutations in $\text{Av}(24153, 31524, \ell \cdots 21)$ are contained in $\text{Grid}(M)$ for a finite 0/1 matrix M which does not contain $\begin{pmatrix} 1 & 1 \\ 1 & * \end{pmatrix}$ as a submatrix.*

Proof. Let π be an arbitrary simple permutation in $\text{Av}(24153, 31524, \ell \cdots 21)$. As we observed in Footnote 1, it suffices to show that there are constants a and b such that $\pi \in \text{Grid}(M)$ for an $a \times b$ matrix M satisfying the desired conditions.

By Proposition 4.3, π is contained in $\text{Grid}(N)$ for some 0/1 matrix N , of size (say) $t \times u$. We say that a *bad* rectangle within any specified cell is an axis-parallel rectangle which contains two entries which are split both by points below and to the right. Since π does not contain 24153, no cell can contain more than one independent bad rectangle — see Figure 6. Therefore an independent set of bad rectangles can have size at most tu , so Theorem 4.2 shows that the bad rectangles can all be sliced by $f(tu)$ lines.

In any cell of the original gridding, the additional $f(tu)$ slices that have been added can at most slice these points into $f(tu)+1$ maximal unsliced pieces. In the entire permutation, therefore, these slices (together with the original gridlines from N) divide the points into at most $tu(f(tu)+1)$ maximal unsliced pieces. Let \mathfrak{H} denote the rectangular hulls of these maximal unsliced pieces.

We now check that \mathfrak{H} satisfies the hypotheses of Proposition 4.4. Condition (H1) follows immediately by construction. Next, note that no two hulls of \mathfrak{H} from the same cell of the N -gridding can be dependent, since every cell is monotone, so we may take $k = \max(t, u)$ to satisfy (H2). Finally, no hull can contain a bad rectangle (since all bad rectangles have been sliced), and so no hull can simultaneously be dependent with a hull from a cell below it, and a hull from a cell to its right, as required by (H3).

Now, applying Proposition 4.4 (noting that all the rectangular hulls in \mathfrak{H} contain increasing entries), we have a 0/1 gridding matrix M_π for π , of dimensions at most $v \times w$ for some v, w , which does not contain $\begin{pmatrix} 1 & 1 \\ 1 & * \end{pmatrix}$ as a submatrix. We are now done by our comments at the beginning of the proof. \square

Having proved Proposition 4.5, we merely need to put the pieces together to finish the proof of our main theorem. This proposition shows that there is a finite 0/1 matrix M

with no submatrix of the form

$$\begin{pmatrix} 1 & 1 \\ 1 & * \end{pmatrix}$$

such that the simple permutations of $\text{Av}(24153, 31524, \ell \cdots 21)$ are contained in $\text{Grid}(M)$. It follows that

$$\text{Av}(24153, 31524, \ell \cdots 21) \subseteq \langle \text{Grid}(M) \rangle.$$

Moreover, M is a forest because if it were to contain a cycle, it would have to contain a submatrix of the form $\begin{pmatrix} 1 & 1 \\ 1 & * \end{pmatrix}$. Therefore the permutation class $\text{Av}(24153, 31524, \ell \cdots 21)$ is wqo by Theorem 3.4.

Theorem 4.6. *For every ℓ , the permutation class $\text{Av}(24153, 31524, \ell \cdots 21)$ is well-quasi-ordered. Therefore the class of permutation graphs omitting P_5 and K_ℓ is also well-quasi-ordered.*

5 Permutation Graphs Omitting P_6 , P_7 or P_8

In this section, we establish the following:

Proposition 5.1. *The following three classes of graphs are not wqo:*

- (1) the P_6 , K_6 -free permutation graphs,
- (2) the P_7 , K_5 -free permutation graphs, and
- (3) the P_8 , K_4 -free permutation graphs.

In order to prove these three classes are not wqo, it suffices to show that each class contains an infinite antichain. This is done by showing that the related permutation classes contain “generalised” grid classes, for which infinite antichains are already known. In general, we cannot immediately guarantee that the permutation antichain translates to a graph antichain, but we will show that this is in fact the case for the three we construct here.

We must first introduce *generalised* grid classes. Suppose that \mathcal{M} is a $t \times u$ matrix of permutation classes (we use calligraphic font for matrices containing permutation classes). An \mathcal{M} -*gridding* of the permutation π of length n in this context is a pair of sequences $1 = c_1 \leq \cdots \leq c_{t+1} = n + 1$ (the column divisions) and $1 = r_1 \leq \cdots \leq r_{u+1} = n + 1$ (the row divisions) such that for all $1 \leq k \leq t$ and $1 \leq \ell \leq u$, the entries of π from indices c_k up to but not including c_{k+1} , which have values from r_ℓ up to but not including $r_{\ell+1}$ are either empty or order isomorphic to an element of $\mathcal{M}_{k,\ell}$. The *grid class* of \mathcal{M} , written $\text{Grid}(\mathcal{M})$, consists of all permutations which possess an \mathcal{M} -gridding. The notion of monotone griddability can be analogously defined, but we do not require this.

Here, our generalised grid classes are formed from gridding matrices which contain the monotone class $\text{Av}(21)$, and a non-monotone permutation class denoted $\oplus 21$. This is

formed by taking all finite subpermutations of the infinite permutation $21436587\cdots$. In terms of minimal forbidden elements, we have $\bigoplus 21 = \text{Av}(321, 231, 312)$.

Our proof of Proposition 5.1 requires some further theory to ensure we can convert the permutation antichains we construct into graph antichains. The primary issue is that a permutation graph G can have several different corresponding permutations. With this in mind, let

$$\Pi(G) = \{\text{permutations } \pi : G_\pi \cong G\}$$

denote the set of permutations each of which corresponds to the permutation graph G .

Denote by π^{-1} the (group-theoretic) *inverse* of π , by π^{rc} the *reverse-complement* (formed by reversing the order of the entries of π , then replacing each entry i by $|\pi| - i + 1$), and by $(\pi^{-1})^{\text{rc}}$ the *inverse-reverse-complement*, formed by composing the two previous operations (in either order, as the two operations commute). It is then easy to see that if $\pi \in \Pi(G)$, then $\Pi(G)$ must also contain all of π^{-1} , π^{rc} and $(\pi^{-1})^{\text{rc}}$. However, it is possible that $\Pi(G)$ may contain other permutations, and this depends on the graph-theoretic analogue of the substitution decomposition, which is called the *modular decomposition*.

We also need to introduce the graph analogues of intervals and simplicity, which have different names in that context. A *module* M in a graph G is a set of vertices such that for every $u, v \in M$ and $w \in V(G) \setminus M$, u is adjacent to w if and only if v is adjacent to w . A graph G is said to be *prime* if it has no nontrivial modules, that is, any module M of G satisfies $|M| = 0, 1$, or $|V(G)|$.

The following result, arising as a consequence of Gallai's work on transitive orientations, gives us some control over $\Pi(G)$:

Proposition 5.2 (Gallai [9]). *If G is a prime permutation graph, then, up to the symmetries inverse, reverse-complement, and inverse-reverse-complement, $\Pi(G)$ contains a unique permutation.*

We now extend Proposition 5.2 to suit our purposes. Consider a simple permutation σ of length $m \geq 4$, and form the permutation π by inflating two² of the entries of σ each by the permutation 21. Note that $G_{21} = K_2$, and $\Pi(K_2) = \{21\}$. In the correspondence between graphs and permutations, modules map to intervals and vice versa, so prime graphs correspond to simple permutations, and there is an analogous result to Proposition 3.5 for (permutation) graphs.

Thus, for any permutation $\rho \in \Pi(G_\pi)$, it follows that ρ must be constructed by inflating two entries of some simple permutation τ by the permutation 21. Moreover, τ must be one of σ , σ^{-1} , σ^{rc} or $(\sigma^{-1})^{\text{rc}}$, and the entries of τ which are inflated are determined by which entries of σ were inflated, and which of the four symmetries of σ is equal to τ . In other words, we still have $\Pi(G_\pi) = \{\pi, \pi^{-1}, \pi^{\text{rc}}, (\pi^{-1})^{\text{rc}}\}$.

We require one further easy observation:

Lemma 5.3. *If G and H are permutation graphs such that $H \leq G$, then for any $\pi \in \Pi(G)$ there exists $\sigma \in \Pi(H)$ such that $\sigma \leq \pi$.*

²It is, in fact, possible to inflate more than two entries and establish the same result, but we do not require this here.




Figure 7: The first three elements and the general structure of the infinite antichain A in $\text{Grid}(\oplus 21 \oplus 21)$. The grey lines indicate the sequence used to place the entries.

Proof. Given any $\pi \in \Pi(G)$, let σ denote the subpermutation of π formed from the entries of π which correspond to the vertices of an embedding of H as an induced subgraph of G . Clearly $G_\sigma \cong H$, so $\sigma \in \Pi(H)$ and $\sigma \leq \pi$, as required. \square

We are now in a position to prove Proposition 5.1. Since the techniques are broadly similar for all three cases, we will give the details for case (2), and only outline the key steps for the other two cases.

Proof of Proposition 5.1 (2). First, the graph P_7 corresponds to two permutations, namely 3152746 and 2416375 respectively. Thus the class of P_7 , K_5 -free permutation graphs corresponds to the permutation class $\text{Av}(3152746, 2416375, 54321)$. This permutation class contains the grid class

$$\text{Grid}(\oplus 21 \oplus 21),$$

because this grid class avoids the permutations 241635 (contained in both 3152746 and 2416375), and 54321.

We now follow the recipe given by Brignall [7] to construct an infinite antichain which lies in $\text{Grid}(\oplus 21 \oplus 21)$. Call the resulting antichain A , the first three elements and general term of which are illustrated in Figure 7. This antichain is related to the ‘‘parallel’’ antichain in Murphy’s thesis [15].

Now set $G_A = \{G_\pi : \pi \in A\}$, and note that G_A is contained in the class of permutation graphs omitting P_7 and K_5 by Lemma 5.3. If G_A is an antichain of graphs, then we are done, so suppose for a contradiction that there exists $G, H \in G_A$ with $H \leq G$. Take the permutation $\pi \in A$ for which $G_\pi = G$. Applying Lemma 5.3, there exists $\sigma \in \Pi(H)$ such that $\sigma \leq \pi$. We cannot have $\sigma \in A$ since A is an antichain, so σ must be some other permutation with the same graph. Choose $\tau \in A \cap \Pi(H)$.

It is easy to check that the only non-trivial intervals in any permutation in A are the two pairs of points in Figure 7 which are circled. Thus, τ is formed by inflating two entries of a simple permutation each by a copy of 21. By the comments after Proposition 5.2, we conclude that σ is equal to one of τ^{-1} , τ^{rc} or $(\tau^{-1})^{\text{rc}}$.

Note that every permutation in A is closed under reverse-complement, so $\tau = \tau^{\text{rc}}$ and thus $\sigma \neq \tau^{\text{rc}}$. If $\sigma = \tau^{-1}$, by inspection every element of A contains a copy of 24531, which means that σ contains $(24531)^{-1} = 51423$. Since $\sigma \leq \pi$, it follows that π must also contain a copy of 51423. However this is impossible, because 51423 is not in $\text{Grid}(\oplus 21 \oplus 21)$.




Figure 8: The first three elements of an infinite antichain in $\text{Grid} \left(\begin{array}{cc} \oplus_{21} & \text{Av}(21) \\ & \oplus_{21} \end{array} \right)$.

Thus, we must have $\sigma = (\tau^{-1})^{\text{rc}}$, in which case σ must contain a copy of $((24531)^{\text{rc}})^{-1} = 34251$. However, this permutation is also not in $\text{Grid} \left(\begin{array}{cc} \oplus_{21} & \oplus_{21} \end{array} \right)$ so cannot be contained in π . Thus $\sigma \notin \pi$, and from this final contradiction we conclude that $H \not\leq G$, so G_A is an infinite antichain of permutation graphs, as required. \square

Sketch proof of Proposition 5.1 (1) and (3). For (1), the class of P_6 , K_6 -free permutation graphs corresponds to $\text{Av}(241635, 315264, 654321)$. This class contains

$$\text{Grid} \left(\begin{array}{cc} \oplus_{21} & \text{Av}(21) \\ & \oplus_{21} \end{array} \right)$$

because this grid class does not contain 23154 (contained in 241635), 31254 (contained in 315264) and 654321.

By [7], this grid class contains an infinite antichain whose first three elements are illustrated in Figure 8. The permutations in this antichain have exactly two proper intervals, indicated by the circled pairs of points in each case, and this means that for any permutation π in this antichain, $\Pi(G_\pi) = \{\pi, \pi^{-1}, \pi^{\text{rc}}, (\pi^{-1})^{\text{rc}}\}$.

Following the proof of Proposition 5.1 (2), it suffices to show that for any permutations σ and π in the antichain, none of σ , σ^{-1} , σ^{rc} , or $(\sigma^{-1})^{\text{rc}}$ is contained in π . This is done by identifying permutations which are not contained in any antichain element π , but which are contained in one of the symmetries σ^{-1} , σ^{rc} , or $(\sigma^{-1})^{\text{rc}}$. We omit the details.

For (3), The P_8 , K_4 -free permutation graphs correspond to the permutation class $\text{Av}(4321, 24163857, 31527486)$, and this contains the (monotone) grid class

$$\text{Grid} \left(\begin{array}{cc} \text{Av}(21) & \text{Av}(21) \\ \text{Av}(21) & \text{Av}(21) \end{array} \right).$$

In this case, we appeal to Murphy and Vatter [16] for an infinite antichain, and the same technique used to prove cases (1) and (2) can be applied. \square

6 Enumeration

One significant difference between studies of the induced subgraph order and the subpermutation order is the research interests of the different camps of investigators. In the




Figure 9: The substitution decomposition tree of 375896214 (of height 3), and a pruned version of the same height which alternates between the labels \oplus and \ominus .

latter area, a great proportion of the work is enumerative in nature. Therefore, having established the structure of permutation graphs avoiding P_5 and cliques, we briefly study the enumeration of the corresponding permutation classes in this section. Our goal is to show that these classes are *strongly rational*, i.e., that they and every one of their subclasses have rational generating functions. Here we refer to

$$\sum_{\pi \in \mathcal{C}} x^{|\pi|}$$

as the generating function of the class \mathcal{C} , where $|\pi|$ denotes the length of the permutation π . Note that by a simple counting argument (made explicit in Albert, Atkinson, and Vatter [3]), strongly rational permutation classes must be wqo.

Proposition 3.5 allows us to associate with any permutation π a unique *substitution decomposition tree*. This tree is recursively defined by decomposing each node of the tree as

- $\sigma[\alpha_1, \dots, \alpha_m]$ where σ is a nonmonotone simple permutation,
- $\alpha_1 \oplus \dots \oplus \alpha_m$ where each α_i is sum indecomposable, or
- $\alpha_1 \ominus \dots \ominus \alpha_m$ where each α_i is skew indecomposable.

See Figure 9 for an example. In particular, note that by our indecomposability assumptions, sum nodes (resp., skew sum nodes) cannot occur twice in a row when reading up a branch of the tree. The *substitution depth* of π is then the height of its substitution decomposition tree, so for example, the substitution depth of the permutation from Figure 9 is 3, while the substitution depth of any simple (or monotone) permutation is 1. As we show in our next result, substitution depth is bounded for the classes we are interested in. This result is a special case of Vatter [18, Proposition 4.2], but we include a short proof for completeness.

Proposition 6.1. *The substitution depth of every permutation in $\text{Av}(\ell \dots 21)$ is at most $2\ell - 3$ for $\ell \geq 2$.*

Proof. We prove the result using induction on ℓ . Only increasing permutations avoid 21 and they have substitution depth 1, so the result holds for $\ell = 2$. Suppose to the contrary

that the result holds for $\ell \geq 2$ but that there is a permutation $\pi \in \text{Av}((\ell + 1) \cdots 21)$ of substitution depth 2ℓ (we may always assume that the depth is precisely this value, because of downward closure). If π is a sum decomposable permutation, express it as $\pi = \pi_1 \oplus \cdots \oplus \pi_m$ and note that at least one π_i must have substitution depth $2\ell - 1$.

Thus we may now assume that there is a sum indecomposable permutation $\pi \in \text{Av}((\ell + 1) \cdots 21)$ of substitution depth $2\ell - 1$. Write $\pi = \sigma[\alpha_1, \dots, \alpha_m]$ where σ is either a simple permutation or decreasing of length at least 2 (to cover the case where π is skew decomposable). At least one of the α_i must have substitution depth $2\ell - 2$, and contains $\ell \cdots 21$ by induction. However, there is at least one entry of σ which forms an inversion with $\sigma(i)$, and thus π itself must contain $(\ell + 1) \cdots 21$, as desired. \square

It was shown in Albert, Atkinson, Bouvel, Ruškuc and Vatter [2, Theorem 3.2] that if M is a forest, then $\text{Grid}(M)$ is a *geometric grid class*. In the same paper, two strong properties of geometric grid classes were established: they are defined by finitely many minimal forbidden permutations and they are strongly rational. We refer the reader to that paper for a comprehensive introduction to geometric grid classes. Our enumerative result follows from the following theorem proved in a subsequent paper.

Theorem 6.2 (Albert, Ruškuc, and Vatter [4, Theorem 7.6]). *The class $\mathcal{C}[\mathcal{U}]$ is strongly rational for all geometrically griddable classes \mathcal{C} and strongly rational classes \mathcal{U} .*

Applying induction on the height of substitution decomposition trees (which are bounded in all of our classes by Proposition 6.1), we immediately obtain the following result.

Theorem 6.3. *For every ℓ , the class $\text{Av}(24153, 31524, \ell \cdots 21)$ is strongly rational.*

7 Concluding Remarks

As shown in Figure 1, there are only three cases remaining: permutation graphs avoiding $\{P_6, K_5\}$, $\{P_6, K_4\}$ and $\{P_7, K_4\}$. Due to the absence of an “obvious” infinite antichain in these cases, we conjecture that they are all wqo. However, all three classes contain (for example) the generalised grid class $\text{Grid}(\bigoplus_{21} \text{Av}(21))$, so these classes all contain simple permutations which are not monotone griddable. Thus our approach would have to be significantly changed to approach this conjecture.

A natural first step might be to show that the corresponding permutation classes are \mathcal{D} -griddable (in the sense of Section 5) for a “nice” class \mathcal{D} . As any proof along these lines would need to use some sort of slicing argument as well (such as we used to prove Proposition 4.5), \mathcal{D} would have to be nice enough to allow for such arguments. Vatter [18, Lemma 5.3] has shown that such slicing arguments can be made to work when \mathcal{D} has only finitely many simple permutations and finite substitution depth (in the sense of Section 6). As we already have finite substitution depth by Proposition 6.1, it would suffice to show that we could take \mathcal{D} to contain only finitely many simple permutations.

Again, though, this would only be an encouraging sign and not a proof, because one would then have to develop more sophisticated tools to prove wqo for such classes. Specifically, it is unlikely that the existing machinery of Brignall [7] would suffice.

References

- [1] ALBERT, M. H., AND ATKINSON, M. D. Simple permutations and pattern restricted permutations. *Discrete Math.* 300, 1-3 (2005), 1–15.
- [2] ALBERT, M. H., ATKINSON, M. D., BOUVEL, M., RUŠKUC, N., AND VATTER, V. Geometric grid classes of permutations. *Trans. Amer. Math. Soc.* 365 (2013), 5859–5881.
- [3] ALBERT, M. H., ATKINSON, M. D., AND VATTER, V. Subclasses of the separable permutations. *Bull. Lond. Math. Soc.* 43 (2011), 859–870.
- [4] ALBERT, M. H., RUŠKUC, N., AND VATTER, V. Inflations of geometric grid classes of permutations. *Israel J. Math.* 205, 1 (2015), 73–108.
- [5] ALBERT, M. H., AND VATTER, V. Generating and enumerating 321-avoiding and skew-merged simple permutations. *Electron. J. Combin.* 20 (2) (2013), Research paper 44, 11 pp.
- [6] ATMINAS, A., AND LOZIN, V. Labelled induced subgraphs and well-quasi-ordering. *Order* (2014), 1–16.
- [7] BRIGNALL, R. Grid classes and partial well order. *J. Combin. Theory Ser. A* 119 (2012), 99–116.
- [8] DING, G. Subgraphs and well-quasi-ordering. *J. Graph Theory* 16, 5 (1992), 489–502.
- [9] GALLAI, T. Transitiv orientierbare Graphen. *Acta Math. Acad. Sci. Hungar* 18 (1967), 25–66.
- [10] GYÁRFÁS, A., AND LEHEL, J. A Helly-type problem in trees. In *Combinatorial theory and its applications, II (Proc. Colloq., Balatonfüred, 1969)*. North-Holland, Amsterdam, 1970, pp. 571–584.
- [11] HIGMAN, G. Ordering by divisibility in abstract algebras. *Proc. London Math. Soc.* (3) 2 (1952), 326–336.
- [12] HUCZYNSKA, S., AND VATTER, V. Grid classes and the Fibonacci dichotomy for restricted permutations. *Electron. J. Combin.* 13 (2006), R54, 14 pp.
- [13] KORPELAINEN, N., AND LOZIN, V. Two forbidden induced subgraphs and well-quasi-ordering. *Discrete Math.* 311, 16 (2011), 1813–1822.
- [14] KORPELAINEN, N., AND LOZIN, V. V. Bipartite induced subgraphs and well-quasi-ordering. *J. Graph Theory* 67, 3 (2011), 235–249.
- [15] MURPHY, M. M. *Restricted Permutations, Antichains, Atomic Classes, and Stack Sorting*. PhD thesis, Univ. of St Andrews, 2002.

- [16] MURPHY, M. M., AND VATTER, V. Profile classes and partial well-order for permutations. *Electron. J. Combin.* 9, 2 (2003), Research paper 17, 30 pp.
- [17] ROBERTSON, N., AND SEYMOUR, P. Graph minors I–XX. *J. Combinatorial Theory Ser. B* (1983–2004).
- [18] VATTER, V. Small permutation classes. *Proc. Lond. Math. Soc. (3)* 103 (2011), 879–921.
- [19] VATTER, V., AND WATON, S. On partial well-order for monotone grid classes of permutations. *Order* 28 (2011), 193–199.