



Open Research Online

The Open University's repository of research publications and other research outputs

Radio identifications in the NEP Deep Field

Journal Item

How to cite:

White, Glenn J.; Barrufet de Soto, Laia; Pearson, Chris; Serjeant, Stephen; Lim, Tanya; Matsuhara, Hideo; Sirothia, S.K.; Pal, S. and Karouzos, Marios (2017). Radio identifications in the NEP Deep Field. Publications of The Korean Astronomical Society, 32(1) pp. 231–233.

For guidance on citations see [FAQs](#).

© 2017 The Korean Astronomical Society

Version: Version of Record

Link(s) to article on publisher's website:

<http://dx.doi.org/doi:10.5303/PKAS.2017.32.1.231>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

RADIO IDENTIFICATIONS IN THE NEP DEEP FIELD

GLENN J. WHITE^{1,2}, LAIA BARRUFET DE SOTO¹, CHRIS PEARSON^{1,2}, STEPHEN SERJEANT², TANYA LIM¹, HIDEO MATSUHARA³, S.K.SIROTHIA⁴, S.PAL⁵, MARIOS KAROUZOS⁶, AND AKARI-NEP TEAM³¹RAL Space, The Rutherford Appleton Laboratory, Chilton, UK²Dept of Physical sciences, The Open University, Milton Keynes, UK³Inst of Space and Astronautical Science, Japan Aerospace Exploration Agency, Sagami-hara, Kanagawa 229-8510, Japan⁴National Centre for Radio Astrophysics/TIFR, Pune University Campus, Pune 411 007, India⁵Indian Centre for Space Physics, Kolkata-700084, India⁶CEOU-Astronomy Program, Department of Physics & Astronomy, Seoul National University, Gwanak-gu, Seoul, Korea*E-mail: glenn.white@stfc.ac.uk**(Received July 22, 2015; Revised October 25, 2016; Accepted October 25, 2016)*

ABSTRACT

We have imaged the AKARI Deep Field with the GMRT radio telescope at 610 MHz, detecting 1224 radio components, which are optically identified with 455 optical galaxies having a mean r' magnitude brighter of 22.5 (to a completeness limit of 25.4 mag), and an average redshift ~ 0.8 .

Key words: NEP— FIRC —radio continuum: galaxies — multiwavelength survey

1. INTRODUCTION

The AKARI Deep Field North Ecliptic Pole survey (NEP - Matsuhara et al., 2007; Hwang et al., 2007) was designed to study the star formation history as a function of redshift, and to understand the mass assembly of early type galaxies. Using the Japanese AKARI infrared satellite, we made a deep far-infrared survey close to the NEP covering the high exposure (AKARI observed the NEP during every orbit) part of the deep field. To support this we have made a sensitive radio survey with the Giant Metre Wave Radio Telescope (GMRT) at 610 MHz covering the AKARI Deep Field (Takagi et al., 2012; White et al., 2015 *in preparation*), which, by cross-correlating the radio components with AKARI source catalogues, is used to identify radio loud galaxies and active galactic nuclei.

2. RADIO SURVEY

GMRT observations were obtained as part of a 2-beam mosaic of the NEP region observed during June 2011, covering an area of 1.5 degree² around the NEP. The synthesised half power beam width of the data was 4",

<http://pkas.kas.org>

the half power beam-width of the primary beam is 43', and the rms noise level was 20 μ Jy per beam. Figure 1 shows the map observed towards the North Ecliptic Pole using the GMRT radio telescope at 610 MHz.

The GMRT data was compared with the 1400 MHz survey of the NEP by White et al. (2010) with the Westerbork Synthesis Radio Telescope (WSRT) survey which had a resolution of $17 \times 15.5''$ and sensitivity 21 μ Jy per beam, and areal coverage of 1.7 degree². The GMRT survey has the advantage of sampling sources a factor ~ 2 times fainter, assuming that the average spectral index for extragalactic radio sources scales as $\propto \nu^{-0.7}$, and benefits from a factor of ~ 4 times higher angular resolution. In this paper, results are presented for the 0.6 degree² area of the NEP Deep region (Takagi et al., 2012).

Figure 2 shows a montage of images obtained with various telescopes and at various wavelengths (for details of the various ancillary data sets used in this Figure please refer to Table 1 of Karouzos et al., 2014). In this example, the displayed WSRT radio source, NEP 175714+662139, has a photometric redshift of 0.604 and r' magnitude of 20.53 (Oi et al., 2014).




Figure 1. GMRT 610 MHz mosaiced image of the North Ecliptic Pole. 1224 discrete radio components were identified to a 5σ threshold, with the brightest source close to the map centre, NGC6543, The Catseye Nebula, having a flux of 27 mJy per beam. Noise increases at the map edges due to primary beam attenuation.




Figure 2. $40'' \times 40''$ images centred on the WSRT radio source NEP 175714+662139. The circle is a nominal $4''$ diameter reference at the position of the WSRT radio source, to show the relative positional accuracy of the various surveys.

3. CROSS-IDENTIFICATIONS

Matching the GMRT catalogue with the photometric redshift catalogue of Oi et al. (2014) for the NEP Deep area. The GMRT source density in the NEP Deep region is 695 radio sources per square degree. We find 455 sources that coincide to within a 2 arc second radius search area. The distribution of the r' magnitudes of the associations is shown in Figure 3.

The mean of the magnitude distributions of r' magnitudes of optically identified galaxies is ~ 22.5 . Based on the r' source density from Oi et al. (2014), the probability that an unrelated and randomly distributed galaxy with $r' = 23$ or brighter lies within the $2''$ diameter radio search box is $< 3\%$. Further work is needed to con-




Figure 3. r' magnitudes of the GMRT radio associations in the NEP Deep field.




Figure 4. Photometric redshifts of the GMRT radio associations in the AKARI NEP Deep field.

firm associations with galaxies whose r' magnitudes are fainter than 23, because of the possibility to mis-identify as a consequence of confusion.

The assigned photo- z or spectroscopic measurements taken from Oi et al. (2014) are shown in Figure 4, where the mean of the redshift distribution peaks at $z \sim 0.8$.

The identified sources from the GMRT radio sample have been examined in Herschel/AKARI colour-colour space as shown in Figure 5.

It is notable that the template colours and redshift tracks follow the data very well, i.e. for the blue star forming track the 10th marker ($z=1$) is around the point between the $z < 1$ and $z > 1$ sources, as might be expected. The data suggest that $\log(350\mu\text{m}/250\mu\text{m})$ colours > -0.2 preferentially select $z > 1$ objects, and appear better represented by the M82 or LIRG templates rather than the archetypal ULIRG Arp220 SED.

4. CONCLUSIONS

The AKARI North Ecliptic Pole deep field has been imaged using the GMRT telescope at 610 MHz, detecting 1224 radio components. By simple cross-matching with a CFHT r' catalogues, we associated 455 of the GMRT radio sources with r' galaxies down to the optical completeness limit $r' = 25.4$ mag, although the




Figure 5. Colour-colour plot for GMRT radio sources. The black squares are sources $z < 1$, red circles $1 < z < 2$ and the (very few), and blue diamonds at $z > 2$. The small dots on the SED tracks along solid lines correspond to increments in $dz = 0.1$ for the given spectral template or either a normal, star-forming (M82), LIRG, or ULIRG (Arp22) galaxy template.

possibility for confusion rises for galaxies fainter than $r' = 23$ mag. The mean value of the r' associations is $r' = 22.5$, and the corresponding mean redshift is 0.8. Using a combination of Herschel and AKARI data we show that $\log(350\mu\text{m}/250\mu\text{m})$ colours > -0.2 preferentially select objects having $z > 1$, and that the SEDs are better fitted by local M82 or LIRG spectral templates.

REFERENCES

- Hwang, N., et al., 2007, An Optical Source Catalog of the North Ecliptic Pole Region, *ApJS*, 172, 583
- Karouzos, M., et al., 2014, A Tale of Two Feedbacks: Star Formation in the Host Galaxies of Radio AGNs, *ApJ*, 784, 137
- Matsuhara, H., et al., 2006, Deep Extragalactic Surveys around the Ecliptic Poles with AKARI (ASTRO-F), *PASJ*, 58, 673
- Oi, N., et al., 2014, Optical - near-infrared catalog for the AKARI north ecliptic pole Deep field, *A&A*, 566, A60
- Takagi, T., et al., 2012, The AKARI NEP-Deep survey: a mid-infrared source catalogue, *A&A*, 537, A24
- White, G. J., et al., 2010, A deep survey of the AKARI north ecliptic pole field . I. WSRT 20 cm radio survey description, observations and data reduction, *A&A*, 517, A54
- White, G. J., et al., 2012, A deep ATCA 20 cm radio survey of the AKARI Deep Field South near the South Ecliptic Pole, *MNRAS*, 427, 1830