Accepted Manuscript

The bogus taste test: Validity as a measure of laboratory food intake

Eric Robinson, Ashleigh Haynes, Charlotte A. Hardman, Eva Kemps, Suzanne Higgs, Andrew Jones

PII: S0195-6663(17)30321-5

DOI: 10.1016/j.appet.2017.05.002

Reference: APPET 3453


To appear in: Appetite

Received Date: 27 February 2017

Revised Date: 27 April 2017 Accepted Date: 1 May 2017

Please cite this article as: Robinson E., Haynes A., Hardman C.A., Kemps E., Higgs S. & Jones A., The bogus taste test: Validity as a measure of laboratory food intake, *Appetite* (2017), doi: 10.1016/j.appet.2017.05.002.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1	The bogus taste test: Validity as a measure of laboratory food intake
2	
3	
4	
5	
6	
7	
8	Eric Robinson ¹ , Ashleigh Haynes ¹ , Charlotte A. Hardman ¹ , Eva Kemps ² , Suzanne Higgs ³ ,
9	Andrew Jones ¹
10	
11	¹ Institute of Psychology, Health & Society, University of Liverpool, UK
12	² School of Psychology, Flinders University, Adelaide, Australia
13	³ School of Psychology, University of Birmingham, Edgbaston, Birmingham, UK
14	sensor of rejenerogy, emiterary of biningham, Eugenstein, biningham, err
15	Common dance Eric Dahiman Institute of Development Health & Contact Heimster of
16	Correspondence: Eric Robinson, Institute of Psychology, Health & Society, University of
17	Liverpool, L69 7ZA, UK. Email: eric.robinson@liv.ac.uk
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	Word Count: 5,992
45	TOTA COMIN 3,772
46	Dunning Head: Validity of the tests test
47	Running Head: Validity of the taste test
48	
49	

51 Abstract

Because overconsumption of food contributes to ill health, understanding what affects how
much people eat is of importance. The 'bogus' taste test is a measure widely used in eating
behaviour research to identify factors that may have a causal effect on food intake. However,
there has been no examination of the validity of the bogus taste test as a measure of food
intake. We conducted a participant level analysis of 31 published laboratory studies that used
the taste test to measure food intake. We assessed whether the taste test was sensitive to
experimental manipulations hypothesized to increase or decrease food intake. We examined
construct validity by testing whether participant sex, hunger and liking of taste test food were
associated with the amount of food consumed in the taste test. In addition, we also examined
whether BMI (body mass index), trait measures of dietary restraint and over-eating in
response to palatable food cues were associated with food consumption. Results indicated
that the taste test was sensitive to experimental manipulations hypothesized to increase or
decrease food intake. Factors that were reliably associated with increased consumption during
the taste test were being male, have a higher baseline hunger, liking of the taste test food and
a greater tendency to overeat in response to palatable food cues, whereas trait dietary restraint
and BMI were not. These results indicate that the bogus taste test is likely to be a valid
measure of food intake and can be used to identify factors that have a causal effect on food
intake.

Key Words: Taste test; food intake; laboratory; appetite; eating behaviour;

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

The bogus taste test: Validity as a measure of laboratory food intake Because of the damaging effects that poor diet and overconsumption of food have on health (Kopelman, 2007; Prentice, 2001), there is a need to understand the factors effecting how much people eat. Moreover, isolating the causal effect that biological, environmental and psychological factors have on food intake enables more nuanced theories of human eating behaviour. A variety of methods exist to measure eating behaviour. A large amount of epidemiological research has measured food and energy intake by using self-report methods, including food frequency questionnaires and dietary recalls. Although widely used and relatively inexpensive, the precision of such measures have long been questioned because of concerns over respondents' ability and motivation to provide highly accurate reports of their eating behaviour (Heitmann & Lissner, 1995; Macdiarmid & Blundell, 1998; Schoeller, 1990; Schoeller et al., 2013). Laboratory measurement of food intake is another approach used to assess human eating behaviour. Unlike self-report measures, the controlled environment of the laboratory allows for objective examination of food intake. One laboratory approach is to examine food intake from test meals. In such studies participants are served a single or multi-item meal at breakfast, lunch and/or dinner, are told to eat until they are comfortably full, and the total amount of ad-libitum energy consumed is calculated (Blundell et al., 2010). The measurement of energy intake from test meals is common in research that examines the underlying physiology of human eating. For example, by assessing food intake at test meals across the day (or even for several days), it is feasible to examine whether pharmaceutical or nutritional interventions increase or decrease energy intake and/or affect food preference (Gibbons, Finlayson, Dalton, Caudwell, & Blundell, 2014; Hill, Rogers, & Blundell, 1995;

Welch et al., 2011). This type of test meal design has been reported to be valid and reliable

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

(Blundell, et al., 2010; Gregersen et al., 2008; Martin et al., 2005). However, it has practical limitations. Test meal methods can be expensive and time consuming for researchers and require specialist research facilities. Furthermore, methods used at present rarely attempt to disguise that the test meal is being used to measure participant food consumption, e.g. (Andrade, Kresge, Teixeira, Baptista, & Melanson, 2012; Yip, Wiessing, Budgett, & Poppitt, 2013). This could be problematic because transparency of the purpose of the test meal may affect the amount of food that participants eat due to self-presentation concerns (Robinson, Hardman, Halford, & Jones, 2015; Robinson, Kersbergen, Brunstrom, & Field, 2014) and this effect may be differential dependent on participant individual differences within or across samples (Robinson, Proctor, Oldham, & Masic, 2016). This line of reasoning is consistent with classic social psychology research on demand characteristics and 'observer' effects, whereby behaviour can be biased by awareness of the purpose of a study (Nichols & Maner, 2008; Orne, 1962). Indeed, for some time there has been concern that commonly used laboratory methods to study eating behaviour are too artificial, and therefore lack ecological validity (de Castro, 2000; Meiselman, 1992). A different laboratory measure of food intake is the bogus taste test. The bogus taste

A different laboratory measure of food intake is the bogus taste test. The bogus taste test typically involves providing participants with one or more food items and unobtrusively measuring the amount of food consumed. In an attempt to disguise that food intake is being measured, participants are led to believe that the purpose of the task is to assess their taste perception of the food(s). Participants are provided with the food, a series of taste ratings to complete (e.g. how sweet is the food?) in a set time period (e.g. 10 minutes) and are normally informed that once they have completed the ratings they are free to eat as they please. The taste test therefore is relatively inexpensive and convenient to use, as well as acting as a 'disguised' and objective measurement of food intake that can be easily implemented in laboratory settings. The taste test has been employed to examine whether a range of

environmental and psychological factors influence food intake, including but not exclusive to; social norms (Robinson, Sharps, Price, & Dallas, 2014), advertisement (Harris, Bargh, & Brownell, 2009), portion size (Spanos, Kenda, & Vartanian, 2015), alcohol intoxication (Christiansen, Rose, Randall-Smith, & Hardman, 2016), stress (Sproesser, Schupp, & Renner, 2013), memory for recent eating (Higgs, 2002), attentional bias (Werthmann et al., 2011), mindfulness (Hooper, Sandoz, Ashton, Clarke, & McHugh, 2012), impulsivity (Guerrieri, Nederkoorn, & Jansen, 2008) and inhibitory control (Houben, 2011). Although the taste test has been employed by researchers for some time, e.g. (Conger, Conger, Costanzo, Wright, & Matter, 1980), unlike other measures of eating behaviour there has been no formal assessment of the validity of the taste test as a measure of food intake. For a recent examination of the bogus taste test in alcohol research see (Jones et al., 2016).

Variables Associated with Food Intake

Here we examine the validity of the bogus taste test as a measure of food intake by making use of participant level data from 31 published studies that adopted the taste test. If the taste test is a valid measure of food intake then factors that have been shown to reliably predict how much food a person consumes using other paradigms would be expected to predict food intake in the taste test. For example, although not all studies show a statistically significant relationship between hunger and food intake, there is now consistent evidence that self-reported hunger measured prior to eating modestly predicts how much a person will subsequently eat during a meal (de Castro & Elmore, 1988; Horner, Byrne, & King, 2014; Sadoul, Schuring, Mela, & Peters, 2014). Likewise, studies have consistently shown that individuals prefer to eat less of foods they dislike and more of a food if they like its taste (Brunstrom & Shakeshaft, 2009; de Graaf et al., 2005; Drewnowski & Hann, 1999). There are also marked sex differences in food intake, whereby men have a higher energy need and

tend to consume more food than women (Rolls, Fedoroff, & Guthrie, 1991). Thus, in the present analyses we predicted that hunger, food liking and being male (as opposed to female) would positively predict taste test food intake and that evidence for these associations would imply support for construct validity of the taste test.

We also examined whether trait dietary restraint and the tendency to over-eat in response to palatable food cues predict taste test food intake. Trait dietary restraint can be defined as the tendency to consciously attempt to restrict food intake in order to prevent weight gain. Based on this definition, we predicted that higher dietary restraint should be predictive of lower taste test food intake. However, we made this prediction tentatively because whether attempts to restrict food intake reliably translate to reduced food intake is questionable, with some research suggesting that dietary restraint can often 'backfire'. Rather than being predictive of lower energy consumption, restraint has in some studies been associated with over-eating (Herman & Mack, 1975; Johnson, Pratt, & Wardle, 2012; Stroebe, van Koningsbruggen, Papies, & Aarts, 2013; Wardle, Steptoe, Oliver, & Lipsey, 2000). In addition, there is observational data which suggest that dietary restraint does not predict restriction of objectively measured food intake in the real world (Stice, Sysko, Roberto, & Allison, 2010).

The tendency to over-eat in response to palatable food cues is a factor that may also predict taste test food intake. In the present research we made use of self-reported data on trait disinhibited eating and trait external eating to characterize 'over-eating in response to palatable food cues'. In particular, trait disinhibition has been implicated in greater food intake and weight gain in multiple studies (Bryant, King & Blundell, 2008; French, Epstein, Jeffery, Blundell, & Wardle, 2012). However, there has been some debate over the accuracy and validity of self-reported trait measures of behaviour (Evers et al., 2011; Bongers & Jansen, 2016; Evers, de Ridder, & Adriaanse, 2009). Based on this we tentatively predicted

that self-reported tendencies to over-eat in response to palatable food cues would be positively associated with taste test food intake.

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

We also know that participants with a higher body mass index (BMI) should on average have a greater energy need and therefore eat more than individuals with a lower BMI. In line with this, in multiple laboratory and epidemiology studies individuals of heavier body weight have demonstrated a greater total energy intake (Forslund, Torgerson, Sjostrom, & Lindroos, 2005; Sadoul, et al., 2014; Trichopoulou, Gnardellis, Lagiou, Benetou, & Trichopoulos, 2000). De Castro et al. (2012) found evidence that a heavier BMI was associated with self-reported energy intake and this relationship was most pronounced when participants were eating outside of the home. Yet, there are studies which report no significant association between BMI and energy intake. For example, Bell and Rolls (2001) found no difference in laboratory measured energy intake between females with normal weight and obesity. Similarly, in addition, although Berg et al. (2009) found that obesity was related to larger self-reported meal size for main meals among a large sample of Swedish adults, there was no significant relationship between BMI and daily energy intake in this study. There are also complex relationships between dietary restraint, over-eating in response to food cues and BMI. Individuals of heavier BMI are more likely to be restrained eaters, but ironically, also more likely to score higher on measures of over-eating (French, et al., 2012). In addition, laboratory taste tests typically involve the consumption of 'unhealthy' energy dense food. Because individuals of heavier body weight may be more likely to present their eating behaviour in a socially desirable way (Hebert, Clemow, Pbert, Ockene, & Ockene, 1995), or eat minimally when they are aware that their food intake is assessed because of self-presentation concerns (Robinson, et al., 2016), heavier BMI may not predict greater food intake. Thus, in the context of a taste test it is not clear whether a heavier BMI would predict

200	greater, limited or equivocal food intake. Because of these considerations we tentatively
201	predicted that a higher BMI would be associated with greater taste test food intake.
202	
203	Sensitivity to Experimental Manipulation
204	A further test of the validity of the taste test is whether the amount of food a participant eats
205	in a taste test is sensitive to experimental manipulations hypothesized to increase or decrease
206	food intake. Although previous research suggests that the taste test is sensitive to
207	experimental manipulation (Conger, et al., 1980; Roth, Herman, Polivy, & Pliner, 2001),
208	there are instances in which taste test methods have been used, and manipulations expected to
209	increase or decrease food intake, did not do so (Blodorn, Major, Hunger, & Miller, 2016;
210	Cavanagh, Vartanian, Herman, & Polivy, 2013). It is difficult to conclude why 'null' findings
211	occur in individual studies; it may be that theoretical predictions are inaccurate, studies lack
212	adequate statistical power and/or the methods used (e.g. the taste test) are not sufficiently
213	sensitive. In the present analyses we were able to formally examine, with more than adequate
214	statistical power, whether manipulations that had been hypothesized to increase or decrease
215	taste test food intake did do so. We predicted that the taste test would be sensitive to
216	manipulations hypothesized to increase or decrease food intake and evidence of this would
217	provide further support for the validity of the taste test.
218	
219	Testing Validity of the Taste Test
220	We reasoned that the taste test being sensitive to experimental manipulation and associated
221	with participant level variables that are reliably associated with food intake in other

paradigms (participant sex, baseline hunger and liking of the food used in a taste test) would

provide strong confirmatory evidence for the validity of the taste test.

222

224

Methods 225 Because our approach required analysis of participant level data, we made use of available 226 data sets from published studies of three research groups based in the UK and Australia that 227 have routinely employed the bogus taste test in laboratory settings over the last 15 years. 228 These studies were performed by, or under the supervision of, at least one of the present 229 article's authors. See https://osf.io/ggkqp/ for preregistration of our methods and a-priori 230 231 analysis strategy. 232 *Inclusion:* In total, 34 independent studies from 27 publications were identified initially. We 233 limited our analysis to 31 studies (from 26 publications) that used between-subjects designs. 234 As the taste test is typically used in between-subjects studies and there would be insufficient 235 236 data to make comparisons between study types (i.e. comparing within, mixed and betweensubjects), we did not include 3 studies that used within or mixed subjects designs. Studies 237 included in the analysis are denoted in the reference list with an asterisk. 238 239 Study procedure: In all studies participants were led to believe that the aim of the taste test 240 was to examine taste perception of the foods in the taste test, rather than to assess food intake. 241 Participants were provided with the taste test food, a questionnaire about taste perceptions 242 (e.g. how crunchy is the food?), before being asked to complete the ratings and were told that 243 244 they were free to eat as much or as little of the foods as desired after completing the ratings. Participants were left alone to do this task, typically for 10 minutes. Hunger was self-reported 245 shortly before the taste test in all studies. Liking of the foods used in the taste test was self-246 247 reported by participants during or immediately after the taste test. Self-reported participant level characteristics (sex, trait dietary restraint, trait over-eating in response to palatable food 248 cues) tended to be measured after the taste test. Weight and height tended to be measured 249

250	after the taste test to calculate BMI, although in a small proportion of studies, weight and
251	height were self-reported. See Supplemental Table 1 for a list of the individual studies
252	included and the variables included in the analyses for each study.
253	
254	Sex: Participants in the 31 studies were predominantly female (2613/2692: 97%), so our main
255	analyses were planned only on women (N=2613). However, we conducted an additional
256	separate analysis to examine sex differences in food intake from studies (N=4) in which both
257	men and women participated.
258	
259	Participant level variables: To assess variables of interest that would have sufficient data for
260	analysis, we first identified variables that were measured and available in the majority of data
261	sets (i.e. > 50% data sets were required to include a measurement of a variable of interest in
262	order to ensure adequate statistical power for analyses). This resulted in us extracting
263	participant level data for baseline hunger (N=2464), taste test food liking (N=1871), trait
264	dietary restraint (N=1640), trait over-eating in response to palatable food cues (N=1546) and
265	BMI ($N=2275$). A total of $N=1071$ participants had data for taste test food intake and all of
266	the above participant level variables. We Z-scored baseline hunger, liking, restraint and over-
267	eating in response to palatable food cues for each individual study because of variability in
268	the way these constructs were measured across studies. BMI was measured consistently in
269	each study (weight/height squared), so we did not Z score BMI.
270	
271	Experimental conditions: Based on the introduction section of each published article, two
272	authors independently coded the experimental conditions in each study as either hypothesised
273	to increase, decrease or have no overall effect on food intake (no effect on food intake
274	'control' condition). Blinded initial agreement between the two coders was high (90%

agreement). In the remaining cases there was some ambiguity in papers about the specific hypotheses for an experimental condition, but the two coders agreed after discussion.

Operationalising taste test food intake: Because the amount of time given, number of taste test ratings required, type of food, number of food items, quantities of food and measurement of intake (e.g. grams, calories) used varied (and was sometimes not reported in detail) across taste tests in each study, to standardize our dependent variable of interest we Z scored food intake in each individual study. In 25/31 studies food intake was coded as total amount of food consumed. In two studies (Kemps et al., 2016a, 2016b), 50% of participants received grapes as the taste test food and 50% received chocolate. We did not include the data from participants receiving grapes, as taste tests typically involve an energy dense food and there were insufficient studies using only grapes to be able to formally compare them to other studies in the analysis. In four studies (Kakoschke, Kemps, & Tiggemann, 2014; Kemps, Tiggemann, & Elford, 2015; Kemps, Tiggemann, Orr, & Grear, 2014; Schumacher, Kemps, & Tiggemann, 2016) there were multiple taste test foods and the authors had experimental hypotheses specific to the intake of one of the foods in the taste test (e.g. chocolate muffin, but not blueberry muffin intake). In these studies, we used food intake data for only the food type that was central to the authors' experimental hypotheses.

Planned primary unadjusted analyses: We first planned to examine our hypotheses using all available data in a set of unadjusted analyses, in which statistical significance was set at p < .05. To assess whether the taste test is sensitive to experimental manipulation, we planned a one way ANOVA, with experimental condition as the between-subjects factor. If a main effect was observed, we planned follow up pairwise comparisons between the three experimental conditions (increase, decrease and control). To assess whether participant level

variables were associated with food intake we planned Pearson's r correlations. To examine sex differences on taste test food intake, we planned an independent samples-test on data from the four studies in which men and women participated.

Planned primary adjusted analyses: Next, we planned to assess the extent to which experimental conditions and participant level variables independently predicted food intake using stepwise regression. The first step included experimental design (i.e. dummy coded experimental conditions). The second step included participant level variables (hunger, restraint, over-eating in response to palatable food cues, BMI). Because taste test food liking in the studies was measured during the taste test, or immediately after, we reasoned that its association with food intake may be inflated due to reverse causality. According to self-perception theory (Bem, 1972), people base their beliefs in part on their prior behaviour (e.g., 'I ate a lot of cookies, so I must really like the taste of cookies'), so it is plausible that a participant who ate a lot of food in the taste test would assigned a higher liking rating to it.

Because of this, we planned to enter liking separately in a final step of the regression model.

Planned secondary analyses: We planned to test whether results were similar in the UK vs Australian studies. If any participant level variables were predictive of food intake, we planned to assess whether these associations were observed consistently across UK vs Australian studies by computing interactions between country of origin and the participant level variables and entering them into the above regression model at a further step. We also planned to examine whether the associations between taste test food intake and trait measures of restraint and over-eating in response to palatable food cues differed dependent on the trait questionnaire used; restraint and disinhibition subscales of the TFEQ (Stunkard & Messick, 1985) vs. the restraint and external eating subscales of the DEBQ (Van Strien, Frijters,

Bergers, & Defares, 1986), by computing interactions between trait measure type and scale score, and entering them into the regression model at a further step.

Statistical power: Sample sizes provided us with adequate statistical power to detect statistically small effects ($f^2 = 0.02$, > 80% power, p < .05) in our planned primary and secondary analyses.

332 Results

In our unadjusted analyses we made use of data from 2613 female participants, with a mean age of 20.7 years (SD = 4.6) and a mean BMI (kg/metres²) of 22.8 (SD = 4.4).

Experimental manipulations of food intake

There was a significant effect of experimental condition on food intake (F (2, 2610) = 26.10, p < .001, partial eta sq = 0.02). Pairwise comparisons indicated that participants in conditions that were hypothesized to increase food intake ate significantly more (p = .016, d = 0.11) than did the participants in 'control' conditions that were not hypothesized to have an effect on food consumption, and participants in conditions that were hypothesized to decrease food intake ate significantly less (p < .001, d = 0.27) than did participants in 'control' conditions that were not hypothesized to affect food consumption. The difference in food intake between participants in the conditions hypothesized to increase vs. decrease food intake was also statistically significant (p < .001, d = 0.38). See Table 1.

Table 1. Effect of experimental conditions on taste test food intake

Condition	N	Z scored food intake
Decrease intake	689	22 (0.89)
Control	1180	.04 (0.99)
Increase intake	744	.15 (1.06)

348 Z scored food intake values are means (standard deviations in brackets)

Unadjusted associations between participant level variables and food intake

Baseline hunger, liking of taste test food and trait over-eating in response to palatable food cues were all significantly positively correlated with taste test food intake. Trait dietary restraint was significantly negatively correlated with taste test food intake, whereas BMI was not significantly correlated with taste test food intake. See Table 2.

Table 2. Unadjusted associations between taste test food intake and participant level variables

	1	1			1
	Baseline	Body mass	Liking of	Trait	Trait over-
	hunger	index	test food	dietary	eating
	S			restraint	O
Food	r = .19	r = .03	r = .27	r =05	r = .13
intake	p < .001	p = .18	p < .001	p = .04	p < .001
	N = 2464	N = 2275	N = 1871	N = 1640	N = 1546
Baseline		r =04	r = .20	r =05	r = .10
hunger		p = .09	p < .001	p = .06	p < .001
		N = 2126	N = 1871	N = 1640	N = 1546
Body mass			r = .02	r = .10	r = .08
index			p = .53	p < .001	p = .002
			N = 1735	N = 1528	N = 1463
Liking of				r =07	r = .22
test food				p = .016	p < .001
				N = 1248	N = 1155
Trait					r = .10
dietary		Y			p < .001
restraint					N = 1543

Sex and food intake

An independent samples t-test indicated that male participants (N = 79, M Z scored intake =

.23, SD = 1.10) consumed significantly more food (t (258) = 2.50, p = .013, d = 0.34) than

did female participants (N = 181, MZ scored intake = -.10, SD = 0.93).

Predictors of taste test food intake using stepwise regression

The final model was statistically significant (F = 37.05, p < .001, Adjusted $R^2 = .12$) and included the following predictor variables; experimental manipulations hypothesized to decrease food intake, baseline hunger, over-eating in response to palatable food cues and taste test food liking. See Table 3. Manipulations hypothesized to increase food intake, BMI and restraint were not significant predictors in any steps of the model. Over-eating in response to palatable food cues was a significant predictor in all steps, but became non-significant in the final step in which taste test food liking was included. Experimental manipulations hypothesized to increase food intake approached significance as a predictor variable in a number of the steps of the model, but was not included in the final model.

Table 3. Stepwise linear regression model results

Predictor variables	Model (step one) Adjust $R^2 = .02$	Model (step two) Adjust $R^2 = .07$	Final model Adjust R ² = .12
Exp. condition	$B =14, p < .001^a$	$B =14, p < .001^{a}$	$B =14, p < .001^{a}$
decrease intake		y	
Exp. condition	$B = .05, p = .15^{b}$	$B = .06, p = .06^{b}$	B = .04, p = .15
increase intake	\wedge		
Baseline	-	$B = .21, p < .001^{a}$	$B = .16, p < .001^a$
hunger			
Trait	-	$B = .09, p = .002^{a}$	$B = .05, p = .12^{a}$
over-eating			
Trait dietary	<u> </u>	$B =02, p = .57^{b}$	$B =01, p = .80^{b}$
restraint			
Body mass	-	$B = .02, p = .58^{b}$	$B = .01, p = .62^{b}$
index			
Taste test food	_	-	$B = .23, p < .001^a$
liking)		h

B refers to standardized Beta values. ^a indicates predictor variable was included in model step. ^b indicates predictor variable was not included in model step.

Generalizability of findings

Of the 31 included studies, 18 were conducted in the UK and 13 in Australia. Study country of origin did not interact significantly with participant liking of the taste test food or trait

over-eating in response to palatable food cues to predict food intake. However, there was a small but significant interaction between study country and baseline hunger (B = .09, p = .04, R^2 change = .004). To examine the direction of the interaction we conducted our planned main regression models separately in studies conducted in the UK and Australia. In line with our main findings, baseline hunger was a modest significant positive predictor of food intake in both countries, although the strength of association between hunger and food intake was stronger in UK studies (N participants = 439, B = .25, p < .001) than Australian studies (N participants = 631, B = .11, p = .006). We also found a significant interaction between trait over-eating in response to palatable food cues and measure type (i.e., TFEQ disinhibition versus DEBQ external eating) (B = .07, p = .04, R^2 change = .003). To follow up this interaction we conducted our planned main regression models separately using data from studies that measured trait over-eating in response to palatable food cues using the TFEQ vs. the DEBQ. Over-eating in response to palatable food cues was a significant predictor of food intake in studies that used the TFEQ disinhibition scale (N = 324, B = .15, p = .005), but was not a significant predictor of food intake in studies that used the DEBQ external eating scale (N = 746, B = .002, p = .95). By contrast, there was no significant interaction between trait dietary restraint score and restraint measure type (i.e, TFEQ versus DEBQ).

398

399

400

401

402

403

404

405

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

Post-hoc analyses

As we found no correlation between BMI and taste test food intake we examined whether consistent results were observed when categorizing participants according to World Health Organization BMI categories; underweight (BMI < 18.5, N = 163), normal weight (BMI 18.5-24.9, N = 1642), overweight (BMI 25-29.9, N = 330) and obese (BMI \geq 30, N = 140). In line with the correlational analyses, there was no significant effect of BMI category on food intake tested using a one way ANOVA (F (3, 2271) = 1.27, p = .28, partial eta sq = 0.002).

406 Discussion

The aim of the present study was to examine the validity of the bogus taste test as a laboratory measure of food intake. We made use of data from over 2500 participants across 31 published laboratory studies from three research groups in the UK and Australia that have used the taste test paradigm. To assess validity we examined whether the taste test was sensitive to manipulations hypothesized to decrease or increase food intake and the extent to which participant level characteristics reliably associated with food intake in other paradigms predicted taste test food intake. By finding that the taste test was sensitive to experimental manipulation and all variables identified as being reliably associated with food intake in other paradigms (hunger, sex, liking of food) were associated with taste test food intake, we provide evidence for the validity of the taste test. When examining other participant level characteristics that tend not to be reliably associated with food intake in other paradigms, we found less consistent results; neither BMI or trait dietary restraint were reliably associated with taste test food intake, although trait tendencies to over-eat in response to palatable food cues were predictive of taste test food intake.

Is the taste test sensitive to experimental manipulation?

We found that experimental manipulations hypothesized to increase taste test food intake were associated with increased consumption, and manipulations hypothesized to decrease food intake were associated with reduced taste test food intake. In both instances, the overall effects of the experimental manipulations on taste test food intake were statistically small. Moreover, although a statistically significant predictor of food intake in unadjusted analyses, the effect of manipulations hypothesized to increase food intake on taste test intake was not statistically significant in an adjusted analysis with a smaller sample size. These relatively small effects are perhaps not too surprising because these manipulations were only

431	hypothesized to increase food intake. For example, in Robinson et al., (2014a) a condition
432	was hypothesized to increase food intake because it would make participants feel less self-
433	aware, but the manipulation did not successfully alter self-awareness. Unsurprisingly taste
434	test food intake was also unaffected in this study. The present analyses alongside a range of
435	other studies (Conger, et al., 1980; Oldham-Cooper, Hardman, Nicoll, Rogers, & Brunstrom,
436	2010; Van Strien et al., 2013) indicate that the taste test is a sensitive enough measure to be
437	able to examine the causal effect of a manipulated variable on food intake.
438	
439	Hunger and taste test food liking
440	In the present analyses we found that hungry participants tended to eat more during the taste
441	test than did less hungry participants, and that the extent to which participants liked the food
442	used in the taste test positively predicted food intake. We observed these results in our
443	unadjusted analyses and in an analysis which included other participant level predictors of
444	taste test food intake. We found this pattern of results irrespective of the country (UK vs.
445	Australia) that studies were conducted in, although there was a tendency for baseline hunger
446	to be more strongly associated with taste test food intake in studies conducted in the UK. This
447	result was not predicted and could reflect differences between UK and Australian study
448	methodologies. Overall, these findings are in line with other research which has shown that
449	hunger (Sadoul, et al., 2014) and food liking (de Graaf, et al., 2005) are predictors of food
450	intake, and thus confirm the construct validity of the taste test.
451	
452	Sex
453	In a small sub-analysis we also examined whether there are sex differences in taste test food
454	intake. Based on the notion that men have a higher energy need than women (Rolls, et al.,
455	1991), we hypothesized that men would consume significantly more than women in the taste

test. In line with this hypothesis, men consumed significantly more than women and this was a small to medium sized effect. This result is in support of the taste test having good construct validity.

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

456

457

458

Trait eating behaviour measures

We found evidence that self-reported over-eating in response to palatable food cues predicted food intake in the taste test, whereby participants with a greater tendency to overeat in response to palatable foods consumed significantly more in the taste test than participants with lower scores. However, this association was dependent on the measure used, whereby responses on the TFEQ disinhibition subscale (Stunkard & Messick, 1985), but not DEBQ external eating subscale (Van Strien, et al., 1986) were reliable predictors of taste test food intake. The present finding may reflect that the items on the DEBQ external eating subscale tend to ask participants about the influence that external cues have on stimulating overeating, whereas the TFEQ disinhibition subscale is a more general measure of 'overeating' or loss of control over eating (e.g. scale item: 'Sometimes when I start eating, I just can't seem to stop'). This may results in it being more predictive of taste test food intake because taste test procedures promote initial consumption of food in order to complete taste ratings. We found little evidence that trait dietary restraint predicted taste test food intake. In an unadjusted analysis, there was a very small (r = -.05, p = .04) negative association between restraint and food intake that was close to the threshold for statistical significance. However, in the adjusted analysis this association was no longer statistically significant (p = .80) and was close to zero. Restraint was also correlated with other participant level characteristics that did significantly predict taste test intake which indicates that the small unadjusted association between restraint and taste test food intake may have been caused by confounding. Although we made a tentative hypothesis that dietary restraint would be

associated with lower taste test food intake, other studies outside of the laboratory have suggested that there is a lack of reliable relationship between dietary restraint and energy intake (Johnson, Pratt, & Wardle, 2012; Stice, et al., 2010). However, in the context of a laboratory taste test the association between dietary restraint and food intake may be determined by the extent to which a test food is perceived as being 'forbidden' by a participant. This is a hypothesis we were not able to test in the present study. Moreover, in line with restraint theory (Herman & Mack, 1975), dietary restraint may interact with certain types of experimental manipulation to predict taste test food intake, rather than having a direct association with intake as was tested in the present study. Thus, more sophisticated tests of when dietary restraint does/does not predict food intake may uncover an association between dietary restraint and taste test food intake.

BMI

We found no evidence of a significant relationship between BMI and taste test food intake, irrespective of whether this relationship was examined with BMI as a continuous variable or when BMI was grouped according to weight status (e.g. normal weight, overweight, obese). We had predicted that there would be a positive association because a higher BMI should be associated with a larger energy intake requirement. Both Acosta et al. (2015) and Meyer-Gerspach et al. (2014) report data which indicates that participants with severe obesity have a higher energy intake in the laboratory than participants with normal weight. In the present study we had relatively few participants with obesity and most were of class I obesity (30-34.9 kg/m²). Thus, we may have found a relationship between BMI and taste test food intake if we had a wider BMI range in the present study. In the context of a taste test it is also plausible that individuals of heavier body weight do not eat more than their slimmer counterparts because overconsumption of the foods commonly used in taste tests (high

calorie snack food) may invoke self-presentation concerns. Moreover, there is some debate whether individuals of heavier BMI eat larger meal sizes in the real world and it has instead been argued that eating frequency may be more reliably associated with BMI (Mattes, 2014). Thus, the lack of association between BMI and taste test food intake in the present study may reflect this.

Limitations and Methodological Considerations

The present project involved participant level data and because of this it was not feasible to review and analyze data from all published studies that have adopted the taste test. Thus, it is important to note that our conclusions are based on findings from three research groups. However, we did make use of a relatively large number of studies that had been conducted in two countries and this increases confidence in the generalizability of our findings. A limitation of the present study was that a lack of data from male participants resulted in our main analysis being limited to young women. Although a smaller sub-analysis showed that the taste test is sensitive to sex differences in food intake, we do not know whether our results regarding the sensitivity of the taste test to experimental manipulations and participant level predictors of taste test food intake apply to men. We are not aware of any convincing rationale why for example, taste test food intake in men would not be predicted by baseline hunger, but further work assessing the validity of the taste test in male samples would be informative.

Based on our findings we recommend that the use of the taste test in laboratory eating behaviour research to identify that affect food intake is valid. However, there are caveats to this recommendation. Given that baseline hunger and taste test food liking predicted food intake in our analyses, ensuring that these variables are standardized and/or measured in taste test studies is recommended. All of the included studies in the present analyses adopted cover

stories to attempt to ensure that participants were not aware of the aims of the study or experimental hypotheses. It has been shown in a number of studies that when participants believe their food intake is being measured this tends to affect the amount of food they eat (Robinson et al, 2015). Thus, we would argue that studies which adopt the taste test should a) attempt to ensure that participants are unaware of study hypotheses and b) attempt to conceal that food intake is being measured. The present studies also all used between-subjects designs, as opposed to participants attending several laboratory sessions, being exposed to different manipulations and completing multiple taste tests. Thus, our conclusions are limited to between-subjects designs. It is feasible that with repeated use of the taste test (e.g. a crossover design) the purpose of the taste test may become more apparent to a participant. A final point is that the predictor variables in our analyses combined explained only 12.5% of taste test food intake. Thus, identifying and understanding other factors that explain how much participants consume during a taste test would now be of interest.

545 Conclusions

The results of our analyses indicate that the bogus taste test is likely to be a valid measure of food intake and can be used to identify whether experimental manipulations have a causal effect on food intake.

Funding Funding

This research received no external funding. ER's salary was part funded by the Medical Research Council and the Economic Social Research Council. CAH and ER have received research funding from the American Beverage Association.

Declaration of Conflicting Interests 556 The authors declare that they had no conflicts of interest with respect to their authorship or 557 the publication of this article. 558 559 560 References Acosta, A., Camilleri, M., Shin, A., Vazquez-Roque, M. I., Iturrino, J., Burton, D., ... & 561 Zinsmeister, A. R. (2015). Quantitative gastrointestinal and psychological traits 562 563 associated with obesity and response to weight-loss therapy. Gastroenterology, 148(3), 537-546. 564 Andrade, A. M., Kresge, D. L., Teixeira, P. J., Baptista, F., & Melanson, K. J. (2012). Does 565 eating slowly influence appetite and energy intake when water intake is controlled? 566 Int J Behav Nutr Phys Act, 9, 135. 567 Bell, E. A., & Rolls, B. J. (2001). Energy density of foods affects energy intake across 568 multiple levels of fat content in lean and obese women. Am J Clin Nutr, 73(6), 1010-569 570 1018. Bem, D. J. (1972). Self-perception theory. Advances in Experimental Social Psychology, 6, 1-571 572 573 Berg, C., Lappas, G., Wolk, A., Strandhagen, E., Torén, K., Rosengren, A., ... & Lissner, L. (2009). Eating patterns and portion size associated with obesity in a Swedish 574 population. Appetite, 52(1), 21-26. 575 Berteus Forslund, H., Torgerson, J. S., Sjostrom, L., & Lindroos, A. K. (2005). Snacking 576 frequency in relation to energy intake and food choices in obese men and women 577 compared to a reference population. Int J Obes Relat Metab Disord, 29(6), 711-719. 578 Blodorn, A., Major, B., Hunger, J., & Miller, C. (2016). Unpacking the psychological weight 579 of weight stigma: A rejection-expectation pathway. Journal of Experimental Social 580 Psychology, 63, 69-76. 581 Blundell, J., De Graaf, C., Hulshof, T., Jebb, S., Livingstone, B., Lluch, A., . . . Van Der 582 Knaap, H. (2010). Appetite control: methodological aspects of the evaluation of 583 foods. *Obesity Reviews*, 11(3), 251-270. 584 Bongers, P., & Jansen, A. (2016). Emotional Eating Is Not What You Think It Is and 585 Emotional Eating Scales Do Not Measure What You Think They Measure. Frontiers 586 in Psychology, 7, 1932. 587 Brunstrom, J. M., & Shakeshaft, N. G. (2009). Measuring affective (liking) and non-affective 588 589 (expected satiety) determinants of portion size and food reward. Appetite, 52(1), 108-590 Bryant, E. J., King, N. A., & Blundell, J. E. (2008). Disinhibition: its effects on appetite and 591 weight regulation. Obesity Reviews, 9(5), 409-419. 592 Cavanagh, K., Vartanian, L. R., Herman, C. P., & Polivy, J. (2013). The effect of portion size 593 on food intake is robust to brief education and mindfulness exercises. Journal of 594 595 Health Psychology, 19(6), 730-739. *Christiansen, P., Rose, A., Randall-Smith, L., & Hardman, C. A. (2016). Alcohol's acute 596 effect on food intake is mediated by inhibitory control impairments. Health 597 598 Psychology, 35(5), 518-522.

- Conger, J. C., Conger, A. J., Costanzo, P. R., Wright, K. L., & Matter, J. A. (1980). The effect of social cues on the eating behavior of obese and normal subjects. *J Pers*, 48(2), 258-271.
- de Castro, J. M. (2000). Eating behavior: lessons from the real world of humans. *Nutrition*, 16(10), 800-813.
- de Castro, J. M., & Elmore, D. K. (1988). Subjective hunger relationships with meal patterns in the spontaneous feeding behavior of humans: Evidence for a causal connection. *Physiology & Behavior*, 43(2), 159-165.
- de Castro, J. M., King, G. A., Duarte-Gardea, M., Gonzalez-Ayala, S., & Kooshian, C. H. (2012). Overweight and obese humans overeat away from home. *Appetite*, 59(2), 204-609 211.
- de Graaf, C., Kramer, F. M., Meiselman, H. L., Lesher, L. L., Baker-Fulco, C., Hirsch, E. S.,
 & Warber, J. (2005). Food acceptability in field studies with US army men and
 women: relationship with food intake and food choice after repeated exposures.
 Appetite, 44(1), 23-31.
- Drewnowski, A., & Hann, C. (1999). Food preferences and reported frequencies of food consumption as predictors of current diet in young women. *The American Journal of Clinical Nutrition*, 70(1), 28-36.
- Evers, C., de Ridder, D. T., & Adriaanse, M. A. (2009). Assessing yourself as an emotional eater: Mission impossible? *Health Psychology*, 28(6), 717.
- Evers, C., Stok, F. M., Danner, U. N., Salmon, S. J., de Ridder, D. T., & Adriaanse, M. A. (2011). The shaping role of hunger on self-reported external eating status. *Appetite*, 57(2), 318-320.
- French, S. A., Epstein, L. H., Jeffery, R. W., Blundell, J. E., & Wardle, J. (2012). Eating behavior dimensions. Associations with energy intake and body weight. A review. *Appetite*, *59*(2), 541-549.
- Gibbons, C., Finlayson, G., Dalton, M., Caudwell, P., & Blundell, J. E. (2014). Metabolic
 phenotyping guidelines: studying eating behaviour in humans. *Journal of Endocrinology*, 222(2), G1-G12.
- Gregersen, N. T., Flint, A., Bitz, C., Blundell, J. E., Raben, A., & Astrup, A. (2008).
 Reproducibility and power of ad libitum energy intake assessed by repeated single meals. *Am J Clin Nutr*, 87(5), 1277-1281.
- Guerrieri, R., Nederkoorn, C., & Jansen, A. (2008). The interaction between impulsivity and a varied food environment: its influence on food intake and overweight. *Int J Obes* (*Lond*), 32(4), 708-714.
- *Hardman, C. A., Rogers, P. J., Dallas, R., Scott, J., Ruddock, H. K., & Robinson, E. (2015).

 "Food addiction is real". The effects of exposure to this message on self-diagnosed food addiction and eating behaviour. *Appetite*, *91*, 179-184.
- Harris, J. L., Bargh, J. A., & Brownell, K. D. (2009). Priming effects of television food advertising on eating behavior. *Health Psychology*, 28(4), 404.
- *Haynes, A., Kemps, E., Moffitt, R., & Mohr, P. (2014). Resisting temptation of unhealthy food: interaction between temptation-elicited goal activation and self-control.

 *Motivation and Emotion, 38(4), 485-495.
- *Haynes, A., Kemps, E., Moffitt, R., & Mohr, P. (2015a). Reduce temptation or resist it? Experienced temptation mediates the relationship between implicit evaluations of unhealthy snack foods and subsequent intake. *Psychology & Health*, 30(5), 534-550
- *Haynes, A., Kemps, E., & Moffitt, R. (2015b). Inhibitory self-control moderates the effect of changed implicit food evaluations on snack food consumption. *Appetite*, *90*, 114-122.

- *Haynes, A., Kemps, E., & Moffitt, R. (2015c). The moderating role of state inhibitory control in the effect of evaluative conditioning on temptation and unhealthy snacking.

 *Physiology & Behavior, 152, 135-142.
- *Haynes, A., Kemps, E., & Moffitt, R. (2016). Too Depleted to Try? Testing the Process Model of Ego Depletion in the Context of Unhealthy Snack Consumption. *Appl Psychol Health Well Being*, 8(3), 386-404.
- Hebert, J. R., Clemow, L., Pbert, L., Ockene, I. S., & Ockene, J. K. (1995). Social desirability
 bias in dietary self-report may compromise the validity of dietary intake measures.
 International Journal of Epidemiology, 24(2), 389-398.
- Heitmann, B. L., & Lissner, L. (1995). Dietary underreporting by obese individuals--is it specific or non-specific? *BMJ*, *311*(7011), 986-989.
- 659 Herman, C. P., & Mack, D. (1975). Restrained and unrestrained eating. *J Pers*, *43*(4), 647-660 660.
- *Higgs, S. (2002). Memory for recent eating and its influence on subsequent food intake.

 *Appetite, 39(2), 159-166.
- *Higgs, S. (2015). Manipulations of attention during eating and their effects on later snack intake. *Appetite*, *92*, 287-294.
- *Higgs, S., & Donohoe, J. E. (2011). Focusing on food during lunch enhances lunch memory and decreases later snack intake. *Appetite*, *57*(1), 202-206.
- *Higgs, S., & Jones, A. (2013). Prolonged chewing at lunch decreases later snack intake.

 *Appetite, 62, 91-95.
- *Higgs, S., Williamson, A. C., & Attwood, A. S. (2008). Recall of recent lunch and its effect on subsequent snack intake. *Physiology & Behavior*, *94*(3), 454-462.

671

672673

674

675

676

689

- Hill, A. J., Rogers, P. J., & Blundell, J. E. (1995). Techniques for the experimental measurement of human eating behaviour and food intake: a practical guide. *Int J Obes Relat Metab Disord*, *19*(6), 361-375.
- Hooper, N., Sandoz, E. K., Ashton, J., Clarke, A., & McHugh, L. (2012). Comparing thought suppression and acceptance as coping techniques for food cravings. *Eat Behav*, 13(1), 62-64.
- Horner, K. M., Byrne, N. M., & King, N. A. (2014). Reproducibility of subjective appetite ratings and ad libitum test meal energy intake in overweight and obese males.
 Appetite, 81, 116-122. d
- Houben, K. (2011). Overcoming the urge to splurge: influencing eating behavior by manipulating inhibitory control. *J Behav Ther Exp Psychiatry*, 42(3), 384-388.
- Johnson, F., Pratt, M., & Wardle, J. (2012). Dietary restraint and self-regulation in eating behavior. *International Journal of Obesity*, *36*(5), 665-674.
- Johnson, F., Pratt, M., & Wardle, J. (2012). Dietary restraint and self-regulation in eating behavior. *International Journal of Obesity*, *36*(5), 665-674.
- Jones, A., Button, E., Rose, A. K., Robinson, E., Christiansen, P., Di Lemma, L., & Field, M. (2016). The ad-libitum alcohol 'taste test': secondary analyses of potential confounds and construct validity. *Psychopharmacology*, 233, 917-924.
 - *Kakoschke, N., Kemps, E., & Tiggemann, M. (2014). Attentional bias modification encourages healthy eating. *Eating behaviors*, 15(1), 120-124.
- *Kakoschke, N., Kemps, E., & Tiggemann, M. (2015). Combined effects of cognitive bias for food cues and poor inhibitory control on unhealthy food intake. *Appetite*, 87, 358-364.
- *Kemps, E., Herman, C. P., Hollitt, S., Polivy, J., Prichard, I., & Tiggemann, M. (2016a).

 Contextual cue exposure effects on food intake in restrained eaters. *Physiology & Behavior*, *167*, 71-75.

- *Kemps, E., Herman, C. P., Hollitt, S., Polivy, J., Prichard, I., & Tiggemann, M. (2016b).

 The role of expectations in the effect of food cue exposure on intake. *Appetite*, 103, 259-264.
- *Kemps, E., Tiggemann, M., & Elford, J. (2015). Sustained effects of attentional re-training
 on chocolate consumption. *Journal of Behavior Therapy and Experimental Psychiatry*, 49, 94-100.
 - *Kemps, E., Tiggemann, M., Orr, J., & Grear, J. (2014). Attentional retraining can reduce chocolate consumption. *J Exp Psychol Appl*, 20(1), 94-102.
 - Kopelman, P. (2007). Health risks associated with overweight and obesity. *Obesity Reviews*, 8(s1), 13-17.
- Macdiarmid, J., & Blundell, J. (1998). Assessing dietary intake: Who, what and why of under-reporting. *Nutrition Research Reviews*, 11(2), 231-253.

703

704

705706

728 729

734735

- 709 Martin, C. K., Williamson, D. A., Geiselman, P. J., Walden, H., Smeets, M., Morales, S., & Redmann, S., Jr. (2005). Consistency of food intake over four eating sessions in the laboratory. *Eating Behaviors*, *6*(4), 365-372.
- Mattes, R. (2014). Energy intake and obesity: ingestive frequency outweighs portion size. *Physiol Behav*, *134*, 110-118.
- Meiselman, H. L. (1992). Methodology and theory in human eating research. *Appetite*, *19*(1), 49-55.
- Meyer-Gerspach, A. C., Wölnerhanssen, B., Beglinger, B., Nessenius, F., Napitupulu, M.,
 Schulte, F. H., ... & Beglinger, C. (2014). Gastric and intestinal satiation in obese and
 normal weight healthy people. *Physiology & Behavior*, 129, 265-271.
- Nichols, A. L., & Maner, J. K. (2008). The good-subject effect: investigating participant demand characteristics. *Journal of General Psychology*, *135*(2), 151-165.
- Oldham-Cooper, R. E., Hardman, C. A., Nicoll, C. E., Rogers, P. J., & Brunstrom, J. M. (2010). Playing a computer game during lunch affects fullness, memory for lunch, and later snack intake. *The American Journal of Clinical Nutrition*.
- Orne, M. T. (1962). On the social psychology of the psychological experiment: With particular reference to demand characteristics and their implications. *American Psychologist*, 17(11), 776.
- Prentice, A. M. (2001). Overeating: the health risks. *Obesity Research*, 9(S11), 234S-238S.
 - *Robinson, E., & Field, M. (2015). Awareness of social influence on food intake. An analysis of two experimental studies. *Appetite*, 85, 165-170.
- Robinson, E., Hardman, C. A., Halford, J. C., & Jones, A. (2015). Eating under observation: a systematic review and meta-analysis of the effect that heightened awareness of observation has on laboratory measured energy intake. *American Journal of Clinical Nutrition*, 102(2), 324-337.
 - *Robinson, E., Kersbergen, I., Brunstrom, J. M., & Field, M. (2014a). I'm watching you. Awareness that food consumption is being monitored is a demand characteristic in eating-behaviour experiments. *Appetite*, 83(0), 19-25.
- *Robinson, E., Kersbergen, I., & Higgs, S. (2014b). Eating 'attentively' reduces later energy consumption in overweight and obese females. *British Journal of Nutrition*, 112(4), 657-661.
- *Robinson, E., Oldham, M., Cuckson, I., Brunstrom, J. M., Rogers, P. J., & Hardman, C. A.
 (2016). Visual exposure to large and small portion sizes and perceptions of portion size normality: Three experimental studies. *Appetite*, *98*, 28-34.
- Robinson, E., Proctor, M., Oldham, M., & Masic, U. (2016). The effect of heightened awareness of observation on consumption of a multi-item laboratory test meal in females. *Physiology & Behavior*, *163*, 129-135.

- *Robinson, E., Sharps, M., Price, N., & Dallas, R. (2014c). Eating like you are overweight: the effect of overweight models on food intake in a remote confederate study. *Appetite*, 82, 119-123.
- Rolls, B. J., Fedoroff, I. C., & Guthrie, J. F. (1991). Gender differences in eating behavior and body weight regulation. *Health Psychology*, *10*(2), 133.
- *Rose, A. K., Hardman, C. A., & Christiansen, P. (2015). The effects of a priming dose of alcohol and drinking environment on snack food intake. *Appetite*, *95*, 341-348.
- Roth, D. A., Herman, C. P., Polivy, J., & Pliner, P. (2001). Self-presentational conflict in social eating situations: a normative perspective. *Appetite*, *36*(2), 165-171.

755

756

757

758

759

760

761

762

763

764 765

766

767 768

769

770 771

772

773

774

775

776

- *Ruddock, H. K., Christiansen, P., Jones, A., Robinson, E., Field, M., & Hardman, C. A. (2016). Believing in food addiction: Helpful or counterproductive for eating behavior? *Obesity*, 24(6), 1238-1243.
- Sadoul, B. C., Schuring, E. A., Mela, D. J., & Peters, H. P. (2014). The relationship between appetite scores and subsequent energy intake: an analysis based on 23 randomized controlled studies. *Appetite*, *83*, 153-159.
- Schoeller, D. A. (1990). How accurate is self-reported dietary energy intake? *Nutrition Research Reviews*, 48(10), 373-379.
- Schoeller, D. A., Thomas, D., Archer, E., Heymsfield, S. B., Blair, S. N., Goran, M. I., . . . Allison, D. B. (2013). Self-report-based estimates of energy intake offer an inadequate basis for scientific conclusions. *American Journal of Clinical Nutrition*, 97(6), 1413-1415.
- *Schumacher, S. E., Kemps, E., & Tiggemann, M. (2016). Bias modification training can alter approach bias and chocolate consumption. *Appetite*, *96*, 219-224.
- Spanos, S., Kenda, A. S., & Vartanian, L. R. (2015). Can serving-size labels reduce the portion-size effect? A pilot study. *Eating Behaviors*, 16, 40-42.
- Sproesser, G., Schupp, H. T., & Renner, B. (2013). The bright side of stress-induced eating eating more when stressed but less when pleased. *Psychological Science*.
- Stice, E., Sysko, R., Roberto, C. A., & Allison, S. (2010). Are Dietary Restraint Scales Valid Measures of Dietary Restriction? Additional Objective Behavioral and Biological Data Suggest Not. *Appetite*, *54*(2), 331-339.
- Stroebe, W., van Koningsbruggen, G. M., Papies, E. K., & Aarts, H. (2013). Why most dieters fail but some succeed: A goal conflict model of eating behavior. *Psychological Review*, *120*(1), 110-138.
- Stunkard, A. J., & Messick, S. (1985). The three-factor eating questionnaire to measure
 dietary restraint, disinhibition and hunger. *Journal of Psychosomatic Research*, 29(1),
 71-83.
- 782 Trichopoulou, A., Gnardellis, C., Lagiou, A., Benetou, V., & Trichopoulos, D. (2000). Body 783 mass index in relation to energy intake and expenditure among adults in Greece. 784 *Epidemiology*, 11(3), 333-336.
- van Strien, T., Cebolla, A., Etchemendy, E., Gutierrez-Maldonado, J., Ferrer-Garcia, M.,
 Botella, C., & Banos, R. (2013). Emotional eating and food intake after sadness and
 joy. *Appetite*, 66, 20-25.
- Van Strien, T., Frijters, J. E., Bergers, G., & Defares, P. B. (1986). The Dutch Eating
 Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external
 eating behavior. *International Journal of Eating Disorders*, 5(2), 295-315.
- Wardle, J., Steptoe, A., Oliver, G., & Lipsey, Z. (2000). Stress, dietary restraint and food intake. *Journal of Psychosomatic Research*, 48(2), 195-202.
- Welch, R. W., Antoine, J. M., Berta, J. L., Bub, A., de Vries, J., Guarner, F., . . . Woodside, J.
 V. (2011). Guidelines for the design, conduct and reporting of human intervention

	ACCEPTED MANUSCRIPT
795 796	, 11
797	•
798	Can(not) take my eyes off it: attention bias for food in overweight participants. Health
799	
800	
801	11 U
802	women. Appetite, 69, 102-107.
803	
804	