



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

An updated view of hypothalamic-vascular-pituitary unit function and plasticity

Citation for published version:

Le Tissier, P, Campos, P, Lafont, C, Romanò, N, Hodson, DJ & Mollard, P 2016, 'An updated view of hypothalamic-vascular-pituitary unit function and plasticity' *Nature reviews endocrinology*. DOI: 10.1038/nrendo.2016.193

Digital Object Identifier (DOI):

[10.1038/nrendo.2016.193](https://doi.org/10.1038/nrendo.2016.193)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Nature reviews endocrinology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



An updated view of hypothalamic–vascular–pituitary unit function and plasticity

Paul Le Tissier¹, Pauline Campos^{2,3,4}, Chrystel Lafont^{2,3,4}, Nicola Romanò¹, David J. Hodson^{5,6} and Patrice Mollard^{2,3,4}

¹Centre for Integrative Physiology, University of Edinburgh, George Square, Edinburgh, EH8 9XD, UK. ²CNRS, UMR-5203, Institut de Génomique Fonctionnelle, rue de la Cardonille, F-34000 Montpellier, France; ³INSERM, U661, rue de la Cardonille, F-34000 Montpellier, France; ⁴Université de Montpellier, rue de la Cardonille, UMR-5203, F-34000 Montpellier, France; ⁵Institute of Metabolism and Systems Research and Centre of Membrane Proteins and Receptors (COMPARE), University of Birmingham, Edgbaston, B15 2TT, UK. ⁶Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, B15 2TH, UK.

Correspondence to

Paul.LeTissier@ed.ac.uk

patrice.mollard@igf.cnrs.fr

Abstract

The discoveries of novel functional adaptations of the hypothalamus and anterior pituitary gland for physiological regulation have transformed our understanding of their interaction. The activity of a small proportion of hypothalamic neurons can control complex hormonal signalling, which is disconnected from a simple stimulus and subsequent hormone secretion relationship and is dependent on physiological status. The interrelationship of the terminals of hypothalamic neurons and pituitary cells with the vasculature has an important role in determining the pattern of neurohormone exposure. Cell in the pituitary gland form networks with distinct organizational motifs that are related to the duration and pattern of output. modification of these occurs in different physiological states, can persist after cessation of demand and results in enhanced function. Consequently, the hypothalamus and pituitary can no longer be considered as having a simple stratified relationship: with the vasculature they form a tripartite system, which must function in concert for appropriate hypothalamic regulation of physiological processes, for example reproduction. An improved understanding of the mechanisms underlying these regulatory features has implications for current and future therapies that correct defects in the hypothalamic–pituitary axes. In addition, recapitulating proper network organization will be an important challenge for regenerative stem cell treatment.

[H1] Introduction

To maximize reproductive success, via the appropriate timing of ovulation, lactation, or body growth, the outputs of several hypothalamic–pituitary axes are dramatically altered. These adaptive changes occur over differing time scales, with varying frequencies and levels of predictability. For example, the occurrence of the increase in growth hormone (GH) output at puberty is largely predictable. On a relatively short time scale (days), the surge in luteinizing hormone (LH) secretion required for oestrus is an acute change that occurs regularly once every reproductive cycle and, in humans, continues for years in the absence of pregnancy. On a longer time scale (months to years depending on the species), the increase in prolactin required for lactation is maintained for a variable time (which depends on when offspring are weaned) and recurs at each pregnancy, but is unpredictable before gestation. These large changes in the output of the various pituitary axes require modification of both hypothalamic and pituitary function, but whether this effect is reversed on cessation of physiological demand likely reflects the expectation that increased output will recur. A mechanistic understanding of these alterations in hypothalamic–pituitary function is fundamental to interpret and treat defects that lead to endocrine diseases resulting from hormone deficiencies (for example, dwarfism) or excess (such as polycystic ovarian syndrome). In this Review, we will focus on three pituitary axes that have roles in driving changes in physiology; the gonadatropin, prolactin and GH axes. The level of our understanding varies for each of these axes and the features that might serve as general principles of hypothalamic-pituitary and other endocrine organ function will be highlighted in the text.

[H1] Beyond stimulus-secretion coupling

The path of communication between the brain and peripheral tissues is exemplified by muscle contraction, which requires the transfer of electrical signals from axons via the neuromuscular junction. This sequence of events, known as excitation-contraction coupling¹, takes <1 s in mammals and adapts to altered functional demand. Similarly, in the hypothalamus just a few thousand neurons can also send signals to the periphery, in this case towards the median eminence (ME) via a specialized neurohaemal junction. In this junction, nerve terminal depolarization, either originating from the perikarya² or the terminals themselves³, allows the sufficiently rapid entry of calcium ions to trigger exocytosis of neurohormones towards the first

loop of the portal fenestrated capillaries^{4,5}. This rapid (<1 s) sequence of events was termed ‘stimulus-secretion coupling’ due to the clear similarities with excitation-contraction coupling^{5,6}. Soon after release, neurohormones pervade the second loop of fenestrated capillaries within the pituitary gland, before binding to cognate receptors on endocrine cells to induce pituitary hormone exocytosis through a second ‘stimulus-secretion coupling’ event (FIG. 1)⁷⁻¹⁰.

In the second half of the twentieth century (and building on Harris’ work on the hypothalamus–pituitary axis¹¹) the analogy between excitation-contraction and stimulus-secretion coupling was developed further¹². However, important and fundamental differences exist between the two processes. Specifically, in the hypothalamus, endogenous rhythms with axis-dependent frequencies exist and the time scale for pituitary hormone release is measured in minutes to several hours¹³. Nevertheless, the analogy with neural control of locomotor activity led to a generally accepted model of hypothalamic regulation of pituitary function (FIG. 1).. Specifically, the excitation of specific hypothalamic neuron populations, determined by higher brain centres and peripheral feedback, is relayed as an unmodified series of signals to drive balanced pituitary hormone output¹¹. The release of neurohormones and subsequent transportation and the effects on target cells were considered to be passive events in the regulation of pituitary hormone secretion, with only variation in the number of endocrine cells seeming to affect response levels^{11,14}. Similarly, the alterations in gene expression and cell proliferation, which support maintenance of hormone output, were simply considered a correlated response to hypothalamic regulation of secretion¹⁵.

In the early twenty-first century, a series of paradigm shifts in our understanding of the hypothalamic–pituitary system was established as a consequence of newly developed tools and techniques, including calcium imaging, fluorescent protein identification of specific cell types and two-photon microscopy, for use in genetically modified mice (eg¹⁶⁻¹⁸). The use of these methods have shown that both the pituitary gland and portal system can no longer be considered as static structures simply responding to neurohormonal regulation (BOX 1), although they have confirmed the pre-eminence of the hypothalamus as the driver of pituitary hormone output. In addition, hypothalamic neuron function has been found to be more dynamic than

initially thought, which might contribute to modifications in its regulation of the pituitary under different physiological states (FIG. 1)².

[H1] Gonadotroph axis

The reproductive system is critically dependent upon pulsatile secretion of gonadotrophin-releasing hormone (GnRH) and LH; however, the understanding of pulse generation has been hampered by the complexity of the regulatory mechanisms, many of which will clearly be lost in *in vitro* preparations. Investigators working in the late 1980s using pituitary portal bleeding and microdialysis documented the pulsatile nature of GnRH release into the portal vasculature of the sheep, monkey and rat¹⁹⁻²⁵, and showed a strong correlation between GnRH and LH pulses^{22,26,27}. However, the scattered distribution and relative paucity of GnRH cell bodies limited the investigation of the cellular events that lead to pulsatile secretion of LH *in vivo*. In the past few years, the development of optogenetic techniques in rats and mice and an ultrasensitive ELISA capable of measuring LH levels in whole blood microsamples²⁸ has enabled investigators to dissect the GnRH neuron excitation parameters that generate LH pulses²⁹. In these studies, the stimulation of just 60 GnRH neurons (~5% of the GnRH neurons) can trigger short-lived increases in LH secretion that resemble endogenous pulses²⁹. Given the critical importance of GnRH neurons to the survival of all mammalian species, a degree of functional redundancy within this cell population is expected. This finding is consistent with studies in which just 10% of the GnRH neuron population is sufficient to maintain pulsatile LH secretion^{30,31}. Varying the timing and frequency of stimulation has demonstrated that a brief (2 min) optogenetic stimulation at high frequency (10 Hz) evokes an LH pulse, whereas shorter periods and lower frequencies cannot elicit LH output that resembles endogenous pulses²⁹. A bursting pattern of stimulation, which had been assumed to be most effective for pulse generation and the focus of many previous studies³², also failed to increase LH secretion. Whether such a stimulatory signal exists endogenously *in vivo* and where its origin might be is unknown, although a ‘GnRH pulse driver’ might be located in the mediobasal–hypothalamus, specifically at the level of neurons co-expressing kisspeptin, neurokinin B, as well as dynorphin A (so-called ‘KNDy’ neurons)^{33,34}.

The generation of GnRH pulses, requires a coordinated release from multiple neurons through the synchronization of GnRH neuron population excitation. The cell bodies

of GnRH neurons are scattered throughout the basal forebrain, while their projections have dendrodendritic bundling and/or shared synapses, and become highly concentrated around the ME³⁵. Fascinatingly, these projections simultaneously receive and integrate synaptic inputs — they possess both axonal and dendritic characteristics, leading to their description as ‘dendrons’, before finally acquiring an axonal morphology within the ME and ramifying into numerous terminals that appose blood vessels³⁶. Dendrons might be an ideal location for putative afferent axons to modulate the excitability of multiple GnRH neuron dendrites, and for multiple GnRH neurons to align their firing pattern, which thereby provides a potential mechanism for their synchronized activity directly in the mediobasal hypothalamus³⁶. An additional source of pulse synchronization is in the ME, where hypophysiotropic GnRH neurons terminate within the external zone close to endothelial cells of the portal vasculature³⁷. Endothelial cells in the ME might modulate GnRH release through nitric oxide secretion³⁸⁻⁴⁰ (reviewed elsewhere⁴¹). At the ME, nitric oxide is spontaneously released from an endothelial source and follows a pulsatile and cyclic pattern of secretion³⁸, and inhibition of nitric oxide synthesis within the ME can disrupt reproductive cyclicity³⁹. Conversely, in the GnRH neuron perikarya, basal nitric oxide synthase activity might provide the tonic inhibition of the GnRH neural system required to maintain nadir levels of LH⁴⁰.

Once released into the ME, the transport of GnRH to the pituitary, and the pattern of gonadotroph exposure to the neurohormone, have been largely assumed to represent a simple linear process²². However, the use of fluorescent tracking using 4 kDa dextran, which mimics the size of most hypothalamic neurohormones, has shown that the diffusion processes, both at the level of the ME and the pituitary capillaries, are complex and non-linear⁸. Consequently, the portal vessel network might function as a ‘physical integrator’, enabling neurohormones to be transferred from the ME to the gonadotroph within a few seconds⁴². Once in the blood stream, the moderately rapid clearance rate (which varies between species) of LH generates the specific asymmetric pulse shape of this hormone, which is characterized by a fast increase immediately followed by a slower decrease²⁸. Importantly, a faithful delivery of the pulsatile pattern of GnRH secretion to the pituitary is crucial for gonadotroph function⁴³⁻⁴⁶. For example, high GnRH pulse frequencies (>1 pulse per h) activate LH production, whereas low frequencies (<1 pulse per 2–3 h) preferentially induce

follicle-stimulating hormone (FSH) synthesis and release⁴⁷. Overall, the intricate relationships between pulsatile GnRH release, secretory competency of the pituitary gonadotrophs and regulatory mechanisms within the vasculature generate the rhythmic fluctuations in LH secretion.

[H3] GnRH and LH surge generation

The GnRH/LH surge mechanism is sex specific and normally occurs only in women^{48,49}. During the oestrous cycle, increasing concentrations of plasma oestrogen alter feedback to the GnRH neuronal afferent networks and gonadotrophs from negative to positive to induce the gonadotrophin surge⁵⁰⁻⁵². That the oestrogen-responsive kisspeptin neurons in the rostral periventricular area of the third ventricle have a critical role in enabling ovulation in rodents by activating GnRH neurons is now well accepted⁵³. Importantly, the relative contribution of the hypothalamic and pituitary levels to the oestrogen-induced gonadotropin surge seems to be species-dependent, with the latter the predominant mechanism in human and non-human primates⁵⁴. In the female sheep, the GnRH surge is composed of high-frequency pulsatile events superimposed on a constantly elevated level of GnRH release, although whether the surge is driven by a fundamentally altered pattern of GnRH secretion⁵⁵, or by a simple increase in the frequency of pulsatile secretion is unclear⁵⁶. This huge increase in GnRH secretion continues for a period of 24 h, considerably longer than the duration of the LH surge it induces, before returning to a strictly episodic pattern of release^{25,55,57}. The firing pattern of GnRH neurons needed to generate the GnRH/LH surge is unknown. However, it is reasonable to assume that the prolonged firing of an increased number of GnRH neurons is required for the secretion of surge levels, compared with that required for a pulse. Indeed, in transgenic mice with a dose-dependent reduction in GnRH neuron migration, 10% of the normal GnRH neuronal content failed to rescue ovulation, but cyclicity was restored when approximately 30% of the GnRH population was present³¹.

In addition to the putative change in GnRH population electrical activity, anatomical changes are found within the external zone of the ME where GnRH nerve terminals are ensheathed by tanycytes^{58,59} (Fig. 2). The cellular conformation changes with fluctuating oestrogen profiles throughout the oestrous cycle. For example, in rats, semaphorin-7a-dependent structural remodelling of tanycytes occurs during the

preovulatory surge, resulting in release of the engulfed axons and direct access of GnRH nerve terminals to the portal vasculature⁶⁰ (FIG. 2). By contrast, fenestrated rat endothelial cells of the hypothalamic–hypophyseal portal vessels release semaphorin-3A, which is thought to induce GnRH neuron axonal growth and sprouting within the ME as a function of the oestrous cycle⁶¹. These mechanisms are likely to enable the generation of high concentrations of GnRH, which evoke the GnRH/LH surge, to be released into the pituitary portal circulation^{62,63}.

Within the pituitary, the distinct network organization of gonadotrophs⁶⁴ and their large scale reorganization during puberty⁶⁵ suggests that communication mediated by cell contact between gonadotrophs might have a functional role in the regulation of gonadotrophin output (Box 1). However, this organisation has not been studied in the same detail as that of other pituitary axes to date. Although the dynamic gonadotroph responses at the time of the proestrous surge have not yet been described *in vivo*, sampling of pituitaries at single time points in rats and sheep suggest that changes occur in gonadotrophin subunit expression, granule distribution and GnRH receptor abundance^{66,67}. Sequencing of mRNA isolated from gonadotrophs of the anterior pituitary glands from female mice reveals that genes regulating the secretory process, blood pressure and cell adhesion were also enriched during proestrus⁶⁸. Likewise, immortalized cell lines and cells in pituitary slices increase their cellular movement following GnRH stimulation⁶⁹, and extend cellular processes and increase their cellular movement at puberty⁶⁵ (FIG 3 and Box 1). These findings suggest that changes in the relationship of the gonadotroph network with the vasculature might modify the secretory response of gonadotrophs⁷⁰.

[H3] Clinical relevance

The mechanisms that underlie both pulsatile secretion and surge generation of LH have important implications for the treatment of infertility in women. For example, polycystic ovarian syndrome (PCOS), the most common anovulatory cause of infertility⁷¹ affecting >100 million women worldwide, is associated with a dysregulation of the normal pattern of LH secretion⁷². Whether the origin of this multi-factorial disorder is at the level of the hypothalamic–pituitary axis is unknown⁷³, but PCOS is characterized by increases in GnRH pulse frequency and sensitivity of the pituitary gland to the neurohormone^{74,75}. Consequently, potential interventions that modify the dynamics of GnRH output, its transport to the ME or its

actions in the pituitary might have implications for the treatment of PCOS. This is also the case for congenital hypogonadotropic hypogonadism, which results from a pituitary or a hypothalamic defect with or without anosmia⁷⁶. Several novel gene mutations that are associated with this disorder have been identified, including those encoding neuropeptides (such as kisspeptin), transcription factors (chromodomain helicase DNA-binding 7) and G-protein coupled receptors (GnRHR)⁷⁷. To induce female fertility, hypogonadotropic hypogonadism of pituitary origin can be reversed by subcutaneous injections of FSH followed by human chorionic gonadotrophin or LH to trigger ovulation⁷⁶. Conversely, hypogonadotropic hypogonadism of hypothalamic origin can be treated using GnRH pumps to restore pituitary hormone secretion⁷⁶. Pulsatile GnRH has the advantage of decreasing the risk of multiple pregnancy and ovarian hyperstimulation syndrome⁷⁸. In both situations, the pulsatility of GnRH or the rhythmic secretion of LH and/or FSH is required to obtain sufficient follicular maturation and proper ovulation⁷⁶. Advances in understanding of GnRH secretion and its interactions with LH are essential for designing novel, and indeed modifying existing, therapies for hypogonadotropic hypogonadism. For example, an estimated 22% of patients with this disease, who are yet to undergo treatment, have transient phases of normal fertility⁷⁹. The underlying mechanisms and relevant therapeutic interventions to maintain this phenomenon might be elucidated by further investigation of pulsatility and rhythmicity.

[H1] The prolactin axis

The prolactin axis is unique among the pituitary hormonal systems, as in men and non-lactating women it can be considered a system primed for activation but tonically inhibited by hypothalamic dopamine⁸⁰. In this situation, low concentrations (<25 ng/ml) of circulating prolactin are maintained by short-loop feedback, with prolactin receptor-mediated stimulation of dopamine neuron firing rate leading to an increase in catecholamine production⁸¹ and output². The timescale of the feedback response to prolactin (~10–20 min) could be explained by coordinated release of dopamine from multiple neurons². One possible mechanism is the coordinated changes in firing rates of a subset of tuberoinfundibular dopamine neurons over tens of minutes; these correlate with episodic dopamine secretion recorded from multiple terminals at the median eminence in mice². Gap junctions and local dendritic dopamine release have

been proposed to mediate this activity^{82,83}, and integration of single cell firing rates seems to be involved in the generation of longer dopamine release output events (N. Romano and P. Mollard, unpublished data).

Variations in prolactin output occur in virgin female rats as a surge at proestrus, which coincides with that of LH⁸⁴. Prolactin also increases following vaginal stimulation of both rats⁸⁵ and mice⁸⁶ as twice daily surges. These surges are coordinated by signals from the suprachiasmatic nucleus⁸⁷, most likely through the actions of vasoactive-intestinal peptide⁸⁸. At the level of the pituitary, lactotrophs form a network of honeycomb motifs (Box 1) that allow the congregation of cells along the fine pituitary capillary network (FIG. 3)⁸⁹. This organization supports low levels of cell-cell coordination, with a small proportion (~1–10%) of cells acting as coordinating nodes by functionally connecting distant ensembles⁹⁰. In addition to synchronizing Ca²⁺ activity, cellular organization also mediates the coordination of gene transcription, with gap junction signalling enabling local correlation of bursts of transcriptional activity that are otherwise randomly timed^{91,92}. This mechanism resembles quorum-sensing where apparently random systems display complex activity as long as the components (in this case the cells) can interact, and might contribute to hormone gene expression and cell proliferation⁹²⁻⁹⁴. Precisely how gap junctions might orchestrate this mechanism remains unknown.

[H3] Increased prolactin output during lactation

The long-term requirement for large increases of circulating prolactin in lactation is associated with a decrease in dopamine output, which begins in late pregnancy and is coincident with a surge of pituitary prolactin secretion⁹⁵ (FIG. 2). The dopamine tone needs to be strongly decreased throughout lactation to enable the necessary increase in circulating prolactin, and is mediated by a decrease in phosphorylation of tyrosine hydroxylase, the rate-limiting enzyme for dopamine synthesis⁹⁵. This mechanism is not the result of a reduced feedback of prolactin on dopamine neurons, which remain electrically responsive at the level of the cell body, but rather, neuronal firing becomes uncoupled from dopamine secretion². Remarkably, the reduction in dopamine tone is accompanied by the production of opioids, which might enable these neurons to stimulate prolactin secretion^{96,97}.

In concert with changes in the hypothalamic inhibition of prolactin secretion, substantial alterations occur in the pituitary to support the 10–50-fold increase in prolactin secretion that is required for milk production in mammals⁹⁸. In humans and rats, this increased hormone secretion is generally accompanied by proliferation and hypertrophy of lactotrophs, although the studies describing hypertrophy are based on 2D histological studies⁹⁸. By contrast, in lineage tracing and FACS studies in mice, lactotrophs become hypertrophied during lactation and increase their volume threefold without the accompanying increase in number⁹⁹. Other investigators have confirmed these findings, and also showed that the lactotroph network *in situ* becomes highly-connected during lactation, which is associated with the strength of the suckling stimulus¹⁰⁰. This increase in structural connectivity leads to an ~100% increase in the proportion of the subpopulation of lactotrophs that function as coordinating nodes and orchestrate increased output of prolactin¹⁰⁰.

[H3] Memory of prolactin demand after weaning

At weaning, a rapid decrease in prolactin secretion occurs as a result of a return of dopamine inhibition¹⁰¹. In rodents^{101,102} and humans¹⁰³ basal prolactin secretion is reduced below that of virgin animals, which might reflect an enhanced pituitary response to dopamine inhibition¹⁰⁴. Strikingly, and despite this reduction in basal prolactin secretion, lactotrophs remain enlarged and well-connected with each other at both the structural and functional levels, with a twofold increased number of nodes, which persists for many months after lactation has ceased¹⁰⁰. Such hard-wiring or ‘memory’ of previous stimuli, which was previously thought to only exist for neurons and immune cells, leads to augmented network-mediated lactotroph calcium activity during suckling of subsequent litters, which drives even higher concentrations of prolactin¹⁰⁰. This mechanism is independent of reproductive experience *per se*, since it can be prevented by reducing the suckling demand¹⁰⁰.

[H3] Clinical relevance

The dysregulation of the prolactin axis, owing to either pituitary adenomas¹⁰⁵ or as an adverse effect of treatment with antipsychotic drugs¹⁰⁶, leads to impaired fertility. Hyperprolactinaemia has a prevalence of ~10 to 30 per 100,000 in men and women, respectively, and is the second most common cause of infertility in women after PCOS¹⁰⁷. Whilst treatment with commonly-used dopamine receptor agonists is an

effective treatment of the majority of hyperprolactinaemic patients, side effects of these drugs including nausea, headaches and postural hypotension leads to compliance problems¹⁰⁸. Clearly, an understanding of the interactions that lead to altered dopamine output and the response of the pituitary might help to identify novel treatment strategies for this disease. In rodent studies, prolactin seems to affect multiple neuroendocrine axes, including those regulating fertility, body weight and appetite, stress and maternal behaviour⁸⁰ and these warrant further study to determine the potential effects of its over-secretion in humans. For example, hyperprolactinaemia might lead to changes in GnRH neuron activity via interactions with the GPR54/kisspeptin pathway in mice¹⁰⁹ and GnRH pulsatility has been reinstated in mice with physiological hyperprolactinaemia by administration of kisspeptin¹¹⁰. By contrast with the study in mice where an acute high dose of prolactin was delivered peripherally¹⁰⁹, in studies treating sheep with a chronic centrally administered low (10,000-fold lower) prolactin dose, no effects of prolactin on hypothalamic kisspeptin expression have been seen¹¹¹. An improved understanding of these pathways could aid the development of treatments for women with hyperprolactinaemia that is resistant to dopamine agonists¹⁰⁸.

[H1] The GH axis

In humans and animals, such as cattle and horses, in which the measurements can be taken, pulsatile GH output is present from birth¹¹²⁻¹¹⁴. However, the output is markedly increased at puberty when sexually-dimorphic body growth occurs¹¹⁵. Since the discoveries of GH releasing hormone (GHRH) and somatostatin that control GH secretion from pituitary somatotrophs¹¹⁵⁻¹¹⁷, a remarkable advancement our understanding of GH pulse generation during critical physiological windows has taken place.

[H3] Pulsatile GHRH output

Using genetically-modified mouse models with GHRH neurons marked with green fluorescent protein¹¹⁸, several investigators have defined the mechanisms that underlie pulse generation using *ex vivo* slices of brain. Before puberty, GHRH neurons are excitable neuroendocrine neurons with complex synaptic inputs¹¹⁹. These early stages of hypothalamic development ensure appropriate regulation of the somatotroph axis, as shown in the Ames dwarf mice in which loss of GH leads to a compensatory

increase in GHRH cell number¹²⁰. Additionally, steroid exposure in young animals can have programming effects on the GHRH neuron population, with testosterone exposure in neonates permanently increasing adult GHRH cell number and GHRH gene transcription¹²¹. Modification of these synaptic inputs and electrical properties over the first 6 [weeks of postnatal life](#) correlates with and likely drives, at least in part, increased pituitary GH output and sexual dimorphism¹²². The intrinsic hourly rhythms of GHRH neuronal activity predicted by simulation studies of *in vivo* GH pulsatility have not been identified^{119,123}, however, somatostatin can generate GHRH neuron pulsatile output by delaying oscillations of action potential firing via a recurring inhibition of inhibitory GABAergic interneurons (that is, inhibition of inhibition)¹²³. Consequently, somatostatin can both acutely inhibit the excitability of GHRH neurons and also promote their patterned output together with more sustained GHRH neuron stimulation in response to other stimuli in the brain (for example, acetylcholine¹²²) and peripheral tissues (such as, ghrelin)¹¹⁷.

[H3] Modification of pituitary somatotroph output

No full description of the *in vivo* dynamics of GHRH and somatostatin neurons and their regulation of pituitary somatotrophs exist. This event can be viewed as a three-step process: delivery of neurohormone to target cells; cellular secretory responses to regulation; and entry of pituitary hormone into the peripheral circulation⁸. *In vivo* imaging of the mouse portal system and somatotroph network have provided insights into the first step in this process and the role of the vasculature in shaping the pattern of exposure of the pituitary to hypothalamic neuropeptides⁸. In this study, delivery of neuropeptides such as GHRH to the somatotroph network, which extends throughout the pituitary gland, follows specific vascular/capillary routes and results in specific temporal patterned regional regulation rather than a homogenous exposure of the whole pituitary to the secretagogue. In addition, the initial stimulation by GHRH evokes a coordinated enhancement of oxygen supply to the stimulated somatotroph network via increased capillary blood flow, which provides fuel for energy-depleting secretory responses. Indeed, this study also demonstrated that capillaries closely line the clusters and strings of cells that form the GH cell, which suggests an important role of local oxygen regulation on GH release (Figure 3). The dynamic association of gonadotrophs with the vasculature, which varies through the ovarian cycle⁶⁵, suggests that this situation might also be the case for other pituitary axes. The second step in

pituitary regulation has been characterized using *ex vivo* data from acute pituitary slices where the somatotroph network organization is preserved, in which the homotypic network mediates coordination of stimulation, triggering long-lasting GH secretion¹²⁴ (Box 1) These studies have also shown that network organization is likely to have a major role in the increased GH output at puberty. In particular, using males the study showed that the volume of the GH network undergoes large changes relative to its surface area that coincides with the onset of puberty, before gradually returning to normal prepubertal levels by day 100 in mice¹⁶. These changes occur with a timing that is coincident with the increased pulse output of GH associated with puberty in males and are blocked by castration, which also prevents pubertal changes in GH output¹⁶. A direct effect of sex steroids on organisation of pituitary cells into homotypic networks is apparent from the rapid and dramatic increased motility of somatotrophs in *ex vivo* pituitary cultures treated with oestradiol¹²⁵. These findings highlight the importance of somatotroph network organization and its plasticity in the generation of pituitary somatotroph output. The vasculature also has an important role in somatotroph output, where *in vivo* imaging shows that capture of secreted GH is a controlled event where the perivascular space acts as a gate-keeper for hormone entry into the capillary lumen⁸. The relationship between cellular network organization and the vasculature in the pituitary is, therefore, central to the delivery of incoming hypothalamic signals, and the build-up of GH pulses within capillaries.

[H3] Clinical relevance

GH deficiency resulting from congenital defects or acquired following traumatic brain injury, pituitary tumours or cranial irradiation^{126,127}, is commonly treated with a daily subcutaneous dose of recombinant GH in childhood to increase growth rate¹²⁸. However, considerable uncertainty exists regarding the optimal dosage or regime, and current treatments by injection of GH do not fully recapitulate the physiological pattern of GH secretion¹¹⁵. One potential therapy is repopulation of the pituitary with stem cells, which have been identified in the mouse¹²⁹. However, such approaches are likely to require the recapitulation of the normal cellular organization to achieve normal pulsatility and homeostatic regulation.

Patients with acromegaly, which results from a GH secreting pituitary adenoma, frequently have glycaemic disorders: a lack of GH pulsatility modifies lipolysis,

whereas overall GH hypersecretion can induce insulin resistance¹³⁰. Consequently, an improved understanding of the mechanisms that determine the pattern of GH output might help to define new therapeutic options for dyslipidaemia or diabetes mellitus. In addition, GH pulsatility also has an important role in a subgroup of patients who have clinical acromegaly with increased insulin-like growth factor 1, but unaltered mean 24-h GH concentration compared with healthy controls^{131,132}. Altered GH pulsatility might explain the clinical presentation of this sub-group of patients, and our new understanding of the mechanisms underlying patterning of pituitary output might explain the abnormal GH axis function in these individuals and warrants further investigation.

[H1] Conclusions

The examples in this Review provide new insights into regulation of three hypothalamic–pituitary axes and demonstrate that these mechanisms are not a simple relay of stimulus-secretion coupled events. The disconnection or modulation of hypothalamic excitation and neurohormone release, and an active role of the vasculature and pituitary in the network-mediated modification of responses, demonstrates that the previous stimulus-secretion coupled view of the hypothalamic–pituitary system is over-simplistic. Advances in imaging technologies are allowing us to understand more about the organization and function of this axis (FIG. 4). Given that the hypothalamus contains no more than a few thousand parvocellular neurons, the rapid development of techniques for interrogating neuronal function should enable the characterization of this structure's regulation and output. Such studies will be invaluable for the deeper understanding of mammalian physiology, as this region controls a much larger panel of known body functions than any other brain region.

The regulation of pulsatile pituitary secretion must now be considered as an integration of hypothalamic, vasculature and pituitary regulation, which has further implications for the understanding of disease. For example, the identification of kisspeptin has provided an exciting new target for the treatment of infertility¹³³. The uncoupling of neuronal excitation and hormone output also has deeper implications such as in the case of ageing. For example, the reduction in GH output with age might be due to a failure of neurohormone secretion from GHRH neurons without any change in their excitation¹²⁹ (Figure 2). The human pituitary gland can be accessed by

transphenoidal surgery, which makes this structure an ideal target organ for regenerative therapy¹³⁴. The pituitary cell networks and their relationship with the vasculature must be considered for such therapy and the microenvironment clearly has an important role in the regulation of the pituitary gland, which might also affect the development of tumours¹³⁵. Finally, as pituitary networks are sensitive to peripheral regulation and their modification can persist for extended periods, they are a potential target for endocrine disrupting chemicals (EDCs). Indeed, the identification of bisphenol A-mediated reduction of expression of *ICAM5* in the pituitary¹³⁶ has led us to speculate that some EDC effects might be mediated by changes in pituitary organization¹³⁷. These possibilities require further investigation for the understanding of both the aetiology and treatment of diseases associated with pituitary hormones.

Competing interests statement

The authors declare no competing interests

Acknowledgements

We thank F. Castinetti (Aix-Marseille Universite, France) for helpful comments and suggestions. P.L.T. was supported by a grant from the Biotechnology and Biological Sciences Research Council, UK (BB/N007026/1). N.R. was supported by a Medical Research Council, UK project grant (MR/J008893/1). D.J.H. was supported by an R.D. Lawrence Fellowship, Diabetes UK (12/0004431), European [Foundation](#) for the Study of Diabetes /Novo Nordisk Rising Star and Birmingham Fellowships, a Medical Research Council, UK project grant (MR/N00275X/1), Imperial Confidence in Concept and Wellcome Trust Institutional Support Awards, both UK, and an [ERC Starting Grant](#) (OptoBETA; 715884). P.M. was supported by funding from the Agence Nationale de la Recherche (ANR 12 BSV1 0032-01, ANR-15-CE14-0012-01), Institut National de la Santé et de la Recherche Médicale, Centre National de la Recherche Scientifique, Université de Montpellier, Fondation pour la Recherche Médicale (DEQ20150331732), and IPAM-Biocampus of Montpellier and France-Bioimaging, all in France.

Author contributions

The authors contributed equally to all aspects of the preparation of this article

Key points

- The activity of hypothalamic neurons is modified by inputs leading to heterogeneous activity; a small proportion of the total population can drive pituitary hormone pulsatility
- Neurohormone output can vary following neuron excitation according to physiological status, which might also lead to declining neuroendocrine output with age
- The release of hypothalamic factors into the blood is modified by alterations in the juxtaposition of nerve terminals with the vasculature and tanycytes in the median eminence
- Cells in the pituitary gland form homotypic networks, the organization of these relationships with the vasculature are distinct for each endocrine axis, which modifies responses to regulatory factors and patterns of output in response to demand
- Reorganisation of the pituitary network can store long-term memories of increased output and learn to increase function on repeated challenge
- Understanding the importance of coordinated hypothalamic–vasculature–pituitary function provides new understanding of a range of endocrine axes defects and targets for novel therapies

Figure 1 |A dynamic interplay between hypothalamic neuron output, the vasculature and the pituitary response alters hormone output. a) The concept of stimulus-secretion coupling considered pituitary hormone output to result from a cascade of events dictated by the pattern of regulatory neuron excitation. **b)** Recent studies have shown that this view (a) is too simplistic: alterations at the level of neuron response to excitation, release of hypothalamic factors in the median eminence and pituitary cell responses are all capable of adapting output of hormone to the periphery.

Figure 2 |Output of hypothalamic regulatory factors to the median eminence is mediated by a number of processes. a) A simplified view of tanycyte regulation of gonadotrophin releasing hormone (GnRH) release. Access of GnRH nerve terminals to capillaries at diestrus is blocked by tanycyte ensheathment, which retracts of at proestrus, possibly in response to oestrogen-driven secretion of nitric oxide (NO), leading to formation of a direct neurohaemal junction and augmented release of

GnRH. **b)** Feedback inhibition of prolactin secretion is reduced in lactation by an uncoupling of dopamine production and release from neuronal electrical activity and possible switch to production and secretion of opioids. **c)** Growth hormone releasing hormone (GHRH) secretion is reduced in late adulthood, with an altered localisation of nerve terminals and redistribution of secretory vesicles to autophagic vacuoles, both resulting in reduced neuropeptide release to the portal circulation.

Figure 3 Pituitary cells form homotypic networks, with distinct organisational motifs and relationship with the vasculature, which alter with physiological status. **a)** Gonadotroph cells are arranged as strings of cells which align with capillaries and make direct contact with the vasculature through protrusions at proestrus. **b)** The organisation of lactotroph into honeycomb structures in virgin mice becomes more pronounced in animals which are lactating, with an alteration of cell-cell contacts, increase in cell size, and appearance of cells which act as hubs of functional connectivity (dark orange). This change in structural and functional organisation is maintained for months following cessation of physiological demand at weaning. **c)** The homotypic network of somatotrophs changes dramatically at puberty, with the formation of large clusters which are linked by strands of cells in young males. The strands of cells found in juvenile and mature animals are closely associated with capillaries, which also line the post-pubertal somatotroph clusters

Figure 4 | Updated view of the hypothalamic–vascular–pituitary unit. In the past decade, a more complex relationship between the hypothalamus and pituitary than previously appreciated has emerged, in part owing to new imaging techniques which now allow high-resolution optical imaging of the hypothalamic-vascular-pituitary unit in living animal models. **a** | At the level of the parvocellular neurons, complex inputs modify the excitation of neurons, which can vary coupling with neurohormone output at terminals of the median eminence (ME) through modification of intracellular pathways. **b** | Alterations in tanycyte ensheathment and the anatomical location of neuron terminals modify their interactions with the vasculature, changing the dynamics of neurohormone release. **c** | In the pituitary gland, changes in blood flow in the portal circulation alter the pattern of exposure of pituitary cells to neurohormone and nutrient supply to facilitate secretion. **d** | Cells of the pituitary gland, such as somatotrophs, are organised into intermingled networks with distinct morphological features, which can be altered to meet physiological demand, and relationships with the vasculature.

Box 1 | Organization of pituitary cells into homotypic networks

Somatotrophs, corticotrophs, gonadotrophs and lactotrophs form networks with the following features^{16,64,65,100,138,139}

- *The lineages have distinct developmental programmes*
 - Placement of each endocrine cell network occurs at distinct stages of pituitary organogenesis, before expansion in early postnatal life^{16,64}
- *Networks have distinct motifs and interact with the vasculature*
 - Somatotrophs organise as clusters linked with strands¹⁶ along the capillary network¹³⁸, while lactotrophs form a honeycomb structure¹⁰⁰ aligned with these vessels¹³⁸. Gonadotrophs connect to one or more blood vessels via their protrusions⁶⁵, whereas the corticotroph network has a loose arrangement⁶⁴. Cell-type specific homotypic network organisation and their distinct relationships with the vasculature are likely to affect levels and timing of hormone release^{8, 16, 70, 100, 142}.
 - The mechanisms underlying the distinct endocrine cell-vascular relationship are unclear, although one factor shown to be important is Prop1, since its loss leads to failure of organ vascularization¹⁴³
- *Hypothalamic and steroidal factors generate network motifs*
 - Loss of growth hormone-releasing hormone (GHRH) leads to isolated somatotrophs, whereas somatotroph ablation with intact GHRH stimulation results in clusters of cells that are isolated from each other¹⁴⁰. Gonadal steroids also influence somatotroph cell clustering and cell movement leading to reorganisation^{124,125}
- *Pituitary networks have functional relevance*
 - Pituitary networks integrate, amplify and propagate hypothalamic signals that arrive from the median eminence. For example, the male somatotroph network responds to GHRH input with large, coordinated, oscillatory Ca²⁺ increases that outlast the stimulus to drive large excursions in hormone secretion¹²⁴.
- *Endocrine and non-endocrine homotypic networks interact*
 - Communication between each pituitary hormonal cell-type, as well as the non-hormonal folliculostellate cells, via gap-junctions mediated coupling^{91,100} and paracrine and autocrine interactions^{138,141}, can modify output of each of the pituitary hormones and allow cross-talk between pituitary axes.
 - Loss of specific pituitary hormonal cell types influences the network organisation of heterotypic cells. For example, changes in the gonadotroph network occur following alteration of corticotroph terminal differentiation in mice⁶⁴.

Author biographies

Paul Le Tissier is a Senior Lecturer in the Centre for Integrative Physiology, University of Edinburgh, UK. His primary research interests are the understanding of hypothalamic regulation of the anterior pituitary gland, how this mechanism maintains and alters the function of the different cell populations to ensure appropriate output throughout life and modelling how dysregulation leads to pathology.

Pauline Campos has recently become a Postdoctoral Fellow in the laboratory of Patrice Mollard, at the Institute of Functional Genomics of Montpellier, France. She completed her PhD in the Centre for Neuroendocrinology at the University of Otago, New Zealand, and has been focused on developing cutting-edge tools to study the electrical activity of hypothalamic neurons underlying pituitary hormone pulsatility. Her current work aims at understanding the central control of thyroid function in conscious animals.

Chrystel Lafont is an INSERM Engineer in the Institute of Functional Genomics of Montpellier, France. She is the technical leader of a small animal *in vivo* imaging facility that develops cellular *in vivo* imaging techniques. Her research interests look at the role of the vasculature and pituitary cell networks in the build-up of hormone pulses.

Nicola Romanò is a Lecturer at the University of Edinburgh, UK. His main interest is the study of the cellular and systemic mechanisms of pulsatile hormone secretion in neuroendocrine axes, and how they correlate and influence physiological function. His research interests span from analysis of activity patterns of hypothalamic neurons to structural and functional analysis of pituitary cell networks, such as, the spatio-temporal analysis of cell activity.

David Hodson is a Professor in Cellular Metabolism at the University of Birmingham, UK. His laboratory is developing innovative optical methodologies that enable the mapping and manipulation of endocrine cells directly within their tissue context, focusing on the pituitary gland and pancreas as tractable systems.

Patrice Mollard is the Head of the Department of Physiology and a CNRS Research Director at the Institute of Functional Genomics, Montpellier, France. Dr Mollard's laboratory pioneered the study of large-scale functional organization of endocrine cell types within the pituitary gland in the early 2000s. His research is now exploring how pituitary cell networks function together with hypothalamic inputs and vasculature in health and disease.

- 1 Sandow, A. Excitation-contraction coupling in skeletal muscle. *Pharmacol Rev* **17**, 265-320, (1965).
- 2 Romano, N. *et al.* Plasticity of hypothalamic dopamine neurons during lactation results in dissociation of electrical activity and release. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **33**, 4424-4433, (2013).
- 3 Smith, J. T. *et al.* Kisspeptin is essential for the full preovulatory LH surge and stimulates GnRH release from the isolated ovine median eminence. *Endocrinology* **152**, 1001-1012, (2011).
- 4 Lim, N. F., Nowycky, M. C. & Bookman, R. J. Direct measurement of exocytosis and calcium currents in single vertebrate nerve terminals. *Nature* **344**, 449-451, (1990).

- 5 Drouva, S. V., Epelbaum, J., Laplante, E. & Kordon, C. Calmodulin involvement on the Ca⁺⁺-dependent release of LHRH and SRIF in vitro. *Neuroendocrinology* **38**, 189-192, (1984).
- 6 Douglas, W. W. Stimulus-secretion coupling: the concept and clues from chromaffin and other cells. *Br J Pharmacol* **34**, 451-474, (1968).
- 7 Tse, A., Tse, F. W., Almers, W. & Hille, B. Rhythmic exocytosis stimulated by GnRH-induced calcium oscillations in rat gonadotropes. *Science* **260**, 82-84, (1993).
- 8 Lafont, C. *et al.* Cellular in vivo imaging reveals coordinated regulation of pituitary microcirculation and GH cell network function. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 4465-4470, (2010).
- 9 Thomas, P., Wong, J. G., Lee, A. K. & Almers, W. A low affinity Ca²⁺ receptor controls the final steps in peptide secretion from pituitary melanotrophs. *Neuron* **11**, 93-104, (1993).
- 10 Tse, A. & Lee, A. K. Voltage-gated Ca²⁺ channels and intracellular Ca²⁺ release regulate exocytosis in identified rat corticotrophs. *J Physiol* **528 Pt 1**, 79-90, (2000).
- 11 Raisman, G. An urge to explain the incomprehensible: Geoffrey Harris and the discovery of the neural control of the pituitary gland. *Annual review of neuroscience* **20**, 533-566, (1997).
- 12 Stojilkovic, S. S., Tabak, J. & Bertram, R. Ion channels and signaling in the pituitary gland. *Endocrine reviews* **31**, 845-915, (2010).
- 13 Keenan, D. M. & Veldhuis, J. D. Pulsatility of Hypothalamo-Pituitary Hormones: A Challenge in Quantification. *Physiology (Bethesda)* **31**, 34-50, (2016).
- 14 Moenter, S. M. Leap of Faith: Does Serum Luteinizing Hormone Always Accurately Reflect Central Reproductive Neuroendocrine Activity? *Neuroendocrinology* **102**, 256-266, (2015).
- 15 Petersenn, S. & Schulte, H. M. Structure and function of the growth-hormone-releasing hormone receptor. *Vitam Horm* **59**, 35-69, (2000).
- 16 Bonnefont, X. *et al.* Revealing the large-scale network organization of growth hormone-secreting cells. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 16880-16885, (2005).
- 17 Han, S. K., Todman, M. G. & Herbison, A. E. Endogenous GABA release inhibits the firing of adult gonadotropin-releasing hormone neurons. *Endocrinology* **145**, 495-499, (2004).
- 18 Han, S. K., Lee, K., Bhattarai, J. P. & Herbison, A. E. Gonadotrophin-releasing hormone (GnRH) exerts stimulatory effects on GnRH neurons in intact adult male and female mice. *Journal of neuroendocrinology* **22**, 188-195, (2010).
- 19 Dierschke, D. J., Bhattacharya, A. N., Atkinson, L. E. & Knobil, E. Circular oscillations of plasma LH levels in the ovariectomized rhesus monkey. *Endocrinology* **87**, 850-853, (1970).
- 20 Carmel, P. W., Araki, S. & Ferin, M. Pituitary stalk portal blood collection in rhesus monkeys: evidence for pulsatile release of gonadotropin-releasing hormone (GnRH). *Endocrinology* **99**, 243-248, (1976).
- 21 Knobil, E. The neuroendocrine control of the menstrual cycle. *Recent Prog Horm Res* **36**, 53-88, (1980).
- 22 Clarke, I. J. & Cummins, J. T. The temporal relationship between gonadotropin releasing hormone (GnRH) and luteinizing hormone (LH) secretion in ovariectomized ewes. *Endocrinology* **111**, 1737-1739, (1982).
- 23 Levine, J. E., Pau, K. Y., Ramirez, V. D. & Jackson, G. L. Simultaneous measurement of luteinizing hormone-releasing hormone and luteinizing hormone release in unanesthetized, ovariectomized sheep. *Endocrinology* **111**, 1449-1455, (1982).
- 24 Caraty, A., Locatelli, A. & Martin, G. B. Biphasic response in the secretion of gonadotrophin-releasing hormone in ovariectomized ewes injected with oestradiol. *The Journal of endocrinology* **123**, 375-382, (1989).
- 25 Moenter, S. M., Brand, R. C. & Karsch, F. J. Dynamics of gonadotropin-releasing hormone (GnRH) secretion during the GnRH surge: insights into the mechanism of GnRH surge induction. *Endocrinology* **130**, 2978-2984, (1992).
- 26 Levine, J. E. New concepts of the neuroendocrine regulation of gonadotropin surges in rats. *Biol Reprod* **56**, 293-302, (1997).
- 27 Karsch, F. J., Bowen, J. M., Caraty, A., Evans, N. P. & Moenter, S. M. Gonadotropin-releasing hormone requirements for ovulation. *Biol Reprod* **56**, 303-309, (1997).
- 28 Steyn, F. J. *et al.* Development of a methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile and adult male mice. *Endocrinology* **154**, 4939-4945, (2013).
- 29 Campos, P. & Herbison, A. E. Optogenetic activation of GnRH neurons reveals minimal requirements for pulsatile luteinizing hormone secretion. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 18387-18392, (2014).

- 30 Kokoris, G. J., Lam, N. Y., Ferin, M., Silverman, A. J. & Gibson, M. J. Transplanted gonadotropin-releasing hormone neurons promote pulsatile luteinizing hormone secretion in congenitally hypogonadal (hpg) male mice. *Neuroendocrinology* **48**, 45-52, (1988).
- 31 Herbison, A. E., Porteous, R., Pape, J. R., Mora, J. M. & Hurst, P. R. Gonadotropin-releasing hormone neuron requirements for puberty, ovulation, and fertility. *Endocrinology* **149**, 597-604, (2008).
- 32 Jasoni, C. L., Romano, N., Constantin, S., Lee, K. & Herbison, A. E. Calcium dynamics in gonadotropin-releasing hormone neurons. *Front Neuroendocrinol* **31**, 259-269, (2010).
- 33 Lehman, M. N., Coolen, L. M. & Goodman, R. L. Minireview: kisspeptin/neurokinin B/dynorphin (KNDy) cells of the arcuate nucleus: a central node in the control of gonadotropin-releasing hormone secretion. *Endocrinology* **151**, 3479-3489, (2010).
- 34 Navarro, V. M. *et al.* Role of neurokinin B in the control of female puberty and its modulation by metabolic status. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **32**, 2388-2397, (2012).
- 35 Campbell, R. E., Gaidamaka, G., Han, S. K. & Herbison, A. E. Dendro-dendritic bundling and shared synapses between gonadotropin-releasing hormone neurons. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 10835-10840, (2009).
- 36 Herde, M. K., Iremonger, K. J., Constantin, S. & Herbison, A. E. GnRH neurons elaborate a long-range projection with shared axonal and dendritic functions. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **33**, 12689-12697, (2013).
- 37 Prevot, V. *et al.* Evidence that members of the TGFbeta superfamily play a role in regulation of the GnRH neuroendocrine axis: expression of a type I serine-threonine kinase receptor for TGRbeta and activin in GnRH neurones and hypothalamic areas of the female rat. *Journal of neuroendocrinology* **12**, 665-670, (2000).
- 38 Knauf, C. *et al.* Evidence for a spontaneous nitric oxide release from the rat median eminence: influence on gonadotropin-releasing hormone release. *Endocrinology* **142**, 2343-2350, (2001).
- 39 De Seranno, S. *et al.* Vascular endothelial cells promote acute plasticity in ependymogial cells of the neuroendocrine brain. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **24**, 10353-10363, (2004).
- 40 Hanchate, N. K. *et al.* Kisspeptin-GPR54 signaling in mouse NO-synthesizing neurons participates in the hypothalamic control of ovulation. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **32**, 932-945, (2012).
- 41 Bellefontaine, N. *et al.* Nitric oxide as key mediator of neuron-to-neuron and endothelia-to-glia communication involved in the neuroendocrine control of reproduction. *Neuroendocrinology* **93**, 74-89, (2011).
- 42 Page, R. B. Pituitary blood flow. *Am J Physiol* **243**, E427-442, (1982).
- 43 Belchetz, P. E., Plant, T. M., Nakai, Y., Keogh, E. J. & Knobil, E. Hypophysial responses to continuous and intermittent delivery of hypothalamic gonadotropin-releasing hormone. *Science* **202**, 631-633, (1978).
- 44 Wildt, L. *et al.* Frequency and amplitude of gonadotropin-releasing hormone stimulation and gonadotropin secretion in the rhesus monkey. *Endocrinology* **109**, 376-385, (1981).
- 45 Pohl, C. R., Richardson, D. W., Hutchison, J. S., Germak, J. A. & Knobil, E. Hypophysiotropic signal frequency and the functioning of the pituitary-ovarian system in the rhesus monkey. *Endocrinology* **112**, 2076-2080, (1983).
- 46 McArdle, C. A. & Roberson, M. S. in *Knobil and Neill's Physiology of Reproduction (Fourth Edition)* 335-397 (Academic Press, 2015).
- 47 Tsutsumi, R. & Webster, N. J. GnRH pulsatility, the pituitary response and reproductive dysfunction. *Endocrine journal* **56**, 729-737, (2009).
- 48 Cooke, B., Hegstrom, C. D., Villeneuve, L. S. & Breedlove, S. M. Sexual differentiation of the vertebrate brain: principles and mechanisms. *Front Neuroendocrinol* **19**, 323-362, (1998).
- 49 Simerly, R. B. Organization and regulation of sexually dimorphic neuroendocrine pathways. *Behavioural brain research* **92**, 195-203, (1998).
- 50 Levine, J. E. & Ramirez, V. D. Luteinizing hormone-releasing hormone release during the rat estrous cycle and after ovariectomy, as estimated with push-pull cannulae. *Endocrinology* **111**, 1439-1448, (1982).
- 51 Sarkar, D. K., Chiappa, S. A., Fink, G. & Sherwood, N. M. Gonadotropin-releasing hormone surge in pro-estrous rats. *Nature* **264**, 461-463, (1976).
- 52 Park, O. K. & Ramirez, V. D. Spontaneous changes in LHRH release during the rat estrous cycle, as measured with repetitive push-pull perfusions of the pituitary gland in the same female rats. *Neuroendocrinology* **50**, 66-72, (1989).

- 53 Herbison, A. E. Control of puberty onset and fertility by gonadotropin-releasing hormone
neurons. *Nat Rev Endocrinol*, (2016).
- 54 Plant, T. M. A comparison of the neuroendocrine mechanisms underlying the initiation of the
preovulatory LH surge in the human, Old World monkey and rodent. *Front Neuroendocrinol*
33, 160-168, (2012).
- 55 Moenter, S. M., Caraty, A., Locatelli, A. & Karsch, F. J. Pattern of gonadotropin-releasing
hormone (GnRH) secretion leading up to ovulation in the ewe: existence of a preovulatory
GnRH surge. *Endocrinology* **129**, 1175-1182, (1991).
- 56 Clarke, I. J. Variable patterns of gonadotropin-releasing hormone secretion during the
estrogen-induced luteinizing hormone surge in ovariectomized ewes. *Endocrinology* **133**,
1624-1632, (1993).
- 57 Caraty, A. *et al.* Nature and bioactivity of gonadotropin-releasing hormone (GnRH) secreted
during the GnRH surge. *Endocrinology* **136**, 3452-3460, (1995).
- 58 Kozlowski, G. P. & Coates, P. W. Ependymoneuronal specializations between LHRH fibers
and cells of the cerebroventricular system. *Cell and tissue research* **242**, 301-311, (1985).
- 59 King, J. C. & Rubin, B. S. Dynamic alterations in luteinizing hormone-releasing hormone
(LHRH) neuronal cell bodies and terminals of adult rats. *Cellular and molecular neurobiology*
15, 89-106, (1995).
- 60 Parkash, J. *et al.* Semaphorin7A regulates neuroglial plasticity in the adult hypothalamic
median eminence. *Nat Commun* **6**, 6385, (2015).
- 61 Giacobini, P. *et al.* Brain endothelial cells control fertility through ovarian-steroid-dependent
release of semaphorin 3A. *PLoS Biol* **12**, e1001808, (2014).
- 62 King, J. C. & Letourneau, R. J. Luteinizing hormone-releasing hormone terminals in the
median eminence of rats undergo dramatic changes after gonadectomy, as revealed by
electron microscopic image analysis. *Endocrinology* **134**, 1340-1351, (1994).
- 63 Prevot, V. *et al.* Definitive evidence for the existence of morphological plasticity in the
external zone of the median eminence during the rat estrous cycle: implication of neuro-glio-
endothelial interactions in gonadotropin-releasing hormone release. *Neuroscience* **94**, 809-
819, (1999).
- 64 Budry, L. *et al.* Related pituitary cell lineages develop into interdigitated 3D cell networks.
Proceedings of the National Academy of Sciences of the United States of America **108**, 12515-
12520, (2011).
- 65 Alim, Z. *et al.* Gonadotrope plasticity at cellular and population levels. *Endocrinology* **153**,
4729-4739, (2012).
- 66 Thomas, S. G., Takahashi, M., Neill, J. D. & Clarke, I. J. Components of the neuronal
exocytotic machinery in the anterior pituitary of the ovariectomised ewe and the effects of
oestrogen in gonadotropes as studied with confocal microscopy. *Neuroendocrinology* **67**, 244-
259, (1998).
- 67 Barkan, A. L., Regiani, S. R., Duncan, J. A. & Marshall, J. C. Pituitary gonadotropin-releasing
hormone receptors during gonadotropin surges in ovariectomized-estradiol-treated rats.
Endocrinology **112**, 1042-1048, (1983).
- 68 Qiao, S. *et al.* Molecular Plasticity of Male and Female Murine Gonadotropes Revealed by
mRNA Sequencing. *Endocrinology* **157**, 1082-1093, (2016).
- 69 Navratil, A. M., Knoll, J. G., Whitesell, J. D., Tobet, S. A. & Clay, C. M. Neuroendocrine
plasticity in the anterior pituitary: gonadotropin-releasing hormone-mediated movement in
vitro and in vivo. *Endocrinology* **148**, 1736-1744, (2007).
- 70 Schaeffer, M., Hodson, D. J., Lafont, C. & Mollard, P. Endocrine cells and blood vessels work
in tandem to generate hormone pulses. *J Mol Endocrinol* **47**, R59-66, (2011).
- 71 Padmanabhan, V. Polycystic ovary syndrome--"A riddle wrapped in a mystery inside an
enigma". *J Clin Endocrinol Metab* **94**, 1883-1885, (2009).
- 72 McCartney, C. R., Eagleson, C. A. & Marshall, J. C. Regulation of gonadotropin secretion:
implications for polycystic ovary syndrome. *Semin Reprod Med* **20**, 317-326, (2002).
- 73 Francou, M. *et al.* Characterization of pituitary cell populations in rats with induced polycystic
ovaries. *Cells Tissues Organs* **188**, 310-319, (2008).
- 74 Roland, A. V. & Moenter, S. M. Reproductive neuroendocrine dysfunction in polycystic
ovary syndrome: insight from animal models. *Front Neuroendocrinol* **35**, 494-511, (2014).
- 75 Cardoso, R. C., Puttabyatappa, M. & Padmanabhan, V. Steroidogenic versus Metabolic
Programming of Reproductive Neuroendocrine, Ovarian and Metabolic Dysfunctions.
Neuroendocrinology **102**, 226-237, (2015).

- 76 Boehm, U. *et al.* Expert consensus document: European Consensus Statement on congenital hypogonadotropic hypogonadism--pathogenesis, diagnosis and treatment. *Nat Rev Endocrinol* **11**, 547-564, (2015).
- 77 Pitteloud, N., Durrani, S., Raivio, T. & Sykiotis, G. P. Complex genetics in idiopathic hypogonadotropic hypogonadism. *Front Horm Res* **39**, 142-153, (2010).
- 78 Christin-Maitre, S., de Crecy, M. & Groupe Francais des pompes a Gn, R. H. [Pregnancy outcomes following pulsatile GnRH treatment: results of a large multicenter retrospective study]. *J Gynecol Obstet Biol Reprod (Paris)* **36**, 8-12, (2007).
- 79 Sidhoum, V. F. *et al.* Reversal and relapse of hypogonadotropic hypogonadism: resilience and fragility of the reproductive neuroendocrine system. *J Clin Endocrinol Metab* **99**, 861-870, (2014).
- 80 Grattan, D. R. & Kokay, I. C. Prolactin: a pleiotropic neuroendocrine hormone. *Journal of neuroendocrinology* **20**, 752-763, (2008).
- 81 Arbogast, L. A. & Voogt, J. L. Hyperprolactinemia increases and hypoprolactinemia decreases tyrosine hydroxylase messenger ribonucleic acid levels in the arcuate nuclei, but not the substantia nigra or zona incerta. *Endocrinology* **128**, 997-1005, (1991).
- 82 Stagkourakis, S., Kim, H., Lyons, David J. & Broberger, C. Dopamine Autoreceptor Regulation of a Hypothalamic Dopaminergic Network. *Cell reports* **15**, 735-747, (2016).
- 83 Lyons, D. J., Horjales-Araujo, E. & Broberger, C. Synchronized network oscillations in rat tuberoinfundibular dopamine neurons: switch to tonic discharge by thyrotropin-releasing hormone. *Neuron* **65**, 217-229, (2010).
- 84 Freeman, M. E., Reichert, L. E., Jr. & Neill, J. D. Regulation of the proestrus surge of prolactin secretion by gonadotropin and estrogens in the rat. *Endocrinology* **90**, 232-238, (1972).
- 85 Butcher, R. L., Fugo, N. W. & Collins, W. E. Semicircadian rhythm in plasma levels of prolactin during early gestation in the rat. *Endocrinology* **90**, 1125-1127, (1972).
- 86 Larsen, C. M. & Grattan, D. R. Prolactin-induced mitogenesis in the subventricular zone of the maternal brain during early pregnancy is essential for normal postpartum behavioral responses in the mother. *Endocrinology* **151**, 3805-3814, (2010).
- 87 Mai, L. M., Shieh, K. R. & Pan, J. T. Circadian changes of serum prolactin levels and tuberoinfundibular dopaminergic neuron activities in ovariectomized rats treated with or without estrogen: the role of the suprachiasmatic nuclei. *Neuroendocrinology* **60**, 520-526, (1994).
- 88 Egli, M., Bertram, R., Sellix, M. T. & Freeman, M. E. Rhythmic secretion of prolactin in rats: action of oxytocin coordinated by vasoactive intestinal polypeptide of suprachiasmatic nucleus origin. *Endocrinology* **145**, 3386-3394, (2004).
- 89 Hodson, D. J. & Mollard, P. Navigating pituitary structure and function - defining a roadmap for hormone secretion. *Journal of neuroendocrinology* **25**, 674-675, (2013).
- 90 Hodson, D. J. *et al.* Coordination of calcium signals by pituitary endocrine cells in situ. *Cell calcium* **51**, 222-230, (2012).
- 91 Featherstone, K. *et al.* Spatially coordinated dynamic gene transcription in living pituitary tissue. *eLife* **5**, (2016).
- 92 Harper, C. V. *et al.* Dynamic organisation of prolactin gene expression in living pituitary tissue. *Journal of Cell Science* **123**, 424-430, (2010).
- 93 Long, T. *et al.* Quantifying the integration of quorum-sensing signals with single-cell resolution. *PLoS Biol* **7**, e68, (2009).
- 94 Weber, W. *et al.* Streptomyces-derived quorum-sensing systems engineered for adjustable transgene expression in mammalian cells and mice. *Nucleic Acids Res* **31**, e71, (2003).
- 95 Andrews, Z. B., Kokay, I. C. & Grattan, D. R. Dissociation of prolactin secretion from tuberoinfundibular dopamine activity in late pregnant rats. *Endocrinology* **142**, 2719-2724, (2001).
- 96 Ciofi, P. *et al.* Plasticity in expression of immunoreactivity for neuropeptide Y, enkephalins and neurotensin in the hypothalamic tubero-infundibular dopaminergic system during lactation in mice. *Journal of neuroendocrinology* **5**, 599-602, (1993).
- 97 Merchantaler, I. Induction of enkephalin in tuberoinfundibular dopaminergic neurons during lactation. *Endocrinology* **133**, 2645-2651, (1993).
- 98 Le Tissier, P. R., Hodson, D. J., Martin, A. O., Romano, N. & Mollard, P. Plasticity of the prolactin (PRL) axis: mechanisms underlying regulation of output in female mice. *Advances in experimental medicine and biology* **846**, 139-162, (2015).

- 99 Castrique, E., Fernandez-Fuente, M., Le Tissier, P., Herman, A. & Levy, A. Use of a prolactin-Cre/ROSA-YFP transgenic mouse provides no evidence for lactotroph transdifferentiation after weaning, or increase in lactotroph/somatotroph proportion in lactation. *Journal of Endocrinology* **205**, 49-60, (2010).
- 100 Hodson, D. J. *et al.* Existence of long-lasting experience-dependent plasticity in endocrine cell networks. *Nat Commun* **3**, 605, (2012).
- 101 Guillou, A. *et al.* Assessment of Lactotroph Axis Functionality in Mice: Longitudinal Monitoring of PRL Secretion by Ultrasensitive-ELISA. *Endocrinology* **156**, 1924-1930, (2015).
- 102 Byrnes, E. M. & Bridges, R. S. Lactation reduces prolactin levels in reproductively experienced female rats. *Horm Behav* **48**, 278-282, (2005).
- 103 Musey, V. C., Collins, D. C., Musey, P. I., Martino-Saltzman, D. & Preedy, J. R. Long-term effect of a first pregnancy on the secretion of prolactin. *The New England journal of medicine* **316**, 229-234, (1987).
- 104 Byrnes, E. M. & Bridges, R. S. Reproductive experience and expression of dopamine D(2) receptor mRNA: a possible mechanism for reduced prolactin secretion in primiparous rats. *Journal of neuroendocrinology* **19**, 773-778, (2007).
- 105 Wong, A., Eloy, J. A., Couldwell, W. T. & Liu, J. K. Update on prolactinomas. Part 1: Clinical manifestations and diagnostic challenges. *J Clin Neurosci* **22**, 1562-1567, (2015).
- 106 Holt, R. I. & Peveler, R. C. Antipsychotics and hyperprolactinaemia: mechanisms, consequences and management. *Clin Endocrinol (Oxf)* **74**, 141-147, (2011).
- 107 Melmed, S. *et al.* Diagnosis and treatment of hyperprolactinemia: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* **96**, 273-288, (2011).
- 108 Wong, A., Eloy, J. A., Couldwell, W. T. & Liu, J. K. Update on prolactinomas. Part 2: Treatment and management strategies. *J Clin Neurosci* **22**, 1568-1574, (2015).
- 109 Brown, R. S., Herbison, A. E. & Grattan, D. R. Prolactin regulation of kisspeptin neurones in the mouse brain and its role in the lactation-induced suppression of kisspeptin expression. *Journal of neuroendocrinology* **26**, 898-908, (2014).
- 110 Sonigo, C. *et al.* Hyperprolactinemia-induced ovarian acyclicity is reversed by kisspeptin administration. *J Clin Invest* **122**, 3791-3795, (2012).
- 111 Li, Q., Rao, A., Pereira, A., Clarke, I. J. & Smith, J. T. Kisspeptin cells in the ovine arcuate nucleus express prolactin receptor but not melatonin receptor. *Journal of neuroendocrinology* **23**, 871-882, (2011).
- 112 Adcock, C. J. *et al.* The use of an automated microsampling system for the characterization of growth hormone pulsatility in newborn babies. *Pediatr Res* **42**, 66-71, (1997).
- 113 Coxam, V., Davicco, M. J., Robelin, J. & Barlet, J. P. Growth hormone secretory pattern and somatomedin C plasma concentrations in newborn calves. *J Dev Physiol* **9**, 113-121, (1987).
- 114 Davicco, M. J. *et al.* Growth hormone (GH) secretory pattern and GH response to GH-releasing factor (GRF) or thyrotropin-releasing hormone (TRH) in newborn foals. *J Dev Physiol* **19**, 143-147, (1993).
- 115 Robinson, I. C. A. F. & Hindmarsh, P. C. in *Comprehensive Physiology* (John Wiley & Sons, Inc., 2010).
- 116 Tannenbaum, G. S. Genesis of episodic growth hormone secretion. *J Pediatr Endocrinol* **6**, 273-282, (1993).
- 117 Steyn, F. J., Tolle, V., Chen, C. & Epelbaum, J. Neuroendocrine Regulation of Growth Hormone Secretion. *Compr Physiol* **6**, 687-735, (2016).
- 118 Balthasar, N. *et al.* Growth hormone-releasing hormone (GHRH) neurons in GHRH-enhanced green fluorescent protein transgenic mice: a ventral hypothalamic network. *Endocrinology* **144**, 2728-2740, (2003).
- 119 Gouty-Colomer, L. A. *et al.* Specific involvement of gonadal hormones in the functional maturation of growth hormone releasing hormone (GHRH) neurons. *Endocrinology* **151**, 5762-5774, (2010).
- 120 Romero, M. I. & Phelps, C. J. Identification of growth hormone-releasing hormone and somatostatin neurons projecting to the median eminence in normal and growth hormone-deficient Ames dwarf mice. *Neuroendocrinology* **65**, 107-116, (1997).
- 121 Chowen, J. A., Frago, L. M. & Argente, J. The regulation of GH secretion by sex steroids. *Eur J Endocrinol* **151 Suppl 3**, U95-100, (2004).
- 122 Baccam, N. *et al.* Dual-level afferent control of growth hormone-releasing hormone (GHRH) neurons in GHRH-green fluorescent protein transgenic mice. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **27**, 1631-1641, (2007).

- 123 Osterstock, G. *et al.* Somatostatin triggers rhythmic electrical firing in hypothalamic GHRH
neurons. *Sci Rep* **6**, 24394, (2016).
- 124 Sanchez-Cardenas, C. *et al.* Pituitary growth hormone network responses are sexually
dimorphic and regulated by gonadal steroids in adulthood. *Proceedings of the National
Academy of Sciences of the United States of America* **107**, 21878-21883, (2010).
- 125 Schaeffer, M. *et al.* Influence of estrogens on GH-cell network dynamics in females: a live in
situ imaging approach. *Endocrinology* **152**, 4789-4799, (2011).
- 126 Alatzoglou, K. S., Webb, E. A., Le Tissier, P. & Dattani, M. T. Isolated growth hormone
deficiency (GHD) in childhood and adolescence: recent advances. *Endocrine reviews* **35**, 376-
432, (2014).
- 127 Pekic, S. & Popovic, V. Alternative causes of hypopituitarism: traumatic brain injury, cranial
irradiation, and infections. *Handb Clin Neurol* **124**, 271-290, (2014).
- 128 Hindmarsh, P. C. & Dattani, M. T. Use of growth hormone in children. *Nat Clin Pract
Endocrinol Metab* **2**, 260-268, (2006).
- 129 Andoniadou, C. L. *et al.* Sox2(+) stem/progenitor cells in the adult mouse pituitary support
organ homeostasis and have tumor-inducing potential. *Cell stem cell* **13**, 433-445, (2013).
- 130 Olarescu, N. C. & Bollerslev, J. The Impact of Adipose Tissue on Insulin Resistance in
Acromegaly. *Trends in endocrinology and metabolism: TEM* **27**, 226-237, (2016).
- 131 Dimaraki, E. V., Jaffe, C. A., DeMott-Friberg, R., Chandler, W. F. & Barkan, A. L.
Acromegaly with apparently normal GH secretion: implications for diagnosis and follow-up. *J
Clin Endocrinol Metab* **87**, 3537-3542, (2002).
- 132 Holly, J. M. *et al.* Inter-relations between growth hormone, insulin, insulin-like growth factor-
I (IGF-I), IGF-binding protein-1 (IGFBP-1) and sex hormone-binding globulin in acromegaly.
Clin Endocrinol (Oxf) **34**, 275-280, (1991).
- 133 Narayanaswamy, S. *et al.* Subcutaneous infusion of kisspeptin-54 stimulates gonadotrophin
release in women and the response correlates with basal oestradiol levels. *Clin Endocrinol
(Oxf)*, (2015).
- 134 Castinetti, F., Davis, S. W., Brue, T. & Camper, S. A. Pituitary stem cell update and potential
implications for treating hypopituitarism. *Endocrine reviews* **32**, 453-471, (2011).
- 135 Andoniadou, C. L. *et al.* Identification of novel pathways involved in the pathogenesis of
human adamantinomatous craniopharyngioma. *Acta neuropathologica* **124**, 259-271, (2012).
- 136 Eckstrum, K. S., Weis, K. E., Baur, N. G., Yoshihara, Y. & Raetzman, L. T. Icam5 expression
exhibits sex differences in the neonatal pituitary and is regulated by estradiol and bisphenol A.
Endocrinology, en20151521, (2016).
- 137 Le Tissier, P. R. & Mollard, P. Bisphenol A Effects on Gonadotroph Function: Disruption of
Pituitary Cell-Cell Communication? *Endocrinology* **157**, 1324-1325, (2016).
- 138 Le Tissier, P. R. *et al.* Anterior pituitary cell networks. *Front Neuroendocrinol* **33**, 252-266,
(2012).
- 139 Mollard, P., Hodson, D. J., Lafont, C., Rizzoti, K. & Drouin, J. A tridimensional view of
pituitary development and function. *Trends in endocrinology and metabolism: TEM* **23**, 261-
269, (2012).
- 140 Waite, E. *et al.* Different degrees of somatotroph ablation compromise pituitary growth
hormone cell network structure and other pituitary endocrine cell types. *Endocrinology* **151**,
234-243, (2010).
- 141 Deneff, C. Paracrinicity: the story of 30 years of cellular pituitary crosstalk. *Journal of
neuroendocrinology* **20**, 1-70, (2008).
- 142 Schaeffer, M., Hodson, D. J., Lafont, C. & Mollard, P. Functional importance of blood flow
dynamics and partial oxygen pressure in the anterior pituitary. *Eur J Neurosci* **32**, 2087-2095,
(2010).
- 143 Ward, R. D., Stone, B. M., Raetzman, L. T. & Camper, S. A. Cell proliferation and
vascularization in mouse models of pituitary hormone deficiency. *Molecular endocrinology*
20, 1378-1390, (2006).

Author notes

Please check these figures carefully and return any comments/amendments that you might have to me as soon as possible. In particular, we would like you to check the following:

- Do the figures convey the intended message?
- Are all the labels accurate and in the right place?
- Are all the arrows in the right place?
- Are any chemical structures correct?
- Have shapes and colours been used consistently and accurately throughout the figures?
- Have any of the figures been previously published, or have they been supplied by a colleague(s) who is not a named author on the article?

To mark up any corrections, please use the commenting tools in the PDF, or print and draw by hand, rather than directly editing the PDFs.

Fig 1




Fig 2




Fig 3




Fig 4

