27

145

ALLERGIC BRONCHO-PULMONARY HELMINTHIASIS (ABPH)

VIJAYAN VK

Cardio-Pulmonary Medicine Unit, Tuberculosis Research Centre, Indian Council of Medical Research Madras 600 031 (Received original May 1991; revised October 1991)

ABSTRACT

Based on current literature, an aetiological classification of pulmonary eosinophilia due to helminths including tropical eosinophilia is proposed.

Introduction

Infestations with helminths (cestodes, nematodes and trematodes) are the commonest causes of pulmonary eosinophilia in tropical countries (1-3). Other causes of pulmonary eosinophilia include Allergic Broncho-Pulmonary Aspergillosis (ABPA) (4), chronic eosinophilic pneumonia (5), cryptogenic pulmonary fibrosis (6), Wegener's granulomatosis (7), lymphomatoid granulomatosis eosinophilic granuloma of lung (9), the (8) Churg-Strauss syndrome (10), drug hypersensitivity reactions (11) and infections such as brucellosis (12), coccidioidomycosis (13), corynebacterium pseudotuberculosis (14) and tuberculosis (15).

Pulmonary eosinophilia due to helminths :

One of the commonest forms of pulmonary in tropical countries is due to hyeosinophilia persensitivity reaction to an occult form of filariasis termed as tropical (filarial) eosinophilia (16) or Tropical Eosinophilia (TE) (17-20). TE presents with symptoms suggestive of bronchial asthma such as cough, dysphoea. nocturnal wheezing discomfort, occassionally associated and chest with such constitutional symptoms as weight loss, anorexia and fever (17,21). Physical examination reveals rhonchi and rales (21). Chest radiographs show a variety of abnormalities, including diffuse miliary mottling, pleural effusions, and uncommonly cavitation and large areas of consolidation (22). High levels of blood and lung eosinophils with elevated serum and lavage fluid levels of IgE and filaria - specific IgE and IgG antibodies (17, 23, 24) had also been demonstrated. Pulmonary function studies had shown airway obstruction in addition to reduced lung volumes and reduced diffusing capacity (21, 25-27). Following a three -week course of diethylcarbamazine (DEC), there was incomplete reversal of clinical, haematological, radiological and physiological changes

one month after starting treatment (25). In conformity with these findings, a chronic respiratory tract inflammation as evidenced by a persistent eosinophilic alveolitis with lower respiratory tract inflammatory cells spontaneously releasing exaggerated amounts of 0, and H,O, has also been demonstrated (28). It had also been shown that, at one year, patients with eosinophilic alveolitis had a significantly reduced diffusing capacity despite three weeks treatment with DEC (21). It had also been reported that bronchiectasis could occur during and persist after clinical attacks of TE (29). Therefore, there is an urgent need to undertake controlled clinical trials in TE to assess the role of corticosteroids in preventing the occurence of irreversible radiological and lung function changes.

A similar syndrome of pulmonary eosinophilia with respiratory symptoms can result from infestations with other helminths such as Ascaris (30), Ancylostoma species Strongyloides ster-(31), coralis (32) and Toxocara species (33, 34). Currently available immunological investigations of antibody assays (35) may not differentiate the eosinophilic syndromes due to various helminthic infections because of antigenic similarities. Because of the probable broad-spectrum anti-helminthic action of DEC, patients with pulmonary eosinophilia due to various helminthic infections respond to treatment with DEC (Personal observation). Thus it appears that the syndrome of tropical eosinophilia is an end result of a common pathogenesis resulting from infestations with various helminths. Though the syndrome of Tropical eosinophilia is common in tropical and sub-tropical countries, it had also been described in subjects from temperate countries on exposure to appropriate infections in endemic areas (36). Therefore, it will be ideal to have a better name for this syndrome based on aetiology and research should be directed to identify tests that may differentiate various types of pulmonary

146

eosinophilia.

Allergic broncho-pulmonary mycosis :

A similar disease resulting from a hypersensitivity reaction to fungal antigens (Aspergillus species) has been termed Allergic Broncho-Pulmonary Aspergillosis (ABPA)(4). ABPA also manifests as asthma with elevated serum levels of total IgE and specific antibodies to aspergillus antigens (37). Pulmonary function tests had revealed both obstructive and restrictive ventilatory defects (38). Irreversible lung changes including proximal bronchiectasis can occur if ABPA is left untreated (39). Since other fungi such as Candida (40, 41), Helminthosporium (42), Stemphyllium (43) and Curvaluria (44) can cause a similar syndrome, it was suggested that the term "Allergic Broncho-Pulmonary Fungal diseases" was a better name for the disease (45) (Table 1).

TABLE 1

Allergic Broncho-Pulmonary Mycoses (45)

1.	Allergic	Broncho-Pulmonary	Aspergillosis (4j.
2.	Allergic	Broncho-Pulmonary	Candidiasis (40).
3.	Allergic	Broncho-Pulmonary	Helminthosporiasis (42).
4.	Allergic	Broncho-Putmonary	Stemphyliosis (43).
5.	Allergic	Broncho-Pulmonary	Curvulariasis (44).

Members in parenthesis refer to articles quoted in reference

Comparison of ABPA and TE :

Thus, there appears to be a similarity in pathogenesis of pulmonary eosinophilia resulting from hypersensitivity reactions to fungi and helminths. A comparison of two diseases viz. ABPA and TE which are prototypes of pulmonary eosinophilia due to fungi and helminths are shown in Table 2.

Comparison of ABPA and TE

		ABPA	TE
1.	Features of Asthma	+	+
2.	Blood and Sputum eosidophila	a + .	+
3.	Lung Eosinophilia	+	+
4.	Chest Radiographic infiltration	15 + .	/ +
5.	Total IgE	Ť	Î
6.	Specific IgG and IgE	° ↑	Ť
7.	Pulmonary Function	Obstruc-	Obstruc-
		tion	tion
		+	+
		Restriction	Restriction
8.	Irreversible lung damage		
	including bronchiectasis	Can occur	Can occur
9.	Antigens	Fungus	Helminths
10.	Treatment	Corticosteroids	DEC
			? Corticoteroids

LUNG INDIA (1991) IX, NO.4 (P.145-148)

Both result from exaggerated immune-responses to respective antigens as evidenced by elevated levels of total IgE and specific antibodies (17,35,37). Both have bronchial and lung parenchymal abnormalities as revealed by similar pulmonary function abnormalities (25,38). These may be due to type I and type III hypersensitivity reactions. Lung eosinophils in both diseases can cause irreversible lung changes if left untreated (17,29,39). Therefore, it is suggested that the term "Allergic Broncho-Pulmonary Helminthiasis" is a better term, for pulmonary eosinophilia due to helminthic infections. Terminologies that are appropriate for pulmonary eosinophilia resulting from various helminthic infections are given in Table 3.

TABLE 3

Allergic Broncho-Pulmonary Helminthiasis

Suggested list of Infestations

1.	Allergic	Broncho-Pulmonary	Filariasis
2.	Allergic	Broncho-Pulmonary	Ascariasis
3.	Allergic	Broncho-Pulmonary	Ancylostomiasis
4.	Allergic	Broncho-Pulmonary	Strongyloidiasis
5.	Allergic	Broncho-Pulmonary	Toxocariasis
-			

Tropical Eosinophilia can then be termed as "Allergic Broncho-Pulmonary Filariasis".

In case of an incorrect diagnosis in a patient with pulmonary eosinophilia due to strongyloidiasis and if this patient is treated with steroids, the parasite may become disseminated (46). This emphasises the need for an aetiological diagnosis in patients with pulmonary eosinophilia.

REFERENCES

- 1. Crofton JW, Livingstone JL, Oswald NC and Roberts ATM : Pulmonary eosinophilia. Thorax, 1952; 7: 1-35.
- Spry CJF : Lung Diseases associated with eosinophils : In : Current perspectives in Allergy. Edinburgh. Churchill Livingston, 1982.67-77.
- 3. Sharma O.P. Pulmonary eosinophilia. Lung India, 1988; 6:39-47.
- Hinson KFW, Moon AJ; and Plummer NS : Bronchopulmonary aspergillosis : a review and a report of eight new cases. Thorax, 1952; 7: 317-33.
- Carrington C, Addington W, Goff A, Madoff L, Marks A, Schwaber J and Gaensler E. : Chronic eosinophilic penumonia. New Engl. J.Med. 1969; 280: 737-98.
- Turner-Warwick M and Haslam PL : The value of serial broncho-alveolar lavagas in assessing the clinical progress of patients with cryptogenic fibrosing alveolitis. Am.Rev.Respir.Dis. 1987, 135: 26-34.

Vijayan : Allergic Bronchopulmonary Helminthiasis

- Hunninghake GW and Fauci AS : State of the art : Pulmonary involvement in the collagen vascular diseases. Am.Rev.Respir.Dis. 1979, 119: 471-503.
- 8. Leibow AA, Carrington CRB and Friedman PJ : Lymphomatoid granulomatosis. Hum.Pathol, 1972, *3*: 457-558.
- Case Records of the Massachusettes General Hospital. Weekly Clinico-Pathological exercises. Case 8-1986. A 39year old woman with bilateral pulmonary infiltrates. New.Engl.J.Med. 1986, 314: 564-74.
- Churg J and Strauss L.: Allergic granulomatosis, allergic angiitis, and polyarteritis nodosa. Am J. Pathol, 1951, 27: 277-301,
- 11. Rosenow EC III : The spectrum of drug-induced pulmonary disease. Ann. Intern. Med. 1972, 77: 977-91.
- 12. Elsom KA, and Ingelfinger FJ : Eosinophilia and pneumonitis in chronic brucellosis. A report of two cases. Ann.Intern.Med. 1942, *16*: 995-1002.
- Lombard CM, Tazelaar HD and Krasne DL : Pulmonary eosinophilia in Coccidioidal infections. Chest, 1987, *91*: 734-36.
- Keslin MH, McCoy EL, McCusker JJ, and Lutch JS : Corynebacterium pseudotuberculosis. A new cause of infectious and eosinophilic penumonia. Am.J.Med. 1979, 67. 228-302.
- Vijayan VK, Reetha AM, Jawahar MS, Sankaran K and Prabhakar R: Pulmonary eosinophilia in pulmonary tuberculosis. Chest. (In press.)
- 16. Neva FA and Ottesan EA.: Tropical (filarial) eosinophilia. New Engl. J. Med, 1978, *298*: 1129-31.
- Udwadia FE : Tropical Eosinophilia : In : Pulmonary eosinophilia : Progress in Respiration Research. Vol 7. Basel : Karger, 1975; 35-155.
- Viswanathan R. : Pulmonary eosinophiliosis : Natural history, aetiology. arid pathogenesis. Ind J Chest Dis. 1963, 5: 213-30.
- Kamat SR, Warner NVU, Store SD, Kavandikar ED, and Hoskote VR. : Clinical studies in pulmonary eosinophilia.I. Comparative study of response to diethylcarbamazine and corticosteroid drugs. Ind.J. Chest Dis. 1976, 18: 221-32.
- 20. Ray D. : Relapsing tropical eosinsphilia. Ind.J. Chest Dis. 1977, 19:65-71:
- Vijayan VK : Tropical Eosinophilia : Bronchoalveolar lavage and Pulmonary pathophysiology in relation to treatment. Ph D Thesis. University of Madras, 1990.
- 22. Jain VK and Beniwal OP. : Unusual presentation of tropical pulmonary eosinophilia. Thorax; 1984, 39: 634-5.
- Pinkston P, Vijayan VK, Nutman TB, Rom WN, O'Donnell KM, Cornelius MJ, Kumaraswamy V, Ferrans VJ, Yenokida G, Thiruvengadam KV, Tripathy SP, Ottesen EA and Crystal RG : Acute tropical eosinophilia : characterisation of the lower respiratory tract inflammation and its response to therapy. J.Clin Invest. 1987, 80: 216-25.
- Nutman TB, Vijayan VK, Pinkston P, Kumaraswamy V, Steel C, Crystal RG and Ottesen EA : Tropical Pulmonary eosinophilia : Analysis of Antifilarial Antibody localised to the lung. J. Infect Dis, 1989, 160: 1042-50.

- Vijayan VK, Kuppurao KV, Sankaran K, Venkatesan P and Prabhakar R.: Tropical eosinophilia : Clinical and Physiological response to diethylcarbamazine. Respiratory Medicine, 1991, 85: 17-20.
- 26. Poh SC : The course of lung function in treated tropical eosinophilia. Thorax. 1974, 29: 710-12.
- Mann JM and Heurich AE : Response of diffusion capacity in the treatment of Tropical Esoinophilia. Chest, 1991, 99: 776-77.
- Rom WN, Vijayan VK, Cornelius MJ, Kumaraswamy V. Prabhakar R, Ottesen EA and Crystal RG: Persistent Lower respiratory tract inflammation associated with interstitial lung disease in patients with tropical pulmonary eosinophilia following coventional treatment with diethylcarbamazine. Am.Rev.Respir.Dis. 1990, 142: 1088-92.
- Rab SM, Nurul Haq AQM and Chaudhary GM: Tropical eosinophilia and bronchiectasis. Brit J Dis Chest 1963, 57: 103-6.
- Loffler W: Zur Differential Diagnose der Lungeninfiltrierugen 1. Fruhinfiltrate unter besonderer Beruchsichtigung der Ruckbildungzeten. 11. Uber fluschitge succedan-infiltrate (mit eosinophilie). Beitrage zur klinik der Tuberkulose and Spezifischen Tuberkulose Forschung. 1932, 79: 338-82.
- Butland. RJ, and Coulson IH.: Pulmonary eosinophilia associated with cutaneous larva migrans. Thorax, 1985. 40: 76-7.
- deMessias IT, Telles FQ. Boaretti AC, Silva S, Guimarres LM and Genti RM: Clinical, immunological and epidemiological aspects of strongyloidiasis in an endemic area of Brazil. Ailergol Immunopathol. (Madr). 1987, 15: 37-41.
- 33. Chaudhari RN and Saha TK; Tropical eosinophilia: Experiments with toxocara canis. Lancet 1959, 2: 493-4.
- Beaver PC: Toxocariasis (visceral larva migrans) in relation to tropical eosinophilia. Bull Soc. Patho. Extol. 1963, 55: 555-76.
- **35.** Saha K, Sarkar. N, Paul DN, and Ray D.: Serum immunoglobulins and complement profile in parasitic diseases. Ind J Med Res. 1979, *70*: 22-32.
- **36.** Hodes PJ and Wood R.D.: Eosinophilic Lung (Tropical eosinophilia): Am.J.Med. Sci. 1945, *210*. 288-95.
- **37.** Rosenberg M, Patterson R, Mintzer R, Cooper BJ, Roberts M, and Harris KE : Clinical and immunologic. criteria for the diagnosis of allergic bronchopulmonary aspergillosis. Ann. Intern Med. 1977, *86*: 405-14.
- Malo, JL, Hawkins R, and Pepys J: Studies in chronic Allergic bronchopulmonary aspergillosis. 1. Clinical and Physiological findings. Thorax. 1977; 32 254-61.
- Chapman BJ, Capewell S, Gibson R, Greining AP and Crompton GK: Pulmonary eosinophilia with and without allergic bronchopulmonary aspergillosis. Thorax, 1989. 44: 919-24.
- Pepys J.: Hypersensitivity disease of the lungs due to fungi and organic dusts. Monogr. Allergy 1969; *4*: 63-3.
- Slavin RG, Stanzyk DJ, Lonigro AJ, and Broun GO: Allergic bronchopulmonary aspergillosis - a North Americal rarity. Clinical and Immunologic characteristics. Am J Med, 1969, 47. 306-13.

- 42. Dolan CT, Wood LA, and Dines DE: Bronchopulmonary helmintnosporiosis. Am J Clin Path 1970. 53: 235-42.
- Benatar SR, Allan B, Hewitson RP and Don PA. : Allergic bronchopulmonary stemphyliosis. Thorax, 1980, 35: 515-18.
- 44. Mcheer R., Kroenert DB, Elder JL and Fondist JH: Allegic bronchopulmonary disease caused by Curvularia lunata and Drechslera hawaiiensis. Thorax 1981. *36*: 338-44.
- 45. Glancy JJ, Elder JL and McAleer R.: Allergic bronchopulmonary fungal disease without- clinical asthma. Thorax,

1981, 36: 345-49.

- Case records of Massachusetts General Hospital. Weekly Clinico-Pathological exercises. Case 47-1987. A 79-year old Cuban native with asthma, weight loss, vomitting eosinophilia and past meningitis. New Engl.J. Med, 1987, 317: 1332-42.
 - Correspondence/ request for reprints : Dr. V.K.Vijayan Tuberculosis Research Centre, Spurtank Road, Madras 600 031.