

ROLE OF HUMAN LEUCOCYTE ANTIGEN (HLA) AND NON-HLA GENES IN SUSCEPTIBILITY OR RESISTANCE TO PULMONARY TUBERCULOSIS

Human beings live in an environment endemic to various diseases. Under such conditions, some individuals are susceptible to certain diseases while most are resistant to those diseases. This raises many questions. One such question is whether the person who is susceptible is genetically different from the person who is resistant to the disease? Similarly, in a family, one of the parents may be affected with a particular disease (*e.g.* diabetes); among the children born to the parents, if one child (younger one) develops the disease and the other child (elder one) does not, the question arises why did the younger one develop the disease and the elder did not, when both of them were living together with the affected parent. This raises the possibility of the disease being inherited from parents. If genetic factors play a major role in disease susceptibility or resistance, in what way are they associated with the disease development? Do they control the immune response to an antigen and/or pathogen? Do they play a role in the immune mechanism of susceptibility or resistance to disease? This article gives an account of disease susceptibility or resistance in pulmonary tuberculosis in the context of the host genetic factors, *i.e.* HLA (Human Leucocyte Antigens) and non-HLA genes and their gene products.

HLA and disease association - a hypothesis

For considering genetic susceptibility to various diseases and disease association in the context of HLA, several hypotheses exist for explaining MHC (Major Histocompatibility Complex) associated diseases. The main hypotheses are:-

- (1) That HLA A, B, C (Class-I) and DR, DQ and DP (Class-II) antigens could act directly as disease susceptibility agents. For this, three possible mechanisms have been suggested: (a) There could be antigenic cross reactivity or mimicry between infectious organisms and a given HLA antigen. This phenomenon is termed as "molecular mimicry". Serological cross-reaction between HLA 827 antigen and the bacterial strains *Klebsiella* and *Shigella* has been

identified^{1,2}. This means that common antigenic determinants are shared by HLA B27 and the bacteria; (b) HLA antigens could act as receptors for microorganism; and (c) HLA antigens could influence particular immune responses, acting as immune response (Ir) genes. It has been shown that Ir genes regulate the immune response to any antigen or pathogen).

- (2) The MHC genes may be physically close to the chromosome region that carries a gene conferring susceptibility or resistance to a particular disease. This hypothesis may explain the lack of complete association, and geographical variation in the association, due to linkage disequilibrium.
- (3) Genetic susceptibility due to non-HLA genes: Though classical genetic studies in humans and experimental models have clearly documented the primary contribution of the MHC genes, these genes themselves appear to be insufficient in conferring susceptibility or resistance to disease.

Susceptibility or resistance to tuberculosis

Pulmonary tuberculosis is a granulomatous lung disease caused by *Mycobacterium tuberculosis*. Susceptibility to tuberculosis has been suggested to be multifactorial. Though environmental and socio-economic factors are primarily related, numerous studies have emphasised the importance of host resistance and hereditary susceptibility^{4,5}. Susceptibility or resistance to tuberculosis may be under the control of the genetic make up of an individual. The host genetic factors such as HLA (Human Leucocyte Antigen) and non-HLA genes may be associated with susceptibility or resistance to tuberculosis.

HLA gene polymorphism

Finding out of HLA and non-HLA genes/gene products (antigens) which are associated with susceptibility or resistance to tuberculosis will serve

to provide HLA genetic markers to development of the disease. Moreover, studying the role of these markers in the immune mechanism underlying susceptibility or resistance to tuberculosis will be useful for understanding the immunopathogenesis of the disease. Studies of HLA and tuberculosis susceptibility have been carried out, in populations as well as in families by several groups in different part of the world. The association of various HLAs with disease susceptibility in different ethnic population groups has been shown in these studies⁶⁻¹².

The HLA association shows some geographic variation. It seems likely that evolutionary selection pressures have given rise to frequent polymorphisms in genes involved in resisting infectious pathogens and contributed to marked allele frequency differences at the same loci. When geographic variation in pathogen polymorphism is superimposed on host genetic heterogeneity, considerable variation may occur in detectable allelic association. Gene-environmental interactions are likely to introduce another layer of complexity¹³. The genes involved in defense against infectious pathogens evolve more rapidly than others.¹⁴ and excessive polymorphism in the human genome may result from selection pressures exerted by infectious diseases. Similarly, the causative organism *M. tuberculosis* also has genetic variation. During evolution, all these polymorphic forms might have evolved due to the host-parasite interaction.

Studies in non-Asian countries

A large number of HLA association studies have been carried out in non-Asian countries⁶⁻¹¹. One of the first reports of an association between HLA and tuberculosis showed an increased frequency of HLA B8 in Canada⁶. Other studies showed an increased frequency of HLA B5, B15 and DR5 in the North American blacks^{7,8}, HLA A2 and B5 in the Egyptian population and B27 in the Greek population¹¹. A negative association has been reported for DR6 in American blacks⁹. Since more interesting findings have been reported from the Asian region, the discussion is being restricted to the Asian populations.

Studies in Asian populations

Several studies of HLA association with pulmonary tuberculosis have been carried out in

Chinese^{12,15}, Korean¹⁶, Indonesian¹⁷, Indian¹⁸⁻²⁴ and Russian patients²⁵. A significantly increased frequency of HLA-DR2 was seen in the major studies^{20,21,23,24,17,25} which have revealed HLA DR2 association with higher susceptibility to tuberculosis.

Of the numerous Indian studies on HLA association with pulmonary tuberculosis¹⁸⁻²⁴, the earlier ones focussed on association of HLA Class-I antigens^{18,19} in tuberculosis patients and later, in family studies in north India^{21,23} and south India^{20,24}. An increased frequency of HLA DR2 in pulmonary tuberculosis patients compared with the control subjects was found. Molecular study has revealed that the allele DRB1 * 1501 of HLA-DR2 was higher compared with DRB1 *1502 in north Indian patients²³. While both DRB1 * 1501 and * 1502 were increased in south Indian patients²⁷, no deviation was seen between * 1501 and *1502 alleles. Association of HLA DP B102 and DQ1 with susceptibility to pulmonary tuberculosis has also been shown in patients in south India²⁷. On the other hand, a decreased frequency of HLA-DR6 in pulmonary tuberculosis patients suggests that DR6 may be associated with resistance to tuberculosis^{21,24}.

Previous investigations showed a varied picture of the association in different populations mainly because of geographical distribution of antigens and the ethnic background of the populations. However, studies carried out in India, Indonesia and Russia have shown a definite association of HLA DR2 and DQ1 with pulmonary tuberculosis. It is well established that DR2 is in linkage disequilibrium with DQ1.

In a recent study²⁷, HLA DRB1 * 1501, DRB1 *08 HLA DQB1 *0601 (a subtype of HLA-DQ1) and DPB1 *02 were found to be positively associated with susceptibility to pulmonary tuberculosis while a negative association (preventive fractions associated with resistance) has also been identified (DRB1 *11(5), DRB1 *10, DQB1 *0501 and DPB1 *08). Haplotype analysis also supports the DRB1 * 1501 -DQB1 *0601 association with susceptibility to pulmonary tuberculosis.

HLA DR2 and immunity to tuberculosis

The association of HLA DR2 with severity of pulmonary tuberculosis has also been emphasised in Indian population. In south Indian patients, DR2 has been shown to be more strongly associated with

far-advanced, smear positive tuberculosis than with minimal and moderately advanced disease²⁰.

In Russian patients, DR2 was associated with chronic fibrocavitary cases who failed to react to PPD²⁵. Our study has revealed that HLA DR2 is associated with both low and high tuberculin test response²⁸. Further, among north Indian patients a high frequency of DR2 was seen in treatment failures compared with healthy controls and treatment responders²³. Though severity of the disease is associated with HLA DR2, follow up of patients treated with short course chemotherapy did not reveal any difference in DR2 frequency between relapse and quiescent patients²⁹.

Patients bearing DR2 antigen have higher level of antibodies to *M. tuberculosis* compared with other patients. Further, there is evidence favouring a spectrum wherein cell-mediated and humoral immunity are at the opposing ends of the spectrum. Our recent study suggests that in active tuberculosis, patients having DR2 and a trend towards higher antibody response to *M. tuberculosis* antigens possibly show that higher antibody titre may have suppressed the cell mediated immune response²⁶. It has been suggested that patients and controls having DR2 give a high reaction to PPD²⁰. Further study has revealed that DRB1 * 1501 (a subtype of DR2) is responsible for greater tuberculin induration compared with DRB1 * 1502³⁰. Similarly, HLA DR2 is found associated with lymphocyte response to *M. tuberculosis* culture filtrate antigen both in active and cured (quiescent) tuberculosis patients³¹. A decreased level of plasma lysozyme, a phagocytic enzyme (lysosomal enzyme) involved in microbicidal activity has been found in DR2 positives as compared with non-DR2 active pulmonary tuberculosis patients³². This suggests that susceptibility to pulmonary tuberculosis is not only influenced by HLA DR2 genes/gene products but by non-HLA gene factors as well.

Recently, association of multi candidate genes has been suggested for various infectious diseases¹³. In north Indian pulmonary tuberculosis patients, compared with control subjects, the "Transporter" associated with antigen processing gene 2 (TAP2) has been found associated with active tuberculosis, along with HLA DR2³³. Studies on non-HLA gene polymorphism have revealed that functional mutant

homozygotes of mannose binding protein genes are associated with pulmonary tuberculosis independent of HLA DR2³⁴. Moreover, study of Vitamin D receptor (VDR) gene polymorphism has revealed a significant increase of VDR mutant homozygotes (tt) in female patients compared with female contacts while a significant increase of wild type homozygotes (TT) was observed among female contacts compared with female cases³⁵, suggesting that HLA DR2 gene/gene products in combination with other HLA and/or non-HLA genes, play a major role in susceptibility to pulmonary tuberculosis.

Non-HLA gene polymorphism

Non-HLA gene polymorphism studies in tuberculosis have revealed definite association between tuberculosis and the haptoglobin 2-2 phenotype³⁶, except in Indonesia³⁷ and India¹⁹.

Recently, human genome analysis has revealed several candidate non-HLA genes, including cytokine genes, but due to point mutations most of these non-HLA genes occur as diallelic polymorphic forms. Such polymorphic genes have been shown to be associated with susceptibility to a number of infectious and non-infectious diseases. Some of the important non-HLA genes are as follows:

Mannose-binding protein

Mannose-binding protein (MBP), also known as mannose-binding lectin (MBL) is an acute phase protein secreted by liver. It binds mannose and N-acetylglucosamine terminated glycoproteins and plays an important role in host defence against pathogens. Upon binding with certain carbohydrate moieties, such as terminal N-acetyl glucosamine or mannose, on various pathogens, MBP activates complement via specific protease and acts directly as an opsonin using the C1q receptor on macrophages. Mutations are found at the coding regions of the MBP genes *i.e.* at codons 52, 54 and 57 that lead to low or near absent serum MBP levels in heterozygotes and homozygotes, respectively. Low serum level of MBP is associated with a common opsonic defect and is frequent in recurrent infections of infancy and possibly infections in adult life.

Studies carried out in Gambia have revealed that MBP is not associated with pulmonary

tuberculosis³⁸. But in an Indian population, an increased genotype frequency of MBP functional mutant homozygote (including 52, 54 and 57) was found in pulmonary tuberculosis (10.9%) compared with control subjects (1.8%). Analysis of association of MBP genes and HLA DR2 has revealed that these genes are associated with susceptibility to pulmonary tuberculosis, independent of each other³⁹.

Vitamin D receptor

Vitamin D regulates calcium metabolism. Vitamin D3 is an immunoregulatory hormone and activates monocytes and stimulates cell-mediated immunity. Further, Vitamin D3 is one of the few mediators which have been shown to impair the growth of *M. tuberculosis* in macrophages³⁹. Vitamin D effects are exerted through interaction with Vitamin D receptor (VDR). It has been suggested that the mutant allele of the VDR gene may be associated with reduced VDR mRNA expression. Allelic variants of Vitamin D receptor appear to be associated with differential susceptibility to several infectious and non-infectious diseases¹³. Vitamin D receptor gene polymorphism studied in Gambian tuberculosis patients suggested that the tt genotype (mutant) is less frequent in cases of pulmonary tuberculosis⁴⁰ but in Indian leprosy patients the tt genotype has been shown to be associated with tuberculoid leprosy whereas the opposite TT genotype is associated with lepromatous leprosy⁴¹. Our study of VDR gene polymorphism in pulmonary tuberculosis patients revealed a significant increase in VDR mutant homozygotes (tt) in female patients as compared with female contacts (spouses of male patients). Similarly, a significant increase of wild type homozygotes (TT) was observed in female contacts compared with female patients. This suggests that the mutant genotype variant of VDR gene (tt) is associated with susceptibility to pulmonary tuberculosis in female patients and the wild type genotype (TT) with resistance in female contacts, in India³⁵.

Natural resistance-associated-macrophage-protein-1 (NRAMP-1)

Studies in mice of acute susceptibility to infection with *Leishmania donovani* and *Salmonella typhimurium* led to the mapping of a major gene denoted as *Lsh* or *lty* on mouse chromosome-1. Subsequently, this was found to affect susceptibility

to some strains of BCG (*Bacille Calmette Guerin*). This gene is expressed in macrophages and appears to influence macrophage activation. The equivalence for the BCG gene is the NRAMP-1 gene, which encodes for NRAMP-1 protein localised in the phagosome membrane. Several polymorphisms in the human homologue NRAMP-1 have been defined. Genetic linkage and association of tuberculosis and leprosy have been reported⁴². Recently, Bellamy *et al*⁴³ have identified variants in the NRAMP-1 gene that are associated with a significantly increased risk of pulmonary tuberculosis in Gambian population. Our study of one of the NRAMP-1 CA repeat microsatellite polymorphism at 199 and 201 region of the human NRAMP-1 gene revealed no such association in Indian patients (Selvaraj *et al.*, unpublished data) but other gene polymorphisms of the NRAMP-1 gene are under study.

Cytokine genes and receptors

An analysis of the course of infection in gene-knock-out mice has provided examples of the potential relevance of polymorphism in cytokine and cytokine receptor genes to infectious disease susceptibility in humans.

1) Tumor necrosis factor - α (TNF - α)

Increased production of inflammatory cytokines, such as tumour necrosis factor- μ (TNF- μ) has been found in tuberculosis and various other infectious diseases. The TNF - α is mainly produced by monocytes and macrophages and variant genes of TNF - α are associated with increased production of TNF - α . Association studies have been carried out on polymorphisms in and near the tumor necrosis factor (TNF) gene located in class III region of MHC. Studies of gene polymorphism, at -308 promoter region in Cambodian patients have revealed no association with TNF - α in tuberculosis patients⁴⁴. Our study of TNF- μ and -238 and -308 gene polymorphisms in Indian pulmonary tuberculosis patients also revealed no association either with susceptibility or resistance³⁵. However, studies in Indian leprosy patients have revealed that the -308 variant is associated with lepromatous leprosy⁴⁵.

2) IL-1 receptor antagonist (IL-1 RA)

Interleukin-1 receptor antagonist (IL-1 RA) is

another cytokine which competes for the IL-1 binding site. The production of IL-1 μ and IL-1 β is controlled by IL-1 RA. The association of IL-1 RA gene polymorphism has been studied in various diseases⁴⁶. Macrophages from carriers of IL-1 RA allele 2 have been shown to produce more IL-1 RA and less IL-1 μ than other genotypes⁴⁷. Our study of IL-1 RA gene polymorphism in Indian pulmonary tuberculosis patients revealed no association with any of the genotypes³⁵ but spinal tuberculosis patients showed an increased frequency of genotype 22 compared with the control subjects (Selvaraj *et al.*, unpublished data).

Non-HLA gene polymorphisms such as Interferon γ receptor/gene polymorphism have shown recessive susceptibility to atypical mycobacterial species such as *M. fortuitum*, *M. chelonii* and *M. avium*. A different mutation was identified in a child with fatal disseminated BCG infection.

Over the last few years, interest has increased in both MHC and non-MHC genes that are important to the immunogenetics of human infectious diseases. Susceptibility is mainly determined by a large or very large number of polymorphic host genes, each contributing a relatively small effect. Polygenic determination/candidate gene studies in infectious disease have been more rewarding than other multifactorial diseases. With better knowledge of the mechanism of immunity, variants are being found that are likely to affect resistance to many infectious pathogens¹³. Finding out new MHC and non-MHC genes which are associated with susceptibility or resistance to disease will serve as genetic markers for predetermining development of disease. Moreover, the role of these multi-candidate genes on the immune mechanism of disease susceptibility or resistance will throw light on the mechanism of susceptibility or resistance to tuberculosis.

ACKNOWLEDGEMENTS

The author sincerely thanks Dr.P.R.Narayanan, Director, Tuberculosis Research Centre, Chennai for encouragement in writing this article and Mrs.V.Shanthi for typing the manuscript.

P.Selvaraj

*Tuberculosis Research Centre (ICMR),
Chennai*

REFERENCES

1. Van Boheman-OG, Grument FC, Zanen HC. Identification of HLA B27MI and M2 cross-reactive antigens in *Klebsiella*, *Shigella* and *Yersinia*. *Immunol* 1984, 51, 607
2. Schwimmbeck PL, Yu DTY. Autoantibodies to HLA-B27 in the sera of HLA-B27 patients with ankylosing spondylitis and Reiter's syndrome. Molecular mimicry with *Klebsiella pneumoniae* as potential mechanism of autoimmune disease. *J Exp Med* 1987, 166, 173
3. Mc Devitt H. Ir genes : towards the year 2000. *Res Immunol* 1991, 142, 509
4. Puffer RR (ed). Family susceptibility to tuberculosis. Harvard University Press, Cambridge, p. 80. 1994
5. Comstock GW. Tuberculosis in twins : a re-analysis of the Proffit survey. *Am Rev Respir Dis* 1978, 117, 621
6. Selby R, Barnard JM, Buchler SK, Crumley J, Larsen B, Marshall WH. Tuberculosis associated with HLA-B8, Bfs in a Newfoundland community study. *Tissue Antigens* 1978, 11, 403
7. Al-Arif LI, Goldstein RA, Affronti LF, Janicki BW. HLA-BW15 and tuberculosis in a North American Black population. *Am Rev Respir Dis* 1979, 120, 1275
8. Cox RA, Arnold DR, Cook D. *et al.* HLA phenotypes in Mexican Americans with tuberculosis. *Am Rev Respir Dis* 1982, 126, 653
9. Hwang CH, Khan S, Ende N, Mangura BT, Reichman LB, Chou J. The HLA-A, B and DR phenotypes and tuberculosis. *Am Rev Respir Dis* 1985, 132, 382
10. Hafez M, EL\I-Salab SH, El-Shennawy F, Bassiony MR. HLA antigens and tuberculosis in the Egyptian population. *Tubercle* 1985, 66, 35
11. Zervas J, Constantopoulos C, Toubis M, Anagnostopoulos D, Cotsovouloce V. HLA-A and -B antigens and pulmonary tuberculosis in Greeks. *Br J Dis Chest* 1987, 81, 147
12. Hawkins BR, Higgins DA, Chan SL, Lowrie DB, Mitchison DA, Girling DJ. HLA typing in the Hong Kong Chest Service/British Medical Research Council study of factors associated with the breakdown to active tuberculosis of inactive pulmonary lesions. *Am Rev Respir Dis* 1988, 138, 1616
13. Hill AVS. The immunogenetics of human infectious diseases. *Annu Rev Immunol* 1998, 16, 593
14. Murphy PM. Molecular mimicry and the generation of host defense protein diversity. *Cell* 1993, 72, 823
15. Jiang ZF, An JB, Sun YP, Mittal KK, Lee TD. Association of HLA BW35 with tuberculosis in the Chinese. *Tissue Antigens* 1983, 22, 86
16. Chung TH, Jung TH, Kim JC *et al.* HLA antigens in Korean patients with tuberculosis. In : HLA in Asian Oceania. *Proceedings of the Third Asia-Oceania Histocompatibility Workshop Conference* (eds). Hokkaido University Press. Sappora, Japan. page 644 1986
17. Bothamley GH, Beck JS, Schreuder GMT. *et al.* Association of tuberculosis and *M. tuberculosis* - specific antibody levels with HLA. *J Infect Dis* 1989, 159, 549

18. Papiha SS, Wentzed J, Behjati F. Human leukocyte antigens and circulating immunoglobulin levels in Indian patients with pulmonary tuberculosis. *Tubercle* 1985, 66, 25
19. Papiha SS, Singh BN, Lanchbury JSS, *et al.* Association of HLA and other genetic markers in south Indian patients with pulmonary tuberculosis. *Tubercle* 1987, 68, 159
20. Brahmajothi V, Pitchappan RM, Kakkanaiah VN, *et al.* Association of pulmonary tuberculosis and HLA in south India. *Tubercle* 1991, 72, 123
21. Singh SPN, Mehra NK, Dingley HB, Pandey JN, Vaidya MC. Human leukocyte antigen (HLA)-linked control of susceptibility to pulmonary tuberculosis and association with HLA-DR types. *J Infect Dis* 1983, 148, 676
22. Sanjeevi CB, Narayanan PR, Prabhakar R, *et al.* No association or linkage with HLA-DR or -DQ genes in south Indians with pulmonary tuberculosis. *Tuberc Lung Dis* 1992, 73, 280
23. Rajalingam R, Mehra NK, Jain RC, Mynedu VP, Pande JN. Polymerase chain reaction based sequence-specific oligonucleotide hybridisation analysis of HLA-Class II antigens in pulmonary tuberculosis: Relevance to chemotherapy and disease severity. *J Infect Dis* 1996, 173, 669
24. Selvaraj P, Uma H, Reetha AM, *et al.* HLA antigen profile in pulmonary tuberculosis patients and their spouses. *Ind J Med Res* 1998, 107, 155
25. Khomenko AG, Litvinov VI, Chukanova VP, Pospelov LE. Tuberculosis in patients with various HLA phenotypes. *Tubercle* 1990, 71, 187
26. Selvaraj P, Uma H, Reetha AM, Xavier T, Prabhakar R, Narayanan PR. Influence of HLA-DR2 phenotype on humoral immunity and lymphocyte response to *M. tuberculosis* culture filtrate antigens in pulmonary tuberculosis. *Ind J Med Res* 1998, 107, 208
27. Pitchappan RM, Dheenadhayalan, Ravikumar M, *et al.* MHC and Th1/Th2 phenotypes in pulmonary tuberculosis in south India. *International MHC Symposium MHC in Medicine: 21st century*. 29th October - 1st November, 1998. New Delhi
28. Selvaraj P, Reetha AM, Uma H, *et al.* Influence of HLA-DR and -DQ phenotypes on tuberculin reactive status in pulmonary tuberculosis patients. *Tuberc Lung Dis* 1996, 77, 369
29. Selvaraj P, Uma H, Reetha AM, Xavier T, *et al.* Association of HLA-Class I antigens and haplotypes with relapse of pulmonary tuberculosis in patients treated with short course chemotherapy. *Ind J Tub* 1997, 44, 9
30. Pitchappan RM, Dheenadayalan V. Host genetics and Tuberculosis. *Proceedings of the Workshop on Tuberculosis Research into the 21st century*. 3-5th February. 1998. Organised by Tuberculosis Research Centre (ICMR)
31. Uma H, Selvaraj P, Reetha AM, Xavier T, Prabhakar R, Narayanan PR. Influence of HLA-DR antigens on lymphocyte response to *Mycobacterium tuberculosis* culture filtrate antigens and mitogens in pulmonary tuberculosis. *Tuberc Lung Dis* 1999 (in Press)
32. Selvaraj P, Kannapiran M, Reetha AM, Uma H, Xavier T, Narayanan PR. HLA-DR2 phenotype and plasma lysozyme, β -glucuronidase and acid phosphatase levels in pulmonary tuberculosis. *Int J Tuberc Lung Dis* 1997, 1, 265.
33. Rajalingam R, Singal DP, Mehra NK. Transporter associated with antigen-processing (TAP) genes and susceptibility to tuberculoid leprosy and pulmonary tuberculosis. *Tissue Antigens* 1997, 49, 168
34. Selvaraj P, Narayanan PR, Reetha AM. Association of functional mutant homozygotes of the mannose binding protein gene with susceptibility to pulmonary tuberculosis in India. *Tuberc Lung Dis* 1999, 79.3 (in Press)
35. Selvaraj P, Narayanan PR, Reetha AM. Association of Vitamin-D receptor genotypes with the susceptibility to pulmonary tuberculosis in female patients and resistance in female contacts. *Tuberc Lung Dis* (submitted)
36. Kharakter ZZ, Mazhak KD, Pavlenko AV. The role of genetically determined haptoglobin phenotypes in patients with destructive pulmonary tuberculosis. *Probl Tuberk* 1990, 7, 50
37. Grange JM, Kardjito T, Beck JS, Ebeid O, Kohler W, Prokop O. Haptoglobin: an immunoregulatory role in tuberculosis? *Tubercle* 1985, 66, 41
38. Bellamy R, Ruwende C, McAdam *et al.* Mannose binding protein deficiency is not associated with increased susceptibility to malaria, hepatitis-B nor tuberculosis in Africans. *Quarterly Journal of Med* 1998, 91, 13
39. Rook GAW. The role of Vitamin-D in tuberculosis. *Am Rev Respir Dis* 1998, 138, 768
40. Bellamy R, Ruwende C, Corrah T, *et al.* Resistance to tuberculosis and persistent hepatitis-B virus infection in Africans are associated with variation in the Vitamin-D receptor gene. *J Infect Dis* 1999, 179, 721
41. Roy S, Frodsham A, Saha B, Hazra SK, Mascie-Taylor CGN and Hill AVS. Association of vitamin-D receptor genotype with leprosy type. *J Infect Dis* 1999, 179, 187
42. Liu J, Fujiwara TM *et al.* Identification of polymorphisms and sequence variants in the human homologue of the mouse natural resistance associated macrophage protein gene. *Am J. Hum Genet* 1995, 56, 845
43. Bellamy R, Ruwende C, Corrah T, *et al.* Variations in the NRAMP-1 gene and susceptibility to tuberculosis in West Africans. *N Engl J Med* 1998, 338, 640
44. Goldfeld AE, Delgado JC *et al.* Association of an HLA-DQ allele with clinical tuberculosis. *JAMA* 1998, 279, 226
45. Roy S, McGuire W, Mascie-Taylor CGN, *et al.* Tumor necrosis factor promoter polymorphism and susceptibility to lepromatous leprosy. *J Infect Dis* 1997, 176, 530
46. Arend WP, Maylak M, Guthridge CJ, Gabay C. Interleukin-1 Receptor antagonist: Role in Biology. *Annu Rev Immunol* 1998, 16, 27
47. Danis VA, Millington M, Hyland VJ, Grennan D. Cytokine production by normal human monocytes: inter-subject variation and relationship to an IL-1 receptor antagonist (IL-1RA) gene polymorphism. *Clin Exp Immunol* 1995, 99, 303.