

# Flip Distance to a Non-crossing Perfect Matching

Édouard Bonnet \*

Tillmann Miltzow †

## Abstract

A *perfect straight-line matching*  $M$  on a finite set  $P$  of points in the plane is a set of segments such that each point in  $P$  is an endpoint of exactly one segment.  $M$  is *non-crossing* if no two segments in  $M$  cross each other. Given a perfect straight-line matching  $M$  with at least one crossing, we can remove this crossing by a flip operation. The flip operation removes two crossing segments on a point set  $Q$  and adds two non-crossing segments to attain a new perfect matching  $M'$ . It is well known that after a finite number of flips, a non-crossing matching is attained and no further flip is possible. However, prior to this work, no non-trivial upper bound on the number of flips was known. If  $g(n)$  (resp.  $k(n)$ ) is the maximum length of the longest (resp. shortest) sequence of flips starting from any matching of size  $n$ , we show that  $g(n) = O(n^3)$  and  $g(n) = \Omega(n^2)$  (resp.  $k(n) = O(n^2)$  and  $k(n) = \Omega(n)$ ).

Van Leeuwen and Schoone showed with the same argument and the same definition of flip how to transform a Hamilton cycle to a non-crossing Hamilton cycle on a set of  $n$  points within  $O(n^3)$  flips [17]. Therefore, we do not consider the main result (our upper bound on  $g(n)$ ) as a new contribution, because the used technique is exactly the same.

We want to use these proceedings to draw attention again on this old problem and hope to stimulate research that will close the gap between the upper and lower bound.

## 1 Introduction




Figure 1: Two crossing segments are replaced by two non-crossing segments. There are two ways to flip.

\*Institute for Computer Science and Control, Hungarian Academy of Sciences (MTA SZTAKI), [edouard.bonnet@lamsade.dauphine.fr](mailto:edouard.bonnet@lamsade.dauphine.fr)

†Institute for Computer Science and Control, Hungarian Academy of Sciences (MTA SZTAKI), [t.miltzow@gmail.com](mailto:t.miltzow@gmail.com)

Given  $2n$  points in the plane in general position (no three points on a line), we define a *perfect straight-line non-crossing matching* as a set of  $n$  segments such that each point is incident to exactly one segment and no two segments intersect. Given  $2n$  points in the plane, it is well-known that a perfect straight-line non-crossing matching always exists. One elegant argument to see this is to start with any perfect straight-line matching, potentially self-intersecting, and remove any crossing by a flip (see Figure 1). Although the total number of crossings might increase (see Figure 5), the sum of the length of all the segments decreases (see Figure 2). Thus, the process will eventually end with a perfect non-crossing matching.




Figure 2: The two new edges (dotted) are shorter than the old edges (solid) since the dashed part to the left (resp. to the right) of the crossing is longer than the dotted segment on the left (resp. on the right).

A simpler argument is to take the first two points with lowest  $x$ -coordinate and connect them with a segment and continue with the remaining points by induction. Contrary to the first argument, this does not carry over to the bichromatic setting. In the bichromatic setting, we are given  $n$  red and  $n$  blue points and a *bichromatic matching* is a matching as above with the additional property that only segments linking points of different colors are allowed.

Motivated by this old folklore result, we investigate the question on the maximum and minimum number of flips that are necessary and sufficient to reach a straight-line non-crossing matching.

### 1.1 Preliminaries

From here on,  $P$  always denotes a set of  $2n$  points in the plane and  $M$  a perfect straight-line matching on  $P$ . Given two points  $a$  and  $b$  we denote by  $\text{seg}(a, b)$  the segment with endpoints  $a$  and  $b$ . Given a perfect straight-line matching  $M$  with at least one crossing, we can remove this crossing by a flip operation. The flip operation removes any two crossing segments

$\text{seg}(a, b)$  and  $\text{seg}(c, d)$  on a point set  $Q = \{a, b, c, d\}$  and adds two non-crossing segments ( $\text{seg}(a, c)$  and  $\text{seg}(b, d)$ ) or ( $\text{seg}(a, d)$  and  $\text{seg}(b, c)$ ) to attain a new matching  $M'$ . Matching  $M$  is a *successor* of matching  $M'$  if we can construct  $M$  from  $M'$  by a single flip. We say that  $\mathcal{M} = (M_0, \dots, M_k)$  is a valid sequence of matchings, if each matching  $M_{i+1}$  is a successor of  $M_i$  and  $M_k$  is non-crossing. The number  $k$  denotes the length of  $\mathcal{M}$ . Given a set  $P$  of  $2n$  points in the plane, we define:

$$f(M) = \max\{k : \exists \mathcal{M} \text{ of length } k \text{ with } M = M_0\};$$

$$h(M) = \min\{k : \exists \mathcal{M} \text{ of length } k \text{ with } M = M_0\}.$$

Consequently functions  $g(n)$  and  $k(n)$  are defined as:

$$g(n) = \max\{f(M) : M \text{ is a matching on } 2n \text{ points}\};$$

$$k(n) = \max\{h(M) : M \text{ is a matching on } 2n \text{ points}\}.$$

### 1.2 Results

We establish the following result:

**Theorem 1 ([17])** *Let  $n$  be a large enough natural number. Then it holds:*

$$\frac{n^2 - n}{2} = \binom{n}{2} \leq g(n) \leq n^3.$$

This result immediately carries over to bichromatic matchings. We conjecture that  $g(n) = \Theta(n^2)$ .

**Theorem 2** *Let  $n$  be a large enough natural number. Then it holds:*

$$n - 1 \leq k(n) \leq \frac{n^2}{2},$$

for some constants  $C$ .

Our proof of Theorem 2 does *not* carry over to the bichromatic case. However, we will see that the upper bound further holds if the crossing to flip is imposed at each step by an adversary and we may only choose which of the two flips (see Figure 1) we perform.

### 1.3 Related Work

The most relevant work is by van Leeuwen and Schoone, who showed the same upper bound on  $g(n)$  for Hamilton cycles instead of matchings [17]. The study of Hamilton cycles is motivated as a post-processing step for algorithms that find a traveling salesman tour on a set of points. They show that they can improve the solution to be non-crossing in  $O(n^3)$  steps.

For points in convex position, Oda and Watanabe established linear upper and lower bounds on  $g(n)$ , again on Hamilton cycles instead of matchings [15].

The combinatorial work on flip graphs of geometric structures is fairly large. See the survey by Bose and Hurtado [11] for an overview and some motivations.

Matchings, triangulations and spanning trees are commonly studied in recent work [1–9, 12, 13, 16]. Particularly interesting are triangulations of points that are in convex positions as they correspond to Catalan structures. Another interesting application comes from Lawson flips, which can be used to reach the Delaunay triangulation in  $O(n^2)$  flips [14]. This can be used for the reverse search technique to enumerate triangulations [10].

## 2 Lower Bounds

We start with the lower bound for Theorem 1 and 2. Let  $\ell$  and  $\ell'$  be two parallel horizontal lines and let  $P$  be a set of  $2n$  points,  $n$  of which are on  $\ell$  and  $\ell'$  respectively. In the following, we consider only matchings that connect points from  $\ell$  to  $\ell'$  (see Figure 3). Every




Figure 3: Matching corresponding to cycle (123).

such matching  $M$  can be interpreted as a permutation  $\pi_M$  and  $M$  is crossing free if and only if  $\pi_M$  is the identity. We can always do flips that correspond to an elementary step in bubble sort. Bubble sort on permutation  $\pi$  needs as many steps as the number of inversions of  $\pi$ . And, the number of inversions is at most  $\binom{n}{2}$ . A small perturbation of the point set ensures general position.

For the lower bound of Theorem 2, we define a *Schoone matching* on  $2n$  points in convex position denoted by  $p_1, p_2, \dots, p_{2n}$  in counterclockwise order as follows. The points  $p_1$  is linked to  $p_{n+1}$  and for each  $i \in [2, n]$ ,  $p_i$  to  $p_{2n+2-i}$  (see Figure 4). Observe that




Figure 4: An initial configuration guaranteeing the lower bound of Theorem 2, and a possible flip.

a Schoone matching decomposes into two Schoone matchings after any flip. By this the total number of

crossings decreases by exactly one. This implies by induction that  $h(M)$  equals the number of crossings minus one, for all Schoone matchings. Thus  $n-1 \leq k(n)$ .

### 3 Upper Bounds

Before we prove the upper bound, observe in Figure 5 that the number of crossings might increase after a flip. It is also possible that a segment that has dis-




Figure 5: After the depicted flip, the number of crossings goes from 1 to 3.

appeared after a flip reappear after some more flips (see Figure 6). These two observations suggest that




Figure 6: Segment  $A$  disappears and reappears.

there is no straightforward way of getting a good upper bound.

For the upper bound of Theorem 1, we define a potential function  $\Phi_{\mathcal{L}}(M)$  that depends on a well-chosen set of lines  $\mathcal{L}$ . We show that  $\Phi_{\mathcal{L}}(M) \leq 4n^3$  and that  $\Phi_{\mathcal{L}}$  decreases by at least four after any flip. The potential function  $\Phi_{\mathcal{L}}(M)$  is defined as the number of intersections between a line of  $\mathcal{L}$  and a segment of  $M$ . We define  $\mathcal{L}$  as follows. Given two points  $p, q \in P$  let  $\ell$  be the supporting line of  $p$  and  $q$ . We add to  $\mathcal{L}$  the two lines slightly above and below  $\ell$  (see Figure 7).




Figure 7: Construction of  $\mathcal{L}$ .

It holds that  $|\mathcal{L}| = 2\binom{2n}{2} \leq 4n^2$ . As any line and segment can cross at most once it follows  $\Phi_{\mathcal{L}}(M) \leq |\mathcal{L}| \cdot |M| = 4n^3$ . It remains to show that the number of segment-line intersections decreases by at least four in any flip. Consider two crossing segments  $A$  and  $B$




Figure 8: Flipping  $A$  and  $B$  yields fewer segment-line intersections.

on points  $Q = \{p_1, \dots, p_4\}$  as in Figure 8. Note that there are only three combinatorial types of lines intersecting the convex hull of  $Q$ . Either a line separates  $p_1$  and  $p_2$  from  $p_3$  and  $p_4$  as  $\ell_1$ ; a line separates  $p_1$  and  $p_4$  from  $p_2$  and  $p_3$  as  $\ell_2$ ; or a line separates one point from the other three as  $\ell_3$ . For every type of lines the number of intersections does not increase after flipping  $A$  and  $B$ . It is also easy to see that the number of intersections decreases by two for lines of type  $\ell_1$  or  $\ell_2$  after flipping  $A$  and  $B$ . By definition of  $\mathcal{L}$ , there exists for every crossing of two segments at least two lines of type  $\ell_1$  and at least two lines of type  $\ell_2$ . Thus  $\Phi_{\mathcal{L}}(M)$  decreases by at least four as claimed.

For the upper bound of Theorem 2, we define a different set of lines  $\mathcal{K}$ , which contains one vertical line between any two consecutive points ordered in  $x$ -direction, see Figure 9. It follows that  $\Phi_{\mathcal{K}}(M) \leq$




Figure 9: The set  $\mathcal{K}$ .

$n^2$  since  $|\mathcal{K}| = n - 1$ . We have to show that  $\Phi_{\mathcal{K}}$  decreases by at least two after each flip. Let  $A$  and  $B$  be two crossing segments on the points  $p_1, p_2, p_3, p_4$  ordered by  $x$ -coordinate. Then we replace  $A$  and  $B$  by  $\text{seg}(p_1, p_2)$  and  $\text{seg}(p_3, p_4)$ , see Figure 10. It is clear that at least one line  $\ell$  between  $p_2$  and  $p_3$  is not crossed after the flip and was crossed twice before the flip.




Figure 10: The number of crossings between  $\ell$  and the segments of the matchings decreases by 2.

## Acknowledgments

We want to thank anonymous reviewers for the hint to the paper of van Leeuwen and Schoone [17] and a hint to the paper by Oda and Watanabe [15]. Both authors are supported by the ERC grant PARAMTIGHT: "Parameterized complexity and the search for tight complexity results", no. 280152. The second author thanks the organizers and participants of the Emléktábla Workshop in the Summer 2015 for a nice event and interesting discussions respectively. (There are too many to list names and we do not want to forget someone.)

## References

- [1] Oswin Aichholzer, Andrei Asinowski, and Tillmann Miltzow. Disjoint compatibility graph of non-crossing matchings of points in convex position. *Electr. J. Comb.*, 22(1):P1.65, 2015.
- [2] Oswin Aichholzer, Franz Aurenhammer, Clemens Huemer, and Hannes Krasser. Transforming spanning trees and pseudo-triangulations. *Inf. Process. Lett.*, 97(1):19–22, 2006.
- [3] Oswin Aichholzer, Sergey Bereg, Adrian Dumitrescu, Alfredo García Olaverri, Clemens Huemer, Ferran Hurtado, Mikio Kano, Alberto Márquez, David Rappaport, Shakhbar Smorodinsky, Diane L. Souvaine, Jorge Urrutia, and David R. Wood. Compatible geometric matchings. *Comput. Geom.*, 42(6-7):617–626, 2009.
- [4] Oswin Aichholzer, Thomas Hackl, David Orden, Alexander Pilz, Maria Saumell, and Birgit Vogtenhuber. Flips in combinatorial pointed pseudo-triangulations with face degree at most four. *Int. J. Comput. Geometry Appl.*, 24(3):197–224, 2014.
- [5] Oswin Aichholzer, Thomas Hackl, David Orden, Pedro Ramos, Günter Rote, André Schulz, and Bettina Speckmann. Flip graphs of bounded degree triangulations. *Graphs and Combinatorics*, 29(6):1577–1593, 2013.
- [6] Oswin Aichholzer, Wolfgang Mulzer, and Alexander Pilz. Flip distance between triangulations of a simple polygon is np-complete. *Discrete & Computational Geometry*, 54(2):368–389, 2015.
- [7] Greg Aloupis, Luis Barba, Stefan Langerman, and Diane L. Souvaine. Bichromatic compatible matchings. *Comput. Geom.*, 48(8):622–633, 2015.
- [8] Andrei Asinowski, Tillmann Miltzow, and Günter Rote. Quasi-parallel segments and characterization of unique bichromatic matchings. *Journal on Computational Geometry*, 6(1):185–219, 2015.
- [9] Andrei Asinowski and Günter Rote. Point sets with many non-crossing matchings. *CoRR*, abs/1502.04925, 2015.
- [10] David Avis and Komei Fukuda. Reverse search for enumeration. *Discrete Applied Mathematics*, 65(1):21–46, 1996.
- [11] Prosenjit Bose and Ferran Hurtado. Flips in planar graphs. *Comput. Geom.*, 42(1):60–80, 2009.
- [12] Michael E. Houle, Ferran Hurtado, Marc Noy, and Eduardo Rivera-Campo. Graphs of triangulations and perfect matchings. *Graphs and Combinatorics*, 21(3):325–331, 2005.
- [13] Mashhood Ishaque, Diane L Souvaine, and Csaba D. Tóth. Disjoint compatible geometric matchings. *Discrete & Computational Geometry*, 49(1):89–131, 2013.
- [14] Charles L Lawson. Properties of n-dimensional triangulations. *Computer Aided Geometric Design*, 3(4):231–246, 1986.
- [15] Yoshiaki Oda and Mamoru Watanabe. The number of flips required to obtain non-crossing convex cycles. In *Computational Geometry and Graph Theory*, pages 155–165. Springer, 2008.
- [16] Alfredo García Olaverri, Clemens Huemer, Ferran Hurtado, and Javier Tejel. Compatible spanning trees. *Comput. Geom.*, 47(5):563–584, 2014.
- [17] Jan van Leeuwen, Anneke A. Schoone. *Untangling a traveling salesman tour in the plane*. Rijksuniversiteit. Vakgroep Informatica, 1980.