

The Sum and Product of Finite Sequences of Complex Numbers

Keiichi Miyajima
 Ibaraki University
 Faculty of Engineering
 Hitachi, Japan

Takahiro Kato
 Graduate School of Ibaraki University
 Faculty of Engineering
 Hitachi, Japan

Summary. This article extends the [10]. We define the sum and the product of the sequence of complex numbers, and formalize these theorems. Our method refers to the [11].

MML identifier: RVSUM_2, version: 7.11.07 4.156.1112

The notation and terminology used in this paper have been introduced in the following papers: [5], [7], [6], [4], [8], [13], [9], [2], [3], [15], [10], [12], and [14].

1. AUXILIARY THEOREMS

Let F be a complex-valued binary relation. Then $\text{rng } F$ is a subset of \mathbb{C} .

Let D be a non empty set, let F be a function from \mathbb{C} into D , and let F_1 be a complex-valued finite sequence. Note that $F \cdot F_1$ is finite sequence-like.

For simplicity, we adopt the following rules: i, j denote natural numbers, x, x_1 denote elements of \mathbb{C} , c denotes a complex number, F, F_1, F_2 denote complex-valued finite sequences, and R, R_1 denote i -element finite sequences of elements of \mathbb{C} .

The unary operation sqrcomplex on \mathbb{C} is defined as follows:

(Def. 1) For every c holds $(\text{sqrcomplex})(c) = c^2$.

Next we state two propositions:

- (1) sqrcomplex is distributive w.r.t. $\cdot_{\mathbb{C}}$.
- (2) $\cdot_{\mathbb{C}}$ is distributive w.r.t. $+_{\mathbb{C}}$.

2. SOME FUNCTORS ON THE i -TUPLES OF COMPLEX NUMBERS

Let us consider F_1, F_2 . Then $F_1 + F_2$ is a finite sequence of elements of \mathbb{C} and it can be characterized by the condition:

$$\text{(Def. 2)} \quad F_1 + F_2 = (+_{\mathbb{C}})^{\circ}(F_1, F_2).$$

Let us observe that the functor $F_1 + F_2$ is commutative.

Let us consider i, R_1, R_2 . Then $R_1 + R_2$ is an element of \mathbb{C}^i .

The following propositions are true:

$$(3) \quad (R_1 + R_2)(s) = R_1(s) + R_2(s).$$

$$(4) \quad \varepsilon_{\mathbb{C}} + F = \varepsilon_{\mathbb{C}}.$$

$$(5) \quad \langle c_1 \rangle + \langle c_2 \rangle = \langle c_1 + c_2 \rangle.$$

$$(6) \quad i \mapsto c_1 + i \mapsto c_2 = i \mapsto (c_1 + c_2).$$

Let us consider F . Then $-F$ is a finite sequence of elements of \mathbb{C} and it can be characterized by the condition:

$$\text{(Def. 3)} \quad -F = -_{\mathbb{C}} \cdot F.$$

Let us consider i, R . Then $-R$ is an element of \mathbb{C}^i .

The following propositions are true:

$$(7) \quad -\langle c \rangle = \langle -c \rangle.$$

$$(8) \quad -i \mapsto c = i \mapsto (-c).$$

$$(9) \quad \text{If } R_1 + R = R_2 + R, \text{ then } R_1 = R_2.$$

$$(10) \quad -(F_1 + F_2) = -F_1 + -F_2.$$

Let us consider F_1, F_2 . Then $F_1 - F_2$ is a finite sequence of elements of \mathbb{C} and it can be characterized by the condition:

$$\text{(Def. 4)} \quad F_1 - F_2 = (-_{\mathbb{C}})^{\circ}(F_1, F_2).$$

Let us consider i, R_1, R_2 . Then $R_1 - R_2$ is an element of \mathbb{C}^i .

The following propositions are true:

$$(11) \quad (R_1 - R_2)(s) = R_1(s) - R_2(s).$$

$$(12) \quad \varepsilon_{\mathbb{C}} - F = \varepsilon_{\mathbb{C}} \text{ and } F - \varepsilon_{\mathbb{C}} = \varepsilon_{\mathbb{C}}.$$

$$(13) \quad \langle c_1 \rangle - \langle c_2 \rangle = \langle c_1 - c_2 \rangle.$$

$$(14) \quad i \mapsto c_1 - i \mapsto c_2 = i \mapsto (c_1 - c_2).$$

$$(15) \quad R - i \mapsto 0_{\mathbb{C}} = R.$$

$$(16) \quad -(F_1 - F_2) = F_2 - F_1.$$

$$(17) \quad -(F_1 - F_2) = -F_1 + F_2.$$

$$(18) \quad \text{If } R_1 - R_2 = i \mapsto 0_{\mathbb{C}}, \text{ then } R_1 = R_2.$$

$$(19) \quad R_1 = (R_1 + R) - R.$$

$$(20) \quad R_1 = (R_1 - R) + R.$$

Let us consider F, c . We introduce $c \cdot F$ as a synonym of cF .

Let us consider F, c . Then $c \cdot F$ is a finite sequence of elements of \mathbb{C} and it can be characterized by the condition:

(Def. 5) $c \cdot F = \cdot_c \cdot F$.

Let us consider i, R, c . Then $c \cdot R$ is an element of \mathbb{C}^i .

One can prove the following four propositions:

(21) $c \cdot \langle c_1 \rangle = \langle c \cdot c_1 \rangle$.

(22) $c_1 \cdot (i \mapsto c_2) = i \mapsto (c_1 \cdot c_2)$.

(23) $(c_1 + c_2) \cdot F = c_1 \cdot F + c_2 \cdot F$.

(24) $0_{\mathbb{C}} \cdot R = i \mapsto 0_{\mathbb{C}}$.

Let us consider F_1, F_2 . We introduce $F_1 \bullet F_2$ as a synonym of $F_1 F_2$.

Let us consider F_1, F_2 . Then $F_1 \bullet F_2$ is a finite sequence of elements of \mathbb{C} and it can be characterized by the condition:

(Def. 6) $F_1 \bullet F_2 = (\cdot_{\mathbb{C}})^{\circ}(F_1, F_2)$.

Let us note that the functor $F_1 \bullet F_2$ is commutative.

Let us consider i, R_1, R_2 . Then $R_1 \bullet R_2$ is an element of \mathbb{C}^i .

Next we state four propositions:

(25) $\varepsilon_{\mathbb{C}} \bullet F = \varepsilon_{\mathbb{C}}$.

(26) $\langle c_1 \rangle \bullet \langle c_2 \rangle = \langle c_1 \cdot c_2 \rangle$.

(27) $i \mapsto c \bullet R = c \cdot R$.

(28) $i \mapsto c_1 \bullet i \mapsto c_2 = i \mapsto (c_1 \cdot c_2)$.

3. FINITE SUM OF FINITE SEQUENCE OF COMPLEX NUMBERS

One can prove the following propositions:

(29) $\sum(\varepsilon_{\mathbb{C}}) = 0_{\mathbb{C}}$.

(30) $\sum\langle c \rangle = c$.

(31) $\sum(F \wedge \langle c \rangle) = \sum F + c$.

(32) $\sum(F_1 \wedge F_2) = \sum F_1 + \sum F_2$.

(33) $\sum(\langle c \rangle \wedge F) = c + \sum F$.

(34) $\sum\langle c_1, c_2 \rangle = c_1 + c_2$.

(35) $\sum\langle c_1, c_2, c_3 \rangle = c_1 + c_2 + c_3$.

(36) $\sum(i \mapsto c) = i \cdot c$.

(37) $\sum(i \mapsto 0_{\mathbb{C}}) = 0_{\mathbb{C}}$.

(38) $\sum(c \cdot F) = c \cdot \sum F$.

(39) $\sum(-F) = -\sum F$.

(40) $\sum(R_1 + R_2) = \sum R_1 + \sum R_2$.

(41) $\sum(R_1 - R_2) = \sum R_1 - \sum R_2$.

4. THE PRODUCT OF FINITE SEQUENCES OF COMPLEX NUMBERS

One can prove the following propositions:

- (42) $\prod(\varepsilon_{\mathbb{C}}) = 1.$
- (43) $\prod(\langle c \rangle \wedge F) = c \cdot \prod F.$
- (44) For every element R of \mathbb{C}^0 holds $\prod R = 1.$
- (45) $\prod((i + j) \mapsto c) = \prod(i \mapsto c) \cdot \prod(j \mapsto c).$
- (46) $\prod((i \cdot j) \mapsto c) = \prod(j \mapsto \prod(i \mapsto c)).$
- (47) $\prod(i \mapsto (c_1 \cdot c_2)) = \prod(i \mapsto c_1) \cdot \prod(i \mapsto c_2).$
- (48) $\prod(R_1 \bullet R_2) = \prod R_1 \cdot \prod R_2.$
- (49) $\prod(c \cdot R) = \prod(i \mapsto c) \cdot \prod R.$

5. MODIFIED PART OF [1]

We now state several propositions:

- (50) For every complex-valued finite sequence x holds $\text{len}(-x) = \text{len } x.$
- (51) For all complex-valued finite sequences x_1, x_2 such that $\text{len } x_1 = \text{len } x_2$ holds $\text{len}(x_1 + x_2) = \text{len } x_1.$
- (52) For all complex-valued finite sequences x_1, x_2 such that $\text{len } x_1 = \text{len } x_2$ holds $\text{len}(x_1 - x_2) = \text{len } x_1.$
- (53) For every real number a and for every complex-valued finite sequence x holds $\text{len}(a \cdot x) = \text{len } x.$
- (54) For all complex-valued finite sequences x, y, z such that $\text{len } x = \text{len } y = \text{len } z$ holds $(x + y) \bullet z = x \bullet z + y \bullet z.$

REFERENCES

- [1] Kanchun and Yatsuka Nakamura. The inner product of finite sequences and of points of n -dimensional topological space. *Formalized Mathematics*, 11(2):179–183, 2003.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [3] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [5] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(1):175–180, 1990.
- [6] Czesław Byliński. Binary operations applied to finite sequences. *Formalized Mathematics*, 1(4):643–649, 1990.
- [7] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [8] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [9] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [10] Czesław Byliński. The sum and product of finite sequences of real numbers. *Formalized Mathematics*, 1(4):661–668, 1990.
- [11] Keith E. Hirst. *Numbers, Sequences and Series*. Butterworth-Heinemann, 1984.

- [12] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Formalized Mathematics*, 1(3):477–481, 1990.
- [13] Andrzej Trybulec. Binary operations applied to functions. *Formalized Mathematics*, 1(2):329–334, 1990.
- [14] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers. *Formalized Mathematics*, 1(3):445–449, 1990.
- [15] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

Received January 12, 2010
