

# Antioxidants & Redox Signaling

Antioxidants & Redox Signaling: <http://mc.manuscriptcentral.com/liebert/ARS>

## The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

Journal:	<i>Antioxidants &amp; Redox Signaling</i>
Manuscript ID:	Draft
Manuscript Type:	Forum Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Szarka, András; Budapest University of Technology and Economics, Department of Applied Biotechnology and Food Science, Laboratory of Biochemistry and Molecular Biology Bánhegyi, Gábor; Semmelweis University, Dept. of Medical Chemistry Asard, Han; University of Antwerp, Department of Biology
Keyword:	Free Radicals, Mitochondria, Plants

SCHOLARONE™  
Manuscripts

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

Forum Review Article

**The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration**

**András Szarka<sup>1</sup>, Gábor Bánhegyi<sup>2</sup>, Han Asard<sup>3</sup>**

<sup>1</sup> Department of Applied Biotechnology and Food Science, Laboratory of Biochemistry and Molecular Biology, Budapest University of Technology and Economics, Budapest, Hungary 1111; <sup>2</sup> Department of Medical Chemistry, Molecular Biology and Pathobiochemistry, Semmelweis University, Budapest, Hungary 1444; <sup>3</sup> Department of Biology, University of Antwerp, Antwerp, Belgium 2020

Running head: Ascorbate, between mitochondria and chloroplasts

Corresponding author: András Szarka, Department of Applied Biotechnology and Food Science, Laboratory of Biochemistry and Molecular Biology, Budapest University of Technology and Economics, 1111 Szent Gellért tér 4. Budapest, Hungary, Telephone: +36-1-463-3858; Fax: +36-1-463-3855, e-mail: [szarka@mail.bme.hu](mailto:szarka@mail.bme.hu);

Words: 4319

References: 60

Color illustration: 5 (online 5 and hardcopy 0)

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

## Abstract

Significance: Ascorbate, this multifaceted small molecular weight carbohydrate derivative, plays important roles in a range of cellular processes in plant cells from, the regulation of cell cycle, through cell expansion and senescence. Beyond these physiological functions, ascorbate has a critical role in responses to abiotic stresses such as high light, high salinity or drought. The biosynthesis, recycling and intracellular transport are important elements of the balancing of ascorbate level to the always-changing conditions and demands. Recent Advances: A bidirectional tight relationship was described between ascorbate biosynthesis and the mitochondrial electron transfer chain, since L-galactono-1,4-lactone dehydrogenase (GLDH), the enzyme catalyzing the ultimate step of ascorbate biosynthesis, uses oxidized cytochrome c as the only electron acceptor and has a role in the assembly of Complex I. A similar bidirectional relationship was revealed between the photosynthetic apparatus and ascorbate biosynthesis since the electron flux through the photosynthetic electron transfer chain affects the biosynthesis of ascorbate and the level of ascorbate could affect photosynthesis. Critical Issues: The details of this regulatory network of photosynthetic electron transfer, respiratory electron transfer and ascorbate biosynthesis are still not clear, as are the potential regulatory role and the regulation of intracellular ascorbate transport and fluxes. Future Directions: The elucidation of the role of ascorbate as an important element of the network of photosynthetic, respiratory electron transfer chains and tricarboxylic acid cycle will contribute to understanding plant cell responses to different stress conditions.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

## Introduction

Ascorbate. The molecule that raises new issues and supplies new findings continuously. Our knowledge about its conventional but crucial role in antioxidant defense as well as in stress protection has deepened (14, 47). Furthermore, the cellular and physiological role of the protagonist of this review has notably broadened in the last two decades. This repertoire includes functions in nearly every aspect of plant biology from mitosis and cell expansion (48), through programmed cell death (15), as well as protection against environmental stresses including ozone, UV radiation (14), high temperatures (26) and high light intensity (35) to growth (40), hormone responses, flowering, senescence (5) and defense against pathogens (39).

Probably the paramount advances were the discovery of the link between the mitochondrial electron transport chain and ascorbate biosynthesis (7, 34, 41, 43), the regulatory role of photosynthetic electron transfer to ascorbate biosynthesis and accumulation (8, 55) and the possible role for ascorbate in coordinating the rates of respiration, the tricarboxylic acid (TCA) cycle and photosynthesis (38, 52, 16, 49).

Ascorbate. The molecule, a (plant) biologist cannot lose interest in.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

### Plant mitochondrion: ascorbate biosynthesis

The first complete pathway of ascorbate biosynthesis in plant cells, the so called D-Man/L-Gal or Smirnoff-Wheeler pathway, was only unraveled in 1998 (53) (Fig. 1). In the mean time, all the genes and their products, involved in the pathway in *Arabidopsis thaliana* have been fully characterized (30). Shortly after the description of the Smirnoff-Wheeler pathway the existence of alternative ascorbate biosynthetic pathways became apparent with the cloning and characterization of other ascorbate biosynthetic enzymes. It was shown that GDP-Mannose-3',5'-epimerase catalyzes at least two distinct epimerization reactions and releases GDP-L-Gulose (GDP-L-Gul) in addition to the well known, Smirnoff-Wheeler pathway intermediate, GDP-L-galactose (GDP-L-Gal) (54). GDP-L-Gul can be channeled directly into the proposed L-Gulose branch of the biosynthetic pathway, in which the last step is the oxidation of L-gulonono-1,4-lactone to ascorbate (54) (Fig. 1). The possible contribution of myo-inositol as a precursor of ascorbate biosynthesis via the formation of UDP-glucuronate by myo-inositol oxidase was also suggested in *Arabidopsis* (31). This route – similar to the L-Gulose branch one – provides L-gulonono-1,4-lactone (Fig. 1). Finally, evidence was provided that D-galacturonic acid, a principal component of cell wall pectins, is produced by D-galacturonate reductase and contributes to the ascorbate pool in strawberry fruit (1). The latter route – similar to the classic D-Man/L-Gal pathway – provides L-galactono-1,4-lactone to the biosynthetic pathway (Fig. 1). Mutants in the D-Man/L-Gal pathway, e.g. *vtc-1* (an *Arabidopsis* mutant with defective GDP-mannose pyrophosphorylase activity) and *vtc-2* (an *Arabidopsis* mutant with defective GDP-L-galactose

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

phosphorylase activity), demonstrated that ascorbate derived from alternative pathways, could not compensate for the low ascorbate levels (13, 17).

Both the D-Man/L-Gal and the Galacturonate routes converge to L-galactono-1,4-lactone. The oxidation of L-galactono-1,4-lactone to ascorbate was attributed to mitochondrial GLDH already more than fifty years ago (32). However, the intimate link between the mitochondrial electron transfer chain (mETC) and the activity of this enzyme became clear only more recently (7). Subfractionation of mitochondria demonstrated that GLDH is largely associated with the inner-membrane fraction (7). Also, the latency of the enzyme is similar to that of cytochrome c oxidase (CCO) suggesting that the GLDH localization is similar to CCO. GLDH delivers electrons to oxidized cytochrome c (7, 28). The absolute requirement of the enzyme for oxidized cytochrome was proved by the stimulatory effect of antimycin A, an inhibitor of cytochrome c reductase (7) and by the observation of enhanced GLDH activity in the Arabidopsis ppr40-1 mutant, which has a block of electron flow at complex III (60). Therefore, L-galactono-1,4-lactone can be defined as a respiratory substrate in plants (7, 34) (Fig. 2).

Although three membrane spanning regions were predicted previously (7), no transmembrane regions were found in the mature AtGLDH in a recent study (28). A tight link between complex I and GLDH was predicted by the fact, that rotenone, a specific inhibitor of electron flow through complex I, inhibited ascorbate biosynthesis (34). Indeed, GLDH was present in the minor higher-mobility form (850 kDa) of complex I (22, 41). The 58 kDa GLDH protein did not attach to the 1000 kDa holo complex, but only to a slightly smaller version of complex I (34, 41). Interestingly, complex I and its associated NADH dehydrogenase activity were lacking in leaf and root mitochondria of *gldh* mutant Arabidopsis, although the other respiratory complexes

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

1  
2  
3  
4 were unaffected. This suggests that GLDH might represent an assembly factor in complex I  
5  
6 biogenesis (41). Consistent with the reduction of cytochrome c, GLDH is believed to be located  
7  
8 in the intermembrane space (IMS), associated with the membrane arm of complex I (25). A  
9  
10 recent study described two new low abundance GLDH containing protein complexes with  
11  
12 molecular mass of 470 and 420 kDa in Arabidopsis mitochondria (43). These complexes  
13  
14 represent parts of the membrane arm of complex I and are assembly intermediates. The authors  
15  
16 suggest that GLDH binds to the 420 and 470 kDa complex I assembly intermediates, which at a  
17  
18 later stage form the 850 kDa intermediate. Formation of the 1000 kDa holo complex I is  
19  
20 preceded by detachment of GLDH (43).  
21  
22  
23  
24  
25

26  
27 The dual role of GLDH in ascorbate biosynthesis and in the assembly of complex I is  
28  
29 indisputable, however the connection between these two processes is still a mystery. GLDH  
30  
31 belongs to the vanillyl-alcohol oxidase flavoprotein family, including L-gulono-lactone oxidase.  
32  
33 The latter enzyme is responsible for ascorbate synthesis in several animal groups, generating  
34  
35 H<sub>2</sub>O<sub>2</sub> as a by-product. However, in contrast to related aldonolactone oxidoreductases, GLDH  
36  
37 contains an alanine (Ala-113), instead of the conserved glycine or proline, at an essential position  
38  
39 near the isoalloxazine flavin-ring of the cofactor. This substitution possibly prevents molecular  
40  
41 oxygen from reaching the reduced flavin, avoiding the generation of superoxide or H<sub>2</sub>O<sub>2</sub> (27, 28).  
42  
43 GLDH contains non-covalently bound FAD, as it also lacks the conserved histidine involved in  
44  
45 covalent FAD binding (28). The dehydrogenase activity of the enzyme – instead of a possible  
46  
47 oxidase activity – may contribute to the reduction of the H<sub>2</sub>O<sub>2</sub> load of mitochondria and avoid  
48  
49 the possible inactivation of GLDH by the oxidation of its redox sensitive thiol (Cys-340) (29).  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

In addition to L-galactono-1,4-lactone, recombinant Arabidopsis GLDH, also oxidizes the isomer, L-gulono-1,4-lactone, at significant rates, although with a much higher  $K_m$  value (0.17 mM vs. 13.1 mM) (28). Hence it is possible that GLDH also plays a role in the oxidation of L-gulono-1,4-lactone, in an alternative, minor, ascorbate synthesizing pathway (Fig. 1). (Although L-gulono-lactone oxidase activities have also been described in plants (33).)

Tobacco plants overexpressing GLDH do not have elevated ascorbate content (23), and silencing of GLDH in tomato did not reduce ascorbate levels or its redox status, despite decreased mRNA, protein levels and enzyme activity (2). These observations strongly suggest that the mitochondrial GLDH-catalyzed step is not rate limiting in ascorbate biosynthesis. (The GDP-L-galactose phosphorylase (encoded by VTC2 and VTC5) catalyzed conversion of GDP-L-galactose to L-galactose-1-P (Fig. 1) is supposed to be the rate-limiting step of ascorbate synthesis (17, 47).) The metabolic changes that were observed in GLDH-silenced plants (2) are probably explained by decreased functioning of the mitochondrial holo complex I in the absence of GLDH (41).

### **Plant mitochondrion: ascorbate regeneration**

The role of ascorbate recycling (i.e. reduction of its oxidized forms) in the maintenance of a sufficient level of ascorbate is generally accepted.

Chloroplasts are the major source of reactive oxygen species (ROS) in plant cells and ascorbate plays multiple roles in their elimination (4, 20, 35, 36, 47). It is therefore perhaps not surprising



Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

that enzymes for the reduction of oxidized forms of ascorbate, i.e. monodehydroascorbate (MDHA) and dehydroascorbate (DHA), using GSH and NADPH (from PSI) were first described in chloroplasts (i.e. the ascorbate-glutathione or Foyer-Halliwel-Asada cycle) (4, 20). Although the mitochondrial ROS production is considerably less than the ROS production in illuminated chloroplasts or in peroxisomes, however in the dark, or in non-green tissues, mitochondria are major sources of ROS (47).

After scattered reports on presence in mitochondria of the individual elements of the ascorbate-glutathione cycle (3, 18), all four enzymes, ascorbate peroxidase (APX), monodehydroascorbate reductase (MDHAR), dehydroascorbate reductase (DHAR), and glutathione reductase (GR), were described in mitochondria and peroxisomes (24). The mitochondrial location for these proteins was confirmed by green fluorescent protein fusions (12). Furthermore, a dual import assay showed targeting of APX, MDHAR, and GR gene products to mitochondria and chloroplasts, while a putative DHAR was only imported into mitochondria (12). The subcellular localization and the topology of the enzymes were also determined by preparing submitochondrial fractions and performing enzyme latency measurements. APX and MDHAR activities were associated mainly with the inner membrane fraction. The low latency suggested a predominantly IMS side localization for APX, while the high latency of MDHAR suggested a predominantly matrix side localization (Fig. 2.) (24). GR and DHAR activities were prominent in the matrix, but could also be found in other submitochondrial fractions, suggesting they are partitioned between compartments and can be partially membrane associated (12). The predominant matrix location of MDHAR, DHAR, and GR is presumably related to the supply of reductants NADH and NADPH to fuel these reactions (from the TCA cycle). In this way the

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

MDHA and DHA generated by the action of APX in plant mitochondria must pass through the inner mitochondrial membrane to the matrix for reduction back to ascorbate.

Hence the DHA transporter – located in the inner mitochondrial membrane – is an important member of the ascorbate regeneration machinery (45). Interestingly uptake of glucose by tobacco mitochondria was also observed in the same study. The strong inhibition of DHA and glucose uptake by the known GLUT inhibitor genistein and the inhibitory effect of glucose on DHA transport suggested that the two molecules could be ligands for the same transporter. The uptake of both compounds is independent of mitochondrial respiration. Since the uptake of the reduced form, ascorbate – in contrast to the DHA transport – showed a rather low affinity it cannot be excluded that only DHA is taken up by these organelles. In that case, the intramitochondrial appearance of ascorbate may be merely due to the uptake and concomitant reduction of the oxidized form (45). Although components of the ascorbate-glutathione cycle have been demonstrated in mitochondria, GSH is certainly not the only possible electron donor for DHA reduction in these organelles. Results obtained from animal mitochondria indicate a role for the mETC in the reduction of DHA (46). Indeed, ascorbate generation (upon DHA addition) could be stimulated by the complex II substrate succinate. The involvement of complex II in DHA reduction was further supported by the inhibitory effect of the succinate dehydrogenase competitive inhibitor, malonate. Glucose, a competitive inhibitor of mitochondrial DHA uptake, interrupted the regeneration cycle and decreased ascorbate production in the absence and in the presence of succinate. At the same time the role of complex I and III could be ruled out, since neither the substrates nor the inhibitors of these complexes had any effect on ascorbate generation (46).

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

1  
2  
3  
4 In addition to succinate, KCN showed a strong positive effect on ascorbate accumulation upon  
5  
6 DHA addition, presumably due to the decreased ascorbate oxidation. The strong inhibitory effect  
7  
8 of KCN on ascorbate oxidation could be verified by EPR measurements (46). The assumption  
9  
10 was strengthened in Arabidopsis ppr40-1 mutant. Impaired electron transport through Complex  
11  
12 III in the mitochondria isolated from ppr40-1 mutant is accompanied by enhanced cytochrome c  
13  
14 oxidase activity and ascorbate consumption (59). CCO activity and ascorbate consumption were  
15  
16 increased in same manner. Furthermore ascorbate consumption could be inhibited also in this  
17  
18 case by KCN treatment (59). These observations support that ascorbate can act as an alternative  
19  
20 electron donor *in vivo* to sustain oxidative respiration in states with impaired electron flow  
21  
22 through complex III (Fig. 3).  
23  
24  
25  
26  
27

28  
29 Based on these observations the following model can be proposed: DHA, formed in the IMS due  
30  
31 to the activity of APX, crosses the inner membrane through a glucose transporter. DHA is  
32  
33 subsequently reduced to ascorbate in the mitochondrial matrix via the ascorbate-glutathione  
34  
35 cycle or at complex II. A fraction of ascorbate can leave the mitochondrial matrix by a presently  
36  
37 unidentified mechanism. Ascorbate in the IMS can be oxidized by supplying electrons to  
38  
39 complex IV (Fig. 3). The less well characterized step of this hypothesis is the efflux of ascorbate  
40  
41 from the matrix. This process is potentially rate-limiting, since the ascorbate produced in the  
42  
43 mitochondria appeared in the incubation medium with a significant delay compared to the matrix  
44  
45 (Szarka A, unpublished). This electron route represented by the DHA/ascorbate redox couple  
46  
47 may also provide an alternative route in case of complex III damage by taking up electrons at  
48  
49 complex II (by the reduction of DHA to ascorbate) and providing electrons through a bypass to  
50  
51 complex IV (by the oxidation of ascorbate to DHA).  
52  
53  
54  
55  
56  
57  
58  
59  
60

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

In conclusion, the plant mETC, and mitochondria, play an important role not only in the synthesis of ascorbate but also in its regeneration.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

### Ascorbate metabolism and the chloroplast - Ascorbate as a connection between mitochondrial respiration and photosynthesis

Ascorbate levels are dramatically increased in response to light (58), which is undoubtedly related to its multiple roles in photoprotection. On one hand ascorbate has an important role in non-photochemical quenching, since violaxanthin de-epoxidase uses ascorbate as its reductant *in vitro* and the availability of ascorbate could also limit the activity of the enzyme *in vivo* in *vtc-2* mutants (35). On the other hand ascorbate is also a key component of the water-water cycle in chloroplast (4) (Fig. 4).

Since the ultimate step of ascorbate biosynthesis occurs in the mitochondria ascorbate must be transported into the chloroplasts. The chloroplast envelope contains an ascorbate translocator (9) (Fig. 5). In intact spinach chloroplasts ascorbate transport occurs through facilitated diffusion and has a relatively low affinity ( $K_M \sim 20$  mM) (9). In contrast, thylakoid membranes appear to have no transport system for ascorbate and uptake may result from diffusion alone (21). Recently the subcellular distribution of ascorbate was determined by ascorbate-specific immunogold labeling (58). In both *Arabidopsis* and *Nicotiana*, highest levels of ascorbate were found in the cytosol (21.7 mM) whereas mitochondria (10.4 mM) and chloroplasts (10.8 mM) contained intermediate levels (The concentrations refer to *Arabidopsis*.) (58). Subcellular concentrations strengthened the idea that ascorbate uptake occurs by facilitated diffusion through the chloroplast envelope. Consistent with the transport experiments (21), ascorbate was only detected in the chloroplast stroma and mitochondrial matrix, but not within the lumen of thylakoids or cristae in

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

1  
2  
3  
4 non-stressed plants, ascorbate-specific labeling could be found in the lumen of thylakoids only  
5  
6 during high light conditions (58) (Fig. 5).  
7  
8  
9

10 The tight interaction of photosynthetic and respiratory electron transfer chains is a key feature of  
11 plant cells (see for reviews 36, 37, 56). During high light conditions, NADP-malate  
12 dehydrogenase, the key enzyme of the malate valve, converts oxaloacetate to malate using  
13 NADPH, facilitating the regeneration of the  $\text{NADP}^+$  pool in the chloroplast. Malate then can be  
14 exported to the cytosol via the malate/oxaloacetate shuttle (51). The exported malate is oxidized  
15 by NAD-malate dehydrogenases (MDH) of the cytosol, peroxisomes and mitochondria  
16 generating NADH. Finally NADH may be consumed by mitochondrial respiration (42). During  
17 photorespiration a large amount of NADH is also produced by the glycine decarboxylase  
18 complex (GDC) in the mitochondria. Although a part of the generated NADH is transported to  
19 the peroxisomes through the malate/oxaloacetate shuttle of mitochondria, the rest of the  
20 photorespiratory NADH is oxidized by the respiratory chain (36). The induction of the leaf  
21 alternative oxidase (AOX) and internal-type II NAD(P)H dehydrogenase ( $\text{ND}_{\text{in}}$ ) genes by light  
22 (19, 57) presume that mitochondrial respiration and especially the non-phosphorylating pathways  
23 can be important elements of the dissipating system of reductants in the illuminated leaf (Fig. 4).  
24 The suppressed rate of photosynthesis in leaves of uncoupling protein-1 (UCP1)-deficient  
25 Arabidopsis suggests that UCP also allows for more efficient NADH consumption and can be a  
26 part of this dissipation system (44).  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 Increased capacities of the cytochrome c and AOX pathways, in Arabidopsis leaves grown under  
51 high light, were accompanied by an increase of the capacity for ascorbate synthesis and GLDH  
52 activities (8) (Fig. 4). However, high light had no detectable effects on GLDH transcript levels or  
53  
54  
55  
56  
57  
58  
59  
60

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

1  
2  
3  
4 on day/night variations in protein abundance (8). Similarly, leaves of AOX1a overexpressing  
5  
6 lines also had higher rates of ascorbate production and accumulation, without affecting GLDH  
7  
8 activity (Fig. 4). Elevated ascorbate production in these lines may be enhanced by an increased  
9  
10 availability of oxidized cytochrome c, which is the electron acceptor in the last step of ascorbate  
11  
12 synthesis (see above). On the other hand, an increased rate of the cytochrome pathway and an  
13  
14 increased amount of cytochrome c, in high light, may also contribute to higher CCO activities  
15  
16 and enhanced ascorbate synthesis (8) (Fig. 2, 4, 5).  
17  
18  
19

20  
21  
22 In contrast to the results of Bartoli et al. (8) the transcript level of GLDH (and GDP-mannose  
23  
24 pyrophosphorylase, L-galactose 1-P phosphatase, and VTC2) changed in parallel with changes in  
25  
26 the leaf ascorbate pool during dark and light periods in the study of Yabuta et al (55). Overall,  
27  
28 the correlation between the GLDH expression and activity, and foliar ascorbate concentrations  
29  
30 seems to be quite loose. No significant changes were observed in the transcript levels of GLDH  
31  
32 in tobacco leaves during the day-night cycles, whereas foliar ascorbate increased nearly 4-fold in  
33  
34 the light (40), and no correlation was found in the abundance and activity of GLDH and foliar  
35  
36 ascorbate levels in a range of plant species (6). At the same time an approximately 2-fold  
37  
38 increase of the GLDH transcript level was observed in young *Arabidopsis* rosette leaves during  
39  
40 the daytime (50).  
41  
42  
43  
44

45  
46 Since AOX is induced by light, and an AOX overexpression line showed elevated ascorbate  
47  
48 production, the effects of photosynthesis inhibitors on ascorbate levels in leaves, was  
49  
50 investigated. DCMU (3-(3,4-dichlorophenyl)-1,1-dimethylurea) and atrazine, which block  
51  
52 electron flow to plastoquinones in PSII, mimic the effect of darkness by causing the down-  
53  
54 regulation of GLDH, and prevented the increase in leaf ascorbate (55) (Fig. 4). Unfortunately in  
55  
56  
57  
58  
59  
60



Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

these experiments the transcription level and the activity of AOX were not determined. The importance of photosynthetic electron flow and of the plastoquinone redox status was further strengthened by the transfer of plants to a sucrose containing medium, which resulted in lower photosynthetic activity and leaf ascorbate levels, even under exposure to light (55). Together, these results underline the role of the redox status of the plastoquinone pool and photosynthesis itself in the control of ascorbate level in illuminated leaves.

For continuous scavenging of  $H_2O_2$  generated in chloroplasts, ascorbate needs to be regenerated. With an estimated maximum production rate of  $H_2O_2$  of  $450 \mu M \cdot s^{-1}$ , and 30 mM ascorbate in chloroplasts (without regeneration), it was calculated that ascorbate would be consumed within 60 s (4). Hence it is not surprising that the existence of an ascorbate regenerating system in chloroplasts was demonstrated earlier than its complete biosynthetic pathway (4, 12, 20). It was also shown that the capacity to regenerate ascorbate from its oxidized forms is enhanced in the high light-grown leaves, suggesting that amelioration of the recycling systems backs up the enhanced capacity for synthesis to ensure that the cellular ascorbate pool remains in the reduced state (8).

Plants show light inhibition of the TCA cycle and TCA cycle-linked respiration (52) (Fig. 4). The induction of GDC in mitochondria of illuminated leaves (during photorespiration) results in a more reduced redox status and increased adenylate phosphorylation level in the mitochondrial matrix (10). These changes cause restrictions at the pyruvate dehydrogenase complex and at other reactions within the TCA cycle. This way the reductant overload of the mETC can be reduced during light conditions. Hence the elevated rates of photosynthesis, over a range of light intensities, could be reached by the generation of mitochondrial MDH antisense (38) or aconitase



Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

mutant (*Aco1*) (11) tomato plants. A dramatic increase in ascorbate could be observed in both transgenic lines (38). These observations further strengthened the link between the photosynthetic electron flow and ascorbate biosynthesis, accumulation (Fig. 4). However a recent report of Tomaz et al. (52) reduces the clarity of this picture somewhat. These authors found higher mitochondrial respiratory rates in the double mutant mitochondrial MDH *Arabidopsis* line (52). Intriguingly the amount of GLDH was lower, but the ascorbate content was doubled in the mitochondrial MDH deficient plant compared to the wild type (52).

The photoprotective role of ascorbate can be shown more pronounced if the electron sink of mitochondria is plugged in, thus the elimination of reductants is diminished. Both Antimycin A (inhibitor of the cytochrome *c* pathway) and SHAM (inhibitor of the AOX pathway) caused marked reduction in photosynthetic carbon assimilation with concomitant rise in total cellular ROS. Furthermore the restriction of mitochondrial electron transport decreased the ratio of the reduced form of both ascorbate and GSH (16). Accordingly, the application of these respiratory inhibitors in *vtc-1* mutant *Arabidopsis* caused an extreme level of photoinhibition (83%) (49). However the photoinhibition – even in the presence of antimycin A or SHAM – could be decreased significantly by the treatment with L-galactono-1,4-lactone. As was expected the protective effect of L-galactono-1,4-lactone was maximal in *vtc-1* mutants in the presence of SHAM (49).

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

### Concluding remarks

Ascorbate was initially considered to be a simple antioxidant or cofactor for hydroxylation reactions. However, this view has been radically changed by observations of the last two decades.

The enzyme, catalyzing the ultimate step of ascorbate biosynthesis, is deeply embedded into the mETC (Fig. 2). This tight relationship seems to be bidirectional, since GLDH has a role in the assembly of Complex I (41, 43). It is still an enigma whether the ascorbate generating function of the GLDH protein is a requirement for its assembly function. Despite this question, it is more than interesting that the electron flow through complex I has a regulatory role on the ascorbate biosynthesis capacity (34) and the enzyme responsible for ascorbate biosynthesis has a role in the assembly of complex I holo complex (43). It may have a role in balancing the electron flows under different (environmental) circumstances.

It became clear that the regeneration of ascorbate is as important in the maintenance of its level (redox status) as its biosynthesis. Furthermore, the intensity of the regeneration (beside the biosynthesis) is also influenced by the changing (redox) environment both in the mitochondria (60) and in the chloroplast (8) or rather its regulation seemed to be coupled in these two compartments (12). However, the contribution of mitochondrial recycling of ascorbate to the whole cell ascorbate pool is still a matter of future research.

Ascorbate can be an important element of the photosynthetic ETC, mETC and TCA cycle regulatory network since i) the level of ascorbate can influence the intensity of photosynthesis

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

(35, 38, 49); ii) the electron flux through the photosynthetic ETC influences the biosynthesis of ascorbate (55); iii) ascorbate is synthesized tightly coupled to the mETC by providing electrons to complex IV (7) in case of the inhibition of TCA cycle (38) and iv) the activity of AOX also influences the biosynthetic rate of ascorbate (8). Any changes in either of these elements of this network influence the others (Fig. 4). However, the details of this transmitter function of ascorbate between the plastidic and mitochondrial functions so far remains a mystery and needs further investigations.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

### Acknowledgments

This work was supported by the Hungarian Scientific Research Fund (77826). This work is connected to the scientific program of the “Development of quality-oriented and harmonized R + D + I strategy and functional model at BME” project. The project was supported by the New Széchenyi Plan (Project ID: TÁMOP-4.2.1/B-09/1/KMR-2010-0002 and TÁMOP-4.2.1/B-09/1/KMR-2010-0001). Thanks are due to the János Bolyai Research Scholarship of the Hungarian Academy of Sciences for supporting A. Sz. This work was also supported by the Flemish Fund for Scientific Research, FWO (grant G.0118.07N to H.A.).

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

**Author Disclosure Statement**

No competing financial interests exist.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

### List of abbreviations

AOX, alternative oxidase; APX, ascorbate peroxidase; CCO, cytochrome c oxidase; DHA, dehydroascorbate; DHAR, dehydroascorbate reductase; GDC, glycine decarboxylase complex; GDP-L-Gal, GDP-L-galactose; GDP-L-Gul, GDP-L-gulose; GLDH, L-galactono-1,4-lactone dehydrogenase; GR, glutathione reductase; IMS, intermembrane space; MDH, malate dehydrogenase; MDHA, monodehydroascorbate; MDHAR, monodehydroascorbate reductase; mETC, mitochondrial electron transfer chain; ND<sub>in</sub>, internal type II NAD(P)H dehydrogenases; ROS, reactive oxygen species ; SHAM, salicylhydroxamic acid; TCA, tricarboxylic acid; UCP, uncoupling protein

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

## References

1. Agius F, González-Lamothe R, Caballero JL, Muñoz-Blanco J, Botella MA, Valpuesta V. Engineering increased vitamin C levels in plants by overexpression of a D-galacturonic acid reductase. *Nat Biotechnol* 21: 177-181, 2003.
2. Alhagdow M, Mounet F, Gilbert L, Nunes-Nesi A, Garcia V, Just D, Petit J, Beauvoit B, Fernie AR, Rothan C, Baldet P. Silencing of the mitochondrial ascorbate synthesizing enzyme L-galactono-1,4-lactone dehydrogenase affects plant and fruit development in tomato. *Plant Physiol* 145: 1408-1422, 2007.
3. Arrigoni O, Dipierro S, Borraccino G. Ascorbate free radical reductase: a key enzyme of the ascorbic acid system. *FEBS Lett* 125: 242-244, 1981.
4. Asada K. THE WATER-WATER CYCLE IN CHLOROPLASTS: Scavenging of Active Oxygens and Dissipation of Excess Photons. *Annu Rev Plant Physiol Plant Mol Biol* 50: 601-639, 1999.
5. Barth C, De Tullio M, Conklin PL. The role of ascorbic acid in the control of flowering time and the onset of senescence. *J Exp Bot* 57: 1657-1665, 2006
6. Bartoli CG, Guamet JJ, Kiddle G, Pastori G, Di Cagno R, Theodoulou FL, Foyer CH. The relationship between L-galactono-1,4-lactone dehydrogenase (GalLDH) and ascorbate content in leaves under optimal and stress conditions. *Plant, Cell and Environment* 28: 1073-1081, 2005.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

- 1  
2  
3  
4  
5 7. Bartoli CG, Pastori GM, Foyer CH. Ascorbate biosynthesis in mitochondria is linked to  
6  
7 the electron transport chain between complexes III and IV. *Plant Physiol* 123: 335-344,  
8  
9 2000.
- 10  
11  
12 8. Bartoli CG, Yu J, Gómez F, Fernández L, McIntosh L, Foyer CH. Inter-relationships  
13  
14 between light and respiration in the control of ascorbic acid synthesis and accumulation  
15  
16 in *Arabidopsis thaliana* leaves. *J Exp Bot* 57: 1621-1631, 2006.
- 17  
18  
19 9. Beck E, Burkert A, Hofmann M. Uptake of l-Ascorbate by Intact Spinach Chloroplasts.  
20  
21 *Plant Physiol* 73: 41-45, 1983.
- 22  
23  
24  
25  
26 10. Bykova NV, Keerberg O, Pärnik T, Bauwe H, Gardeström P. Interaction between  
27  
28 photorespiration and respiration in transgenic potato plants with antisense reduction in  
29  
30 glycine decarboxylase. *Planta* 222: 130-140, 2005.
- 31  
32  
33  
34 11. Carrari F, Nunes-Nesi A, Gibon Y, Lytovchenko A, Loureiro ME, Fernie AR. Reduced  
35  
36 expression of aconitase results in an enhanced rate of photosynthesis and marked shifts in  
37  
38 carbon partitioning in illuminated leaves of wild species tomato. *Plant Physiol* 133:  
39  
40 1322-1335, 2003.
- 41  
42  
43  
44 12. Chew O, Whelan J, Millar AH. Molecular definition of the ascorbate-glutathione cycle in  
45  
46 *Arabidopsis* mitochondria reveals dual targeting of antioxidant defenses in plants. *J Biol*  
47  
48 *Chem* 278: 46869-46877, 2003.



Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

13. Conklin PL, Norris SR, Wheeler GL, Williams EH, Smirnoff N, Last RL. Genetic evidence for the role of GDP-mannose in plant ascorbic acid (vitamin C) biosynthesis. *Proc Natl Acad Sci U S A* 96: 4198-4203, 1999.
14. Conklin PL, Williams EH, Last RL. Environmental stress sensitivity of an ascorbic acid-deficient *Arabidopsis* mutant. *Proc Natl Acad Sci U S A* 93: 9970-9974, 1996.
15. de Pinto MC, Paradiso A, Leonetti P, De Gara L. Hydrogen peroxide, nitric oxide and cytosolic ascorbate peroxidase at the crossroad between defence and cell death. *Plant J* 48: 784-795, 2006.
16. Dinakar C, Abhaypratap V, Yearla SR, Raghavendra AS, Padmasree K. Importance of ROS and antioxidant system during the beneficial interactions of mitochondrial metabolism with photosynthetic carbon assimilation. *Planta* 231: 461-474, 2010.
17. Dowdle J, Ishikawa T, Gatzek S, Rolinski S, Smirnoff N. Two genes in *Arabidopsis thaliana* encoding GDP-L-galactose phosphorylase are required for ascorbate biosynthesis and seedling viability. *Plant J* 52: 673-689, 2007.
18. Edwards EA, Rawsthorne S, Mullineux PM. Subcellular distribution of multiple forms of glutathione reductase in leaves of pea (*Pisum sativum* L.). *Planta* 180: 278-284, 1990.
19. Escobar M.A, Franklin K.A, Svensson A.S, Salter M.G, Whitlam G, Rasmusson A.G. Light regulation of the *Arabidopsis* respiratory chain. Multiple discrete photoreceptor

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

responses contribute to induction of type II NAD(P)H dehydrogenase genes. *Plant Physiol* 136: 2710–2721, 2004.

20. Foyer CH, Halliwell B. The presence of glutathione and glutathione reductase in chloroplasts: A proposed role in ascorbic acid metabolism. *Planta* 133: 21-25, 1976.
21. Foyer CH, Lelandais M. A Comparison of the Relative Rates of Transport of Ascorbate and Glucose Across the Thylakoid, Chloroplast and Plasmalemma Membranes of Pea Leaf Mesophyll Cells. *J Plant Physiol* 148: 391-398, 1996.
22. Heazlewood JL, Howell KA, Millar AH. Mitochondrial complex I from Arabidopsis and rice: orthologs of mammalian and fungal components coupled with plant-specific subunits. *Biochim Biophys Acta* 1604: 159-169, 2003.
23. Imai T, Niwa M, Ban Y, Hirai M, Oba K, Moriguchi T. Importance of the L-galactonolactone pool for enhancing the ascorbate content revealed by L-galactonolactone dehydrogenase-overexpressing tobacco plants. *Plant Cell Tiss Organ Cult* 96:105–112, 2009.
24. Jimenez A, Hernandez JA, Del Rio LA, Sevilla F. Evidence for the Presence of the Ascorbate-Glutathione Cycle in Mitochondria and Peroxisomes of Pea Leaves. *Plant Physiol* 114: 275-284, 1997.
25. Klodmann J, Sunderhaus S, Nimtz M, Jansch L, Braun HP. Internal architecture of mitochondrial complex I from Arabidopsis thaliana. *Plant Cell* 22: 797-810, 2010.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
26. Larkindale J, Hall JD, Knight MR, Vierling E. Heat stress phenotypes of Arabidopsis mutants implicate multiple signaling pathways in the acquisition of thermotolerance. *Plant Physiol* 138: 882-897, 2005.
27. Leferink NG, Fraaije MW, Joosten HJ, Schaap PJ, Mattevi A, van Berkel WJ. Identification of a gatekeeper residue that prevents dehydrogenases from acting as oxidases. *J Biol Chem* 284: 4392-4397, 2009.
28. Leferink NG, van den Berg WA, van Berkel WJ. l-Galactono-gamma-lactone dehydrogenase from Arabidopsis thaliana, a flavoprotein involved in vitamin C biosynthesis. *FEBS J* 275: 713-726, 2008.
29. Leferink NG, van Duijn E, Barendregt A, Heck AJ, van Berkel WJ. Galactonolactone dehydrogenase requires a redox-sensitive thiol for optimal production of vitamin C. *Plant Physiol* 150: 596-605, 2009.
30. Linster CL, Clarke SG. L-Ascorbate biosynthesis in higher plants: the role of VTC2. *Trends Plant Sci* 13: 567-573, 2008.
31. Lorence A, Chevone BI, Mendes P, Nessler CL. myo-inositol oxygenase offers a possible entry point into plant ascorbate biosynthesis. *Plant Physiol* 134: 1200-1205, 2004.
32. Mapson LW, Breslow E. Biological synthesis of ascorbic acid: L-galactono-gamma-lactone dehydrogenase. *Biochem J* 68: 395-406, 1958.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
33. Maruta T, Ichikawa Y, Mieda T, Takeda T, Tamoi M, Yabuta Y, Ishikawa T, Shigeoka S. The contribution of Arabidopsis homologs of L-gulonolactone oxidase to the biosynthesis of ascorbic acid. *Biosci Biotechnol Biochem* 74: 1494-1497, 2010.
34. Millar AH, Mittova V, Kiddle G, Heazlewood JL, Bartoli CG, Theodoulou FL, Foyer CH. Control of ascorbate synthesis by respiration and its implications for stress responses. *Plant Physiol* 133: 443-447, 2003.
35. Müller-Moulé P, Conklin PL, Niyogi KK. Ascorbate deficiency can limit violaxanthin de-epoxidase activity in vivo. *Plant Physiol* 128: 970-977, 2002.
36. Noguchi K, Yoshida K. Interaction between photosynthesis and respiration in illuminated leaves. *Mitochondrion* 8: 87-99, 2008.
37. Nunes-Nesi A, Araújo WL, Fernie AR. Targeting mitochondrial metabolism and machinery as a means to enhance photosynthesis. *Plant Physiol* 155: 101-107, 2011.
38. Nunes-Nesi A, Carrari F, Lytovchenko A, Smith AM, Loureiro ME, Ratcliffe RG, Sweetlove LJ, Fernie AR. Enhanced photosynthetic performance and growth as a consequence of decreasing mitochondrial malate dehydrogenase activity in transgenic tomato plants. *Plant Physiol* 137: 611-622, 2005.
39. Pavet V, Olmos E, Kiddle G, Mowla S, Kumar S, Antoniow J, Alvarez ME, Foyer CH. Ascorbic acid deficiency activates cell death and disease resistance responses in Arabidopsis. *Plant Physiol* 139: 1291-303, 2005.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
40. Pignocchi C, Fletcher JE, Barnes J, Foyer CH. The function of ascorbate oxidase (AO) in tobacco (*Nicotiana tabacum* L.). *Plant Physiology* 132: 1631–1641, 2003.
41. Pineau B, Layoune O, Danon A, De Paepe R. L-galactono-1,4-lactone dehydrogenase is required for the accumulation of plant respiratory complex I. *J Biol Chem* 283: 32500-32505, 2008.
42. Scheibe R, Backhausen J.E, Emmerlich V, Holtgreffe S. Strategies to maintain redox homeostasis during photosynthesis under changing conditions. *J Exp Bot* 56: 1481–1489, 2005.
43. Schertl P, Sunderhaus S, Klodmann J, Gergoff Grozeff GE, Bartoli CG, Braun HP. L-galactono-1,4-lactone dehydrogenase (GLDH) forms part of three subcomplexes of mitochondrial complex I in *Arabidopsis thaliana*. *J Biol Chem* 287:14412-14419, 2012.
44. Sweetlove LJ, Lytovchenko A, Morgan M, Nunes-Nesi A, Taylor NL, Baxter CJ, Eickmeier I, Fernie AR. Mitochondrial uncoupling protein is required for efficient photosynthesis. *Proc Natl Acad Sci U S A* 103: 19587-19592, 2006.
45. Szarka A, Horemans N, Bánhegyi G, Asard H. Facilitated glucose and dehydroascorbate transport in plant mitochondria. *Arch Biochem Biophys* 428: 73-80, 2004.
46. Szarka A, Horemans N, Kovács Z, Gróf P, Mayer M, Bánhegyi G. Dehydroascorbate reduction in plant mitochondria is coupled to the respiratory electron transfer chain. *Physiol Plant* 129: 225–232, 2007.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
47. Szarka A, Tomasskovics B, Bánhegyi G. The ascorbate-glutathione- $\alpha$ -tocopherol triad in abiotic stress response *Int J Mol Sci* 13: 4458-4483, 2012.
48. Tabata K, Oba K, Suzuki K, Esaka M. Generation and properties of ascorbic acid-deficient transgenic tobacco cells expressing antisense RNA for L-galactono-1,4-lactone dehydrogenase. *Plant J* 27: 139-148, 2001.
49. Talla S, Riazunnisa K, Padmavathi L, Sunil B, Rajsheel P, Raghavendra AS. Ascorbic acid is a key participant during the interactions between chloroplasts and mitochondria to optimize photosynthesis and protect against photoinhibition. *J Biosci* 36: 163-173, 2011.
50. Tamaoki M, Mukai F, Asai N, Nakajima N, Kubo A, Aono M, Saji H. Light-controlled expression of a gene encoding L-galactono- $\gamma$ -lactone dehydrogenase which affects ascorbate pool size in *Arabidopsis thaliana*. *Plant Sci* 164: 1111–1117, 2003.
51. Taniguchi M, Taniguchi Y, Kawasaki M, Takeda S, Kato T, Sato S, Tabata S, Miyake H, Sugiyama T. Identifying and characterizing plastidic 2-oxoglutarate/malate and dicarboxylate transporters in *Arabidopsis thaliana*. *Plant Cell Physiol* 43: 706–717, 2002.
52. Tomaz T, Bagard M, Pracharoenwattana I, Lindén P, Lee CP, Carroll AJ, Ströher E, Smith SM, Gardeström P, Millar AH. Mitochondrial malate dehydrogenase lowers leaf respiration and alters photorespiration and plant growth in *Arabidopsis*. *Plant Physiol* 154: 1143-1157, 2010.
53. Wheeler GL, Jones MA, Smirnoff N. The biosynthetic pathway of vitamin C in higher plants. *Nature* 393: 365-369, 1998.

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

- 1  
2  
3  
4  
5 54. Wolucka BA, Van Montagu M. GDP-mannose 3',5'-epimerase forms GDP-L-gulose, a  
6  
7 putative intermediate for the de novo biosynthesis of vitamin C in plants. *J Biol Chem*  
8  
9 278: 47483-47490, 2003.  
10  
11  
12  
13 55. Yabuta Y, Mieda T, Rapolu M, Nakamura A, Motoki T, Maruta T, Yoshimura K,  
14  
15 Ishikawa T, Shigeoka S. Light regulation of ascorbate biosynthesis is dependent on the  
16  
17 photosynthetic electron transport chain but independent of sugars in Arabidopsis. *J Exp*  
18  
19 *Bot* 58: 2661-2671, 2007.  
20  
21  
22  
23 56. Yoshida K, Terashima I, Noguchi K. How and why does mitochondrial respiratory chain  
24  
25 respond to light? *Plant Signal Behav* 6: 864-866, 2011.  
26  
27  
28  
29 57. Yoshida K, Watanabe CK, Hachiya T, Tholen D, Shibata M, Terashima I, Noguchi K.  
30  
31 Distinct responses of the mitochondrial respiratory chain to long- and short-term high-  
32  
33 light environments in Arabidopsis thaliana. *Plant Cell Environ* 34: 618-628, 2011.  
34  
35  
36  
37 58. Zechmann B. Subcellular distribution of ascorbate in plants. *Plant Signal Behav* 6: 360-  
38  
39 363, 2011.  
40  
41  
42  
43 59. Zsigmond L, Rigó G, Szarka A, Székely G, Otvös K, Darula Z, Medzihradzky KF,  
44  
45 Koncz C, Koncz Z, Szabados L. Arabidopsis PPR40 connects abiotic stress responses to  
46  
47 mitochondrial electron transport. *Plant Physiol* 146: 1721-1737, 2008.  
48  
49  
50  
51 60. Zsigmond L, Tomasskovics B, Deák V, Rigó G, Szabados L, Bánhegyi G, Szarka A.  
52  
53 Enhanced activity of galactono-1,4-lactone dehydrogenase and ascorbate – glutathione  
54  
55  
56  
57  
58  
59  
60

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

cycle in mitochondria from Complex III deficient Arabidopsis. *Plant Physiol Biochem*

49: 809-815, 2011.



Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

## Figure legends

### FIG. 1. Biosynthetic pathways to L-ascorbate in higher plants

The major ascorbate biosynthetic pathway is the D-Man/L-Gal or Smirnoff-Wheeler pathway (yellow). The minor L-gulose pathway is branched at GDP-Man 3',5'-epimerase (purple). The last step of this alternative route is the oxidation of L-gulono-1,4-lactone to ascorbate. The myo-Inositol branch (blue); also provides L-gulono-1,4-lactone for the biosynthesis of ascorbate via the formation of UDP-glucuronate by myo-Inositol oxidase. The D-galacturonic acid route (green), uses D-galacturonic acid, a principal component of cell wall and similar to the classical D-Man/L-Gal pathway provides galactono-1,4-lactone via the formation of L-galactonate by D-galacturonate reductase. Question marks indicate reactions where the gene and the specific enzyme have not been identified yet. The conversion of L-galactono-1,4-lactone to ascorbate by GLDH coupled to mitochondrial electron transfer chain is shown in red (mitochondria-like shape) circle.

(To see this illustration in color the reader is referred to the web version of this article at [www.liebertonline.com/ars](http://www.liebertonline.com/ars)).

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

**FIG. 2. Ascorbate generation and regeneration in plant mitochondria**

GLDH, embedded in the respiratory electron transfer chain, delivers electrons to oxidized cytochrome c by oxidizing L-galactono-1,4-lactone (GL) to ascorbate. The enzyme shows absolute requirement for oxidized cytochrome c, hence L-galactono-1,4-lactone can be defined as an alternative respiratory substrate in plants.

DHA is generated in the intermembrane space by the action of APX. DHA can reach the matrix by the aid of a GLUT-like transporter, where a glutathione-dependent ascorbate regenerating system (Foyer-Halliwel-Asada cycle) operates to reduce it to ascorbate. The supply of reductants is ensured in the form of NADH and NADPH to fuel these reactions from the tricarboxylic acid cycle.

(To see this illustration in color the reader is referred to the web version of this article at [www.liebertonline.com/ars](http://www.liebertonline.com/ars)).

**FIG. 3. The complex II coupled reduction of DHA – a possible bypass of complex III by function of the Asc/DHA redox couple**

DHA formed in the IMS, due to the activity of APX and cytochrome c oxidase, crosses the inner membrane with the mediation of a GLUT-like transporter. DHA is reduced to ascorbate in the matrix at complex II with exposure to the inner surface of the inner mitochondrial membrane (or by the ascorbate-glutathione cycle, see Fig. 3). A portion of ascorbate regenerated in the matrix

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

leave the mitochondria. The efflux of ascorbate has not been characterized yet at molecular level.

This electron route represented by the DHA/ascorbate redox couple may also gives survival value in case of complex III damage by taking up electrons at complex II (by the reduction of DHA to ascorbate) and providing electrons through a bypass way to complex IV (by the oxidation of ascorbate to DHA).

(To see this illustration in color the reader is referred to the web version of this article at [www.liebertonline.com/ars](http://www.liebertonline.com/ars)).

#### **FIG. 4. The photosynthesis - respiration - TCA-cycle - ascorbate network**

The electron flux through the photosynthetic ETC enhances the mitochondrial cytochrome c, AOX pathways and the biosynthesis of ascorbate. Furthermore, the activity of AOX itself enhances the biosynthetic rate of ascorbate. Since ascorbate is synthesized tightly coupled to the mitochondrial ETC, the oxidation of L-galactono-1,4-lactone to ascorbate provides electrons to complex IV in the case of the inhibition of TCA cycle. The reductant overload of the respiration can be reduced by light inhibition of TCA cycle and TCA cycle-linked respiration during light conditions. The inhibition of TCA cycle (by genetic approaches) accompanies by the dramatic increase in ascorbate levels. Ascorbate can elevate the intensity of photosynthesis by its multiple roles in photoprotection. Any changes in either of these elements of this network influence the others.


(To see this illustration in color the reader is referred to the web version of this article at [www.liebertonline.com/ars](http://www.liebertonline.com/ars)).

Szarka et al. : The inter-relationship of ascorbate transport, metabolism and mitochondrial, plastidic respiration

**FIG. 5. The ascorbate transporters and ascorbate fluxes of mitochondria and chloroplast**


The ultimate step of ascorbate biosynthesis is localized in the mitochondria. Hence ascorbate must be transported into the other compartments of the plant cell. The chloroplast envelope contains a low affinity ( $K_M \sim 20$  mM) ascorbate translocator, which mediates the facilitated diffusion of ascorbate from the cytosol to the stroma. The thylakoid membranes appear to have no transport system for ascorbate. This system is supported by the recently determined intracellular distribution of ascorbate. In the investigated plants highest levels of ascorbate specific labeling were found in the cytosol (21.7 mM) whereas mitochondria (10.4 mM) and chloroplasts (10.8 mM) contained intermediate levels. In concordance with the transport experiments within chloroplasts and mitochondria, ascorbate was only detected in the stroma and matrix, respectively, but not within the lumen of thylakoids or cristae (58).

(To see this illustration in color the reader is referred to the web version of this article at [www.liebertonline.com/ars](http://www.liebertonline.com/ars)).




447x224mm (300 x 300 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




254x190mm (300 x 300 DPI)

Peer Review Only



254x190mm (300 x 300 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




254x190mm (300 x 300 DPI)

Peer Review Only



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



199x252mm (300 x 300 DPI)

ew Only