

Charmed Bottom Baryon Spectroscopy

Zachary S. Brown*

The College of William & Mary

E-mail: zsbrown@email.wm.edu

William Detmold

Massachusetts Institute of Technology

E-mail: wdetmold@mit.edu

Stefan Meinel

Massachusetts Institute of Technology

E-mail: smeinel@mit.edu

Kostas Orginos

The College of William & Mary

E-mail: kostas@wm.edu

The spectrum of doubly and triply heavy baryons remains experimentally unexplored to a large extent. Although the detection of such heavy particle states may lie beyond the reach of experiments for some time, it is interesting compute this spectrum from QCD and compare results between lattice calculations and continuum theoretical models. Several lattice calculations exist for both doubly and triply charmed as well as doubly and triply bottom baryons. Here, we present preliminary results from the first lattice calculation of doubly and triply heavy baryons including both charm and bottom quarks. We use domain wall fermions for 2+1 flavors (up down and strange) of sea and valence quarks, a relativistic heavy quark action for the charm quarks, and non-relativistic QCD for the heavier bottom quarks. We present preliminary results for the ground state spectrum.

31st International Symposium on Lattice Field Theory - LATTICE 2013

July 29 - August 3, 2013

Mainz, Germany

*Speaker.

1. Landscape of heavy baryon spectroscopy

Experimental observations of charmed and bottom baryons date back to the mid 1970s with the discovery of the Λ_c^+ , and to the early 1990s with the discovery of the Λ_b . Since that time, the experimental spectrum has filled out with many precise measurements of the ground state energies of singly charmed and bottom baryons. In the past 10 years, results from BaBar and Belle, as well as the D0 and CDF experiments at Fermilab have provided more precise measurements of heavy flavor ground and excited states as well as a first glimpse of several new states, most notably the controversial Ω_b at CDF [1] and D0 [2] and the Ξ_{cc} at SELEX [3][4]. Although the former controversy has been resolved by a more recent measurement at LHCb [5], the latter observation has still not received experimental verification to date, and is only marginally in agreement with most current theoretical results. It is controversies like these, as well as the prospect of observations of new states that make this an exciting time for lattice QCD. The opportunities for heavy flavour spectroscopy from the lattice are threefold: 1) Lattice results can help to resolve controversies like the ones mentioned above (providing a model-independent point of comparison for current experimental measurements), 2) Lattice results can help to provide predictions for as yet unobserved states and 3) Spectroscopy calculations can help to pin down values for the low energy constants (LEC's) of chiral perturbation theory (a job that lattice calculations are uniquely suited for as quark masses can be varied), helping to aid in the understanding of hadronic physics at low energies. The predictive power of lattice QCD is especially exciting in the context of the upcoming increase in operating energy at the LHC scheduled for 2014.

The interest in heavy baryon spectroscopy within the lattice community is reflected by the array of calculations of the spectra of singly, doubly, and triply heavy baryons dating back to 2002 with Ref. [6] and more recently with [7],[8] and [9]. The interest in heavy hadron spectroscopy has not been limited to the lattice community, and there has been significant recent progress in theoretical calculations of the charmed and bottom baryon spectra as well using various models [10][11][12][13]. Included among these calculations are several determinations of the doubly heavy baryons which contain charm *and* bottom quarks. With the exception of the early quenched heavy hadron study [6], this subsector of doubly heavy states remains largely unexplored in lattice calculations, and there is no existing experimental data on charmed bottom baryons. This is a void that the present work will seek to fill, while at the same time reinforcing the existing lattice calculations of the charmed and bottom baryon spectra with a calculation using a single action for the light quark sector.

2. Lattice actions and parameters

In the present work, we utilize ensembles of gauge field configurations generated by the RBC/UKQCD collaboration [14]. We work at one lattice volume, with spatial extent $L \approx 2.7$ fm, and two lattice spacings: a fine lattice spacing with $a \approx 0.0849$ fm, and a coarse lattice spacing with $a \approx 0.1119$ fm. These ensembles were generated using the Iwasaki gauge action and include 2+1 flavors of dynamical light quarks implemented with a domain-wall action. Propagators for the light up and down quarks were generated on these ensembles at both the unitary point (where the sea quark mass is equal to the valence quark mass) as well as at several other lighter masses.

Strange quark propagators at an additional strange quark mass were computed on the coarse lattice at the light quark unitary point, allowing for an interpolation to the physical strange quark mass. The current calculation therefore includes sea-sea pion masses in the range $m_\pi^{ss} \approx 295$ MeV - 352 MeV, while the valence-valence pion masses range from $m_\pi^{ss} \approx 227$ MeV - 352 MeV. The kaon mass range is: $m_K^{vv} \approx 523$ MeV - 586 MeV. This partial quenching in both the light (up and down) and strange sectors, requires special care to be taken in ensuring that chiral extrapolations are correctly performed.

For the bottom quarks, we use improved lattice NRQCD [15] accurate through order v^4 and $\Lambda_{\text{QCD}}^2/m_b^2$, with the value for the coefficient c_4 computed specifically for the Iwasaki gauge action by Tom Hammant. As the mass of the charm quark is too light to call for a non-relativistic treatment on the current lattices, we use instead a relativistic heavy quark action in the Fermilab interpretation [16]. The action we use is of the same form as the one in Ref. [17]. To tune this action for the charm quark, the bare action parameters v and m_0 were simultaneously tuned to match the spin-averaged charmonium mass and enforce the dispersion relationship, both of which are given by:

$$M_{SA} = \frac{1}{4}M_{\eta_c} + \frac{3}{4}M_{J/\psi}, \quad c^2 = \frac{E^2(\mathbf{p}^2) - M_0^2}{\mathbf{p}^2}. \quad (2.1)$$

The additional bare parameters in the relativistic heavy quark action, c_E and c_B were set to their tree level values from lattice perturbation theory.

3. Extracting masses

Starting from three quark fields q, q', q'' , we construct two basic types of baryon operators, given by

$$O_5[q, q', q'']_\alpha = \varepsilon_{abc} (C\gamma_5)_{\beta\gamma} q_\beta^a q_\gamma^b (P_+ q'')_\alpha^c, \quad (3.1)$$

$$O_j[q, q', q'']_\alpha = \varepsilon_{abc} (C\gamma_j)_{\beta\gamma} q_\beta^a q_\gamma^b (P_+ q'')_\alpha^c, \quad (3.2)$$

where a, b, c are color indices, α, β, γ are spinor indices, C is the charge conjugation matrix, and P_+ is the positive-parity projector

$$P_+ = \frac{1}{2}(1 + \gamma_0). \quad (3.3)$$

We also compute the energies of heavy pseudoscalar and vector mesons, using two-point functions of the operators

$$O_5^{(M)}[\bar{q}, q'] = \bar{q}\gamma_5 q', \quad (3.4)$$

$$O_j^{(M)}[\bar{q}, q'] = \bar{q}\gamma_j q'. \quad (3.5)$$

It turns out to be useful to form certain linear combinations of baryon and meson energies, in which the quark-mass dependence and some of the other systematic uncertainties are expected to be reduced. For hadrons containing b quarks, such linear combinations can also be used to cancel the NRQCD energy shift, while for hadrons containing c quarks, this can help to eliminate effects introduced by tuning the relativistic heavy quark action for the charm sector. These linear combinations are taken as follows:

$$M_O = M_O^{latt} - \frac{n_c}{2} M_{SA_c}^{latt} - \frac{n_b}{2} M_{SA_b}^{latt} + \frac{n_c}{2} M_{SA_c}^{PDG} + \frac{n_b}{2} M_{SA_b}^{PDG}. \quad (3.6)$$

Here, $n_{c,b}$ is the number of charm or bottom quarks in the baryon, and $M_{SA_{c,b}}$ is the spin averaged charmonium or bottomonium mass.

We construct an operator basis using various levels of Gaussian smearing at the source and sink. We are reusing previously computed domain wall light and strange propagators, therefore we do not have the freedom to change the source smearing on these quarks. For the heavy quarks, we establish a hierarchy of smearing by reducing the level of smearing of the charm quarks in relation to the light quarks, and reducing the level of smearing of the bottom quarks in relation to that of the charm. The reasoning behind this is that the heavier fermions are expected to have a more spatially localized wavefunction. We apply smearing to either all or none of the heavy quarks, leading to 2×2 correlation function matrices for the triply heavy interpolating operators, and a 2×4 correlation function matrices for the doubly and triply heavy interpolating operators. Simultaneous fits of the resulting correlation function matrices were then carried out, allowing for variable fitting ranges for the individual correlation functions. This allows for an optimal fit range to be chosen for correlation functions with varying levels of excited state contamination. This process of choosing the optimal fit ranges was carried out by using initial guesses of the best fit ranges, and then stochastically minimizing the total $\chi^2/d.o.f.$ as a function of the fit ranges. An example of the effects of this minimization procedure can be seen in Figure. 1




Figure 1: An illustration for the effects of the stochastic fitting range determination for the state Ω_{ccb} . Here, the left hand plot corresponds to the initial guesses for fitting ranges, while the right hand plot corresponds to the optimized fitting ranges.

4. Chiral/Continuum extrapolations

As this spectroscopy calculation has been performed using several unphysical pion masses, in order to obtain a final result that is comparable to experimental results (as well as provide useful

predictions for baryons that have yet to be observed) we must perform an extrapolation of our results to the physical pion mass. One complicating factor stems from the fact that the calculation was performed using partial quenching in the light quark sector in order to extend the range of pion masses beyond those offered by the set of gauge field configurations used. To perform the extrapolations, we use partially quenched heavy hadron chiral perturbation theory (PQ HH χ PT) as a guide. PQ HH χ PT constructs the chiral Lagrangian to include heavy quark symmetries in the infinite mass limit. Expressions have been derived for the sea and valence mass dependence of singly [18] and doubly heavy [19] baryon masses, the latter relying on quark-diquark symmetry to relate the properties of singly heavy mesons to doubly heavy baryons. As the chiral expansion for SU(3) is thought to have poor convergence properties, we perform our extrapolations using SU(4|2) HH PQ χ PT. For singly heavy baryons, we extend existing expressions [18] for the baryon masses to include $\mathcal{O}(1/m_Q)$ corrections, while for the doubly heavy sector, we use the expressions presented in Ref. [19]. To include finite volume effects that appear from performing the calculation in a finite sized box, we include corrections to the chiral functions taken from Ref. [20]

In addition to the light quark mass dependence, the dependence on the finite lattice spacing must be accounted for. In order to accomplish this, we parametrize the lattice spacing dependence by adding a term of the form $c_a a^2$ to each of the chiral extrapolations. We then set c_a to zero to obtain the results in the continuum limit.

In order to stabilize the fits, the axial couplings g_1, g_2, g_3 (included in the chiral loop terms) are constrained using Gaussian priors to be near their values as calculated for the infinite mass limit in Ref. [21]. The Gaussian width used in the constraint was taken to be the statistical uncertainty for the g_i from Ref. [21] with an additional increase in the the width for the charmed baryons versus bottom baryons.

An example of the chiral and continuum extrapolations is presented in Figure 2.

5. Results and future outlook

Results for these extrapolations are presented in Tables 1 and 2. The uncertainties presented in these tables are statistical only. Systematic uncertainties are currently being explored, in both the mass extractions and chiral extrapolations. The limited mass range of m_π^{ss} may be inadequate in constraining the chiral extrapolations, therefore we are currently working to supplement the data sets with an additional sea pion mass.

Baryon	Lattice (GeV)	Expt. (GeV)	Baryon	Lattice (GeV)	Expt. (GeV)
Λ_c	2.137(74)	2.286	Λ_b	5.456(114)	5.619
Σ_c	2.444(81)	2.454	Σ_b	5.781(96)	5.811
Σ_c^*	2.518(82)	2.518	Σ_b^*	5.802(97)	5.832
Ξ_c	2.372(58)	2.467	Ξ_b	5.760(80)	5.791
Ξ_c'	2.526(62)	2.575	Ξ_b'	5.947(81)	-
Ξ_c^*	2.600(62)	2.645	Ξ_b^*	5.971(81)	-
Ω_c	2.615(67)	2.685	Ω_b	6.008(80)	6.071
Ω_c^*	2.690(67)	2.765	Ω_b^*	6.036(80)	-

Table 1: Chiral and continuum extrapolated results for all singly heavy states considered in the present work.




Figure 2: Example chiral and continuum extrapolations for the singly charmed states. Here, the open symbols represent partially quenched data while the filled symbols represent unitary points. The dashed line represents the coarse lattice spacing, the dotted line the fine lattice spacing and the solid line represents the continuum limit extrapolation. Both extrapolations are presented with the finite volume effects removed.

Baryon	Lattice (GeV)	Baryon	Lattice (GeV)	Baryon	Lattice (GeV)
Ξ_{cc}	3.558(39)	Ξ_{cb}	6.877(52)	Ξ_{bb}	10.185(53)
Ξ_{cc}^*	3.627(54)	Ξ_{cb}^*	6.915(62)	Ξ_{bb}^*	10.191(56)
Ω_{cc}	3.689(38)	Ω_{cb}	6.973(48)	Ω_{bb}	10.250(51)
Ω_{cc}^*	3.773(38)	Ω_{cb}^*	7.040(48)	Ω_{bb}^*	10.283(51)
Ω_{ccc}	4.794(9)	Ω_{ccb}	7.989(11)	Ω_{ccb}^*	8.012(12)
Ω_{cbb}	11.177(9)	Ω_{cbb}^*	11.206(11)	Ω_{bbb}	14.370(10)

Table 2: Chiral and continuum extrapolated results for all doubly and triply heavy states considered in the present work.

References

- [1] T. Aaltonen *et al.* *Phys.Rev.*, vol. D80, p. 072003, 2009.
- [2] V. Abazov *et al.* *Phys.Rev.Lett.*, vol. 101, p. 232002, 2008.
- [3] M. Mattson *et al.* *Phys.Rev.Lett.*, vol. 89, p. 112001, 2002.
- [4] A. Ocherashvili *et al.* *Phys.Lett.*, vol. B628, pp. 18–24, 2005.
- [5] D. Milanes *EPJ Web Conf.*, vol. 28, p. 04010, 2012.
- [6] N. Mathur, R. Lewis, and R. Woloshyn *Phys.Rev.*, vol. D66, p. 014502, 2002.
- [7] R. A. Briceño, H.-W. Lin, and D. R. Bolton *Phys.Rev.*, vol. D86, p. 094504, 2012.
- [8] C. Alexandrou, J. Carbonell, D. Christaras, V. Drach, M. Gravina, *et al.* *Phys.Rev.*, vol. D86, p. 114501, 2012.
- [9] Y. Namekawa *PoS*, vol. LATTICE2012, p. 139, 2012.
- [10] W. Roberts and M. Pervin *Int.J.Mod.Phys.*, vol. A23, pp. 2817–2860, 2008.
- [11] A. Martynenko *Phys.Lett.*, vol. B663, pp. 317–321, 2008.
- [12] D. Ebert, R. Faustov, V. Galkin, and A. Martynenko *Phys.Rev.*, vol. D66, p. 014008, 2002.
- [13] R. Roncaglia, D. Lichtenberg, and E. Predazzi *Phys.Rev.*, vol. D52, pp. 1722–1725, 1995.
- [14] Y. Aoki *et al.* *Phys.Rev.*, vol. D83, p. 074508, 2011.
- [15] G. P. Lepage, L. Magnea, C. Nakhleh, U. Magnea, and K. Hornbostel *Phys.Rev.*, vol. D46, pp. 4052–4067, 1992.
- [16] A. X. El-Khadra, A. S. Kronfeld, and P. B. Mackenzie *Phys.Rev.*, vol. D55, pp. 3933–3957, 1997.
- [17] L. Liu, H.-W. Lin, K. Orginos, and A. Walker-Loud *Phys.Rev.*, vol. D81, p. 094505, 2010.
- [18] B. C. Tiburzi *Phys.Rev.*, vol. D71, p. 034501, 2005.
- [19] T. Mehen and B. C. Tiburzi *Phys.Rev.*, vol. D74, p. 054505, 2006.
- [20] W. Detmold, C.-J. D. Lin, and S. Meinel *Phys.Rev.*, vol. D84, p. 094502, 2011.
- [21] W. Detmold, C.-J. D. Lin, and S. Meinel *Phys.Rev.Lett.*, vol. 108, p. 172003, 2012.