

Supercritical entanglement in local systems: Counterexample to the area law for quantum matter

Ramis Movassagh^{a,1,2} and Peter W. Shor^{b,c,1,2}

^aIBM Thomas J. Watson Research Center, Yorktown Heights, NY 10598; ^bDepartment of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139; and ^cComputer Science and Artificial Intelligence Laboratory, Massachusetts Institute of Technology, Cambridge, MA 02139

Edited by John Preskill, California Institute of Technology, Pasadena, CA, and approved September 19, 2016 (received for review April 18, 2016)

Quantum entanglement is the most surprising feature of quantum mechanics. Entanglement is simultaneously responsible for the difficulty of simulating quantum matter on a classical computer and the exponential speedups afforded by quantum computers. Ground states of quantum many-body systems typically satisfy an “area law”: The amount of entanglement between a subsystem and the rest of the system is proportional to the area of the boundary. A system that obeys an area law has less entanglement and can be simulated more efficiently than a generic quantum state whose entanglement could be proportional to the total system’s size. Moreover, an area law provides useful information about the low-energy physics of the system. It is widely believed that for physically reasonable quantum systems, the area law cannot be violated by more than a logarithmic factor in the system’s size. We introduce a class of exactly solvable one-dimensional physical models which we can prove have exponentially more entanglement than suggested by the area law, and violate the area law by a square-root factor. This work suggests that simple quantum matter is richer and can provide much more quantum resources (i.e., entanglement) than expected. In addition to using recent advances in quantum information and condensed matter theory, we have drawn upon various branches of mathematics such as combinatorics of random walks, Brownian excursions, and fractional matching theory. We hope that the techniques developed herein may be useful for other problems in physics as well.

quantum matter | local Hamiltonians entanglement entropy | area law | spin chains | Hamiltonian gap

Study of quantum many-body systems (QMBSs) is the study of quantum properties of matter and quantum resources (e.g., entanglement) provided by matter for building revolutionary new technologies such as a quantum computer. One of the properties of the QMBS is the amount of entanglement among parts of the system (1, 2). Entanglement can be used as a resource for quantum technologies and information processing (2–5); however, at a fundamental level it provides information about the quantum state of matter, such as near-criticality (6, 7). Moreover, systems with high entanglement are usually hard to simulate on a classical computer (8). How much entanglement do natural QMBSs possess? What are the fundamental limits on simulation of physical systems?

The area law says that entanglement entropy between two subsystems of a system is proportional to the area of the boundary between them. A generic state does not obey an area law (9); therefore, obeying an area law implies that a QMBS contains much less quantum correlation than generically expected. One can imagine that any given system has inherent constraints such as underlying symmetries and locality of interaction that restrict the states to reside on special submanifolds rendering their simulation efficient (10).

Since the discovery that the Affleck–Kennedy–Lieb–Tasaki (AKLT) model (11) is exactly solvable, and that the density matrix renormalization group method (DMRG) (12) works extremely well on 1D systems, we have come to believe that 1D systems are typically easy to simulate. The DMRG and its natural representation by matrix product states (MPS) (13) gave

systematic recipes for truncating the Hilbert space based on ignoring zero and small singular values in specifying the states of 1D systems. DMRG and MPS have been tremendously successful in practice for capturing the properties of matter in physics and chemistry (14, 15). We now know that generic local Hamiltonians, unlike the AKLT model, are gapless (16). One wonders about the limitations of DMRG.

The rigorous proof of a general area law does not exist; however, it holds for gapped systems in 1D (17). In the condensed matter community it is a common belief that gapped local Hamiltonians of QMBS on a D -dimensional lattice fulfill the area-law conjecture (8). That is, the entanglement entropy of a region of diameter L should scale as the area of the boundary $\mathcal{O}(L^{D-1})$ rather than its volume $\mathcal{O}(L^D)$. In the more general case, when the ground state is unique but the gap vanishes in the thermodynamical limit, it is expected that the area-law conjecture still holds, but now with a possible logarithmic correction, i.e., $S = \mathcal{O}(L^{D-1} \log L)$ (8). In other words, one expects that as long as the ground state is unique, the area law can be violated by at most a logarithmic factor. In particular, in 1D, it is expected that if we cut a chain of n interacting spins in the middle, the entanglement entropy should scale at most like $\log n$. This is based mostly on calculations done in 1+1 conformal field theories (CFTs) (7, 18), as well as in the Fermi liquid theory (19).

This belief has been seriously challenged by both quantum information and condensed matter theorists in recent years. Motivated by hardness results in quantum complexity theory, there are various interesting examples of 1D Hamiltonian constructions (20–22) that can have larger, even linear, scaling of entanglement entropy with the system’s size. In condensed matter physics, nontranslationally invariant models have been proposed and argued to violate the area law maximally (i.e., linearly for a chain) (23); Huijse and Swingle gave a supersymmetric model with some degree of fine-tuning that violates the

Significance

We introduce a class of exactly solvable models with surprising properties. We show that even simple quantum matter is much more entangled than previously believed possible. One then expects more complex systems to be substantially more entangled. For over two decades it was believed that the area law is violated by at most a logarithm in the system’s size for quantum matter (i.e., interactions satisfying physical reasonability criteria clearly stated in the article). In this work we introduce a class of physically reasonable models that we can prove violate the area law by a square root, i.e., exponentially more than the logarithm.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹R.M. and P.W.S. contributed equally to this work.

²To whom correspondence may be addressed. Email: ramis.mov@gmail.com or shor@math.mit.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1605716113/-DCSupplemental.




Fig. 3. Motzkin walk with $s=2$ colors of length $2n=10$. The height m quantifies the degree of correlation between the two halves.

Therefore, the local Hamiltonian, with projectors as interactions, that has the Motzkin state as its unique zero-energy ground state is

$$H = \Pi_{\text{boundary}} + \sum_{j=1}^{2n-1} \Pi_{j,j+1} + \sum_{j=1}^{2n-1} \Pi_{j,j+1}^{\text{cross}}, \quad [2]$$

where $\Pi_{j,j+1}$ implements the local changes discussed above and is defined by

$$\Pi_{j,j+1} \equiv \sum_{k=1}^s \left[|D^k\rangle_{j,j+1} \langle D^k| + |U^k\rangle_{j,j+1} \langle U^k| + |\phi^k\rangle_{j,j+1} \langle \phi^k| \right], \quad [3]$$

with $|D^k\rangle \sim [|0d^k\rangle - |d^k0\rangle]$, $|U^k\rangle \sim [|0u^k\rangle - |u^k0\rangle]$, and $|\phi^k\rangle \sim [|00\rangle - |u^k d^k\rangle]$. The projectors $\Pi_{\text{boundary}} \equiv \sum_{k=1}^s [|d^k\rangle_1 \langle d^k| + |u^k\rangle_{2n} \langle u^k|]$ select out the Motzkin state by excluding all walks that start and end at nonzero heights. Lastly, $\Pi_{j,j+1}^{\text{cross}} \equiv \sum_{k \neq i} |u^k d^i\rangle_{j,j+1} \langle u^k d^i|$ ensures that balancing is well ordered. For example, we want to ensure that the unbalanced sequence of steps $u^3 u^1 u^2$ is balanced by $d^2 d^1 d^3$ and not, say, $d^1 d^3 d^2$. $\Pi_{j,j+1}^{\text{cross}}$ penalizes wrong ordering by prohibiting $00 \leftrightarrow u^k d^i$ when $k \neq i$. These projectors are required only when $s > 1$ and do not appear in ref. 29.

The difference between the ground-state energy and the energy of the first excited state is called the gap. One says a system is gapped when the difference between the two smallest energies is at least a fixed constant in the thermodynamical limit ($n \rightarrow \infty$). Otherwise the system is gapless.

Whether a system is gapped has important implications for its physics. When it is gapless, the scaling by which the gap vanishes as a function of the system's size has important consequences for its physics. For example, gapped systems have exponentially decaying correlation functions (22), and quantum critical systems are necessarily gapless (31). Moreover, systems that obey a CFT are gapless but the gap must vanish as $1/n$ (32). Therefore, to quantify the physics, it is desirable to find new techniques for analyzing the gap that can be applied in other scenarios.

The model proposed here is gapless and the gap scales as n^{-c} where $c \geq 2$ is a constant. We prove this by finding two functions, both of which are inverse powers of n such that the gap is always smaller than one of them (called an upper bound) and greater than the other (called a lower bound). We use techniques from mathematics such as Brownian excursions and universal convergence of random walks to a Brownian motion, as well as other ideas from computer science such as linear programming and fractional matching theory. We describe the ideas and leave the details of the proofs for *SI Appendix*.

To prove an upper bound on the gap, one needs a state $|\phi\rangle$ that has a small constant overlap with the ground state and such that $\langle \phi | H | \phi \rangle \geq \mathcal{O}(n^{-2})$. Take

$$|\phi\rangle = \frac{1}{\sqrt{M_{2n}}} \sum_{m_p} e^{2\pi i \theta \bar{A}_p} |m_p\rangle, \quad [4]$$

where the sum is over all Motzkin walks, M_{2n} is the total number of Motzkin walks on $2n$ steps, \bar{A}_p is the area under the Motzkin walk m_p , and θ is a constant to be determined by the condition of a small constant overlap with the ground state. The overlap with the ground state is defined by $\langle \mathcal{M}_{2n} | \phi \rangle = (1/M_{2n}) \sum_{m_p} e^{2\pi i \theta \bar{A}_p}$. As $n \rightarrow \infty$, the random walk converges to a Wiener process (33) and a random Motzkin walk converges to a Brownian excursion (34). We scale the walks such that they take place on the unit interval. The scaled area is denoted by A and $\theta \rightarrow \theta$. In this limit, the overlap becomes (see Fig. 4 for the density and Fig. 5 for its Fourier transform; $F_A(\theta)$ is the Fourier transform of the probability density function, which is called the characteristic function.)

$$\lim_{n \rightarrow \infty} \langle \mathcal{M}_{2n} | \phi \rangle \approx F_A(\theta) \equiv \int_0^\infty f_A(x) e^{2\pi i x \theta} dx, \quad [5]$$

where $f_A(x)$ is the probability density function for the area of the Brownian excursion (35) shown in Fig. 4. In Eq. 5, taking $\theta \ll \mathcal{O}(1)$, gives $\lim_{n \rightarrow \infty} \langle \mathcal{M}_{2n} | \phi \rangle \approx 1$ because it becomes the integral of a probability distribution. However, taking $\theta \gg \mathcal{O}(1)$ gives a highly oscillatory integrand that nearly vanishes. To have a small constant overlap with the ground state, we take θ to be the standard of deviation of $f_A(x)$. Direct calculation then gives $\langle \phi | H | \phi \rangle = \mathcal{O}(n^{-2})$. This upper bound decisively excludes the possibility of the model being describable by a conformal field theory (18).

Using various ideas in perturbation theory, computer science, and mixing times of Markov chains we obtain a lower bound on the gap that scales as n^{-c} , where $c \gg 1$. Because it might be of independent interest in other contexts, we present a combinatorial and self-contained exposition of the proof in the *SI Appendix*, different in some aspects from that given in ref. 29.

The model above has a unique ground state because the boundary terms select out the Motzkin state among all other walks with different fixed initial and final heights. Without the boundary projectors, all walks that start at height m_1 and end at height m_2 with $-2n \leq m_1, m_2 \leq 2n$ are ground states. For example, when $s = 1$, the ground-state degeneracy grows quadratically with the system's size $2n$ and exponentially when $s > 1$.

For the $s = 1$ case, if we impose periodic boundary conditions, then the superposition of all walks with an excess of k up (down) steps is a ground state. This gives $4n + 1$ degeneracy of the ground state, which includes unentangled product states.




Fig. 4. Plot of the probability density of the area under a Brownian excursion $f_A(x)$ on $[0, 1]$.




Fig. 5. Fourier transform of $f_A(x)$ as defined by Eq. 5.

When $s > 1$, each one of the walks with k excess up (down) steps can be colored exponentially many ways; however, generically they will not be product states. Consider an infinite chain $(-\infty, \infty)$ and take $s > 1$. There is a ground state of this system that corresponds to the balanced state, where on average for each color, the state contains as many u^i as d^i . Suppose we restrict our attention to any block of n consecutive spins. This block contains the sites $j, j+1, \dots, j+n-1$, which is a section of a random walk. Let us assume that it has initial height m_j and final height m_{j+n-1} . Further, let us assume that the minimum height of this section is m_k with $j \leq k \leq j+n-1$. From the theory of random walks, the expected values of $m_j - m_k$ and of $m_{j+n-1} - m_k$ are $\Theta(\sqrt{n})$. The color and number of any unmatched step-ups in this block of n spins can be deduced from the remainder of the infinite walk. Thus, a consecutive block of n spins has an expected entanglement entropy of $\Theta(\sqrt{n})$ with the rest of the chain. A similar argument shows that any block of n spins has an expected half-block entanglement entropy of $\Theta(\sqrt{n})$.

If we take $s = 1$, where the ground state can be a product state, the \sqrt{n} unmatched step-up just mentioned can be matched anywhere on the remaining left and right part of the chain. Two consecutive blocks of n spins can be unentangled because the number of unbalanced steps that are matched in the next block is uncorrelated with the number of unbalanced steps in the first block. However, when $s > 1$ the ordering has to match. Even though the number of unbalanced steps in two consecutive blocks is uncorrelated, the order of the types of unbalanced steps in them agrees.

The Hamiltonian without the boundary terms is truly translationally invariant, yet has a degenerate ground state. We now propose a model with a unique ground state that has the other desirable properties of the model with boundaries, such as the gap and entanglement entropy scalings as before. To do so, we put the system in an external field, where the model is described by the Hamiltonian

$$\begin{aligned} \tilde{H} &\equiv H + \epsilon F \\ F &\equiv \sum_{i=1}^{2n} \sum_{k=1}^s (|d^k\rangle_i \langle d^k| + |u^k\rangle_i \langle u^k|), \end{aligned} \quad [6]$$

where H is as before but without the boundary projectors and $\epsilon = \epsilon_0/n$ with ϵ_0 being a small positive constant. It is clear that F

treats u and d symmetrically; therefore, the change in the energy as a result of applying an external field depends only on the total number of unbalanced steps denoted by m . We denote the change in the energy of m unbalanced steps by ΔE_m . When $s = 1$, the degeneracy after applying the external field will be one for the Motzkin state, twofold when there is a single imbalance, threefold for two imbalances, etc. Because the energies are equal for all m imbalance states, it is enough to calculate the energy for an excited state with m imbalances resulting only from excess step-ups. We denote these states by $|g_m\rangle$, where $0 \leq m \leq 2n$.

The first-order energy corrections, obtained from first-order degenerate perturbation theory, are analytically calculated to be

$$\epsilon \langle g_m | F | g_m \rangle \approx 4\sigma\epsilon n + \frac{m\epsilon}{8\sqrt{s}} \left(\frac{m}{n} \right). \quad [7]$$

The physical conclusion is that the Hamiltonian without the boundary projectors, in the presence of an external field F , has the Motzkin state as its unique ground state with energy $4\sigma\epsilon_0$. Moreover, what used to be the rest of the degenerate zero-energy states acquire energies above $4\sigma\epsilon_0$ that for first elementary excitations scale as $1/n^2$. Moreover, the numerical calculations indicate that the spin-spin correlation functions are flat (36). We leave further investigations for future work.

The energy corrections just derived do not mean that the states with m imbalances will make up for all of the low-energy excitations. For example, when $s > 1$, in the presence of an external field, the energy of states with a single crossed term will be lower than those with large m imbalances and no crossings.

Because $\|\epsilon F\| \ll \|H\|$, the ground state will deform away from the Motzkin state to prefer the terms with more zeros in the superposition. But, as long as ϵ is small, the universality of Brownian motion guarantees the scaling of the entanglement entropy. It is, however, not yet clear to us whether ϵ can be tuned to a quantum critical point where the ground state has a sharp transition from highly entangled to nearly a product state. It is possible that the transition is smooth and that the entanglement continuously diminishes as ϵ becomes larger. For example, in the limit where $|\epsilon| \gg \|H\|/\|F\|$, the effective unperturbed Hamiltonian is approximately F , whose ground state is simply the product state $|0\rangle^{\otimes 2n}$.

Our model shows that simple physical systems can be much more entangled than expected. From a fundamental physics perspective, it is surprising that a 1D translationally invariant quantum spin chain with a unique ground state has about \sqrt{n} entanglement entropy. Moreover, this adds to the collection of exactly solvable models from which further physics can be extracted. Such a spin chain can in principle be experimentally realized, and the large amount of entanglement may be used as a resource for quantum technologies and computation.

ACKNOWLEDGMENTS. We thank Sergey Bravyi and Adrian Feiguin for discussions. R.M. acknowledges the Herman Goldstone Fellowship at IBM Thomas J. Watson Research Center and the support of the National Science Foundation (NSF) through Grant DMS-1312831. P.W.S. was supported by the US Army Research Laboratory's Army Research Office through Grant W911NF-12-1-0486, the NSF through Grant CCF-121-8176, and by the NSF through the Science and Technology Center for Science of Information under Grant CCF0-939370.

1. Tura J, et al. (2014) Quantum nonlocality. Detecting nonlocality in many-body quantum states. *Science* 344(6189):1256–1258.
2. Chuang I, Nielsen M (2000) *Quantum Computation and Quantum Information* (Cambridge Univ Press, Cambridge, UK), 1st Ed.
3. Kimble HJ (2008) The quantum internet. *Nature* 453(7198):1023–1030.
4. Acin A, Cirac JJ, Lewenstein M (2007) Entanglement percolation in quantum networks. *Nat Phys* 3(4):256–259.

5. Brun T, Devetak I, Hsieh M-H (2006) Correcting quantum errors with entanglement. *Science* 314(5798):436–439.
6. Coleman P, Schofield AJ (2005) Quantum criticality. *Nature* 433(7023):226–229.
7. Osterloh A, Amico L, Falci G, Fazio R (2002) Scaling of entanglement close to a quantum phase transition. *Nature* 416(6881):608–610.
8. Eisert J, Cramer M, Plenio MB (2010) Area laws for the entanglement entropy. *Rev Mod Phys* 82:277.

9. Hayden P, Leung D, and Winter A (2006) Aspects of generic entanglement. *Commun Math Phys* 265(1):95–117.
10. Landau Z, Vazirani U, Vidick T (2015) A polynomial time algorithm for the ground state of one-dimensional gapped local Hamiltonians. *Nat Phys* 11:566–569.
11. Affleck I, Kennedy T, Lieb EH, Tasaki H (1998) Finitely correlated states on quantum spin chains. *Commun Math Phys* 115(3):477–528.
12. White SR (1992) Density matrix formulation for quantum renormalization groups. *Phys Rev Lett* 69(19):2863–2866.
13. Perez-Garcia D, Verstraete F, Wolf MM, Cirac JI (2007) Matrix product representation. *Quantum Inf Comput* 7(5):401–430.
14. Kurashige Y, Chan GK-L, Yanai T (2013) Entangled quantum electronic wavefunctions of the Mn_4CaO_5 cluster in photosystem II. *Nat Chem* 5(8):660–666.
15. Yan S, Huse DA, White SR (2011) Spin-liquid ground state of the $S = 1/2$ kagome Heisenberg antiferromagnet. *Science* 332(6034):1173–1176.
16. Movassagh R (2016) Generic local Hamiltonians are gapless. arXiv:1606.09313 [quant-ph].
17. Hastings MB (2007) An area law for one-dimensional quantum systems. *J Stat Phys* 8: 08024.
18. Calabrese P and Cardy J (2009) Entanglement entropy and conformal field theory. *J Phys A Math Theor* 42:504005.
19. Wolf MM (2006) Violation of the entropic area law for fermions. *Phys Rev Lett* 96(1): 010404.
20. Movassagh R (2012) Eigenvalues and low energy eigenvectors of quantum many-body systems. PhD thesis (Massachusetts Institute of Technology, Cambridge). arXiv: 1211.4908 [quant-ph].
21. Irani S (2010) Ground state entanglement in one-dimensional translationally invariant quantum systems. *J Math Phys* 51:022101.
22. Gottesman D, Hastings MB (2010) Entanglement vs. gap for one-dimensional spin systems. *New J Phys* 12:025002.
23. Vitagliano G, Riera A, Latorre JI (2010) Volume-law scaling for the entanglement entropy in spin-1/2 chains. *New J Phys* 12(11):113049.
24. Huijse L, Swingle B (2013) Area law violations in a supersymmetric model. *Phys Rev B* 87(3):035108.
25. Gori G, Paganelli S, Sharma A, Sodano P, Trombettoni A (2014) Bell-paired states inducing volume law for entanglement entropy in fermionic lattices. arXiv: 1405.3616.
26. Shiba N, Takayanagi T (2014) Volume law for the entanglement entropy in non-local QFTs. *J High Energy Phys* 2014(2):1–16.
27. Ramirez G, Rodriguez-Laguna J, Sierra G (2015) Entanglement over the rainbow. arXiv:1503.02695.
28. Swingle B, Todadri S (2013) Universal crossovers between entanglement entropy and thermal entropy. *Phys Rev B* 87(4):045123.
29. Bravyi S, Caha L, Movassagh R, Nagaj D, Shor PW (2012) Criticality without frustration for quantum spin-1 chains. *Phys Rev Lett* 109(20):207202.
30. Verstraete F, Wolf MM, Cirac J (2009) Quantum computation and quantum-state engineering driven by dissipation. *Nat Phys* 5(9):633–636.
31. Sachdev S (2007) *Quantum Phase Transitions* (Wiley Online Library).
32. Di Francesco P, Mathieu P, Sénéchal D (1997) *Conformal Field Theory* (Springer, New York).
33. Prokhorov YV (1956) Convergence of random processes and limit theorems in probability theory. *Theory Probab Appl* 1(2):157–214.
34. Durrett RT, Inglehart DL (1997) Functionals of Brownian meander and Brownian excursion. *Ann Probab* 5(1):130–135.
35. Janson S, et al. (2007) Brownian excursion area, Wrights constants in graph enumeration, and other Brownian areas. *Probability Surveys* 4:80–145.
36. Movassagh R (2016) Entanglement and correlation functions of a recent exactly solvable spin chain. arXiv:1602.07761 [quant-ph].