S1-H03

Efficient Utilization of Plant Materials for Profitable Manufacturing of the Wellness Industry

Umar Isah Abubakar, Chua Lee Suan , Ramlan Aziz

Institute of Bioproduct Development (IBD), Universiti Teknologi Malaysia, 81310 Skudai, Johor, Malaysia

Abstract

Phytochemical compounds are the active ingredients for the wellness industries that are mostly produced from the leaves, flowers, seeds, stems, barks and roots of plant materials through solid-liquid extraction in the upstream section of the industry. The type of solid-liquid extractor and its operation has a crucial influence on the overall performance of the process, and consequently, on the profitability of the whole wellness industry. Many process variables such as solvent type, particle size, solid-liquid ratio, degree of agitation, time and temperature influence the rate of solid-liquid extraction of phytochemical compound (s). The maximum production of the amount of phytochemical compound (s) of interest can be achieved through proper utilization of raw material by clear understanding of the effects of process variables, manipulating and controlling these independent variables during operations. This study has developed a new methodology called extraction selectivity in addition to extraction yield for characterizing the performance of the esolid-liquid extraction of andrographis paniculata for the solid-liquid extraction of andrographolide water extracts and other water soluble phytochemical compounds for the wellness industry. The analysis gives more insights into design, scale-up of solid-liquid extractor, operation planning, and suggests a new way of looking into operation and process optimization.

Keywords: Active ingredients, Raw materials, Process variables, Performance of solid-liquid extraction, Profitability of the wellness industry, Andrographolide water extracts.

ABSTRACT FOR PARALLEL SESSIONS

S2-C03

Optimization Of Polysaccharide Production By Lactobacillus Kefiranofaciens Using Response Surfac Methodology

Dailin, D.J¹, Harun, N¹, Abd Malek, R¹, Othman, N.Z¹, El Enshasy, H.A^{1,2*} Elsayed A. Elsayed^{3,4}

¹Institute of Bioproduct Development (IBD), Universiti Teknologi Malaysia (UTM), Johor Bahru, Malaysia. ²City for Scientific Research and Technology Applications, New Burg Al Arab, Alexandria, Egypt. ³Bioproducts Research Chair, Zoology Department, Faculty of Science, King Saud University, Riyadh, Saudi Arab ⁴Natural and Microbial Product Department, National Research Centre, Dokki, Cairo, Egypt.

Abstract

Kefiran is an exopolysaccharides produced by *Lactobacillus kefiranofaciens* which was isolated from kefir grain Kefiran has wide applications mainly in food and pharmaceutical industries. Growth and kefiran production of *kefiranofaciens* can be significantly enhanced by using mixed culture technique. *Saccharomyces cerevisiae* in th study was used to enhance the kefiran production by reducing lactic acid accumulation in the cultivation mediur. The statistical analysis was used for optimization study by using response surface methodology based c Box-Behnken design. The interaction was studied between three different levels of variables that were lactos yeast extract and phosphate. A second order polynomial model was used to correlate the factors. The model was found to be significant. The optimum concentration of lactose, yeast extract and phosphate obtained usir statistical media optimization is 59.09 g L⁻¹, 8.69 g L⁻¹and 0.5 g L⁻¹ respectively. Kefiran production in optimize medium was 0.97 g L⁻¹ which gave an increase in kefiran production up to 42.65 % compared to the un-optimize medium which was only 0.68 g L⁻¹ of kefiran.

Keywords: Kefiran, exopolysaccharides, medium optimization, *Lactobacillus kefiranofaciens, Saccharomyco cerevisiae*.