



## Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <http://oatao.univ-toulouse.fr/>  
Eprints ID: 16425

**To cite this version:** Alignier, Audrey and Meiss, Helmut and Petit, Sandrine and Reboud, Xavier *Postdispersal weed seed predation ranged between 19 to 84% per week following a species preference rank.* (2008)  
In: 24th German Conference on Weed Biology and Weed Control, 4 March 2008 - 6 March 2008 (Hohenheim, Germany). (Unpublished)

Any correspondence concerning this service should be sent to the repository administrator:  
[staff-oatao@listes-diff.inp-toulouse.fr](mailto:staff-oatao@listes-diff.inp-toulouse.fr)

# Postdispersal weed seed predation ranged between 19 to 84% per week following a species preference rank

Audrey ALIGNIER, Helmut MEISS, Sandrine PETIT & Xavier REBOUD  
INRA Dijon, France, UMR 1210 'Biologie et Gestion des Adventices' (Weed biology & management)

Contact: Audrey.Alignier@toulouse.inra.fr  
Helmut.Meiss@dijon.inra.fr

Postdispersal seed predation may have strong impacts on the demography of annual plants<sup>[1]</sup>.

If this phenomenon is well-known in tropical and woody systems, there is no information about weed seed predation in French agroecosystems.

## Material & Methods

To quantify seed loss due to predation, seeds of 7 contrasted weed species + plastic globules were glued on sandpaper cards<sup>[2]</sup> and pinned on the soil surface of an organic winter wheat.

7 weed + 1 artificial species = 36 unique species pairs  
\* 15 reps. + 2 controls = 612 cards (2\*25 seeds each)

Double **control** for accidental seed losses:

- 1) Seed cards placed under a narrow wire mesh
- 2) Presentation of dummy seeds (plastic globules)



Seed cards




Wire mesh



Dummy seeds

## Results



✓ Seed predation was significantly higher than removals of plastic globules and seed losses under the wire mesh.

✓ Seed predation differed among the tested species (Wilcoxon  $\chi^2 = 381$ ;  $df = 7$ ;  $p < 0.0001$ ). ("species preference rank", Figure)

✓ Seed characteristics like weight, color, lipid or protein content are **not linearly correlated** to the observed predation rate.

✓ **Spatial and temporal variation** could not directly be explained by climatic variables or field border distance.

**H**igh seeds removal rates (19-84%) suggest that:

- 1) Weed seeds may constitute an **important food resource** for animals in the agro-ecosystem.
- 2) Seed predation may be an important factor **shaping weed population dynamics** and **community composition** of arable weeds.

So, to make optimal use of seed predation for reducing weed infestation in arable farming, research should focus on the mechanisms causing variability in predation.

[1] Pullaro, et al. 2006. Effects of killed cover crop mulch on weeds, weeds seeds, and herbivores. *Agriculture, Ecosystems and Environment*, 115, 97-104.

[2] Westerman et al. 2003. Relative importance of vertebrates and invertebrates in epigeic weed seed predation in organic cereals fields. *Agriculture, Ecosystems and Environment*, 95, 417-425.