



CrossMark

Revista Mexicana de Biodiversidad

Revista Mexicana de Biodiversidad 87 (2016) 255–257

Instituto
de Biología
UNAMwww.ib.unam.mx/revista/

Research note

The role of *Phyllocaulis variegatus* (Mollusca: Veronicellidae) in the transmission of digenean parasites

El papel de Phyllocaulis variegatus (Mollusca: Veronicellidae) en la transmisión de parásitos digéneos

Romina Valente^{a,*}, Julia Inés Díaz^b, Oscar Daniel Salomón^a, Graciela Teresa Navone^b^a Instituto Nacional de Medicina Tropical, Ministerio de Salud de la Nación, Jujuy s/n, 3370 Puerto Iguazú, Provincia de Misiones, Argentina^b Centro de Estudios Parasitológicos y de Vectores (CCT La Plata-CONICET), Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, calle 120 e/61 y 62, B1900FWA La Plata, Provincia de Buenos Aires, Argentina

Received 12 November 2014; accepted 14 September 2015

Available online 26 February 2016

Abstract

Ninety-five veronicellid slugs identified as *Phyllocaulis variegatus* were collected in Puerto Iguazú, Misiones Province, Argentina. Specimens were examined for parasites. Metacercariae of the genus *Brachylaima* (Brachylaimidae) were recovered from the body cavity near the female genital pore. The results establish the capability of *P. variegatus* to act as an intermediate host of *Brachylaima* sp. in the area. This study represents the first record of metacercariae of *Brachylaima* sp. in Argentina.

All Rights Reserved © 2015 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access item distributed under the Creative Commons CC License BY-NC-ND 4.0.

Keywords: *Brachylaima* sp.; Metacercariae; *Phyllocaulis variegatus*; Intermediate host; Argentina

Resumen

Noventa y cinco babosas terrestres de la familia Veronicellidae identificadas como *Phyllocaulis variegatus* fueron recolectadas en Puerto Iguazú, provincia de Misiones, Argentina para ser examinadas en busca de parásitos. Las metacercarias del género *Brachylaima* fueron halladas en la cavidad del cuerpo cerca del poro genital femenino. Este resultado establece la capacidad de *P. variegatus* de actuar como hospedador intermediario de *Brachylaima* sp. en el área y representa el primer registro de metacercarias de *Brachylaima* sp. en Argentina.

Derechos Reservados © 2015 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia Creative Commons CC BY-NC-ND 4.0.

Palabras clave: *Brachylaima* sp.; Metacercaria; *Phyllocaulis variegatus*; Hospedador intermediario; Argentina

The family Brachylaimidae Joyeux and Foley, 1930 (Trematoda) comprises 14 genera, including *Brachylaima* Dujardin, 1843 (the type genus). The genus *Brachylaima* contains 72 species that parasitize mammals and birds as definitive hosts around the world, except Antarctica (Butcher, 2003; Gracenea & González-Moreno, 2002; Lunaschi & Drago, 2012; Yamaguti, 1975). Terrestrial snails and slugs are involved as first and

second intermediate hosts (Barger, 2011; Butcher, 2003; Butcher, Parasuramar, Thompson, & Grove, 1998; Cribb, 1990). The specificity at the level of the definitive host greatly differs in the Brachylaimidae (Guisantes, Benito, Estibalez, & Mas-Coma, 1994). Some species of the genus *Brachylaima* show an extensive range of potential definitive hosts, and experimental infections have demonstrated that the same species are able to infect and successfully develop in different species of mammals and birds (Timon-David, 1959). Several cases of human infections by adults of *Brachylaima* sp. causing consistent gastro-intestinal symptoms have been reported in Australia (Butcher & Groove, 2001; Butcher, Talbot, Norton, Kirk,

* Corresponding author.

E-mail address: romina_valente81@hotmail.com (R. Valente).

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

Cribb, & Forsyth, 1996; Butcher et al., 1998; Butcher, Brealey, Grove, & Dymock, 2002). Infections can occur after eating raw mollusks containing metacercariae (Butcher, 2003).

During recent investigations focused on the role of terrestrial mollusks as intermediate hosts of parasites of public health importance in Argentina, terrestrial mollusks were collected (with permission of the Ministerio de Ecología y Recursos Naturales Renovables de la Provincia de Misiones) in twilight hours from Puerto Iguazú (Misiones, Argentina; 25°36'39" S, 54°34'49" W) during spring and summer of 2013 and 2014. Ninety-five specimens of *Phyllocaulis variegatus* Semper 1885 (Mollusca: Veronicellidae) were collected. The slugs were relaxed using menthol crystals dissolved in distilled water and kept in a refrigerator for up to 48 h (Cuezzo, 2013), then preserved in 70% ethanol and examined for parasites under a stereo microscope.

Thirteen of the 95 *P. variegatus* examined were parasitized by trematodes, recovering 235 metacercariae from the body cavity near the female genital pore. Parasites were stained with acetic carmine, dehydrated through ascending ethanol series, cleared with methyl salicylate, and mounted in Canada balsam. Drawings and measurements (range following by mean, in micrometers) were obtained with the aid of a light microscope Olympus BX51 with a camera lucida, and photographs were taken with an Olympus DP 71 camera. Identification of parasites was achieved following keys and specific bibliography (e.g., Butcher, 2003; Gibson, Jones, & Bray, 2001; Yamaguti, 1975). Voucher specimens of parasites were deposited in the Helminthological Collection of the Museo de la Plata, Argentina under the catalog number MLP He 6914.

Description based on 10 specimens of *Brachylaima* sp. (Figs. 1 and 2). Body elongate, 1670–2280 (1976) long, 480–750 (639) wide. Forebody 520–770 (653) long. Oral sucker 300–400 (360) long, 270–380 (335) wide. Ventral sucker 250–350 (291) long, 230–350 (279) wide. Inter-sucker distance 140–350 (262). Pharynx muscular, well developed, 140–180 (155) long, 140–170 (150) wide; sinuous caeca extending almost to the posterior margin of body. Anterior testis 65–100 (76) long, 50–84 (66) wide; posterior testis 66–100 (83) long, 50–93 (67) wide. Ovary intertesticular 55–92 (72) long, 50–80 (64) wide.

Although the specimens were immature, the following morphological characteristics allowed us to assign them to the genus *Brachylaima*: ventral sucker located in the first third of the body; genital pore in the zone of the anterior margin of the anterior testis; gonads located in the posterior part of the body, close together and near the posterior extremity; vitelline glands extending from the anterior margin of the ventral sucker to the anterior zone of the first testis and not surpassing it; uterus extending up to the intestinal bifurcation and thus clearly exceeding the anterior margin of the ventral sucker.

Metacercariae were found near the female genital pore, in contrast to previous studies, which recorded the metacercariae in the pericardial sac (Thiengo & Amato, 1995), or kidney (Barger & Hnida, 2008).

This is the first report of metacercariae of *Brachylaima* sp. in Argentina. In contrast, different species of *Brachylaima* have been reported to use several species of mollusks as intermediate




Figure 1. Metacercariae of *Brachylaima* sp. from the body cavity of *Phyllocaulis variegatus* (ventral view). Measurements are given in micrometers (μm).



Figure 2. Metacercariae of *Brachylaima* sp. from the body cavity of *Phyllocaulis variegatus* (ventral view). Measurements are given in micrometers (μm).

hosts in other regions, including snails of gastronomic interest such as *Otala punctata*, *Helix aspersa*, and *Theba pisana*. In Australia, *Brachylaima cribbi* Cribb, 1990, a parasite of *T. pisana*, was reported to cause human infections (Butcher, 2003; Gracenea & González-Moreno, 2002).

There are a number of previous records of adults of *Brachylaima* from birds and mammals in Argentina (Lunaschi & Drago, 2012; Martínez, 1986). *Brachylaima migrans* Dujardin, 1845 was found in the white-eared opossum *Didelphis albiventris* (Mammalia: Marsupialia) in Corrientes, Chaco, and Formosa Provinces. This mammal has frugivorous-omnivorous habits, although invertebrates (e.g., snails, slugs, worms, and arthropods) constitute an important part of its diet, it has been speculated that snails are involved as intermediate hosts (Santa Cruz, 2006). Particularly in Misiones Province, adult trematodes of the genus *Brachylaima* were observed parasitizing the gut of *Didelphis aurita* (Mammalia: Marsupialia) (GN, pers. comm.). Future experimental or molecular studies may corroborate the identity of the specimens reported here and elucidate their life cycle.

The authors wish to thank Rodrigo Santin for his assistance in the fieldwork, and Kabe Solas and Luis Giambelluca (CEPAVE) for their help with the drawings and photographs, respectively. We specially thank John Mike Kinsella for the English revision of the manuscript. This work was partially supported by UNLP N628 and UNLP N751.

References

- Barger, M. A., & Hnida, J. A. (2008). Survey of trematodes from terrestrial gastropods and small mammals in southeastern Nebraska. *Comparative Parasitology*, 75, 308–314.
- Barger, M. A. (2011). Microhabitat use of two brachylaimid trematodes in terrestrial gastropod second intermediate hosts. *Comparative Parasitology*, 78, 185–187.
- Butcher, A. R. (2003). *Brachylaima cribbi* n. sp. (Digenea: Brachylaimidae): Taxonomy, life-cycle kinetics and infections in animal and humans (Thesis). Australia: Faculty of Science School of Molecular and Biomedical Science, University of Adelaide.
- Butcher, A. R., Brealey, J. K., Grove, D. I., & Dymock, R. B. (2002). *Brachylaima cribbi* (Digenea: Brachylaimidae): scanning electronic microscopical observations of life-cycle stages. *Journal of Helminthology*, 76, 207–215.
- Butcher, A. R., & Grove, D. I. (2001). Description of the life-cycle stages of *Brachylaima cribbi* n. sp. (Digenea: Brachylaimidae) derived from eggs recovered from human feces in Australia. *Systematic Parasitology*, 49, 211–221.
- Butcher, A. R., Parasuramar, P., Thompson, C. S., & Grove, D. I. (1998). First report of the isolation of an adult worm of the genus *Brachylaima* (Digenea: Brachylaimidae), from the gastrointestinal tract of a human. *International Journal for Parasitology*, 28, 607–610.
- Butcher, A. R., Talbot, G. A., Norton, R. E., Kirk, M. D., Cribb, T. H., Forsyth, J. R. L., et al. (1996). Locally acquired *Brachylaima* sp. (Digenea: Brachylaimidae) intestinal fluke infection in two south Australian infants. *Medical Journal of Australia*, 164, 475–478.
- Cribb, T. H. (1990). Introduction of *Brachylaima* species (Digenea: Brachylaimidae) to Australia. *International Journal for Parasitology*, 20, 789–796.
- Cuezzo, M. A. (2013). Mollusca: Gastropoda. In E. Domínguez, & H. Fernández (Eds.), *Macroinvertebrados bentónicos sudamericanos* (pp. 525–629). Buenos Aires: Fundación Miguel Lillo.
- Gibson, D. I., Jones, A., & Bray, R. A. (2001). *Key to the trematoda* (Vol. 1) Wallingford: CABI Publishing.
- Gracenea, M., & González-Moreno, O. (2002). Life cycle of *Brachylaima mascomai* n. sp. (Trematoda: Brachylaimidae), a parasite of rats in the Llobregat Delta (Spain). *Journal of Parasitology*, 88, 124–133.
- Guisantes, J. A., Benito, A., Estibalez, J. J. S., & Mas-Coma, S. (1994). High parasite burdens by *Brachylaima* (*Brachylaima*) sp. (Trematoda: Brachylaimidae) in two dogs in the north of Spain. *Research and Reviews in Parasitology*, 54, 255–258.
- Lunaschi, L. I., & Drago, F. B. (2012). Digenean parasites of *Cariama cristata* (Aves, Gruiformes) from Formosa Province, Argentina, with the description of a new species of the genus *Strigea*. *Acta Parasitologica*, 57, 26–33.
- Martínez, F. A. (1986). Helmintofauna de los mamíferos silvestres. Trematodes. *Veterinaria Argentina*, 3, 544–551.
- Santa Cruz, A. C. M. (2006). *Ecto y endoparásitos de Didelphis albiventris Temmnick, del NEA (Marsupialia, Didelphidae)* Thesis. Corrientes, Argentina: Facultad de Ciencias Veterinarias, Universidad Nacional del Nordeste.
- Timon-David, J. (1959). Recherches sur les kystes et *Brachylaemus* du cyclostome. *Annales de Parasitologie Humaine et Comparée*, 34, 271–287.
- Thiengo, S. C., & Amato, S. B. (1995). *Phyllocaulis variegatus* (Mollusca: Veronicellidae), a new intermediate host for *Brachylaima* sp. (Digenea: Brachylaimidae). *Memorias do Instituto Oswaldo Cruz*, 90, 621–622.
- Yamaguti, S. (1975). *A synoptical review of life histories of digenetic trematodes of vertebrates*. Tokyo: Keigau Publishing.