



IP 80

ENG

Agenda Item: ATCM 15, CEP
7a
Presented by: SCAR
Original: English
Submitted: 21/04/2017

Antarctic Climate Change and the Environment – 2017 Update

Antarctic Climate Change and the Environment – 2017 Update

Information Paper submitted by SCAR

Summary

This paper presents an update on the Antarctic Climate Change and the Environment Report (2009 initial publication and 2013 update), providing an overview of recent science. The update is not meant to be read as a synthesis report, but as a perspective on recent scientific advances.

Introduction

Here we provide an update on recent significant advances in our understanding of climate change across the Antarctic continent and the Southern Ocean, and the impacts on the terrestrial and marine biota. This document builds on the material included in the Antarctic Climate Change and the Environment (ACCE) report, which was published by SCAR in 2009 [Turner *et al.*, 2009], with an update of the key points appearing in 2013 [Turner *et al.*, 2014]. At the request of the ATCM, SCAR agreed to provide regular updates on the original report (e.g. ATCM Resolution 4 (2010)). That activity is coordinated by the SCAR ACCE Expert Group (see <http://www.scar.org/ssg/physical-sciences/acce>), which has provided annual updates to the ATCM. The scope of the group is to keep abreast of recent advances in climate science, with a particular focus on Antarctic climate change and the biological implications of such changes. A recent development has been that the original ACCE report and the updated key points have been made available online as a wiki at http://acce.scar.org/wiki/Antarctic_Climate_Change_and_the_Environment. This online version is being progressively updated by a number of editors with input from many active scientists.

Changes in the Antarctic physical environment

1. In contrast to the Arctic, since the late 1970s the extent of sea ice around Antarctica has been increasing slightly, with record maximum late winter extents observed in 2012, 2013 and 2014. However, on 1 March 2017 Antarctic late summer sea ice extent dropped to just over 2 million square kilometres, which is the smallest amount yet observed since 1979 (Figure 1). The amount of sea ice was particularly low around the coast of West Antarctica. This is an example of the large variability of the Antarctic climate system on annual and decadal timescales. At present, it is difficult to reliably quantify any contribution that increasing greenhouse gas concentrations may have made to the record minimum and whether this represents a more general switch from increasing to decreasing sea ice extent. For more details on the record minimum see www.nsidc.org.
2. The Antarctic Circumpolar Current rings the Antarctic and is the largest wind-driven ocean current on Earth and the only ocean current to travel all the way around the planet. Now researchers have found that the current transports 30% more water than previously thought. The revised estimate is an important update for scientists studying how the world's oceans will respond to a warming climate [Donohue *et al.*, 2016].
3. Since the 1950s the Antarctic Peninsula has experienced a well-publicised warming, which was larger than at any other location in the Southern Hemisphere. However, it is now clear that the warming trend stopped in the late 1990s and that since that time temperatures have been decreasing, especially during the austral summer. The cooling has occurred because of a greater frequency of cold, easterly winds arriving at the Peninsula from the ice-covered Weddell Sea, arising from the development of a

climatological centre of low pressure between the Peninsula and South America, and once again highlighting the large variability of the Antarctic climate [Turner *et al.*, 2016]. Nevertheless, Peninsula temperatures are still higher than in the 1950s and glacier retreat is still ongoing.




Figure 1. The annual sea ice minimum since 1979.

4. Further indications of improvement in stratospheric ozone levels above Antarctica in spring and summer are emerging, although meteorological factors continue to significantly influence the year-to-year depth and size of the ozone hole [Solomon *et al.*, 2016]. The 2016 Antarctic ozone hole was a little smaller than that observed during 2014 and 2015 and was present from early August to mid-November; an earlier finish than in either of the last two years. It's difficult to predict exactly when there will cease to be a springtime ozone hole, but estimates suggest this will be sometime during the second half of this century.
5. Coupled atmosphere-ocean-sea ice models are the main tool we have to predict how the climate of the Antarctic will evolve over the coming decades. Sea ice is one of the most difficult aspects of the environment to incorporate correctly in these models, as it is thin and susceptible to changes in both atmospheric and oceanic conditions. Recently it has been shown that predictions of temperature and precipitation are strongly associated with how these models represent sea ice in their runs over the period since the late 1970s indicating that it is essential to improve the representation of sea ice in climate models [Bracegirdle *et al.*, 2015].
6. Ocean temperatures around the Antarctic continent have increased more rapidly and to greater depth than the global average in recent decades [Roemmich *et al.*, 2015]. Between 500 and 2,000 m warming averaged 0.002 °C/year with a broad intermediate-depth maximum between 700 and 1,400 m. Most of the heat gain (67 to 98%) occurred in the Southern Hemisphere extratropical ocean.
7. West Antarctic glaciers continue to retreat. Konrad *et al.* [2017] used data from five separate satellite missions spanning 1992-2015 to show that the region's glaciers thinned extensively during this period. Thinning was not uniform, but spread upstream along the centre of the Pine Island and Thwaites glaciers twice as fast (13-15 km/yr) as it did elsewhere in the region. The thinning has spread up the Thwaites and Pine Island ice streams into the interior of West Antarctica, likely contributing to an accelerating Antarctic contribution to global sea level rise [Forsberg *et al.*, 2017]. The current thinning episode began in about 1980 in the Pope-Smith-Kohler basin, about 1990 in the Pine Island glacier, and about 2004 on the Thwaites Glacier.

8. A paper by Smith et al. [2017], analysed sediment cores from the seabed beneath the Pine Island ice shelf and suggested that ice shelf retreat began there about 1945, triggered by a period of strong warming of West Antarctica that was associated with El Niño activity in the tropical Pacific. Atmosphere-ocean-ice interactions within the whole Amundsen Sea Embayment have recently been reviewed by Turner et al [2017] who noted that the linkages were not well represented in the current generation of climate models.
9. The water chemistry of the Southern Ocean appears to be changing at a faster rate than previously estimated, particularly in the deep ocean layers. In the cold Southern Ocean, CO₂ is being absorbed at a higher rate than in subtropical waters [Hauri et al., 2016].
10. As a contribution to the SCAR Antarctic Climate 2100 (AntClim21) programme a paper has recently been published that suggests a framework for the development of analogues for understanding past, present and future climates for the Antarctic and Southern Hemisphere [Mayewski et al., 2017], facilitating improved predictions of climate change across the regions. Several future climate scenarios, derived using multiple climate reanalysis data sets, were presented considering the impacts of warming, ozone depletion and changes in the zonal and meridional winds.

Changes in the Antarctic biological environment

1. Climate change is thought to have influenced Southern right whale breeding success in southern Brazil by causing variation in food (krill) availability for the species. Under these circumstances it seems likely that an increase in the frequency of years with reduced krill abundance around South Georgia, may reduce the current rate of recovery of southern right whales from historical overexploitation [Seyboth et al., 2016].
2. A boreal sea-star (*Asterias amurensis*) (Figure 2) that has been transferred to the Southern Hemisphere, most likely in ballast water, has established extensive invasive populations in southern Australia. It is a potentially high-risk invader of the sub-Antarctic and Antarctic, where it could alter community structure and ecosystem functioning because it is a benthic keystone predator [Byrne et al., 2016].



Figure 2. The sea-star *Asterias amurensis*.

3. Fast ice expansion, and the associated increase in the distance Adélie penguins have to forage to find food, have been linked to declines in size and breeding success (Wilson et al., 2016). Another recent

study showed that declining Adélie penguin populations experienced more years with warm sea surface temperature in their foraging areas compared to Adélie penguin populations that are increasing. Based on this relationship, it has been projected that one-third of current Adélie penguin colonies, representing ~20% of their current population, may be in decline by 2060 as Antarctic seas continue to warm. However, climate model projections suggest refugia may exist in continental Antarctica beyond 2099, buffering distribution-wide declines. Climate change impacts on penguins in the Antarctic will likely be highly site-specific based on regional climate trends, but a southward contraction in the range of Adélie penguins is expected over the next century [Cimino *et al.*, 2016].

4. Several recent studies show that response to climate change is highly species specific. This has important implications not only for projecting ecosystem response to climate change, but also for addressing conservation issues. A consequence of these findings is that the tolerance limits to environmental changes of all important key species must be quantified before assessments (up-scaling) of large scale biodiversity shifts and whole ecosystem changes can be made. Recent studies that highlight species responses to climate change include demersal (bottom feeding) fish [Sandersfeld *et al.*, 2017], various marine invertebrates [Clark *et al.*, 2017], amphipods [Schram *et al.*, 2016], sea birds [Grecian *et al.*, 2016]), top predators [Younger *et al.*, 2016], lichens [Bokhorst *et al.*, 2016] and mosses [Ashcroft *et al.*, 2016]
5. A major recent synthesis has assessed the overall levels of protection of Antarctic biodiversity in comparison with other regions of the planet [Chown *et al.*, 2017] concluding that much scope exists for improving the situation. From a protected area perspective, including in response to climate change, readily applicable approaches exist to give effect to such improvements [Coetzee *et al.*, 2017; Hughes & Grant, 2017].
6. The population of wandering albatrosses (*Diomedea exulans*) at South Georgia is decreasing because of bycatch in longline fisheries. Breeding females are at higher risk than males from all the main pelagic longline fleets in the south-west Atlantic. The results have important implications for the management of longline fisheries and the conservation of this highly threatened albatross population [Jiménez *et al.*, 2016].

The SCAR ACCE Expert Group consists of John Turner (Chair), Colin Summerhayes, Tom Bracegirdle, Daniella Liggett, Jenny Baeseman, Paul Mayewski, Pete Convey, Guido di Prisco, Julian Gutt, Dominic Hodgson, Tony Worby, Zhaomin Wang and Alexander Klepikov.

References

- Ashcroft, M.B., *et al.* (2016) Bayesian methods for comparing species physiological and ecological response curve. *Ecological Informatics* **34**, 35-43.
- Bokhorst, S., *et al.* (2016) *Usnea antarctica*, an important Antarctic lichen, is vulnerable to aspects of regional environmental change. *Polar Biology* **39**, 511-21.
- Bracegirdle, T.J., *et al.* (2015) The importance of sea-ice extent biases in 21st century multi-model projections of Antarctic temperature and precipitation. *Geophysical Research Letters* **42**, 10832-10839 doi: 10.1002/2015GL067055.
- Byrne, M., *et al.* (2016) From pole to pole: the potential for the Arctic seastar *Asterias amurensis* to invade a warming Southern Ocean. *Global Change Biology* **22**, 3874-3887.
- Cimino, M.A., (2016) Projected asymmetric response of Adélie penguins to Antarctic climate change. *Scientific Reports* **6**, 28785, doi: 10.1038/srep28785.
- Chown, S.L., *et al.* (2017) Antarctica and the strategic plan for biodiversity. *PLoS Biology* **15**, e2001656. doi: 10.1371/journal.pbio.2001656.
- Clark, M., *et al.* (2017) Biodiversity in marine invertebrate responses to acute warming revealed by a comparative multi-omics approach. *Global Change Biology* **23**, 318-330.

- Coetsee, B.W.T., Convey, P. & Chown, S.L. (2017) Expanding the protected area network in Antarctica is urgent and readily achievable. *Conservation Letters* doi: 10.1111/conl.12342.
- Donohue, K.A., et al. (2016) Mean Antarctic Circumpolar Current transport measured in Drake Passage. *Geophysical Research Letters* **43**, 11760-11767.
- Forsberg, R., Sørensen, L. & Simonsen, S. (2017) Greenland and Antarctica ice sheet mass changes and effects on global sea level. *Surveys in Geophysics* **38**, 89-104.
- Grecian, W.J., et al. (2016) Contrasting migratory responses of two closely related seabirds to long-term climate change. *Marine Ecology-Progress Series* **559**, 231-242.
- Hauri, C., Friedrich, T. & Timmermann, A. (2016) Abrupt onset and prolongation of aragonite undersaturation events in the Southern Ocean. *Nature Climate Change* **6**, 172-176.
- Hughes, K.A. & Grant, S.M. (2017) The spatial distribution of Antarctica's protected areas: A product of pragmatism, geopolitics or conservation need? *Environmental Science and Policy* **72**, 41-51.
- Jiménez, S., et al. (2016) Sex-related variation in the vulnerability of wandering albatrosses to pelagic longline fisheries. *Animal Conservation* **19**, 281-295.
- Konrad, H., et al. (2017) Uneven onset and pace of ice-dynamic imbalance in the Amundsen Sea Embayment, West Antarctica, *Geophysical Research Letters* doi: 10.1002/2016GL070733.
- Mayewski, P.A., et al. (2017) Ice core and climate reanalysis analogs to predict Antarctic and Southern Hemisphere climate changes, *Quaternary Science Reviews* **155**, 50-66.
- Roemmich, D. et al. (2015) Unabated planetary warming and its ocean structure since 2006. *Nature Climate Change* **5**, 240-245.
- Sandersfeld, T., Mark, F.C. & Knust, R. (2017) Temperature-dependent metabolism in Antarctic fish: do habitat temperature conditions affect thermal tolerance ranges? *Polar Biology*, **40**, 141-149.
- Schram, J.B., et al. (2016) Seawater acidification more than warming presents a challenge for two Antarctic macroalgal-associated amphipods. *Marine Ecology-Progress Series* **554**, 81-97.
- Seyboth, E., et al. (2016) Southern right whale (*Eubalaena australis*) reproductive success is influenced by krill (*Euphausia superba*) density and climate. *Scientific Reports* **6**, 28205 doi: 10.1038/srep28205.
- Smith, J.A. et al. (2017) Sub-ice-shelf sediments record history of twentieth century retreat of Pine Island Glacier. *Nature* **541**, 77-80.
- Solomon, S., et al. (2016) Emergence of healing in the Antarctic ozone layer. *Science*, DOI: 10.1126/science.aae0061.
- Turner, J., et al. (2014) Antarctic Climate Change and the Environment - An Update, *Polar Record* **50**, 237-259. doi:10.1017/S0032247413000296.
- Turner, J., et al. (2009) *Antarctic Climate Change and the Environment*. Scientific Committee on Antarctic Research, Cambridge, 526 pp.
- Turner, J. et al. (2016) Absence of 21st century warming on Antarctic Peninsula consistent with natural variability. *Nature* **535**, 411-415.
- Turner, J. et al. (2017) Atmosphere-Ocean-Ice Interactions in the Amundsen Sea Embayment, West Antarctica. *Reviews of Geophysics* **55**, 235-276. doi: 10.1002/2016RG000532.
- Wilson, K.J., et al. (2016) The impact of the giant iceberg B09B on population size and breeding success of Adélie penguins in Commonwealth Bay, Antarctica. *Antarctic Science* **28**, 187-193.
- Younger, J.L., et al. (2016) The influence of historical climate changes on Southern Ocean marine predator populations: a comparative analysis. *Global Change Biology* **22**, 474-493.