

## Male condom use, multiple sexual partners and HIV: a prospective case-control study in Kinshasa (DRC)

Silvia Carlos, PharmD, PhD<sup>a,b,c</sup>  
Cristina López del Burgo, MD, PhD<sup>a,b,c</sup>  
Eduardo Burgueño, MD, PhD<sup>d,e</sup>  
Miguel Ángel Martínez-González, MD, PhD<sup>a,b,f</sup>  
Alfonso Osorio, PhD<sup>b,c,g</sup>  
Adolphe Ndarabu<sup>h</sup>  
Clément Passabosc<sup>i</sup>  
Jokin de Irala, MD, PhD, MPH<sup>a,b,c</sup>

<sup>a</sup>Department of Preventive Medicine and Public Health, School of Medicine, University of Navarra, Pamplona, Spain.

<sup>b</sup>IdisNA Navarra Institute for Health Research, Pamplona, Spain.

<sup>c</sup>Institute for Culture and Society (ICS), Education of Affectivity and Human Sexuality, University of Navarra, Pamplona, Spain.

<sup>d</sup>CEFA-Monkole, Kinshasa, Democratic Republic of the Congo.

<sup>e</sup>Department of Family Medicine and Primary Health Care, UPC-Protestant University in Congo, Kinshasa, Democratic Republic of the Congo.

<sup>f</sup>CIBER Fisiopatología de la Obesidad y Nutrición (CIBER obn), Spanish Government (ISCIII), Madrid, Spain.

<sup>g</sup>School of Education and Psychology, University of Navarra, Pamplona, Spain.

<sup>h</sup>Monkole Hospital, Kinshasa, Democratic Republic of the Congo.

<sup>i</sup>Department of Ophthalmology, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain.

\* Corresponding author:

Silvia Carlos

Department of Preventive Medicine and Public Health, School of Medicine, University of Navarra. Irunlarrea 1, 31008 Pamplona, Navarra, Spain

Tel: (+34) 948 42 56 00 ext.806636 Fax: (+34) 948 42 56 49

E-mail address: [scarlos@unav.es](mailto:scarlos@unav.es)

### Published in:

AIDS Care

<http://www.tandfonline.com/loi/caic20>

### Cite as:

Carlos S, López-del Burgo C, Burgueño E, Martínez-González MA, Osorio A, Ndarabu A, Passabosc C, de Irala J. Male condom use, multiple sexual partners and HIV: a prospective case-control study in Kinshasa (DRC). *AIDS Care* 2016; DOI: 10.1080/09540121.2016.1258450.

**ABSTRACT**

In the Democratic Republic of Congo no previous studies have assessed the factors associated with different patterns of condom use and with multiple sexual partners and the association between condom use simultaneously taking into account multiple sexual partnerships, and HIV infection. We carried out a prospective case-control study. From December 2010 until June 2012, 1,630 participants aged 15-49 getting HIV Voluntary Counseling and Testing in a hospital in Kinshasa were selected. Cases were new HIV diagnosis and controls were HIV-negative participants detected along the study period. We recruited 274 cases and 1,340 controls that were interviewed about HIV-related knowledge, attitudes and behaviours. Among cases there was a high prevalence of multiple lifetime and concurrent sexual partnerships (89.8% and 20.4%, respectively) and most cases never used condoms with only 1.5% using them consistently. Condom use and multiple partnerships were associated with male, single and high-educated participants. An association was found between multiple lifetime partners and 'any condom use' (OR=2.99; 95%CI: 2.14-4.19) but not with consistent use. Both having two or more multiple concurrent sexual partners or not using condoms were variables similarly and highly associated to HIV risk. The association found between having two or more concurrent sexual partners and HIV was slightly higher (OR=3.58, 95%CI:2.31-5.56) than the association found between never condom use and HIV (OR=3.38, 95%CI:1.15-9.93). We found a high prevalence of multiple lifetime sexual partners and an extremely high prevalence of inconsistent condom use, both strongly associated with HIV seropositivity. Local programs would benefit from comprehensive interventions targeting all behavioural and sociocultural determinants.

**KEY WORDS:**

HIV, condom, multiple sexual partners, behaviour, Congo.

## INTRODUCTION

In addition to the new biomedical strategies for HIV prevention that are currently being developed (Marrazzo et al., 2014), the underlying sexual behaviours responsible for most infections are still critical to prevent new transmissions. Thus, HIV prevention programmes must address the avoidance of early sex, the reduction of sexual partners and the correct and consistent condom use among people at risk (Centers for Disease Control and Prevention, 2014; Halperin et al., 2004).

Since the first condom report in 1563 (Fallopian, 1563), its effectiveness in preventing sexually transmitted infections (STIs) has been widely analysed (Centers for Disease Control and Prevention, 2014; Holmes, Levine, & Weaver, 2004; National Institutes of Health, 2001; Vera, Orozco, Soto, & Aburto, 2008; Winer et al., 2006). HIV is the STI with stronger evidence supporting the preventive role of condoms (Centers for Disease Control and Prevention, 2014; Fitch et al., 2002). Right after HIV discovery, condom effectiveness in reducing the HIV risk was evaluated (Conant, Hardy, Sernatinger, Spicer, & Levy, 1986; Davis & Weller, 1999; Judson, Ehret, Bodin, Levin, & Rietmeijer, 1989; Pinkerton & Abramson, 1997; Rietmeijer, Krebs, Feorino, & Judson, 1988; Van de Perre, Jacobs, & Sprecher-Goldberger, 1987). Regarding heterosexual transmission, a 60%-96% relative risk reduction has been estimated (Giannou et al., 2015; Lassi, Imam, Dean, & Bhutta, 2014; Liu et al., 2014; UNAIDS, 2004; Weller & Davis, 2002). A meta-analysis carried out by Giannou *et al.*, 2015 evaluated HIV infection RRs after comparing “always” *versus* “never”, and “always” *versus* “not always” use and estimated that, consistent use provide an approximately 70% relative reduction in HIV risk.

Although most HIV infections occur in sub-Saharan Africa (SSA), local research has been mainly focused on condom availability, use patterns and determinants and barriers of use. Less research has assessed condom effectiveness in Africa (Ahmed et al., 2001; Eaton, & Hoesley, 2014; Hearst, Ruark, Hudes, Goldsmith, & Green, 2013; Hughes et al., 2012;

Medley, Bachanas, Grillo, Hasen, & Amanyeiwe, 2015; Mindel, & Sawleshwarkar, 2013; Peltzer, 2012;). A study carried out at 14 sites in Eastern/Southern Africa, in which serodiscordant couples received free condoms, found a per-act-risk relative reduction of 78% (Hughes et al., 2012). In the Giannou *et al.* review only four out of the twenty-five studies were African (Allen et al., 1992; Hira et al., 1997; Hughes et al., 2012; Kamenga et al., 1991). In all four studies, free condoms were distributed and as much as a 77% relative reduction in HIV risk was observed for consistent use (Giannou et al., 2015).

To achieve the maximum protective effect, condoms must be used consistently and correctly (Centers for Disease Control and Prevention, 2014; Food and Drug Administration; Workowski, & Bolan, 2015) but in SSA use is still low and mostly inconsistent (MacQuarrie, Bradley, Gemmill, & Staveteig, 2014; Mutevedzi, & Newell, 2014; Papo, Bauni, Sanders, Brocklehurst, & Jaffe, 2011; UNAIDS, 2014). Objections to condoms are widespread due to culture, religion misconceptions, stigma and other factors (Abdool Karim et al., 2012; Abdulai et al., 2012; Amo-Adjei, 2012; Carlos et al., 2015; Earnshaw et al., 2014; Exavery et al., 2011; Exavery et al., 2012; Hawkes et al., 2013; Hearst et al., 2013; Kajubi et al., 2005; Mantell et al., 2011; Maticka-Tyndale, 2012; Namisi et al., 2015; Padian, Buvé, Balkus, Serwadda, & Cates, 2008; Rwege, 2013; Skovdal, Campbell, Nyamukapa, & Gregson, 2011; Ukwuani, Tsui, & Suchindran, 2003; UNAIDS, 2004; UNAIDS, 2014b). Moreover, programs that address and provide free condoms, do not always give instructions on their correct use, or about breakage and slippage that can occur (Grasso, et al., 2016; Remis, Alary, Liu, Kaul, & Palmer, 2014; Sanders, et al., 2012; Spaar, et al., 2010; Walsh, et al., 2004).

To develop effective preventive programmes, epidemiological and behavioural evidence within each country in SSA is necessary (Abdool Karim et al., 2012; Kamwi, Kenyon, & Newton, 2006; Ogden, Gupta, Warner, & Fisher, 2011).

The Democratic Republic of Congo (DRC) has an HIV prevalence of 1.2% in the 15-49 aged general population, and of 2.6% in women and 0.3% in men in Kinshasa (Demographic

Health Survey, 2014). No studies have simultaneously evaluated on the general population condom use for HIV prevention also taking into account multiple sexual partnerships. In 1991 (when no antiretrovirals nor Voluntary Counseling and Testing-VCT-services were available in Kinshasa), a study evaluated among 149 serodiscordant couples the effect of a VCT program (which promoted condoms) and collected data on condom effectiveness (Kamenga et al., 1991). Despite an increase in abstinence and in condom use six months after, 6 couples (4%) reporting condom use became HIV concordant due to inconsistent and incorrect use. Subsequent studies in Congo have just shown information about condom availability (which is not nationally ensured) (Bosmans, Cikuru, Claeys, & Temmerman, 2006), condom use (Laga et al., 1994), determinants (Kayembe et al., 2008; Ntumbanzondo, Dubrow, Niccolai, Mwandagalirwa, & Merson, 2006) and barriers for using condoms (Irwin et al., 1991; Kabamba Mulongo, Schirvel, Mukalay Wa Mukalay, & Dramaix Wilmet, 2011; Kayembe, Fatuma, Mapatano, & Mambu, 2006). A recent cross-sectional study among people attending VCT in a rural area found a very low prevalence of consistent use and a non-significant association, in a bivariate analysis, between any condom use and HIV (Kautako-Kiambi, Ekila, Kama-Lemba, Wumba, & Aloni, 2015).

We analyse in a population receiving VCT at a hospital in Kinshasa: 1) the association of sociodemographic characteristics with condom use and consistent use; 2) with multiple sexual partnerships; and 3) the association between condom use, simultaneously taking into account multiple sexual partnerships, and HIV infection.

## **METHODS**

### ***Study design***

Case-control study that evaluates the knowledge, attitudes and sexual behaviours associated with HIV.

### ***Setting***

Outpatient Centre at Monkole referral hospital in Kinshasa that provides VCT and treatment, and also screens blood donors for HIV.

### ***Study participants***

People aged 15-49 attending Monkole to receive VCT or as blood donors to whom VCT was offered. Those never HIV tested or with a previous HIV negative result were accepted.

Cases and controls were defined based on the result of the HIV test. Along the study period (from December 2010 until June 2012) local researchers selected incident HIV+ cases as well as new diagnosed HIV- controls.

### ***Sample size***

When the study was designed, about 1,000 people aged 15-49 annually used Monkole as VCT attendees or blood donors, with around 15% newly HIV+.

Considering a 90% response rate, we estimated that during 18 months about 1,350 people could participate, with around 200 new HIV+ diagnosis (Grimes, & Schulz, 2005). In logistic regressions about 10 events are needed for each parameter in the model (Hosmer, & Lemeshow, 2013), so we were confident to simultaneously adjust for around 20 parameters.

### ***Data collection***

We collected information and compared lifetime exposures that occurred prior to the classification of our participants as cases or controls.

The interviewer-administered questionnaire (available in French and Lingala) was composed of closed questions and few open-ended questions and included 59 items on: sociodemographics, health-related aspects, perceived risk, HIV knowledge, information, attitudes, stigma and sexual behaviours. For this specific paper we evaluated the sociodemographic and behavioural data.

### ***Laboratory analyses***

A serum sample was analysed by the Determine®HIV-1/2 test. If positive, it was analysed by two other rapid tests: DoubleCheckGold® and Unigold®. A confirmatory test was not available and 'undetermined' tests were excluded from the analyses.

### ***Statistical analysis***

Initially we carried out a descriptive analysis of our participants' sociodemographic and behavioural characteristics. Concerning sexual behaviours we first evaluated their use of condoms, which was collected by asking 'Have you ever used a condom?'. The different answer options to this question were: 'never', 'in some sexual relationships', 'in almost all sexual relationships' or 'always, in every sexual relationship'. We also recategorized this variable into 1) 'any' versus 'never'; 2) 'a more consistent use' (almost always/always) versus 'never/sometimes'; and 3) 'consistent use' (always) versus 'never/sometimes/almost always'. With regards to multiple sexual partnerships, the questionnaire included different questions for concurrent sexual partners ('Do you actually have multiple sexual partners' (no/yes), 'how many partners do you have?') and lifetime sexual partners ('how many lifetime partners have you had?'). The number of sexual partners was categorized as a dichotomous categorical variable (multiple sexual partners, yes/no).

We evaluated the association between sociodemographic characteristics and condom use and multiple sexual partnerships and additionally, we evaluated the association between sexual behaviours and HIV simultaneously taking into account both condom use and multiple sexual partnerships.

Multivariate unconditional logistic regressions were carried out with Stata 12.1.

### ***Ethical issues***

The study was approved by the Ethics Committee of the University of Kinshasa and the University of Navarra.

Oral informed consent was obtained, consistent with local clinical practices (Wembonyama, Mpaka, & Tshilolo, 2007; Obermeyer, et al., 2012).

## RESULTS

From December 2010 until June 2012, 274 HIV-positive cases and 1,340 controls were identified. The response rate was 99.9%.

Sixteen HIV-undetermined participants and 59 who had never had sex, were excluded.

**Table 1** shows that overall, our participants were mainly adults (25-49 year old), with a low/medium socioeconomic level and a secondary education level. Among cases the percentage of VCT attendees, women, adults, illiterate and widowed, separated or cohabitating was significantly higher than among controls. Regarding sexual behaviours, among cases the prevalence of multiple sexual partners (lifetime and concurrent) and of never condom use was higher than among controls. Thirty-nine percent of participants never used condoms and only 5.7% of participants reported consistent condom use, which was less frequent for cases than for controls (1.5% and 6.5%;  $p=0.001$ ).

With regards to the specific differences between the study subgroups (data not shown), among VCT attendees there were significantly more women than among blood donors (54.5% vs. 11.6%,  $p<0.001$ ) and a higher percentage of people of a high economic (11.1% vs. 4.2%,  $p<0.001$ ) and education level (21.1% vs. 11.5%,  $p<0.001$ ), divorced (5.3% vs. 0.9%,  $p<0.001$ ) and widowed people (4.9% vs. 0.1%,  $p<0.001$ ). Concerning sexual behaviours, among VCT attendees the percentage of people with multiple concurrent partners was significantly higher than among controls (12.7% vs. 7.7%,  $p=0.001$ ) but there were no significant differences regarding the prevalence of multiple lifetime sexual partners (85.4% vs. 83.8%,  $p=0.382$ ) and of inconsistent condom use (86.1% vs. 83.9%,  $p=0.235$ ).

When we analysed the characteristics of participants associated with condom use and with having multiple sexual partners (**Table 2**) we found that men, people with university studies,



single respondents and participants reporting multiple lifetime partners were more likely to report any condom use and a more consistent use. However, significance for the association between educational level or for reporting two or more lifetime partners and condom use was lost for consistent use. Having concurrent sexual partners was not significantly associated with condom use. Frequency of condom use decreased with age, although the magnitude of this inverse association was low.

VCT attendees, men, and adults were more likely to report multiple sexual partners. Reporting 'any' condom use was associated with having multiple lifetime partners compared to 'never' use.

Both having two or more multiple concurrent sexual partners and not using condoms were variables similarly and highly associated to HIV risk (**Table 3**). The association found between having more than 2 concurrent sexual partners and HIV was slightly higher (OR=3.58, 95%CI:2.31-5.56) than the association found between never condom use and HIV (OR=3.38, 95%CI:1.15-9.93).

The magnitude of the association with HIV positivity was higher for concurrent than for multiple lifetime sexual partners and with regards to inconsistent condom use, the association with HIV was stronger for never use than for sometimes use, compared to always use.

## **DISCUSSION**

To our knowledge, this is the first study in the DRC that has simultaneously analysed the prevalence of different condom use patterns and information on multiple sexual partnerships and related this information to newly diagnosed HIV status.

We found an overall prevalence of 61% of condom use, with HIV+ people using them much less frequently than the negative controls. In the 2013-2014 Demographic Health Survey from the DRC (Demographic Health Survey, 2014) when women were asked about contraception,

only 6.7% reported using male condoms in Kinshasa (where condoms are used more frequently than in rural areas). This was even lower than the 8.7% found in the previous 2007 national survey. In this survey, behavioural information concerning condoms was collected for the last sexual encounter and for people reporting two or more sexual partners in the last 12 months. In that particular high-risk situation, 20% of women and 28% men in Kinshasa reported condom use (14% and 31% in 2007). Among our study participants reporting multiple partnerships, the prevalence of 'any' condom use was close to 65%. We found higher prevalence than the official estimates, however, populations attending healthcare centres are more health-conscious (Wang, Aoumya, & Shanxiao, 2012).

The most important aspect for HIV prevention using condoms is not the overall use but the pattern of use, as inconsistent use can even lead to a higher STI risk (Centers for Disease Control and Prevention, 2014). In the 2013/2014 DRC survey, as in other African surveys, different condom use frequencies are not registered. A comparative publication from several SSA countries that did include information on different condom use patterns, showed that consistent use was rare: less than 1/6 women and 1/3 men reported any condom use at last sex, with consistent use being even lower (Mishra, Praween, Soumya, Yuan, & Shanxiao, 2009). Only 1.5% of our HIV+ cases and 6.5% of controls used condoms consistently. Thus, although there is evidence that in the DRC and other SSA countries condom use may have increased (UNAIDS, 2013), consistent use is still very low, particularly in high-risk sexual relationships.

We found, as many other African studies (Apondi et al., 2011; De Coninck, & Marrone, 2012; Hearst et al., 2013; Kabamba Mulongo et al., 2011; Maticka-Tyndale, 2012; Michielsen et al., 2010; Mishra, Praween, Soumya, Yuan, & Shanxiao, 2009; Oyeyemi, & Oyeyemi, 2012; Shewamene, Legesse, Tsega, Bhagavathula, & Endale, 2015; UNAIDS, 2013), that male, younger and single participants and those with a higher education were more likely to use

condoms. Participants with a higher lifetime number of sexual partners also reported a higher frequency of condom use.

Women in the DRC, as in most SSA countries, are particularly affected by HIV with an HIV incidence and prevalence higher than among men, especially among young women (1.2% and less than 0.1%, respectively) (Demographic Health Survey, 2014; Murray et al., 2014). However, condoms do not seem to be the magic bullet for reducing new infections among women. The DRC has a 75-100% dependency on international sources for HIV response (UNAIDS, 2013; UNAIDS, 2014a). A 2011 USAID audit found that only 30% of male condoms and 0% female condoms were efficiently distributed (USAID, 2011), with important delays in provision and promotion. This can be generalised for many African countries where the donor supply provides around one condom per every ten women per year compared to eight condoms for men. And more critical than this lack of supply, are the traditional gender norms (Exavery et al., 2012; Geary, & Bukusi, 2014; Karim et al., 2014; Mantell et al., 2011; Namisi et al., 2015; Padian et al., 2008). A recent UNAIDS gender-related report, described some women's barriers towards HIV prevention/treatment services, such as cultural beliefs, unequal power gender relations, stigma or mistrust (UNAIDS, 2014b). Thus, gender-related and cultural barriers need to be first overcome to improve the female adherence to any of the available prevention strategies, including condoms (Carlos et al., 2015; Green, 2011; Ogden et al., 2011).

We also found that younger participants were slightly more likely to report condom use than adults. At the 2014 DRC survey, around 30% urban young women and 40% men used condoms in their last sex, 18% and 40% when they had more than one sexual partner and 2% and 0.5% in HIV+ persons (Demographic Health Survey, 2014). When 2014 and 2007 surveys are compared, condom use has not changed much among young people (except for the HIV+ youth in which condom use has in fact decreased). Moreover, usually young people use condoms for contraception and not for HIV prevention (Maticka-Tyndale, 2012). In

addition, the age of first sex (critical for HIV risk), remained similar in 2007 and 2014, with around 12% and 17% single, 15-24 aged, women and men having sex before age 15, and 42% and 55% before age 18. Multiple sexual partnerships among young people have not been reduced since 2007, with around 3% women and 15% men reporting 2 or more sexual partners in the previous year. Since risk avoidance measures offer a higher protection to prevent new HIV infections (Halperin et al., 2004; Okware, Kinsman, Onyango, Opio, & Kaggwa, 2005) and considering that condom availability remains low in these settings, preventive campaigns and VCT information must keep insisting on the benefits of delaying sexual initiation among adolescents and of avoiding multiple sexual partnerships for those sexually active. With regards to adults, the majority of those over 50, even those with multiple partners, do not use condoms (UNAIDS, 2014a). Consequently, youth-friendly and older-age specific HIV information services might have to reconsider how to inform about the preventive strategies for each population group and to increase the adherence to such measures. For example, among young people the uptake and impact can be improved with age-specific training to providers, flexible clinic hours, mobile services or even considering adolescents under 18 years in research so that interventions can be well adapted to them (Dellar, Dlamini, & Karim, 2015).

Our single participants were more likely to report condom use and consistent use. Other SSA studies, including Congolese, have consistently shown that single are significantly more likely to use condoms than married or cohabiting, both for women and men, whether they are HIV+ or not (Ahmed et al., 2001; Anand et al., 2009; Ayiga, 2012; Exavery et al., 2012; Kabamba Mulongo et al., 2011; Reynolds et al., 2011).

Our highly-educated participants were nearly three times more likely to use condoms than participants with no studies or primary studies. The 2007 and 2014 DRC surveys showed that condom use was more frequent for men and women with a highest level of education than for those with no studies. A higher education level has been associated with a higher consistent

use and a lower educational level with not using condoms (Marshall et al., 2015; Pilapil et al., 2016). However, we found the highest prevalence of multiple partnerships in the highly educated, which clearly increases their risk (Hearst et al., 2013; Green, Mah, Ruark, & Hearst, 2009), despite the possible protection due to their education.

Apart from the mentioned socio-ecological factors prevention programs need to address all sexual risk behaviours. The last UNAIDS report found a significant increase in the number of sexual partners for some SSA countries. In Kinshasa the population reporting two or more sexual partners in the last year slightly decreased between 2007 and 2014 but we still found 10% and 88% of our participants reporting multiple concurrent or lifetime sexual partners, respectively. Men were more likely to report both multiple partnerships and since they are potential HIV transmitters to susceptible women, strategies to prevent these behaviours need to involve both sexes.

No matter the type of multiple sexual partnerships, this behaviour was significantly associated with HIV infection in our study, whether condoms were used or not. We also found an association between inconsistent condom use and HIV status, with a higher magnitude for those never using than those using in some sexual relations compared to always use. Consequently, if those with a higher risk are using condoms only in some occasions, the benefits of condom use can be greatly offset since sex with multiple partners is highly prevalent. Today most studies regarding condom use do not take into account that condoms are not always used consistently or that sexual relationships are not always mutually monogamous. Local HIV prevention programs need to therefore consider all these behavioural aspects, together with the underlying sociocultural and structural factors to have a real impact on the infection epidemic.

This study has some limitations. Firstly, it is based on self-report which can be associated with social desirability bias, and could lead to over-reporting condom use and under-reporting multiple partnerships (Hearst et al., 2013). Nevertheless, this bias would likely be present in

both cases and controls, resulting in a non-differential bias of our estimates towards the null. Secondly, only a very low percentage of our participants reported consistent condom use which makes it difficult to adequately analyse the specific predictors of consistent use and the association with HIV. Another constraint is that we did not have information on the correct or incorrect use of condoms. Nevertheless, even if we had considered that participants reporting condom use always used them correctly, the figures for risk reduction would be far away from the goal of optimal protection. Finally, we did not collect information about anal/oral sex and cannot rule out these routes. Today there are still no estimates about the prevalence of these routes among the general Congolese population. We are collecting this information in two prospective cohorts in Kinshasa and will have data on this in the future.

This study also has some strengths. Firstly, participants didn't know their HIV status at the time of the survey so test results could not influence their self-report. Secondly, we evaluated condom use using four categories and not dichotomously, which allows investigating the different use patterns that can be observed in real life (Devine, & Aral, 2004). Thirdly, participants did not receive any intervention pre-interview (i.e. condom counseling) and responses were therefore not influenced. Finally, as far as we know, this is the first study in Congo evaluating condom use patterns in participants with a new HIV diagnosis taking into account the concomitant effect of multiple partners. In future studies we will also consider the different types of sexual partners (i.e. casual, stable) in which condoms are or not used (Lagarde et al., 2001; Omori, & Abu-Raddad, 2016).

In summary, we found a very low prevalence of consistent condom use. Men, single and participants with a higher education were more likely to use them but also more likely to have multiple sexual partners. When these sexual behaviours were analysed altogether we found a significant association between multiple sexual partnerships and HIV infection, whether condoms were used or not. We found a stronger inverse association with HIV for consistent use than for inconsistent use compared to never use. However, the magnitude of the potential

protection of inconsistent use seemed offset by the potential riskier effect of multiple sexual partners. In consequence, local HIV prevention programs in Kinshasa might need to take into account all sexual behaviours from a more comprehensive and holistic approach. In addition, the underlying gender, age-related and sociocultural determinants might also have to be thoroughly addressed if an optimal HIV prevention is targeted.

**Disclosure statement**

No potential conflict of interest was reported by the authors.

**Funding**

This work was supported by University of Navarra [grant number P-2009-05] and received the Dolores Trigo Award from the Alumni Association of the Faculty of Pharmacy of Santiago de Compostela (Spain).

**Table 1. Sociodemographic and behavioural characteristics of cases and controls.**

	<b>Cases (n=274) %</b>	<b>Controls (n=1,340) %</b>	<b>p<sup>a</sup></b>
<b>Study subgroup</b>			
VCT	81.75	39.55	<0.001
Blood donors	18.25	60.45	
<b>Sex (women)</b>	52.92	27.31	<0.001
<b>Age (years)</b>			
15-24	13.50	36.19	<0.001
25-49	86.50	63.81	
<b>Socio-economic status</b>			
Low	47.08	32.54	<0.001
Medium	48.54	59.40	
High	4.38	8.06	
<b>Education</b>			
No studies/Illiterate	12.04	2.84	<0.001
Primary	9.12	2.39	
Secondary	67.52	77.84	
University	11.31	16.94	
<b>Marital status</b>			
Single	41.61	69.48	<0.001
Married	33.58	24.48	
Widowed	11.31	0.52	
Separated/Divorced	10.22	3.88	
Cohabitation	3.28	1.64	
<b>≥2 lifetime sexual partners</b>	89.78	83.51	0.009
<b>≥2 concurrent sexual partners</b>	20.44	7.91	<0.001
<b>Condom use</b>			
Never	61.31	36.72	<0.001
Sometimes	32.12	46.49	
Almost always	5.11	10.52	
Always	1.46	6.27	

<sup>a</sup> p-value obtained by Chi<sup>2</sup> test or Chi<sup>2</sup> for linear trend for ordinal variables.



**Table2. Characteristics of participants associated with condom use and with having multiple sexual partners.**

	Condom use			Multiple sexual partners	
	OR (95%CI) <sup>a,b</sup>			OR (95%CI) <sup>a,b</sup>	
	Any use (n=952)	Almost always/ Always (n=242)	Always (n=88)	Concurrent (n=162)	Lifetime (n=1,362)
<b>Study group (VCT compared to blood donors)</b>	0.58 (0.42-0.69)	-	-	2.80 (1.92-4.09)	1.69 (1.14-2.50)
<b>Sex (men compared to women)</b>	1.48 (1.13-1.93)	1.72 (1.18-2.51)	1.93 (1.05-3.55)	3.73 (2.30-6.03)	2.07 (1.40-3.05)
<b>Age (for each additional year)</b>	0.98 (0.96-0.99)	0.97 (0.95-0.99)	0.93 (0.89-0.97)	-	1.07 (1.04-1.10)
<b>Education (university compared to no studies/primary)</b>	3.14 (1.92-5.13)	2.91 (1.25-6.76)	-	-	-
<b>Marital status (single compared to others)</b>	1.85 (1.38-2.49)	3.12 (1.96-4.97)	2.63 (1.16-5.98)	-	-
<b>≥2 lifetime sexual partners (compared to 1)</b>	2.99 (2.14-4.19)	2.10 (1.24-3.56)	-	NA	NA
<b>≥2 concurrent sexual partners (compared to 1)</b>	-	-	-	NA	NA
<b>Condom use (any compared to never use)</b>	NA	NA	NA	-	2.91 (2.09-4.05)

<sup>a</sup> Adjusted for all the variables listed in the table.

<sup>b</sup> Only statistically significant associations (p<0.05) are shown.

**Table 3. ORs (95%CI) for having a positive HIV test for the different sexual behaviours.**

	OR (95%CI) <sup>a,b</sup>
<b>Multiple concurrent sexual partners</b>	
No concurrent sexual partners	1 (Ref.)
≥2 concurrent sexual partners	3.58 (2.31-5.56)
<b>Multiple lifetime sexual partners</b>	
1 lifetime sexual partners	1 (Ref.)
≥2 lifetime sexual partners	1.86 (1.07-3.23)
<b>Condom use</b>	
Never	3.38 (1.15-9.93)
Sometimes	1.99 (0.68-5.87)
Almost always	1.83 (0.55-6.13)
Always	1 (Ref.)

<sup>a</sup> Adjusted for sex, age, education, marital status, study group (VCT/blood donors), multiple concurrent sexual partners, multiple lifetime sexual partners and condom use.

<sup>b</sup> Interaction between having multiple lifetime sexual partners and condom use was evaluated (p=0.731).

**REFERENCES**

- Abdool Karim, Q, Kharsany, AB, Frohlich, JA, Werner, L, Mlotshwa, M, Madlala, BT, & Abdool Karim, SS. (2012). HIV incidence in young girls in KwaZulu-Natal, South Africa public health imperative for their inclusion in HIV biomedical intervention trials. *AIDS and Behavior*, 16, 1870-6. Retrieved from: PMID: 22618892
- Abdulai, M.A., Baiden, F., Adjei, G., Afari-Asiedu, S., Adjei, K., Tawiah, C., & Newton, S. (2012). An assessment of the likely acceptability of vaginal microbicides for HIV prevention among women in rural Ghana. *BMC Women's Health*, 12, 40. doi: 10.1186/1472-6874-12-40.
- Ahmed, S., Lutalo, T., Wawer, M., Serwadda, D., Sewankambo, N.K., Nalugoda, F...Gray, R. (2001). HIV incidence and sexually transmitted disease prevalence associated with condom use: a population study in Rakai, Uganda. *AIDS*, 15:16 2171-2179. Retrieved from: PMID: 11684937
- Allen, S., Tice, J., Van de Perre, P., Serufulira, A., Hudes, E., Nsengumuremyi, F...Hulley, S. (1992). Effect of serotesting with counselling on condom use and seroconversion among HIV discordant couples in Africa. *BMJ (Clinical research ed.)*, 304,1605-9. Retrieved from: PMID: 1628088
- Amo-Adjei, J. (2012). Age differences and protected first heterosexual intercourse in Ghana. *African Journal of Reproductive Health*, 16, 58-67. Retrieved from: PMID: 23444544
- Anand, A., Shiraishi, R.W., Bunnell, R.E., Jacobs, K., Solehdin, N., Abdul-Quader, A.S...Diaz, T. (2009). Knowledge of HIV status, sexual risk behaviors and contraceptive need among people living with HIV in Kenya and Malawi. *AIDS*, 23, 1565-73. doi: 10.1097/QAD.0b013e32832cb10c
- Apondi, R., Bunnell, R., Ekwaru, J.P., Moore, D., Bechange, S., Khana, K...Mermin, J. (2011). Sexual behavior and HIV transmission risk of Ugandan adults taking antiretroviral therapy: 3 year follow-up. *AIDS*, 25,1317-27. doi: 10.1097/QAD.0b013e328347f775

- Ayiga, N. (2012). Rates and predictors of consistent condom-use by people living with HIV/AIDS on antiretroviral treatment in Uganda. *Journal of Health, Population and Nutrition*, 30, 270-80. Retrieved from: PMID: 23082629
- Bosmans, M., Cikuru, M.N., Claeys, P., & Temmerman, M. (2006). Where have all the condoms gone in adolescent programmes in the Democratic Republic of Congo. *Reproductive Health Matters*, 14, 80-8. Retrieved from: PMID: 17101425
- Carlos, S., Martínez-González, M.A., Burgueño, E., López-Del Burgo C., Ruíz-Canela, M., Ndarabu, A...de Irala, J. (2015). Misconceptions about HIV infection in Kinshasa (Democratic Republic of Congo): a case-control study on knowledge, attitudes and practices. *Sexually Transmitted Infections*, 91, 334-7. doi: 10.1136/sextrans-2014-051734.
- Centers for Disease Control and Prevention. (2014). Condoms and STDs: Fact Sheet for Public Health Personnel. Retrieved from [http:// www.cdc.gov/condomeffectiveness/docs/](http://www.cdc.gov/condomeffectiveness/docs/)
- Conant, M., Hardy, D., Sernatinger, J., Spicer, D., & Levy, J.A. (1986). Condoms prevent transmission of AIDS-associated retrovirus. *The Journal of the American Medical Association*, 255, 1706. Retrieved from: PMID: 3005677
- Davis, K.R., & Weller, S.C. (1999). The effectiveness of condoms in reducing heterosexual transmission of HIV. *Family Planning Perspectives*, 31, 272-9. Retrieved from: PMID: 10614517
- De Coninck, Z., & Marrone, G. (2012). Trends and determinants of condom use in Uganda. *East African Journal of Public Health*, 9,105-11. Retrieved from: PMID: 23136706
- Dellar, R.C., Dlamini, S., & Karim, Q.A. (2015). Adolescent girls and young women: key populations for HIV epidemic control. *Journal of the International AIDS Society*,18 Suppl 1,19408. doi: 10.7448/IAS.18.2.19408. eCollection 2015.
- Demographic Health Survey (DHS). (2014). République Démocratique du Congo Enquête Démographique et de Santé 2013-2014.

Devine, O.J., & Aral, S.O. (2004). The impact of inaccurate reporting of condom use and imperfect diagnosis of sexually transmitted disease infection in studies of condom effectiveness: a simulation-based assessment. *Sexually Transmitted Diseases*, 31, 588-95.

Retrieved from: PMID: 15388995

Earnshaw, V.A., Smith, L.R., Shuper, P.A., Fisher, W.A., Cornman, D.H., & Fisher, J.D. (2014). HIV stigma and unprotected sex among PLWH in KwaZulu-Natal, South Africa: a longitudinal exploration of mediating mechanisms. *AIDS Care*, 26, 1506-13. doi: 10.1080/09540121.2014.938015.

Eaton, E.F., & Hoesley, C.J. (2014). Barrier methods for human immunodeficiency virus prevention. *Infectious Disease Clinics of North America*, 28, 585-99. doi: 10.1016/j.idc.2014.08.006.

Exavery, A., Kanté, A.M., Jackson, E., Noronha, J., Sikustahili, G., Tani, K...Phillips, J.F. (2012). Role of condom negotiation on condom use among women of reproductive age in three districts in Tanzania. *BMC Public Health*, 12, 1097. doi: 10.1186/1471-2458-12-1097.

Exavery, A., Lutambi, A.M., Mubyazi, G.M., Kweka, K., Mbaruku, G., & Masanja, H. (2011). Multiple sexual partners and condom use among 10 - 19 year-olds in four districts in Tanzania: what do we learn? *BMC Public Health*, 11, 490. doi: 10.1186/1471-2458-11-490.

Fallopious, G. (1563). *De Morbo Gallico*. L Gryphium, Padua.

Fitch, J.T., Stine, C., Hager, W.D., Mann, J., Adam, M.B., McIlhaney, J. (2002). Condom effectiveness: factors that influence risk reduction. *Sexually Transmitted Diseases*, 29, 811-7.

Retrieved from: PMID: 12466725

Food and Drug Administration. *Condoms and Sexually Transmitted Diseases*. Retrieved from: <http://www.fda.gov/ForPatients/Illness/HIVAIDS/ucm126372.htm>

Geary, C.W., & Bukusi, E.A. (2014). Women and ARV-based HIV prevention - challenges and opportunities. *Journal of the International AIDS Society*, 17 Suppl 2, 19356. Retrieved from: PMID: 25224621

- Giannou, F.K., Tsiara, C.G., Nikolopoulos, G.K., Talias, M., Benetou, V., Kantzanou, M...Hatzakis, A. (2015). Condom effectiveness in reducing heterosexual HIV transmission: a systematic review and meta-analysis of studies on HIV serodiscordant couples. *Expert Review of Pharmacoeconomics & Outcomes Research*, 2016;16(4):489-99. PMID: 26488070
- Grasso, M.A., Schwarcz, S., Galbraith, J.S., Musyoki, H., Kambona, C., Kellogg, T.A.; KAIS Study Group. (2016). Estimating the Prevalence and Predictors of Incorrect Condom Use Among Sexually Active Adults in Kenya: Results From a Nationally Representative Survey. *Sexually Transmitted Diseases*, 43, 87-93. doi: 10.1097/OLQ.0000000000000393
- Green, E. (2011). *Broken Promises: How the AIDS Establishment has Betrayed the Developing World*, p.59,115.
- Green, E.C., Mah, T.L., Ruark, A., & Hearst, N. (2009). A framework of sexual partnerships: risks and implications for HIV prevention in Africa. *Studies in Family Planning*, 40, 63-70. Retrieved from: PMID: 19397186
- Grimes, D.A., & Schulz, K.F. (2005). Compared to what? Finding controls for case-control studies. *Lancet*, 365,1429–33. PMID: 15836892
- Halperin, D.T., Steiner, M.J., Cassell, M.M., Green, E.C., Hearst, N., Kirby, D...Cates, W. (2004). The time has come for common ground on preventing sexual transmission of HIV. *Lancet*, 364,1913-5. Retrieved from: PMID: 15566989
- Hawkes, M., Sivasivugha, E.S., Ngigi, S.K., Masumbuko, C.K., Brophy, J., & Kibendelwa, Z.T. (2013). HIV and religion in the Congo: a mixed-methods study. *Current HIV Research*, 11, 246-53. Retrieved from: PMID: 23701303
- Hearst, N., Ruark, A., Hudes, E.S., Goldsmith, J., & Green, E.C. (2013). Demographic and health surveys indicate limited impact of condoms and HIV testing in four African countries. *African Journal of AIDS Research*,12, 9-15. doi: 10.2989/16085906.2013.815406.

- Hira, S.K., Feldblum, P.J., Kamanga, J., Mukelabai, G., Weir, S.S., & Thomas, J.C. (1997). Condom and nonoxynol-9 use and the incidence of HIV infection in serodiscordant couples in Zambia. *International Journal of STD & AIDS*, 8, 243-50. Retrieved from: PMID: 9147157
- Holmes, K.K., Levine, R., & Weaver, M. (2004). Effectiveness of condoms in preventing sexually transmitted infections. *Bulletin of World Health Organization*, 82, 454-61. Retrieved from: PMID: 15356939
- Hosmer, D.W., & Lemeshow, S. (2013). Special topics. In: Hosmer DW, Lemeshow S, Sturdivant RX. *Applied Logistic Regression*, 3rd Edition. Wiley. New York; p-407-8.
- Hughes, J.P., Baeten, J.M., Lingappa, J.R., Magaret, A.S., Wald, A., de Bruyn, G...Partners in Prevention HSV/HIV Transmission Study Team. (2012). Determinants of per-coital-act HIV-1 infectivity among African HIV-1-serodiscordant couples. *Journal of Infectious Diseases*, 205, 358-65. doi: 10.1093/infdis/jir747
- Irwin, K., Bertrand, J., Mibandumba, N., Mbuyi, K., Muremeri, C., Mukoka, M...Ryder, R. (1991). Knowledge, attitudes and beliefs about HIV infection and AIDS among healthy factory workers and their wives, Kinshasa, Zaire. *Social Science & Medicine*, 32, 917-30. Retrieved from: PMID: 2031208
- Judson, F.N., Ehret, J.M., Bodin, G.F., Levin, M.J., & Rietmeijer, C.A. (1989). In vitro evaluations of condoms with and without nonoxynol 9 as physical and chemical barriers against *Chlamydia trachomatis*, herpes simplex virus type 2, and human immunodeficiency virus. *Sexually Transmitted Diseases*, 16, 51-6. Retrieved from: PMID: 2545003
- Kabamba Mulongo, L., Schirvel, C., Mukalay Wa Mukalay, A., & Dramaix Wilmet, M. (2011). Understanding couples' attitudes on prenatal HIV testing in the Democratic Republic of Congo. *Revue d'Epidémiologie et de Santé Publique*, 59, 379-83. doi: 10.1016/j.respe.2011.07.004
- Kajubi, P., Kanya, M.R., Kanya, S., Chen, S., McFarland, W., & Hearst, N.(2005). Increasing condom use without reducing HIV risk: results of a controlled community trial in

Uganda. *Journal of Acquired Immune Deficiency Syndromes*, 40, 77-82. Retrieved from: PMID: 16123686

Kamenga, M., Ryder, R.W., Jingu, M., Mbuyi, N., Mbu, L., Behets, F...Heyward, W.L. (1991). Evidence of marked sexual behavior change associated with low HIV-1 seroconversion in 149 married couples with discordant HIV-1 serostatus: experience at an HIV counselling center in Zaire. *AIDS*, 5, 61-67. Retrieved from: PMID: 2059362

Kamwi, R., Kenyon, T., & Newton, G. (2006). PEPFAR and HIV prevention in Africa. *Lancet*, 367,1978-9. Retrieved from : PMID: 16782481

Karim, S.A., Baxter, C., Frohlich, J., & Karim, Q.A. (2014). The need for multipurpose prevention technologies in sub-Saharan Africa. *BJOG: An International Journal of Obstetrics and Gynaecology*, Suppl 5, 27-34.

Kautako-Kiambi, M., Ekila, M.B., Kama-Lemba, S., Wumba, R., & Aloni, M.N. (2015). Voluntary Counseling and Testing for HIV in Rural Area of Democratic Republic of the Congo: Knowledge, Attitude, and Practice Survey among Service Users. *Journal of Tropical Medicine*, 2015, 281093. doi: 10.1155/2015/281093

Kayembe, P.K., Fatuma, A.B., Mapatano, M.A., & Mambu, T. (2006). Prevalence and determinants of the use of modern contraceptive methods in Kinshasa, Democratic Republic of Congo. *Contraception*, 74, 400-6. Retrieved from: PMID: 17046382

Kayembe, P.K., Mapatano, M.A., Busangu, A.F., Nyandwe, J.K., Musema, G.M., Kibungu, J.P...Mayala, GM. (2008). Determinants of consistent condom use among female commercial sex workers in the Democratic Republic of Congo: implications for interventions. *Sexually Transmitted Infections*, 84, 202-6. Retrieved from: PMID: 18055581

Laga, M., Alary, M., Nzila, N., Manoka, A.T., Tuliza, M., Behets, F... Piot, P. (1994). Condom promotion, sexually transmitted diseases treatment, and declining incidence of HIV-1 infection in female Zairian sex workers. *Lancet*, 344,:246-8. Retrieved from: PMID: 7913164



- Lagarde, E., Caraël, M., Glynn, J.R., Kanhonou, L., Abega, S.C., Kahindo, M...Study Group on the Heterogeneity of HIV Epidemics in African Cities. (2001). Educational level is associated with condom use within non-spousal partnerships in four cities of sub-Saharan Africa. *AIDS*, 15,1399-408. Retrieved from: PMID: 11504961
- Lassi, Z.S., Imam, A.M., Dean, S.V., & Bhutta, Z.A. (2014). Preconception care: preventing and treating infections. *Reproductive Health*,11 Suppl 3, S4. doi: 10.1186/1742-4755-11-S3-S4.
- Liu, H., Su, Y., Zhu, L., Xing, J., Wu, J., & Wang, N. (2014). Effectiveness of ART and condom use for prevention of sexual HIV transmission in serodiscordant couples: a systematic review and meta-analysis. *PLoS One*, 9, e111175. doi: 10.1371/journal.pone.0111175. eCollection 2014.
- MacQuarrie, K., Bradley, S., Gemmill, A., & Staveteig, S. (2014). *Contraceptive Dynamics Following HIV Testing*. DHS Analytical Studies No 47. Rockville, Maryland (USA). ICF International.
- Mantell, J.E., Smit, J.A., Beksinska, M., Scorgie, F., Milford, C., Balch, E...Stein, ZA. (2011). Everywhere you go, everyone is saying condom, condom. But are they being used consistently? Reflections of South African male students about male and female condom use. *Health Education Research*, 26, 859-71. doi: 10.1093/her/cyr041.
- Marshall, B.D., Perez-Brumer, A.G., MacCarthy, S., Mena, L., Chan, P.A., Towey, C...Nunn, A.S. (2016). Individual and Partner-Level Factors Associated with Condom Non-Use Among African American STI Clinic Attendees in the Deep South: An Event-Level Analysis. *AIDS and Behavior*, 20, 1334-42. doi: 10.1007/s10461-015-1266-9
- Marrazzo, J.M., del Rio, C., Holtgrave, D.R., Cohen, M.S., Kalichman, S.C., Mayer, K.H...International Antiviral Society-USA Panel. (2014). HIV prevention in clinical care settings: 2014 recommendations of the International Antiviral Society-USA Panel. *JAMA*, 312, 390-409. doi: 10.1001/jama.2014.7999.

- Maticka-Tyndale, E. (2012). Condoms in sub-Saharan Africa. *Sexual Health*, 9, 59-72. doi: 10.1071/SH11033.
- Medley, A., Bachanas, P., Grillo, M., Hasen, N., & Amanyeiwe, U. (2015). Integrating Prevention Interventions for People Living With HIV Into Care and Treatment Programs: A Systematic Review of the Evidence. *Journal of Acquired Immune Deficiency Syndromes*, 68 Suppl 3, S286-S296. doi: 10.1097/QAI.0000000000000520.
- Michielsen, K., Chersich, M.F., Luchters, S., De Koker, P., Van Rossem, R., & Temmerman, M. (2010). Effectiveness of HIV prevention for youth in sub-Saharan Africa: systematic review and meta-analysis of randomized and nonrandomized trials. *AIDS*, 24, 1193-202. doi: 10.1097/QAD.0b013e3283384791
- Mindel, A., & Sawleshwarkar, S. (2013). Physical Barrier Methods: Acceptance, Use, and Effectiveness. In: Lawrence R. Stanberry, Susan L. Rosenthal (Eds.), *Sexually Transmitted Diseases: Vaccines, Prevention, and Control*, (pp. 189-211). London, UK: Elsevier.
- Mishra, V., Prawn, A., Soumya, A., Yuan, G., & Shanxiao, W. (2009). Changes in HIV-related knowledge and behaviors in Sub-Saharan Africa. DHS Comparative Reports No. 24. Calverton, Maryland USA: ICF Macro.
- Murray, C.J., Ortblad, K.F., Guinovart, C., Lim, S.S., Wolock, T.M., Roberts, D.A...Vos T. (2014). Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*, 384, 1005-70. doi: 10.1016/S0140-6736(14)60844-8
- Mutevedzi, P.C., & Newell, M.L. (2014). Review: [corrected] The changing face of the HIV epidemic in sub-Saharan Africa. *Tropical Medicine & International Health*, 19, 1015-28. doi: 10.1111/tmi.12344
- Namisi, F., Aarø, L.E., Kaaya, S., Kajula, L.J., Kilonzo, G.P., Onya, H...Mathews, C. (2015). Adolescents' Communication with Parents, Other Adult Family Members and Teachers on

Sexuality: Effects of School-Based Interventions in South Africa and Tanzania. *AIDS and Behavior*, 19, 2162-76. doi: 10.1007/s10461-015-1019-9

National Institutes of Health (NIH). (2001). Workshop Summary: Scientific Evidence on Condom Effectiveness for Sexually Transmitted Disease (STD) Prevention.

Ntumbanzondo, M., Dubrow, R., Niccolai, L.M., Mwandagalirwa, K., & Merson, M.H. (2006). Unprotected intercourse for extra money among commercial sex workers in Kinshasa, Democratic Republic of Congo. *AIDS Care*, 18, 777-85. Retrieved from: PMID: 16971288

Obermeyer, C.M., Neuman, M., Desclaux, A., Wanyenze, R., Ky-Zerbo, O., Cherutich, P., Hardon, A. (2012). Associations between mode of HIV testing and consent, confidentiality, and referral: a comparative analysis in four African countries. *PLoS Med*, 9, e1001329. doi: 10.1371/journal.pmed.1001329

Ogden, J., Gupta, G.R., Warner, A., & Fisher, W.F. (2011). Revolutionising the AIDS response. *Global Public Health*, 6 Suppl 3, S383-95. doi: 10.1080/17441692.2011.621965.

Okware, S., Kinsman, J., Onyango, S., Opio, A., & Kaggwa, P. (2005). Revisiting the ABC strategy: HIV prevention in Uganda in the era of antiretroviral therapy. *Postgraduate Medical Journal*, 81, 625-8. Retrieved from: PMID: 16210457

Omori, R., & Abu-Raddad, L.J. (2016). Population sexual behavior and HIV prevalence in Sub-Saharan Africa: missing links? *International Journal of Infectious Diseases*, 44, 1-3. doi: 10.1016/j.ijid.2016.01.005

Oyeyemi, A., & Oyeyemi, B. (2012). Young adults and AIDS epidemics: their perception awareness information sources and sexual practices. *East African Journal of Public Health*, 9, 13-8. Retrieved from: PMID: 23120943

Padian, N.S., Buvé, A., Balkus, J., Serwadda, D., & Cates, W. Jr. (2008). Biomedical interventions to prevent HIV infection: evidence, challenges, and way forward. *Lancet*, 372, 585-99. doi: 10.1016/S0140-6736(08)60885-5

- Papo, J.K., Bauni, E.K., Sanders, E.J., Brocklehurst, P., & Jaffe, H.W. (2011). Exploring the condom gap: is supply or demand the limiting factor - condom access and use in an urban and a rural setting in Kilifi district, Kenya. *AIDS*, 25, 247-55. doi: 10.1097/QAD.0b013e328341b9b8
- Peltzer, K. (2012). Correlates of HIV infection among people visiting public HIV counseling and testing clinics in Mpumalanga, South Africa. *African Health Sciences*, 12,8-16. Retrieved from: PMID: 23066414
- Pilapil, M., Morris, L., Saito, K., Kouya, F., Maku, V., Kwalar, R...Jao, J. (2016). Retrospective analysis of the prevalence of and factors associated with condom use among young HIV-infected women in Cameroon. *SAGE Open Medicine*, 4, 2050312115626432. doi: 10.1177/2050312115626432. eCollection 2016.
- Pinkerton, S.D., & Abramson, P.R. (1997). Effectiveness of condoms in preventing HIV transmission. *Social Science & Medicine*, 44,1303-12. Retrieved from: PMID: 9141163
- Remis, R.S., Alary, M., Liu, J., Kaul, R., & Palmer, R.W. (2014). HIV transmission among men who have sex with men due to condom failure. *PLoS One*, 9, e107540. doi: 10.1371/journal.pone.0107540. eCollection 2014.
- Reynolds, S.J., Makumbi, F., Nakigozi, G., Kagaayi, J., Gray, R.H., Wawer, M...Serwadda, D. (2011). HIV-1 transmission among HIV-1 discordant couples before and after the introduction of antiretroviral therapy. *AIDS*, 25, 473-7. doi: 10.1097/QAD.0b013e3283437c2b.
- Rietmeijer, C.A., Krebs, J.W., Feorino, P.M., & Judson, F.N. (1988). Condoms as physical and chemical barriers against human immunodeficiency virus. *JAMA*, 259, 1851-3. Retrieved from: PMID: 2830416
- Rwenge, J.R. (2013). [Sexual behavioral among adolescents and young people in Sub-Saharan Africa and related factors]. *African Journal of Reproductive Health*, 17, 49-66. Retrieved from: PMID: 24069734

Sanders, S.A., Yarber, W.L., Kaufman, E.L., Crosby, R.A., Graham, C.A., & Milhausen, R.R. (2012). Condom use errors and problems: a global view. *Sexual Health*, 9, 81-95. doi: 10.1071/SH11095.

Shewamene, Z., Legesse, B., Tsega, B., Bhagavathula, A.S., & Endale, A. (2015). Consistent condom use in HIV/AIDS patients receiving antiretroviral therapy in northwestern Ethiopia: implication to reduce transmission and multiple infections. *HIV/AIDS (Auckland, N.Z.)*, 7, 119-24. doi: 10.2147/HIV.S79122

Skovdal, M., Campbell, C., Nyamukapa, C., & Gregson, S. (2011). When masculinity interferes with women's treatment of HIV infection: a qualitative study about adherence to antiretroviral therapy in Zimbabwe. *Journal of International AIDS Society*, 14, 29. doi: 10.1186/1758-2652-14-29.

Spaar, A., Graber, C., Dabis, F., Coutoudis, A., Bachmann, L., McIntyre J...ART-LINC of IeDEA. (2010). Prioritising prevention strategies for patients in antiretroviral treatment programmes in resource-limited settings. *AIDS Care*, 22, 775-83. doi: 10.1080/09540120903349102

Ukwuani, F.A., Tsui, A.O., & Suchindran, C.M. (2003). Condom use for preventing HIV infection/AIDS in sub-Saharan Africa: a comparative multilevel analysis of Uganda and Tanzania. *Journal of Acquired Immune Deficiency Syndromes*, 34, 203-13. Retrieved from: PMID: 14526210

UNAIDS. (2004). Making condoms work for HIV prevention.

UNAIDS. (2013). Report on the global AIDS epidemic.

UNAIDS. (2014a). The Gap Report.

UNAIDS. (2014b). Gender matters: overcoming genderrelated barriers to prevent new HIV infections among children and keep their mothers alive.

USAID. (2011). Audit of USAID/Democratic Republic of Congo's HIV/AIDS Program. Audit Report NO. 7-660-12-001-P.

Van de Perre, P., Jacobs, D., & Sprecher-Goldberger, S. (1987). The latex condom, an efficient barrier against sexual transmission of AIDS-related viruses. *AIDS*, 1, 49-52.

Retrieved from: PMID: 3122790

Vera, E.G., Orozco, H.H., Soto, S.S., & Aburto, E.L. (2008). [Condom effectiveness to prevent sexually transmitted diseases]. *Ginecología y Obstetricia de México*, 76, 88-96.

Retrieved from: PMID: 18798401

Walsh, T.L., Frazier, R.G., Peacock, K., Nelson, A.L., Clark, V.A., Bernstein, L., & Wray, B.G. (2004). Effectiveness of the male latex condom: combined results for three popular condom brands used as controls in randomized clinical trials. *Contraception*, 70, 407-13.

Retrieved from: PMID: 15504381

Wang, W., Aoumy, A., & Shanxiao, W. (2012). HIV-related knowledge and behaviors among people living with HIV in eight high HIV prevalence countries in Sub-Saharan Africa. DHS Analytical Studies No.29. Calverton, Maryland, USA: ICF International.

Weller, S., & Davis, K. (2002). Condom effectiveness in reducing heterosexual HIV transmission. *Cochrane Database of Systematic Reviews*, 1, CD003255. Retrieved from: PMID: 11869658

Wembonyama, S., Mpaka, S., & Tshilolo, L. (2007). Medicine and health in the Democratic Republic of Congo: from Independence to the Third Republic. *Med Trop (Mars)*, 67, 447-57.

Retrieved from: PMID: 18225727

Winer, R.L., Hughes, J.P., Feng, Q., O'Reilly, S., Kiviat, N.B., Holmes, K.K., & Koutsky, L.A. (2006). Condom use and the risk of genital human papillomavirus infection in young women. *New England Journal of Medicine*, 354, 2645-54. Retrieved from: PMID: 16790697

Workowski, K.A., & Bolan, G.A. (2015). Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines, 2015. *MMWR Recommendations and Reports*, 64(RR-03):1-137. Retrieved from: PMID: 26042815