



LUND UNIVERSITY

Pathophysiology of extracellular haemoglobin

use of animal models to translate molecular mechanisms into clinical significance

Smeds, Emanuel; Romantsik, Olga; Jungner, Åsa; Erlandsson, Lena; Gram, Magnus

Published in:
ISBT Science Series

DOI:
[10.1111/voxs.12328](https://doi.org/10.1111/voxs.12328)

2017

[Link to publication](#)

Citation for published version (APA):

Smeds, E., Romantsik, O., Jungner, Å., Erlandsson, L., & Gram, M. (2017). Pathophysiology of extracellular haemoglobin: use of animal models to translate molecular mechanisms into clinical significance. *ISBT Science Series*, 12(1), 134-141. <https://doi.org/10.1111/voxs.12328>

Total number of authors:
5

General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Pathophysiology of extracellular haemoglobin: Use of animal models to translate molecular mechanisms into clinical significance

Emanuel Smeds¹, Olga Romantsik², Åsa Jungner², Lena Erlandsson³ and Magnus Gram^{1,2*}

¹Lund University, Department of Clinical Sciences Lund, Infection Medicine, Lund, Sweden

²Lund University, Department of Clinical Sciences Lund, Paediatrics, Lund, Sweden

³Lund University, Department of Clinical Sciences Lund, Obstetrics and Gynaecology, Lund, Sweden

Running Title: Haemoglobin toxicity in animal models

Research Support: This work was supported by the Swedish Research Council, governmental ALF research grants to Lund University and Lund University Hospital, European Commission, the Crafoord Foundation, the Greta and Johan Kock Foundation, the Alfred Österlund Foundation and the Fanny Ekdahl's Foundation.

***Corresponding author**

Dr. Magnus Gram

Lund University, Department of Clinical Sciences Lund, Paediatrics, Infection Medicine

Biomedical Centre, B14, Klinikgatan 10, 221 84 Lund, Sweden

Phone: +46-709378638, Fax: +46-46157756

E-mail: magnus.gram@med.lu.se

Abstract

The blood's major gas-exchange is carried out by haemoglobin, a heme-protein that binds iron and oxygen and can have potentially dangerous side effects due to redox reactions.

Haemoglobin is a very abundant molecule with a concentration of 150 g/L in whole blood, resulting in almost one kg haemoglobin in an adult human body. Normal turn-over of red blood cells results in significant haemoglobin release, and pathological conditions that involve haemolysis can lead to massive haemoglobin levels. To control for the potential threat of extracellular haemoglobin, several protective defence systems have evolved.

Many pathological conditions, diseases as well as iatrogenic conditions, such as infusion of haemoglobin-based oxygen carriers, cerebral intraventricular haemorrhage, extracorporeal circulation and the pregnancy complication preeclampsia, involve abnormal levels of haemolysis and extracellular haemoglobin. Although quite different aetiology, the haemoglobin-induced damage often cause similar clinical sequelae and symptoms. Here we will give an overview of the pathophysiological mechanisms of extracellular haemoglobin and its metabolites. Furthermore, we will highlight the use of animal models in advancing the understanding of these mechanisms and discuss how to utilize the knowledge in the development of new and better pharmaceutical therapies.

Keywords: Haemolysis, extracellular haemoglobin, animal models, organ damage, toxicity, therapeutics

Introduction

Human red blood cells (RBCs) in circulation have a ~120 days life-span, after which they are phagocytized by resident macrophages in the reticuloendothelial system [1, 2]. RBCs have a major role in gas exchange between lungs and tissues as well as in maintaining pH-homeostasis of the body. RBCs are packed with haemoglobin (Hb), a highly reactive protein with numerous chemical and biological properties, including a potent extracellular toxicity, besides its gas-transporting functions [1, 3-5]. While encapsulated within the RBCs, the Hb-molecules are sequestered and the potentially damaging pathways are delimited. However, many clinical conditions result in haemolysis and tissue exposure to extracellular Hb.

Extracellular haemoglobin

A highly reactive and potentially cytotoxic protein

The balance between normal physiology and toxicity is largely dependent on the physical and biochemical barrier, presented by the RBC membrane, separating Hb from the extra-RBC environment. Furthermore, RBCs contain a unique setup of enzymatic mechanisms that preserves intracellular Hb in its functional state and prohibits deleterious side-reactions [6, 7]. Despite this sophisticated system to retain the Hb, significant amounts of Hb escape the RBCs, henceforth denoted as extracellular Hb, during normal RBC turn-over and massive amounts can be released during pathological conditions involving haemolysis. Once Hb escapes the intra-RBC compartment, it is highly reactive and activates cytotoxic, oxidative and inflammatory pathways, eventually leading to tissue damage [8, 9]. Extracellular Hb is rapidly oxidised from ferrous (Fe^{2+} , denoted oxyHb) to ferric (Fe^{3+} , denoted metHb) Hb [10, 11], which readily releases the heme group [12]. Heme is highly lipophilic and binds to lipids intercalating into cell membranes, which results in toxic cytolytic effects [13]. Moreover, free heme is very redox reactive and can damage lipids, proteins and DNA through oxidative

modification, cross-linking and fragmentation [14]. *In vivo*, several physiological and pathophysiological responses are induced by these redox reactions, for instance formation of signalling molecules often derived from oxidation of lipids [15]. The central concept of “redox cell signalling” is that changes in the reductive or oxidative capacity of the cell can lead to posttranslational modifications of proteins [16-18]. This is an emerging field of research in pathologies such as atherosclerosis, ischemia/reperfusion injuries, post-haemolytic injuries, and blood substitutes.

Besides acting as a redox component, extracellular Hb may also bind and react with nitric oxide (NO)[19]. Extracellular Hb thus affects the vasomotor tone, which reflects in the clinical observation that haemolysis and elevated extracellular Hb-concentrations are associated with hypertension (Fig. 1).

Evolutionarily developed protective mechanisms

A number of Hb- and heme-detoxification systems are described in humans. Haptoglobin (Hp) binds extracellular Hb in blood [4, 20], and following binding to the CD163 receptor on macrophages the Hp-Hb complex is removed from the circulation [21]. Free heme in blood is cleared by both albumin and hemopexin (Hpx) [4, 14, 22] and the hepatocyte receptor CD91 removes the Hpx-heme complex from the circulation [23]. A complex network of antioxidation mechanisms (such as superoxide dismutase, glutathione peroxidase and catalase) inhibit and eliminate the oxidative compounds and repair the oxidative damage from uncleared Hb. Intracellularly, heme oxygenase (HO) is the most essential heme catabolic protein, converting heme to free iron, biliverdin and CO [13, 24]. Additionally, the plasma and tissue protein α_1 -microglobulin (A1M) has been described to bind and degrade heme. A series of reports describe that A1M has a physiological role as a protective antioxidant by

clearing extravascular fluids of free radicals and heme-groups, and transporting them to the kidneys for degradation (reviewed in [25])(Fig. 2).

Haemoglobin based oxygen carriers

For more than 70 years, there have been attempts to develop blood substitutes or “artificial blood”. Some of the products, most of them different derivatives of Hb, have undergone clinical trials but so far none have been approved for medical use. As blood has many molecules with functions apart from the transportation of oxygen, the term ‘haemoglobin-based oxygen carrier’ (‘HBOC’) is typically used in this context. There are several important clinical applications that drives the development of HBOCs for medical use. Replacement of blood transfusion products (e.g. stored bank blood), reducing the risk of immunological complications and resolving the issue of a scarcity of bank blood is apparent areas. The development of a clinically approved HBOC would also enable Hb replacement therapy to patients from groups that are concerned in regards to receiving human blood products.

Evaluation of HBOCs in animal models

Although it is possible to make fairly detailed studies of the molecular reactivity of HBOCs using *in vitro* studies, documented safety and efficacy in animal models is an absolute requirement before entering into human clinical trials.

In regards to safety, the toxicology studies that are needed for HBOCs will be similar to other biological compounds. Importantly, as a result of comprehensive human clinical studies during the last decades, a fair amount of data in regards to HBOC related toxicity is nowadays available and can broadly be divided into the following categories: oxidative stress, hypertension and cardiac toxicity (reviewed in Buhler et al [26]). The hypertension is believed to be a result of NO-consumption in the vascular endothelium, whereas the oxidative stress is

a result of the free radicals formed during redox cycling of Hb [26]. The cardiac toxicity have been mimicked in laboratory animals by a NO-synthase inhibitor, suggesting that the resulting toxicity is a result of NO-depletion, but the mechanism may be more complicated [27].

Studies of HBOC efficacy have so far mainly been conducted in bleeding and trauma models. When conducting efficacy studies, it is important that the pathophysiology of the insult is relevant in a clinical context and that the selected animal species can be considered a relevant and transferable species when comparing to humans. A number of different animal species have been used to study HBOC efficacy, including rat, dog, sheep, pig and rhesus monkey [27]. Furthermore, guinea pig is a commonly used animal species for testing HBOCs as guinea pigs, like humans, are incapable of endogenous synthesis of ascorbic acid [28, 29]. This incapability implies a more pronounced susceptibility to Hb toxicity, as has been nicely shown in a comparative trial using rats and guinea pigs [30]. However, guinea pig is a relatively unused species in drug development and it can therefore be speculated that future HBOC animal studies will not use only one single vertebrate species, but several different species. The importance of using more than one species to test HBOC is underscored by the finding that the cardiac toxicity mentioned above was not seen in species like mouse and rat [27].

Next generation HBOCs may require different animal models than the ones used in the past

In the past, HBOCs have typically been evaluated in clinical trials as a blood substitute where the standard therapy would be RBCs from human donors. In fact, all HBOCs entering into clinical trials have been based on either human or bovine Hb. As a result of technical advancement, the development of next generation HBOCs will most likely allow altered properties compared to native forms of Hb. These may be more stable against degradation, have higher oxygen affinity or have an altered binding affinity to Hp or other host molecules.

Consequently, these HBOCs may become interesting for use in clinical indications other than as “blood substitute”. New indications will most likely require development of new or altered animal models, both with regards to safety and efficacy.

Cerebral intraventricular haemorrhage

Cerebral intraventricular haemorrhage (IVH) is a significant problem in neonatal intensive care with an incidence of 25% in very low birth weight infants [31]. Complications of IVH, including periventricular haemorrhagic infarction, posthaemorrhagic ventricular dilatation (PHVD) and the associated cerebellar haemorrhagic injury and periventricular leukomalacia are critical determinants of neonatal morbidity, mortality, and long-term neurodevelopmental sequelae [32]. Approximately 50% to 75% of preterm survivors with IVH/PHVD develop cerebral palsy and mental retardation [33]. Moreover, around a quarter of non-disabled survivors develop psychiatric disorders and problems with executive function [34, 35]. Hence, IVH and its resultant neurologic and psychiatric sequelae continue to be an important public health concern worldwide.

Aetiology and studies of Hb-related brain damage in animal models

The aetiology of IVH is multifactorial, complex, heterogeneous and incompletely understood. At present, IVH is neither preventable nor treatable and hence there is a great need to obtain a better understanding of the molecular mechanisms of brain damage and to develop treatment strategies to prevent or minimize the neurologic consequences of IVH. With regards to this, a number of animal species, including mouse, rat, rabbit, sheep, pig, dog, cat, and primate, have been used to model IVH. These include studies of needle injected blood or collagenase into the cerebral intraventricular space or alterations in haemodynamic parameters, *i.e.* blood pressure, circulating blood volume, serum osmolarity, partial carbon dioxide or oxygen levels

[36]. None of these animal models can be considered ideal, but in combination each of them aid in obtaining a better understanding of the underlying pathophysiological mechanisms of IVH. A generally accepted description of how IVH develops is that following rupture of blood vessels within the germinal matrix, there is a rapid accumulation of blood within the ventricles leading to disruption of normal anatomy and increased local intracerebral pressure. Secondary brain damage is attributable to a number of pathways where the presence of blood and the initial hematoma volume is of great importance [13]. In a preterm rabbit pup model of IVH, it was recently shown that extracellular Hb infiltrates and is widely dispersed in large regions of the cerebral periventricular white matter, especially in areas with high extracellular plasticity [37]. Furthermore, metHb was shown to be a potent inducer of pro-inflammation, displaying a strong correlation with TNF α protein levels in intraventricular CSF and increased mRNA levels for TNF α , IL-1 β and of HO-1 in periventricular brain tissue [10, 38].

Scavenger studies of Hb-metabolites display the importance of Hb in the pathophysiology

To support the hypothesis of extracellular Hb mediated toxicity, several Hb-, heme- and ROS-neutralising systems has been investigated. It has been shown that intracerebroventricular injection of the Hb-scavenger Hp *in vivo* reversed or reduced cellular activation, inflammatory response, structural damage, and cell death following IVH [38](Fig. 3). Deferoxamine, a potent free iron chelator, has been shown to attenuate hippocampal cell loss following IVH [39]. Of further interest, it was recently shown that deletion of Hpx, a natural heme-scavenger, leads to aggravated brain injury following intracranial bleeding and increased iron deposition [40]. Altogether, these data suggest that extracellular Hb plays a crucial role in development of brain damage following IVH.

Extracorporeal circulation and haemolysis

Extracorporeal circulation, *i.e.* extracorporeal membrane oxygenation (ECMO) and cardiopulmonary bypass (CPB), invariably leads to haemolysis and systemic exposure to extracellular Hb as a result of shear forces on the RBCs and, in paediatric extracorporeal circulation, as a result of circuit priming. Furthermore, blood transfusions commonly needed in conditions requiring extracorporeal circulation contribute to the haemolytic burden. Recent studies have shown a clear association between extracellular Hb and the development of postoperative acute kidney injury after surgery on CPB [41-44], suggesting a crucial role in the induction of inflammation and oxidative stress by Hb and its metabolites [45]. In clinical studies on paediatric and adult ECMO patients increased levels of plasma extracellular Hb are associated with increased mortality [43, 46].

Granted the inevitable need for extracorporeal circulation in some clinical settings, and the increased morbidity and mortality associated with intravascular haemolysis, it is essential to define if there is a causal connection between the well-described molecular reactivity of extracellular Hb and clinically relevant outcomes, or if the association remains a numerical association. The use of animal studies might provide the means of establishing a causative link, and ultimately, a possibility to evaluate different treatment options. Full-scale animal models have been developed in primates and non-primates and miniature circuits have been developed for studies on rodents and lagomorphs [47-49]. Importantly, the implementation of extracorporeal circulation requires a circuit that has similar characteristics as the clinical counterparts. As of now, most small-animal circuits are developed at the research site with disparate techniques and disparate performance. A more standardized setting would be required to transfer results between different animal studies, and to translate the results into a clinical context. Up to date, few animal studies exist that specifically intended to delineate the

contribution of extracellular Hb to significant end-organ damage in the context of extracorporeal circulation.

Preeclampsia

Preeclampsia (PE) is one of the most serious pregnancy-related diseases, affecting 3-7 % of pregnant women, making it the leading cause of maternal and foetal morbidity and mortality [50, 51]. Currently, symptomatic blood pressure treatment is the only available treatment for PE and to date the only known cure is delivery.

Animal models support the involvement of foetal Hb in the aetiology

PE is described to have a placental origin, even if the exact aetiology remains unknown, and is believed to evolve in two stages [52]. The first stage occurs during the formation of the placenta, with defective and shallow invasion and remodelling of the spiral arteries resulting in uneven blood perfusion [53]. The second stage include de novo proteinuria and hypertension, appearing as a clinical manifestation after 20 weeks of gestation [54], likely caused by placental-derived material and substances [55-58].

Foetal extracellular Hb (HbF) is believed to play a role in the pathophysiology of PE by inducing damage to the placenta barrier and the subsequent leakage of foetal factors, including HbF into the maternal blood circulation. Clinical studies have shown that there is an increased expression and accumulation of extracellular HbF in the placenta of preeclamptic mothers [59] and results suggest a role for HbF as an important link between stage 1 and 2 in PE [60].

The importance of extracellular Hb in the development of PE has been studied in a few animal models, including rat, rabbit and sheep [61-63]. Following systemic administration of HbF, kidneys displayed an increased glomerular permeability to macromolecules as well as

tissue damage. The importance of HbF-induced kidney damage was further supported by the finding that co-administration of the heme- and radical scavenger A1M restored the structure and function in the kidneys by normalizing both glomerular filtration rate and histology [62]. The importance of Hb in the development of PE was further studied, using a pregnant ewe animal model where PE equivalent symptoms are induced by starvation, causing haemolysis and subsequent release of Hb [63, 64]. In this model, an elimination of the collagen fibres and cellular damage in the placenta was observed. The kidneys displayed structural damage and increased glomerular filtration. Co-administration of A1M was shown to ameliorate the structural tissue damages seen in kidney and placenta and restored the glomerular filtration rate, thus supporting a causal role of extracellular Hb in the development of PE.

In a pregnant rabbit model, administration of species-specific extracellular HbF mimic the human symptoms at stage two of the disease [61]. The rabbits displayed disrupted placental morphology, proteinuria and renal glomerular lesions with a significantly increased glomerular filtration. Co-administration of A1M ameliorated the proteinuria and reversed the increased glomerular filtration, and the animals displayed a significant reduction of the structural and cellular damages in placenta.

Concluding remarks

Extracellular Hb, resulting from haemolysis or exogenous infusion, is shown to be an important pathogenic factor in a growing number of diseases. When using animal models to establish causality of extracellular Hb and evaluate new therapeutic strategies, several pre-requisites needs to be considered: the setting should 1) mirror clinical standards as closely as possible to generate transferrable results, 2) allow long-term follow-up 3) be technically simple and standardized, 4) be relatively inexpensive to allow significant group sizes, and 5) use established and validated outcome measurements. Furthermore, the interpretation of

results derived from animal studies requires caution; animal models often evokes a complex pathophysiology where extracellular Hb toxicity is but one of numerous potential causes of harm.

Acknowledgments

The authors wish to acknowledge the contribution of past and present co-workers and members of the research groups.

All authors wrote, read and approved the manuscript.

Competing Interests

We have read the journal's policy and the authors of this manuscript have the following competing interests: the author MG is co-founder and shareholder of the company A1M Pharma AB.

Abbreviations

A1M, α_1 -microglobulin; CPB, cardiopulmonary bypass; ECMO, extracorporeal membrane oxygenation; Hb, haemoglobin; HbF, foetal haemoglobin; HBOC, haemoglobin based oxygen carrier; HO, heme oxygenase; Hp, haptoglobin; Hpx, hemopexin; IVH, intraventricular haemorrhage; MetHb, ferric (Fe^{3+}) haemoglobin; NO, nitric oxide; OxyHb, ferrous (Fe^{2+}) haemoglobin; PE, preeclampsia; PHVD, post-haemorrhagic ventricular dilatation; RBC, red blood cell.

References

- 1 Buehler PW, D'Agnillo F: Toxicological consequences of extracellular hemoglobin: biochemical and physiological perspectives. *Antioxid Redox Signal* 2010; 12: 275-91.
- 2 Bratosin D, Mazurier J, Tissier JP, et al.: Cellular and molecular mechanisms of senescent erythrocyte phagocytosis by macrophages. A review. *Biochimie* 1998; 80: 173-95.
- 3 Bunn H: Hemoglobin. Weinheim, 1992.
- 4 Ascenzi P, Bocedi A, Visca P, et al.: Hemoglobin and heme scavenging. *IUBMB Life* 2005; 57: 749-59.
- 5 Reeder BJ: The redox activity of hemoglobins: from physiologic functions to pathologic mechanisms. *Antioxid Redox Signal* 2010; 13: 1087-123.
- 6 Rother RP, Bell L, Hillmen P, et al.: The clinical sequelae of intravascular hemolysis and extracellular plasma hemoglobin: a novel mechanism of human disease. *JAMA : the journal of the American Medical Association* 2005; 293: 1653-62.
- 7 Cimen MY: Free radical metabolism in human erythrocytes. *Clin Chim Acta* 2008; 390: 1-11.
- 8 Xi G, Keep RF, Hoff JT: Mechanisms of brain injury after intracerebral haemorrhage. *Lancet Neurol* 2006; 5: 53-63.
- 9 Fang H, Wang PF, Zhou Y, et al.: Toll-like receptor 4 signaling in intracerebral hemorrhage-induced inflammation and injury. *J Neuroinflammation* 2013; 10: 27.
- 10 Gram M, Sveinsdottir S, Ruscher K, et al.: Hemoglobin induces inflammation after preterm intraventricular hemorrhage by methemoglobin formation. *J Neuroinflammation* 2013; 10: 100.
- 11 Umbreit J: Methemoglobin--it's not just blue: a concise review. *Am J Hematol* 2007; 82: 134-44.
- 12 Bunn HF, Jandl JH: Exchange of heme among hemoglobins and between hemoglobin and albumin. *J Biol Chem* 1968; 243: 465-75.
- 13 Wagener FA, Eggert A, Boerman OC, et al.: Heme is a potent inducer of inflammation in mice and is counteracted by heme oxygenase. *Blood* 2001; 98: 1802-11.
- 14 Kumar S, Bandyopadhyay U: Free heme toxicity and its detoxification systems in human. *Toxicol Lett* 2005; 157: 175-88.
- 15 Poli G, Leonarduzzi G, Biasi F, et al.: Oxidative stress and cell signalling. *Curr Med Chem* 2004; 11: 1163-82.
- 16 Cooper CE, Patel RP, Brookes PS, et al.: Nanotransducers in cellular redox signaling: modification of thiols by reactive oxygen and nitrogen species. *Trends Biochem Sci* 2002; 27: 489-92.
- 17 Levonen AL, Patel RP, Brookes P, et al.: Mechanisms of cell signaling by nitric oxide and peroxynitrite: from mitochondria to MAP kinases. *Antioxid Redox Signal* 2001; 3: 215-29.
- 18 Shiva S, Moellering D, Ramachandran A, et al.: Redox signalling: from nitric oxide to oxidized lipids. *Biochem Soc Symp* 2004: 107-20.
- 19 Kim-Shapiro DB, Schechter AN, Gladwin MT: Unraveling the reactions of nitric oxide, nitrite, and hemoglobin in physiology and therapeutics. *Arteriosclerosis, thrombosis, and vascular biology* 2006; 26: 697-705.
- 20 Laurell CB, Nyman M: Studies on the serum haptoglobin level in hemoglobinemia and its influence on renal excretion of hemoglobin. *Blood* 1957; 12: 493-506.
- 21 Kristiansen M, Graversen JH, Jacobsen C, et al.: Identification of the haemoglobin scavenger receptor. *Nature* 2001; 409: 198-201.

- 22 Delanghe JR, Langlois MR: Hemopexin: a review of biological aspects and the role in laboratory medicine. *Clin Chim Acta* 2001; 312: 13-23.
- 23 Hvidberg V, Maniecki MB, Jacobsen C, et al.: Identification of the receptor scavenging hemopexin-heme complexes. *Blood* 2005; 106: 2572-9.
- 24 Tenhunen R, Marver HS, Schmid R: The enzymatic conversion of heme to bilirubin by microsomal heme oxygenase. *Proc Natl Acad Sci U S A* 1968; 61: 748-55.
- 25 Åkerström B, Gram M: A1M, an extravascular tissue cleaning and housekeeping protein. *Free Radic Biol Med* 2014; 74: 274-82.
- 26 Buehler PW, D'Agnillo F, Schaer DJ: Hemoglobin-based oxygen carriers: From mechanisms of toxicity and clearance to rational drug design. *Trends Mol Med* 2010; 16: 447-57.
- 27 Burhop K, Gordon D, Estep T: Review of hemoglobin-induced myocardial lesions. *Artif Cells Blood Substit Immobil Biotechnol* 2004; 32: 353-74.
- 28 Nishikimi M, Fukuyama R, Minoshima S, et al.: Cloning and chromosomal mapping of the human nonfunctional gene for L-gulonogamma-lactone oxidase, the enzyme for L-ascorbic acid biosynthesis missing in man. *J Biol Chem* 1994; 269: 13685-8.
- 29 Nishikimi M, Kawai T, Yagi K: Guinea pigs possess a highly mutated gene for L-gulonogamma-lactone oxidase, the key enzyme for L-ascorbic acid biosynthesis missing in this species. *J Biol Chem* 1992; 267: 21967-72.
- 30 Butt OI, Buehler PW, D'Agnillo F: Differential induction of renal heme oxygenase and ferritin in ascorbate and nonascorbate producing species transfused with modified cell-free hemoglobin. *Antioxid Redox Signal* 2010; 12: 199-208.
- 31 Vermont Oxford Network: Database of very low birth weight infants born in 2012. Burlington, VT, USA, Nightingale Internet Reporting System, Accessed April 4, 2014.
- 32 Sherlock RL, Anderson PJ, Doyle LW, et al.: Neurodevelopmental sequelae of intraventricular haemorrhage at 8 years of age in a regional cohort of ELBW/very preterm infants. *Early Hum Dev* 2005; 81: 909-16.
- 33 Luu TM, Ment LR, Schneider KC, et al.: Lasting effects of preterm birth and neonatal brain hemorrhage at 12 years of age. *Pediatrics* 2009; 123: 1037-44.
- 34 Indredavik MS, Vik T, Evensen KA, et al.: Perinatal risk and psychiatric outcome in adolescents born preterm with very low birth weight or term small for gestational age. *J Dev Behav Pediatr* 2010; 31: 286-94.
- 35 Nosarti C, Giouroukou E, Micali N, et al.: Impaired executive functioning in young adults born very preterm. *J Int Neuropsychol Soc* 2007; 13: 571-81.
- 36 Balasubramaniam J, Del Bigio MR: Animal models of germinal matrix hemorrhage. *J Child Neurol* 2006; 21: 365-71.
- 37 Ley D, Romantsik O, Vallius S, et al.: High Presence of Extracellular Hemoglobin in the Periventricular White Matter Following Preterm Intraventricular Hemorrhage. *Frontiers in Physiology* 2016; 7.
- 38 Gram M, Sveinsdottir S, Cinthio M, et al.: Extracellular hemoglobin - mediator of inflammation and cell death in the choroid plexus following preterm intraventricular hemorrhage. *J Neuroinflammation* 2014; 11: 200.
- 39 Garton TP, He Y, Garton HJ, et al.: Hemoglobin-induced neuronal degeneration in the hippocampus after neonatal intraventricular hemorrhage. *Brain Res* 2016; 1635: 86-94.
- 40 Ma B, Day JP, Phillips H, et al.: Deletion of the hemopexin or heme oxygenase-2 gene aggravates brain injury following stroma-free hemoglobin-induced intracerebral hemorrhage. *J Neuroinflammation* 2016; 13: 26.

- 41 Vermeulen Windsant IC, Hanssen SJ, Buurman WA, et al.: Cardiovascular surgery and organ damage: time to reconsider the role of hemolysis. *J Thorac Cardiovasc Surg* 2011; 142: 1-11.
- 42 O'Neal JB, Shaw AD, Billings FTt: Acute kidney injury following cardiac surgery: current understanding and future directions. *Crit Care* 2016; 20: 187.
- 43 Mamikonian LS, Mamo LB, Smith PB, et al.: Cardiopulmonary bypass is associated with hemolysis and acute kidney injury in neonates, infants, and children*. *Pediatr Crit Care Med* 2014; 15: e111-9.
- 44 Billings FTt, Yu C, Byrne JG, et al.: Heme Oxygenase-1 and Acute Kidney Injury following Cardiac Surgery. *Cardiorenal Med* 2014; 4: 12-21.
- 45 Haase M, Bellomo R, Haase-Fielitz A: Novel biomarkers, oxidative stress, and the role of labile iron toxicity in cardiopulmonary bypass-associated acute kidney injury. *J Am Coll Cardiol* 2010; 55: 2024-33.
- 46 Lou S, MacLaren G, Best D, et al.: Hemolysis in pediatric patients receiving centrifugal-pump extracorporeal membrane oxygenation: prevalence, risk factors, and outcomes. *Crit Care Med* 2014; 42: 1213-20.
- 47 Cook DJ, Orszulak TA, Daly RC: The effects of pulsatile cardiopulmonary bypass on cerebral and renal blood flow in dogs. *J Cardiothorac Vasc Anesth* 1997; 11: 420-7.
- 48 Fujii Y, Shirai M, Inamori S, et al.: A novel small animal extracorporeal circulation model for studying pathophysiology of cardiopulmonary bypass. *J Artif Organs* 2015; 18: 35-9.
- 49 Schnoering H, Arens J, Detering SM, et al.: Development of a rabbit animal model for miniaturized heart-lung machines. *ASAIO J* 2013; 59: 152-6.
- 50 Berg CJ, Mackay AP, Qin C, et al.: Overview of maternal morbidity during hospitalization for labor and delivery in the United States: 1993-1997 and 2001-2005. *Obstetrics and gynecology* 2009; 113: 1075-81.
- 51 MacKay AP, Berg CJ, Atrash HK: Pregnancy-related mortality from preeclampsia and eclampsia. *Obstetrics and gynecology* 2001; 97: 533-8.
- 52 Roberts JM, Hubel CA: The two stage model of preeclampsia: variations on the theme. *Placenta* 2009; 30 Suppl A: S32-7.
- 53 Brosens JJ, Pijnenborg R, Brosens IA: The myometrial junctional zone spiral arteries in normal and abnormal pregnancies: a review of the literature. *Am J Obstet Gynecol* 2002; 187: 1416-23.
- 54 Redman CW: Hypertension in pregnancy: the NICE guidelines. *Heart* 2011; 97: 1967-9.
- 55 Redman CW, Sargent IL: Placental stress and pre-eclampsia: a revised view. *Placenta* 2009; 30 Suppl A: S38-42.
- 56 Roberts JM, Taylor RN, Musci TJ, et al.: Preeclampsia: an endothelial cell disorder. *Am J Obstet Gynecol* 1989; 161: 1200-4.
- 57 Tjoa ML, Cindrova-Davies T, Spasic-Boskovic O, et al.: Trophoblastic oxidative stress and the release of cell-free feto-placental DNA. *The American journal of pathology* 2006; 169: 400-4.
- 58 Cronqvist T, Salje K, Familiari M, et al.: Syncytiotrophoblast vesicles show altered micro-RNA and haemoglobin content after ex-vivo perfusion of placentas with haemoglobin to mimic preeclampsia. *PLoS One* 2014; 9: e90020.
- 59 Centlow M, Carninci P, Nemeth K, et al.: Placental expression profiling in preeclampsia: local overproduction of hemoglobin may drive pathological changes. *Fertil Steril* 2008; 90: 1834-43.

- 60 Hansson SR, Gram M, Åkerström B: Fetal hemoglobin in preeclampsia: a new causative factor, a tool for prediction/diagnosis and a potential target for therapy. *Curr Opin Obstet Gynecol* 2013; 25: 448-55.
- 61 Nääv A, Erlandsson L, Axelsson J, et al.: A1M Ameliorates Preeclampsia-Like Symptoms in Placenta and Kidney Induced by Cell-Free Fetal Hemoglobin in Rabbit. *PLoS One* 2015; 10: e0125499.
- 62 Sverrisson K, Axelsson J, Rippe A, et al.: Extracellular fetal hemoglobin induces increases in glomerular permeability: inhibition with alpha-1-microglobulin and tempol. *Am J Physiol Renal Physiol* 2014; 306: F442-8.
- 63 Wester-Rosenlöf L, Casslen V, Axelsson J, et al.: A1M/alpha-1-microglobulin protects from heme-induced placental and renal damage in a pregnant sheep model of preeclampsia. *PLoS One* 2014; 9: e86353.
- 64 Talosi G, Nemeth I, Nagy E, et al.: The pathogenetic role of heme in pregnancy-induced hypertension-like disease in ewes. *Biochem Mol Med* 1997; 62: 58-64.
- 65 Olsson MG, Allhorn M, Bulow L, et al.: Pathological conditions involving extracellular hemoglobin: molecular mechanisms, clinical significance, and novel therapeutic opportunities for alpha-1-microglobulin. *Antioxid Redox Signal* 2012; 17: 813-46.

Figure Legends

Figure 1. Extracellular Hb-toxicity. While encapsulated within the RBCs, the Hb-molecules are sequestered and the potentially damaging pathways are delimited. Once released into the extracellular compartment, following haemolysis, Hb is highly reactive and is rapidly oxidised. Subsequently, Hb easily loses the heme group and generates free iron and free radicals. All of these components are highly cytotoxic and can generate an oxidative and inflammatory response, eventually leading to tissue damage. Furthermore, extracellular Hb may also bind and react with NO, consequently affecting the vasomotor tone and causing hypertension. In addition, the metabolites formed by the Hb-degradation are haemolytic and thus, oxidative stress accelerates the haemolysis, in a positive feed-back loop. Image modified from Olsson M.G. et al. [65].

Figure 2. Endogenous scavenger systems against Hb-toxicity. Humans have evolved several Hb- and heme-detoxification systems. Hp binds extracellular Hb in blood and the resulting Hp-Hb complex is cleared by binding to the macrophage receptor CD163. Free heme in blood is sequestered mainly by Hpx and the Hpx-heme complex is cleared from the circulation by the hepatocyte receptor CD91. Antioxidation mechanisms inhibit and eliminate the oxidative compounds and repair the oxidative damage caused by uncleared Hb. Intracellularly, HO degrades heme. Extravascular and within tissues, A1M binds and degrades both heme and free radicals. A1M will clear extravascular fluids of free radicals and heme-groups, and transport them to the kidneys for degradation. Image modified from Olsson M.G. et al. [65].

Figure 3. Hp reduces Hb-induced structural damage in the choroid plexus. The importance of Hb-metabolite induced damage in the choroid plexus *in vivo* was investigated by an intraventricular injection of the Hb-scavenger Hp (IVH+Hp) or vehicle solution (IVH+Sham) to rabbit pups with confirmed IVH. Examination of the structural integrity of

the choroid plexus with transmission electron microscopy displayed a clearly reduced tissue damage following Hp injection, as compared to vehicle solution injection. CP= choroid plexus, V= ventricle, Vi= villi. Scale indicate 500 nm. Image modified from Gram M. et al. [38].

Figure 1




Figure 2






Figure 3



IVH+Hp



IVH+Sham