

## Detection and intervention of health problems after stroke – a nurse-led follow-up program

Jönsson, Ann-Cathrin; Pessah-Rasmussen, Hélène

Published in:

Program Book GSA Sixty-Fifth Annual Scientific Meeting

2012

## Link to publication

Citation for published version (APA):

Jönsson, A-C., & Pessah-Rasmussen, H. (2012). Detection and intervention of health problems after stroke – a nurse-led follow-up program. Program Book GSA Sixty-Fifth Annual Scientific Meeting.

Total number of authors:

General rights

Unless other specific re-use rights are stated the following general rights apply: Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

  • You may not further distribute the material or use it for any profit-making activity or commercial gain

  • You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: https://creativecommons.org/licenses/

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**LUND UNIVERSITY** 

## Detection and intervention of health problems after stroke – a nurse-led follow-up program

<sup>1</sup>Ann-Cathrin Jönsson RN PhD & <sup>2</sup>Hélène Pessah-Rasmussen, MD PhD

<sup>1</sup>Department of Health Sciences and Centre of Ageing and Supportive Environments (CASE), Lund University, Sweden

*Introduction:* Secondary prevention among stroke survivors has been reported to be neglected. This randomised controlled trial examines whether a structured nurse-led follow-up program three months after stroke improves health status one year after stroke.

*Method:* During one year, all survivors one month after stroke from a university hospital in Malmö, Sweden, were approached for participation. Those consenting were randomised to intervention group with structured follow-up at three months or to standard care (control group) (mean age 73.8/73.7 years). Three months after stroke, a nurse specialist (NS) followed up the intervention group regarding risk factors and health problems after stroke. The NS sent urgent health problems immediately to a stroke clinician; otherwise patients were referred to a GP for non-urgent or rehabilitative interventions. Patients were informed about stroke and life style factors.

Depression/anxiety was assessed by patient self-report with the EQ-5D scale.

**Results:** After three months, 80% of the intervention group (n=194) needed 1-4 interventions compared with 63% (p<0.001) in the same group after one year, and 74% in the control group (n=197) (p=0.022) after one year, and only 40% in the intervention group experienced depression/anxiety compared with 52% in the control group (p=0.042).

**Conclusions:** A nurse-led follow-up three months after stroke resulted in reduced need for interventions and lower prevalence of self-reported depression/anxiety one year after stroke compared to standard care. The follow-up program may be further enhanced by collaboration between a NS and a stroke clinician at the outpatient clinic regarding all medical interventions, before referring patients to the GP for continued follow-up.

<sup>&</sup>lt;sup>2</sup>Department of Neurology, Skåne University Hospital, Malmö