



LUND UNIVERSITY

Characterization of long-term motor deficits in the 6-OHDA model of Parkinson's disease in the common marmoset.

Santana, M; Palmér, Tobias; Simplício, H; Fuentes, R; Petersson, Per

Published in:
Behavioural Brain Research

DOI:
[10.1016/j.bbr.2015.04.037](https://doi.org/10.1016/j.bbr.2015.04.037)

2015

[Link to publication](#)

Citation for published version (APA):
Santana, M., Palmér, T., Simplício, H., Fuentes, R., & Petersson, P. (2015). Characterization of long-term motor deficits in the 6-OHDA model of Parkinson's disease in the common marmoset. *Behavioural Brain Research*, 290, 90-101. <https://doi.org/10.1016/j.bbr.2015.04.037>

Total number of authors:
5

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

Characterization of long-term motor deficits in the 6-OHDA model of Parkinson's disease in the common marmoset

M. Santana^{a,d,g}, T. Palmér^{b,c}, H. Simplício^{a,e}, R. Fuentes^{a,f}, P. Petersson^{b,*}

a Edmond and Lily Safra International Institute of Neuroscience of Natal, Macaíba, Brazil, 59280-000

b Integrative Neurophysiology and Neurotechnology, Neuronano Research Center, Department of Experimental Medical Sciences, Lund University, BMC F10, S-221 84 Lund, Sweden

c Mathematics LTH Centre for Mathematical Sciences, Faculty of Engineering, Lund University, S-22184 Lund, Sweden

d Psychobiology Program, Federal Univ. of Rio Grande do Norte, Natal, Brazil, 59078-970

e State Univ. of Rio Grande do Norte, Mossoró, Brazil, 59610-210

f Programa de Fisiologia y Biofísica, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, Independencia 1027, Santiago, Chile

g Bacharelado em Ciências Biológicas, Instituto de Ciências e Tecnologia das Águas, Universidade Federal do Oeste do Pará (UFOPA), Santarém, Pará, Brazil.

*) Corresponding author

Per.Petersson@med.lu.se

Ph: +46 46 222 05 78

Medicon Village, Building 404 A2

Scheelevägen 2

223 81 Lund, Sweden

Keywords: Animal models; Parkinson's disease; motor behavior

Research aimed at developing new therapies for Parkinson's disease (PD) critically depend on valid animal models of the disease that allows for repeated testing of motor disabilities over extended time periods. We here present an extensive characterization of a wide range of motor symptoms in the 6-OHDA marmoset model of PD when tested over several months. The severity of motor deficits was quantified in two ways: i) through manual scoring protocols appropriately adapted to include species specific motor behavior and ii) using automated quantitative motion tracking based on image processing of the digital video recordings. We show that the automated methods allow for rapid and reliable characterization of motor dysfunctions, thus complementing the manual scoring procedures, and that robust motor symptoms lasting for several months could be induced when using a two-stage neurotoxic lesioning procedure involving one hemisphere at a time. This non-human primate model of PD should therefore be well suited for long-term evaluation of novel therapies for treatment of PD.

1. Introduction

When investigating new therapeutic approaches for PD, researchers crucially depend on valid animal models of the disease and reliable methods to assess the symptoms displayed. Although several different animal models of PD exist, the preferred choices by many labs today are the 6-hydroxydopamine (6-OHDA) lesioned rat or the MPTP-lesioned non-human primate [1][2], since these models have proven to capture several important features of the disease. However, MPTP is a severe safety hazard to the personnel handling the animals and strict procedures and appropriate laboratory safety equipment are an absolute requirement [3]. Consequently, there have also been a number of studies aimed at developing a primate model of PD based on intracerebral 6-OHDA lesions which would minimize the risk of inadvertent toxic exposure for researchers and animal care taking personnel that is associated with systemic MPTP treatment [4][5][6][7]. In parallel with the ongoing efforts to improve the reliability and validity of PD animal models, more sophisticated and diverse methods to assess severity of PD symptoms in animals has also been a key objective in the methodological development for several labs [8][9]. Given that the relevance of preclinical research ultimately is dictated not only by the validity of the model, but also to a great extent by the reliability and sensitivity of the testing methods used, this work aims towards further improvement of the procedures used to assess symptoms in animal models of PD. In particular, when evaluating new potential therapies, for example, neuromodulatory approaches like deep brain stimulation [10][11], or spinal cord stimulation [12][13], robust testing procedures are needed to allow researchers to repetitively assess the severity of the symptoms displayed over long time periods in response to changes in therapeutic interventions.

To this end, we have here developed new methods for behavioral assessment of PD symptoms in the 6-OHDA lesioned common marmoset (*Callithrix jacchus*). These procedures include manual scoring of PD symptoms according to an adapted PD motor rating scale, and automated movement tracking procedures based on digital video recordings. Using these methods, a thorough characterization of changes in motor behavior in nine 6-OHDA lesioned marmoset monkeys were conducted over a time period of several months. By testing the animals in four different symptomatic stages in a step-wise lesioning procedure, different levels of motor

1
2
3
4 symptom severity could be characterized. The stages evaluated were: 1) intact state prior to
5 lesion; 2) after unilateral lesion; 3) after bilateral lesion; and 4) after bilateral lesion plus
6 treatment with the dopamine synthesis blocker alpha-methyl-p-tyrosine (AMPT). The order of
7 successive lesions and testing procedures in PD model are shown in Fig. 1A.
8
9

10 11 12 13 14 15 **2. Material and Methods** 16 17

18 19 *2.1. Animals and housing conditions* 20 21

22
23 Nine adult male common marmosets (*Callithrix jacchus*) 300-550g were used in the study. The
24 animals were housed in pairs in cages (1.0 x 1.0 x 2.3 m³) in a vivarium with natural light cycle
25 (~12/12 hours). Each cage have cover for rain and direct sun light, and the vivarium has a mobile
26 roof that can be opened or closed according to weather changes such as heavy rain. Common
27 marmosets are endemic to Northeast Brazil where the vivarium is located; thus ensuring suitable
28 temperature, humidity and light conditions. To enrich the housing environment, cages are
29 supplemented with elements such as sticks, tubes, ropes and ladders. Each cage has a small
30 wooden box used as nest for protection and sleeping. Animals are offered two meals a day
31 consisting of primate chow, local fruits, vegetables, mealworm larvae, gum arabic, dairy
32 products, grains, eggs, and meat under the supervision of a veterinarian.
33
34

35
36 All animal procedures were carried out according to approved protocols by AASDAP Ethics
37 Committee and strictly in accordance with the National Institute of Health Guide for the Care
38 and Use of Laboratory Animals (NIH Publications No. 80-23). This project was approved by
39 SISBIO/Brazilian Institute of Environment and Natural Resources (IBAMA) (No. 20795-2).
40
41
42
43
44
45
46
47
48
49
50

51 52 *2.2 Procedure for 6-OHDA injections* 53 54

55
56 The animals were initially sedated with ketamine (10-20 mg/kg ,i.m.) and atropine (0.05 mg/kg
57 i.m.) followed by deep anesthesia with isoflurane inhaled though a nose cone, to be finally
58 intubated with an endotracheal tube and ventilated with artificial ventilator to be maintained with
59
60
61
62
63
64
65

1
2
3
4 isoflurane 1-5% in oxygen at 1-1,5 L/min during the surgery. One mL of 6-OHDA hydrochloride
5 (4 mg/mL dissolved in 0.05% ascorbate/saline solution) was freshly prepared and stored
6 protected from light on ice before use. Five injections (2 μ L each) were made with a 32 gauge
7 Hamilton syringe at 0.5 μ L per minute into the medial forebrain bundle (MFB) in the following
8 locations (Anteroposterior/Mediolateral/Dorsoventral from the interaural midpoint): 6.5/1.2/6.0;
9 6.5/1.2/7.0; 6.5/2.2/6.5; 6.5/2.2/7.5; 6.5/3.2/8.0, which corresponds to a slightly modified version
10 of the protocol used by Annett et al. 1992 [5]. Anteroposterior coordinates were corrected
11 according to the dimensions of the skull of each animal based on the anatomy atlas by Stephan et
12 al., 1980 [14]. After each infusion, the needle was left in place for another 3 min to allow the
13 spread of the solution through the cerebral tissue at the exact area of interest.
14
15

16
17 During the 3-5 days following the surgery, the animals received non-steroid anti-inflammatory
18 analgesic, flunixin meglumine (1 mg/kg, s.c.) and dexamethasone (0.5 – 1 mg/kg, i.m.), and a
19 supplementary high-energy liquid diet. After eight weeks, the same procedures were repeated for
20 the second 6-OHDA lesion in the other hemisphere, as previously described by Mitchell et al.
21 1995 [4]. The second 6-OHDA lesion was made in the contralateral hemisphere to the preferred
22 limb.
23
24
25
26
27
28
29
30
31
32

33 34 35 2.3 *Adaptation to box and tower behavioral testing set-ups* 36 37

38
39 Animals were accustomed to the behavioral testing procedures in a step-wise manner. First,
40 while in their home cage, three times a week for two weeks, the animals were habituated to the
41 food rewards used: 2x5x10 mm³ (~60 mg) marshmallow pieces or mealworm (*Tenebrio molitor*)
42 depending on the preference of each marmoset. Second, animals were accustomed to the
43 transportation box (animals were allowed to explore the transportation box containing food baits
44 while being free to return to their home cage at any time). Once showing interest in the
45 transportation box, the animals were accustomed to a sound signaling entrance and another
46 sound signaling exit from the transportation box. Animals were then trained to exit from the
47 transportation box and explore the two different behavioral testing set-ups used in the study – a
48 transparent cubic acrylic box (0.45 x 0.45 x 0.45 m³) and a vertical tower (width x depth x
49 height: 0.36 x 0.37 x 2.20 m³) with seven horizontal bars located at different distances above
50 ground (0.1, 0.2, 0.4, 0.6, 0.9, 1.25, and 1.75 m; Fig. 1B) [15]. In this training, pieces of
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

marshmallows were placed on the floor of the box or on the bars in the tower testing set-up to encourage the animal to explore the environment. A disposable white sheet of ethylene-vinyl acetate foam covered the floor to preventing the animal from slipping. This training scheme was performed twice a week for two weeks (in a parallel study animals were trained to reach and grasp food rewards through holes in one of the walls - this behavior was not evaluated in the current study and the shelves were not baited). All the procedures were performed either between 10:30 and 12:00 or between 14:00 and 15:30, corresponding to the natural peak of motor activity (cf. Fig. 3C). The food rewards obtained during training of the task replaced the juice portion that the animals would normally receive in their home cages.

2.4 *Automated assessment of motor activity in home cage*

Spontaneous motor activity of two animals were collected using actimeters (Actiwatch Mini, CamNtech) worn in custom made vests inside their home cage. The actimeters collected data every two seconds for three consecutive days (72h) during the baseline, unilateral and bilateral periods. For the panel in Fig. 3B, the average raw motor activity of two consecutive 4am-6pm periods of the 72h-recording session is represented in relation to the date of the second 6-OHDA lesion (except for the unilateral lesion period of Monkey 6 where only one 4am-6pm period was used, since the data from the second period was not available). For the graphics of Fig. 3C, each recording was smoothed with a one-hour (1800 samples) moving average window sliding at every sample, divided in two full 24-hour periods, and the periods finally averaged.

2.5 *Manual PD scoring*

To evaluate the motor disability of the parkinsonian animals, we adapted the Unified Parkinson's Disease Rating Scale developed by Fahn and colleagues for the clinical setting [18] to fit aspects of non-human primate behavior based on previously developed procedures [16], [17].

The adapted scale consists of 16 categories scored from zero to three, which corresponds to absence of altered state to more intense symptomatology, respectively. Some categories involve

1
2
3
4 symptoms that were evaluated for each body part individually (i.e., limbs, trunk, head), each
5 receiving a maximum score of 3, thus, these categories could reach a maximum of 15 (“*Tremor*
6 *at rest*” and “*Tremor in motion*”) or 9 points (“*Fine motor skills*”), and were therefore
7
8 subsequently normalized to 3 in order to facilitate the direct comparison of different categories of
9 symptoms. Hence, the maximum total score of the scale is $16 \times 3 = 48$ points (Table 1).
10
11 The motor examination was performed in the animal’s home cage. Assessments occurred at two
12 times of day: in the morning (~ 9 am) or afternoon (~ 5 pm). All tests were done before meals.
13
14
15
16
17
18

19 The quantified categories are the following:

20 i) Tremor at rest

- 21
22 [0]: Absent
23
24 [1]: Occasional or detected rarely
25
26 [2]: Frequent or easily detected
27
28 [3]: Continuous and intense
29

30
31 ii) Tremor in motion

- 32
33 [0]: Absent
34
35 [1]: Rarely detected, present during action
36
37 [2]: Moderate amplitude, present during action
38
39 [3]: Moderate amplitude, can interfere with feeding
40
41

42 iii) Freezing

- 43
44 [0]: Unhindered to move the body and show normal use of the limbs, e.g., in
45 finding and grasping marshmallows in the reaching task
46
47 [1]: Difficulties in starting to walk, or in the initiation of particular movements. For
48 example, when reaching for a marshmallow, the start of the reaching movement is
49 delayed. In these cases the freezing episodes are short
50
51 [2]: Same as in [1], but the freezing episodes have a longer duration - between 5
52 and 10 seconds
53
54 [3]: Same as in [1], but freezing episodes last over 10 seconds
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 iv) Gait and locomotion
5

6 [0]: Walks normally according to pre-lesion locomotion patterns, with symmetrical
7 limb use
8

9 [1]: Shows reduced walking activity and walks with mild asymmetry
10

11 [2]: Walks slowly, with asymmetry, and occasionally drags a limb (usually a
12 hindlimb)
13

14 [3]: Unable to walk
15
16
17
18

19 v) Fine motor skills (scored for each arm independently)
20

21 [0]: Normal ability to grasp marshmallows
22

23 [1]: Grasps with difficulty
24

25 [2]: Grasps with difficulty and requires one arm to support the stance while using
26 the other to grab the marshmallow
27

28 [3]: Totally unable to grasp marshmallows
29
30
31

32 vi) Bradykinesia (scored independently for limbs and trunk)
33

34 [0]: No difficulty in initiating or performing rapid and precise movements.
35

36 [1]: Difficulties in initiating movements and displays smoother and slower
37 movements when reaching for marshmallows or moving around spontaneously
38

39 [2]: Clear delay in initiating movements and shows a marked slowing of
40 movements in reaching and in spontaneous motor activity
41

42 [3]: Totally immobile
43
44
45

46 vii) Hypokinesia
47

48 [0]: Moves freely and is alert and responsive
49

50 [1]: Reduced activity, moves with less speed
51

52 [2]: Low spontaneous activity, moves when provoked
53

54 [3]: Totally immobile
55
56

57 viii) Rigidity
58

59 [0]: Moves freely; coordinated actions, absence of rigidity
60
61
62
63
64
65

- 1
2
3
4 [1]: Mild rigidity or rigidity apparent only when other body parts are moving
5
6 [2]: Striking stiffness, yet some complete movements are performed easily
7
8 [3]: Severe rigidity, no movements are performed or movements appear incomplete
9

10
11 ix) Body balance (Spontaneous behavior)

- 12 [0]: Normal stance and coordination
13
14 [1]: Compromised coordination, but is able to change from quadrupedalism to
15 bipedalism without falling
16
17 [2]: Compromised coordination, unstable locomotion with occasional falls
18
19 [3]: Face down or lying in supine position unable to maintain any kind of stance
20
21
22
23

24 x) Body balance (Induced behavior elicited by food offering)

- 25 [0]: Normal stance and coordination
26
27 [1]: Compromised coordination but changes from quadrupedalism to bipedalism,
28 without falling
29
30 [2]: Compromised coordination, unstable locomotion with occasional falls
31
32 [3]: Face down or lying in supine position unable to maintain any kind of stance
33
34
35
36

37 xi) Posture

- 38 [0]: Normal posture
39
40 [1]: Somewhat altered posture when standing, such as wider positioning of limbs.
41 Resting with limbs and tail in abnormal body position
42
43 [2]: Hunched posture, abnormal trunk position; abnormal head posture (neck flexed
44 or inclined to one side)
45
46 [3]: Unable to maintain posture, lying in supine or lateral position
47
48
49
50

51 xii) Startle response

- 52 [0]: Immediate, robust threat response
53
54 [1]: Slightly diminished or delayed response, threats with open mouth
55
56 [2]: Minimal or much delayed response, no open mouth threat
57
58 [3]: No response to provocation
59
60
61
62
63
64
65

1
2
3
4
5
6 xiii) Climbing

7 [0]: Normal

8 [1]: Climbs with difficulty. Slow on the branches and home cage mesh. No falling

9 [2]: Very compromised. Climbs branches and cage mesh with great effort. Falling
10 may occurs

11 [3]: Not able to climb
12
13
14
15
16
17
18

19 xiv) Gross motor skills (scored for each arm independently)

20 [0]: Normal limb use when grasping larger objects

21 [1]: Reduced ability to grasp larger objects to support body weight

22 [2]: Rarely is able to grasp larger objects to support body weight

23 [3]: Unable to grasp and hold large objects/structures
24
25
26
27
28
29

30 xv) Facial Expression

31 [0]: Normal

32 [1]: Slightly apparent decrease of facial expression (hypomimia)

33 [2]: Moderate hypomimia with lips separated during brief moments

34 [3]: Fixed face, severe or total loss of facial expression, lips separated in 6 mm or
35 more
36
37
38
39
40
41
42

43 xvi) Vocalization

44 [0]: Normal quantity

45 [1]: Spontaneous vocalization reduced

46 [2]: Induced vocalization reduced

47 [3]: Absent
48
49
50
51
52
53

54 For the categories "*Climbing*", "*Bradykinesia*", "*Fine Motor Skills*" and "*Body Balance*
55 (*Induced*)," mealworms or a piece of marshmallow were offered with tweezers to induce the
56 desired motor behavior. For the evaluation of the category "*Rigidity*", a blunt forceps was
57 presented to the animal. Since the animal associates the forceps with food offering, it would
58
59
60
61
62
63
64
65

1
2
3
4 normally grab it. Following grasping of the forceps, gentle ‘push and pull’ movements were
5 made to evaluate the level of stiffness of the forelimb. The procedure was repeated for both
6 forelimbs.
7
8

9
10 In experiments involving AMPT-treatment, manual PD-scoring was performed six hours after
11 the first injection.
12
13
14
15
16

17 *2.6 Automated tracking procedures*

18
19

20 Digital video recording were performed in the two testing set-ups. Two cameras (AVT –
21 Stingray F033C, 80 fps) were used for digital video recordings in the acrylic box, from top and
22 side views whereas tower activity was recorded using a single front view camera (AVT –
23 Stingray F033C, 80 fps) (Fig. 1B). Motion tracking in the two setups was performed using
24 similar methods. Software tools were developed in MATLAB and included mex-
25 implementations (MATLAB compiled c-code; Mathworks Inc.). Constant light conditions during
26 each recording session eliminated the need for advanced background models. Hence, a simple
27 algorithm where each pixel is modeled as belonging to one of two Gaussian distributions was
28 employed. The two distributions were estimated for each pixel by iterating through a sufficient
29 number of frames of the video and updating the estimated parameters of the most probable
30 distribution. In this case the background is contained in the brighter distributions, as the animals’
31 image in these experiments was always darker than the actual background. After subtracting the
32 background, the resulting foreground images were used in the shape analysis. By assuming that
33 the two-dimensional image of the monkey in each camera plane is approximately elliptically
34 shaped, the position and orientation of the animal could be estimated by the position and
35 orientation of the three-dimensional ellipsoid that best fitted the foreground images. Given a
36 measured foreground image F in a given camera and an estimated foreground image M generated
37 by projection of the ellipsoid onto the camera plane, the matching quality is defined as
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

$$56 \quad q = \frac{\sum_{i,j} \min(F(i,j), M(i,j))}{\sum_{i,j} \max(F(i,j), M(i,j))}$$

57
58
59
60
61
62
63
64
65

1
2
3
4 where i and j are the pixel coordinates. When multiple cameras were used, the combined quality
5 measure was created by multiplying the individual q-scores. Note that the foreground image M is
6 not actually computed, but instead the quality measure was computed using the projected quadric
7 matrix for the ellipsoid. Movement tracking in time was then carried out by using the last known
8 location to initiate estimation for a given frame followed by step-wise improvements of the
9 matching quality by gradual adjustments of the parameters of the estimated ellipsoid. These
10 calculations were performed for every frame in the video, resulting in the vectors $(x, y, z, \theta_x, \theta_y,$
11 $\theta_z)$ describing the position and orientation of the estimated ellipsoid. Each vector will therefore
12 be of the length N , where N is the number of frames in the video. Note that in the tower
13 experiments, the z -coordinate was fixed and not estimated due to the use of only one camera (see
14 <http://homepages.inf.ed.ac.uk/rbf/VAIB14PAPERS/palmer.pdf> for technical details on tracking
15 procedures).

26 27 28 *2.7 Automated extraction of kinematic parameters presented in plots*

29
30
31 Relevant metrics summarizing changes in kinematic parameters over the different experimental
32 conditions were constructed from the tracking data. From the (x,y,z) position vectors, speed was
33 estimated as the Euclidean distance between (x_i, y_i, z_i) and $(x_{i+k}, y_{i+k}, z_{i+k})$, divided by k ,
34 frame number difference, and multiplied by the time resolution. Locomotion bouts were detected
35 by applying a threshold on the acquired speed vectors. A locomotion bout was defined as the
36 period of time where instantaneous speed was uninterruptedly greater than the chosen threshold.
37 To improve robustness, multiple values of k were used for this detection, and all different
38 estimates of the speed at a time have to be greater than the chosen threshold (approximately
39 corresponding to a speed of 0.04 m/s).

40
41 From each locomotion bout, a number of different parameters were obtained: maximum speed,
42 average speed, distance covered, duration and maximal acceleration.
43
44
45
46
47

48 49 50 51 52 53 54 55 *2.8 Tyrosine-hydroxylase staining and quantification*

1
2
3
4 After the period of the experiments, the animals were sacrificed by intracardiac perfusion after
5 deep sedation with ketamine (40 mg/kg i.m.); (xylazine 0.04 mg/kg i.m.) and atropine (0.05
6 mg/kg i.m.). Intracardiac perfusion was performed with 0.9% saline solution and heparin at 37
7 °C, followed by 4% paraformaldehyde in phosphate buffer, 0.1 M (pH 7.4), cooled to 4 °C. The
8 brains were removed and postfixed in the same solution for 2h, washed in 0.1 M phosphate
9 buffer (pH 7.4) at 4 °C for 24 hours, cryoprotected in 20% following 30 % sucrose solution at 4
10 °C, and finally rapidly frozen for cryostat embedding in Tissue-Tek medium. The brains were
11 kept in a freezer at -80 °C until sectioned coronally at 50 µm in a cryostat.
12

13
14
15
16
17
18
19 Immunohistochemical staining was performed free-floating or on sections mounted directly on
20 electrically charged glass slides. The sections were processed for immunohistochemical
21 detection of tyrosine hydroxylase (TH) in substantia nigra and in striatal regions using
22 modifications of the protocol of Eslamboli et al. (2003) [19].
23

24
25
26 The sections were washed in 0.1M phosphate buffer (PB) for 5 minutes. Then, incubated in 1%
27 hydrogen peroxide/methanol solution for 20 minutes to remove endogenous peroxidase activity,
28 and rinsed in 0.05% phosphate buffer-Tween 0.05% (PB-T) for 5 minutes. Thereafter, the
29 sections were confined with the aid of a hydrophobic PAP pen and incubated in 10% goat normal
30 serum diluted in 0,1 PB for 30 minutes. Excess serum was removed and sections were incubated
31 in the primary anti-TH (rabbit polyclonal antibody; 1:500; diluted in normal serum/PB)
32 overnight at room temperature in a humidity chamber to prevent air-drying of the tissue sections.
33 The sections were washed with PB-T (5 min) and incubated in biotinylated goat anti-rabbit
34 secondary antibody (1:200, diluted in PB; Vector Labs) for two hours. After that, the sections
35 were washed again with PB-T (5 min) and incubated in avidin-biotin-peroxidase solution
36 (Vectastain Standard ABC kit, Vector Laboratories) for one hour.
37

38
39
40
41
42
43
44
45
46 After removal of the ABC solution, the sections were washed in PB (5 min) and placed in a
47 solution containing 0.03% 3,3'-diaminobenzidine tetrahydrochloride hydrate (DAB) (Sigma) and
48 0.001% hydrogen peroxide in 0.1M PB. The reaction was monitored in a light microscope. The
49 sections were washed and slides were left to dry overnight. After dehydration through a series of
50 graded alcohols and clearance in xylene, the slides were cover-slipped using Entellan mounting
51 medium.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2.9 Quantification of striatal and nigral tyrosine hydroxylase immunoreactivity

The tissue samples were mounted and photographed using a microscope with the same camera configuration and under identical illumination conditions. TH reactivity in both striatum (caudate and putamen) and in substantia nigra pars compacta (SNc) was assessed by computer densitometry using digital images captured from a camera (CX9000, MBF Bioscience) attached to the microscope (light field Nikon Eclipse 80i - 10x and 20x objectives). TH-reactivity across the striatum was assessed by optical densitometry using ImageJ software (NIH, <http://rsb.info.nih.gov/ij/>). Measurements were obtained using a 0.2 mm² square window positioned in different regions throughout the striatum (60 samples per striatum and animal). To reduce the effects of within-group variability, a normalized scale based on the reactivity for TH of the internal capsule (white matter) was adopted (average over measurements of 10 different sites using the same window). For each animal, a contrast index was calculated according to the equation: $C = (G-W)/(G+W)$ [20], in which G is the average optical density of striatal tissue, and W is the optical density of the white matter (internal capsule). To count TH-labeled cells, we used at least three sections per animal. For the different positions along the rostral-caudal axis (rostral, central and caudal area), the boundaries of the SNc were defined in each section according to the atlas by Paxinos et al. [21] and the area of the SNc was calculated using the sections from control animals (it was not possible to identify the SNc contours in the lesioned animals because of the substantial loss of dopaminergic neurons from 6-OHDA treatment). Cells labeled with TH within the defined areas were subsequently counted (StereoInvestigator system, MBF Bioscience Inc) and the resulting cell densities were expressed as TH⁺ cells/mm².

2.10 Statistical analyses

The statistical tests used in the study are specified in the main text and in the figure legends together with the data used for the respective test. Analyses of significance were performed using either Matlab functions or GraphPad Prism 5.01 software.

3. Results

3.1 Acute effects of 6-OHDA lesions

Immediately following the first lesion, animals showed a rigidity in the limbs and visuospatial neglect contralateral to the lesioned hemisphere and head position deviation ipsilateral to it [22]. In addition, animals showed ipsilateral body rotation while trying to ambulate, and difficulty to use the forelimb contralateral to the lesioned hemisphere. In spite of these evident motor symptoms the animals were still able to feed themselves in their home cages (as indicated by a < 10% weight loss following surgeries). At least eight weeks later, the animals were exposed to a second injection of 6-OHDA in the opposite hemisphere (Fig. 1A). Directly following this second lesion, animals generally showed similar but more severe motor impairments, in some cases requiring special care when animals had difficulties feeding themselves to ensure weight loss would not exceed 10% of total body weight during the first two weeks following surgery [23]. Animals were allowed to recover for two weeks before assessments of PD symptoms commenced.

3.2 Evaluation of motor symptoms using an adapted PD motor disability rating scale

In the manual assessment of PD-symptoms a total of 16 different categories were evaluated: (1) resting tremor, which was not observed in this model; (2) tremor in motion and sporadic postural tremor; (3) episodes of freezing - brief periods of sudden immobility when initiating quadrupedal locomotion or goal directed reaching; (4) uncoordinated gait - inaccurate positioning of the limbs and wobbling of the trunk during locomotion (in the literature referred to as clumsy, poor-balanced gait; Eslamboli, 2003); (5) deficits in fine motor skills - difficulty in using arms to grab any food offered (in some animals the weakness was exacerbated by a worsening of gross motor skills, see below); (6) bradykinesia - noticeable slowing of the execution of movements; (7) hypokinesia - a general reduction in motor activity (motility, grooming, climbing); (8) rigidity - particularly noticeable in forelimbs during extension; (9 and 10) body balance - abnormal body positions and difficulty to rest on branches; (11) hunched posture; (12) a slowed startle response

1
2
3
4 - animals would not respond to alarm vocal signals from mates; (13) slowed climbing; (14) loss
5 of gross motor skills - for example, inability to grasp branches; (15) episodes of hypomimia -
6 reduction of the marmoset's typical behavior of maintaining eye contact and impaired inability to
7 display facial expression in response to interaction with care givers; and (16) lack or decrease of
8 vocalizations (marmosets use vocalizations abundantly to communicate between them).
9

10 For each of these 16 categories, the severity of motor disability was repeatedly evaluated in
11 every individual in a total of eight animals under different degrees of Parkinsonism. Following
12 the first lesion, stable parkinsonian symptoms were observed in all individuals over the more
13 than 8 week long testing period (average score [mean±SEM], week 1-8: 6.9±1.0; Fig. 2A, left).
14 After the second lesion, symptoms were on average more severe compared to the first unilateral
15 lesion during the corresponding assessment period (average score week 1-8: 12.6±0.7, Fig. 2A,
16 right). Animals were then monitored for another few months and persistent symptoms were
17 confirmed. However, during these extended testing periods a certain degree of spontaneous
18 recovery was observed resulting in a gradual decline of the total PD score over a 32 week period
19 (Fig. 2A). Severe Parkinsonism could, however, nevertheless always be transiently reinstated for
20 ~18 h through systemic treatment with the dopamine synthesis blocker AMPT (average score
21 under AMPT effect for week 1-16: 24.6±1.8; week 17-32: 19.7±1.0. Interestingly, the degree of
22 functional recovery varied substantially between different types of motor symptoms. When
23 analyzing the PD-scores for each category of symptoms divided into 8-week periods following
24 the second lesion it became evident that for example symptoms related to locomotion and body
25 balance during spontaneous behavior showed negligible improvements over time (Fig. 2B).
26 These findings indicate that quantitative assessments of spontaneous locomotor behavior could
27 be particularly useful in experiments where testing periods lasting over several months are
28 required.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 *3.3 Twenty-four hour recordings of motility in the home cage*

53
54

55 As a complement to the detailed manual assessments of dysfunctions in motor behavior, the
56 overall spontaneous motor activity during 72h periods in the home cage was also recorded in two
57 animals. It was found that the absolute amount of motor activity was clearly decreased following
58
59
60
61
62
63
64
65

1
2
3
4 the first and second lesion, with a relative decrease after unilateral lesion corresponding to: -44%
5 and -39%, and after bilateral lesion: -78% and -36% for the two monkeys, respectively (Fig. 3A,
6 B). At the same time, the characteristic variations in the relative amount of motor activity
7 displayed throughout the day-night cycle was comparatively preserved also in the parkinsonian
8 state (Fig. 3C).
9

10 11 12 13 14 15 *3.4 Automatic assessment of locomotive activity in the Tower testing set-up* 16 17

18
19 In each testing session, the spontaneous locomotion of the animal was recorded for 5 min in 120
20 testing sessions in a total of 7 animals. The distance travelled during the testing session was
21 subdivided into vertical and horizontal translation (Fig. 4A). It was evident that intact animals
22 were considerably more active than lesioned animals and that the distance travelled successively
23 declined in the more severe PD models (Fig. 4B). On average the distance traveled
24 (horizontal/vertical) in meters per minutes for animals in the four different stages of
25 Parkinsonism were (mean±SD), intact: 1.51±0.52/2.54±1.41, hemilesion: 0.81±0.41/1.16±0.85,
26 bilateral: 0.50±0.28/0.49±0.29, bilateral+AMPT: 0.17±0.06/0.18±0.08 (Fig. 4B; p<0.05,
27 Kruskal-Wallis). Furthermore, healthy individuals preferred staying on the bars positioned
28 relatively higher up in the tower in contrast to the parkinsonian animals resulting in significant
29 differences in mean expectation values in height over ground for the four groups (Fig. 4C;
30 p<0.05, Kruskal-Wallis). Finally, we also observed that when moving between different heights,
31 healthy individuals often displayed longer uninterrupted movement bouts involving multiple
32 transitions between different levels, whereas the lesioned animals moved more frequently one
33 level at a time (fraction of multi-level transitions for the four groups were: 0.23±0.14, 0.09±0.07,
34 0.03±0.03, 0.05±0.10; p<0.05, Kruskal-Wallis Fig. 4D).
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 *3.5 Automatic assessment of locomotive activity in the Box testing set-up* 53 54

55 Spontaneous locomotion in the transparent cubical box was quantified from ~5min recordings in
56 a total of 120 testing sessions in 4 animals. Similarly to the tower test, the distance travelled was
57 subdivided into vertical and horizontal translation and in agreement with the behavior in the
58
59
60
61
62
63
64
65

1
2
3
4 tower the distance travelled was clearly reduced in the more severe PD models. On average the
5 distance travelled (horizontal/vertical) per minute for the four groups were (mean±SD), intact:
6 2.34±0.86/0.55±0.39, hemilesion: 0.96±0.40/0.36±0.41, bilateral: 0.23±0.12/0.04±0.03,
7
8 bilateral+AMPT: 0.14±0.10/0.04±0.05 (Fig. 5A; $p<0.01$ for both horizontal and vertical distance,
9 Kruskal-Wallis). A more detailed analysis of the locomotion bouts revealed further differences in
10 the pattern of locomotion. We found that 1) bout duration, 2) bout maximum speed, 3) bout
11 distance, as well as 4) frequency by which bouts of locomotion were displayed were all reduced
12 in parkinsonian animals (Fig. 5B). Finally, in order to verify that the motor deficits observed
13 were stable over extended time periods, the individual experiments were ordered and analyzed
14 with respect to the time of assessment in relation to the two lesion procedures. To eliminate any
15 inter-individual variability, all the analyzed features of the locomotive behavior were normalized
16 to the motor behavior displayed by each individual during baseline conditions. While slight
17 variations were found between different recording sessions during each of the three conditions, a
18 much greater difference was observed between intact, hemilesioned and bilaterally lesioned
19 animals. Notably, these differences persisted over several months and were found to be
20 particularly evident for bout distance and the frequency by which bouts of locomotion were
21 displayed (Fig. 5C).
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 *3.6 Immunohistochemical verification of 6-OHDA lesions*

40
41
42 Subsequent to these extensive characterizations of behavioral changes during parkinsonian
43 conditions, *post mortem* tissue analyzes were performed. Immunohistochemistry for tyrosine
44 hydroxylase (TH) was used to quantify the extent of the lesions (Fig. 6A, B). A reduction in the
45 number of TH-positive cells of the midbrain dopaminergic neurons projecting to the forebrain in
46 the lesioned hemispheres was confirmed. The cell densities (number of cells/mm²) were
47 (mean±SEM), 57.66 ± 6.23 and 139.01±12.13 in bilaterally lesioned and control animals,
48 respectively ($P<0.0001$, $U=28$, Mann-Whitney U-test; Fig. 6C, bottom panel). The axonal
49 terminal density of TH positive cells projecting to the caudate-putamen was also quantified. A
50 contrast index was used to quantify the TH-staining in relation to background staining (see
51 Methods for detail) showing a significant reduction of TH-immunoreactivity in lesioned animals
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 vs. controls in both the caudate nucleus (0.155 ± 0.01 vs. 0.254 ± 0.02 ; $P<0.05$, $U=112$, Mann-
5
6 Whitney U-test) and in putamen (0.135 ± 0.02 vs. 0.213 ± 0.02 ; $P<0.05$, $U=109$, Mann-Whitney U-
7
8 test; Fig. 6C, top panel). Taken together, the average staining intensity of terminals in the
9
10 caudate-putamen of lesioned animals was 44%, and the density of stained midbrain cells 41%
11
12 compared to intact animals.
13
14

15 16 **4. Discussion** 17

18
19 Non-human primate models have a key role in PD-research aimed at understanding the
20
21 underlying pathophysiology of the disease, as well as for the development of new treatment
22
23 strategies. Whereas experiments in rodents in many cases can provide important insights in the
24
25 early phase of basic PD-research, results are not always transferable to humans. In particular, the
26
27 large difference in overall neuroanatomical complexity between the rodent and primate central
28
29 nervous system can sometimes make findings in rats and mice less clinically relevant. The
30
31 possibility to perform large scale experiments in the MPTP-treated macaque - which by many
32
33 researchers is regarded as the most valid model of PD - is on the other hand very limited due to
34
35 the high costs associated with housing and treating these larger primates and the safety
36
37 precautions required for safe handling of these animals in order to avoid inadvertent neurotoxic
38
39 exposure. In this perspective the 6-OHDA marmoset model of PD, which we here thoroughly
40
41 characterized, may present a valuable complement.

42
43 Investigations aimed at developing prospective treatments for PD generally demand long
44
45 evaluation periods, it was therefore important to systematically evaluate the marmosets with
46
47 respect to a range of motor deficits over a time period of several months following lesions. While
48
49 a recovery of certain motor functions was observed after about three months in the detailed
50
51 manual PD-scoring assessments, other symptoms remained stable also after more than six
52
53 months following lesions, indicating that this model may indeed be useful for the purpose of
54
55 evaluating novel PD therapies under chronic disease conditions. Moreover, in experimental
56
57 situations where severe Parkinsonism is desired, the additional pharmacological treatment with
58
59 the dopamine synthesis inhibitor AMPT reproducibly induced marked motor disability in all
60
61 animals tested. In spite of the comparatively severe symptoms that are transiently induced under
62
63
64
65

1
2
3
4 such conditions, these tests were well tolerated and could be repeated multiple times in all
5
6 animals.

7
8
9 The use of automated procedures for the analysis of spontaneous locomotive behavior provided
10 important information on motor dysfunctions adding to the outcome of the manual scoring of PD
11 symptoms. Both in 24h home cage recordings and in the shorter testing sessions in the Tower
12 and Box set-ups, consistent differences between the different parkinsonian states were observed.
13
14 In fact, in certain respects the automated procedures showed a greater sensitivity than the manual
15 scoring, as evident from the persistent reduction in locomotor-related kinematic parameters such
16 as bout frequency and total distance travelled, which could be established over very long time
17 periods following the second lesion, even when a functional recovery of some other motor
18 functions resulted in a gradual decrease of the total PD-score over time. Taken together, given
19 the complex pattern of motor dysfunctions revealed by the manual scoring procedures, on the
20 one hand, and the robust identification of motor symptoms using the automatic techniques, on
21 the other, the current findings suggest that a combined automatic/manual approach is preferable
22 in order to capture the full range of PD motor symptoms over extended time periods in this
23 model of PD. It can be concluded that, using the methods developed herein, the two-stage 6-
24 OHDA marmoset model of PD provides a robust and reliable primate model of PD lasting for
25 periods of months that can potentially have an important role in the future development of novel
26 therapies.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 **Acknowledgments**

44
45 We are thankful to Miguel Nicolelis for creating the research environment needed to conduct this
46 research and for supporting our work. We would also like to thank Marcelo Carvalho for
47 technical support in histology. This research was supported by The Michael J. Fox Foundation
48 for Parkinson's Research, FINEP 01.06.1092.00, and INCEMAQ – Program of National
49 Institutes of Science and Technology of CNPq/MCT; CAPES; AASDAP - Alberto Santos
50 Dumont Association for Research Support; The Swedish Research Council [#325-2011-6441],
51 the Olle Engkvist, Parkinson Research, Crafoord, Åke Wiberg, Magnus Bergvall, Kockska and
52 Segerfalk Foundation. NIH Transformative award (R01-NS073125-03).
53
54
55
56
57
58

59 **References**

60
61
62
63
64
65

- 1
2
3
4 [1] W. Dauer, S. Przedborski, Parkinson's Disease: Mechanisms and Models, *Neuron*. 39
5 (2003) 889–909. <http://linkinghub.elsevier.com/retrieve/pii/S0896627303005683>.
6
7
8 [2] A. Eslamboli, Marmoset monkey models of Parkinson's disease: Which model, when and
9 why?, *Brain Res. Bull.* 68 (2005) 140–149.
10 doi:<http://dx.doi.org/10.1016/j.brainresbull.2005.08.005>.
11
12 [3] S. Przedborski, V. Jackson-lewis, A.B. Naini, G. Petzinger, R. Miller, M. Akram, The
13 parkinsonian toxin (MPTP): a technical review of its utility and safety, (2001) 1265–
14 1274.
15
16 [4] C. and B.J. Mitchell, I J., Hughes, N., Carroll, Reversal of parkinsonian symptoms by
17 intrastriatal and systemic manipulations of excitatory amino acid and dopamine
18 transmission in the bilateral 6-OHDA lesioned marmoset, *Behav. Pharmacol. Pharmacol.*
19 6 (1995) 492–507.
20
21 [5] L.E. Annett, D.C. Rogers, T.D. Hernandez, S.B. Dunnett, Behavioural analysis of
22 unilateral monoamine depletion in the marmoset., *Brain*. 115 (Pt 3 (1992) 825–56.
23 <http://www.ncbi.nlm.nih.gov/pubmed/1352726>.
24
25 [6] U. Ungerstedt, 6-Hydroxydopamine-Induced Degeneration of the Nigrostriatal Dopamine
26 Pathway: the Turning Syndrome., *Pharmacol. Ther. B.* 2 (1976) 37–40.
27 <http://www.ncbi.nlm.nih.gov/pubmed/772721>.
28
29 [7] R. Klintenber, J. Arts, M. Jongsma, H. Wikström, L. Gunne, P.E. Andrén, Motor effects
30 of a dopamine stabilizer (GMC1111) in primate models of Parkinson and
31 hemiparkinsonism, *Eur. J. Pharmacol.* 459 (2003) 231–237. doi:10.1016/S0014-
32 2999(02)02817-0.
33
34 [8] A. Eslamboli, M. Romero-Ramos, C. Burger, T. Bjorklund, N. Muzyczka, R.J. Mandel, et
35 al., Long-term consequences of human alpha-synuclein overexpression in the primate
36 ventral midbrain, *Brain* . 130 (2007) 799–815.
37 <http://brain.oxfordjournals.org/content/130/3/799.abstract>.
38
39 [9] H. Okano, K. Hikishima, A. Iriki, E. Sasaki, The common marmoset as a novel animal
40 model system for biomedical and neuroscience research applications, *Semin. Fetal*
41 *Neonatal Med.* 17 (2012) 336–340.
42 <http://linkinghub.elsevier.com/retrieve/pii/S1744165X12000844?showall=true>.
43
44 [10] A.L. Benabid, P. Pollak, A. Louveau, S. Henry, J. de Rougemont, Combined
45 (thalamotomy and stimulation) stereotactic surgery of the VIM thalamic nucleus for
46 bilateral Parkinson disease, *Appl Neurophysiol.* 50 (1987) 344–346.
47
48 [11] A.L. Benabid, P. Pollak, D. Hoffmann, C. Gervason, M. Hommel, J.E. Perret, et al., Long-
49 term suppression of tremor by chronic stimulation of the ventral intermediate thalamic
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- nucleus, *Lancet*. 337 (1991) 403–406. doi:[http://dx.doi.org/10.1016/0140-6736\(91\)91175-T](http://dx.doi.org/10.1016/0140-6736(91)91175-T).
- [12] R. Fuentes, P. Petersson, M.A.L. Nicolelis, Restoration of locomotive function in Parkinson's disease by spinal cord stimulation: mechanistic approach., *Eur. J. Neurosci*. 32 (2010) 1100–8. doi:10.1111/j.1460-9568.2010.07417.x.
- [13] R. Fuentes, P. Petersson, W.B. Siesser, M.G. Caron, M.A.L. Nicolelis, Spinal cord stimulation restores locomotion in animal models of Parkinson's disease., *Science*. 323 (2009) 1578–82. doi:10.1126/science.1164901.
- [14] H. (Heinz) Stephan, G. (Georg) Baron, W.K. (Walter K. Schwerdtfeger, *The brain of the common marmoset (Callithrix jacchus) : a stereotaxic atlas / H. Stephan, G. Baron, W.K. Schwerdtfeger, (1980).*
- [15] P.S. Verhave, R. a. P. Vanwersch, H.P.M. van Helden, A.B. Smit, I.H.C.H.M. Philippens, Two new test methods to quantify motor deficits in a marmoset model for Parkinson's disease, *Behav. Brain Res*. 200 (2009) 214–219. doi:10.1016/j.bbr.2009.01.022.
- [16] K.S. Bankiewicz, R. Sanchez-Pernaute, Y. Oiwa, M. Kohutnicka, A. Cummins, J. Eberling, *Preclinical Models of Parkinson's Disease*, in: *Curr. Protoc. Neurosci.*, John Wiley & Sons, Inc., 2001. doi:10.1002/0471142301.ns0904s09.
- [17] M.J. Hansard, L.A. Smith, M.J. Jackson, S.C. Cheetham, P. Jenner, Dopamine, but not norepinephrine or serotonin, reuptake inhibition reverses motor deficits in 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-treated primates., *J. Pharmacol. Exp. Ther*. 303 (2002) 952–8. doi:10.1124/jpet.102.039743.
- [18] S. Fahn, R.L. Elton, U.D. Committee, others, Unified Parkinson's disease rating scale, *Recent Dev. Park. Dis*. 2 (1987) 153–163.
- [19] A. Eslamboli, H.F. Baker, R.M. Ridley, L.E. Annett, Sensorimotor deficits in a unilateral intrastriatal 6-OHDA partial lesion model of Parkinson's disease in marmoset monkeys., *Exp. Neurol*. 183 (2003) 418–29. <http://www.ncbi.nlm.nih.gov/pubmed/14552882> (accessed January 02, 2015).
- [20] M.A.M. Freire, E. Morya, J. Faber, J.R. Santos, J.S. Guimaraes, N.A.M. Lemos, et al., Comprehensive Analysis of Tissue Preservation and Recording Quality from Chronic Multielectrode Implants, *PLoS One*. 6 (2011) e27554. doi:10.1371/journal.pone.0027554.
- [21] H. Paxinos, HG., Watson, ., Petrides, M., Rosa, M., Tokuno, *The Marmoset Brain in Stereotaxic Coordinates*, 2012.
- [22] L.E. Annett, D.C. Rogers, T.D. Hernandez, S.B. Dunnett, Behavioural analysis of unilateral monoamine depletion in the marmoset., *Brain*. 115 (Pt 3 (1992) 825–56.

1
2
3
4 [23] I.J. Mitchell, C.B. Carroll, Reversal of parkinsonian symptoms in primates by antagonism
5 of excitatory amino acid transmission: potential mechanisms of action., *Neurosci.*
6 *Biobehav. Rev.* 21 (1997) 469–75. <http://www.ncbi.nlm.nih.gov/pubmed/9195604>.
7
8
9

10 11 12 13 14 **LEGENDS**

15 16 **Table 1**

17
18
19 Summary of assessments performed in the nine male marmosets included in the study. Animals
20 exposed to bilateral lesions were generally also assessed following the first lesion providing
21 additional data to the hemilesioned group. When multiple tests were performed in the same
22 animal in the Tower/Box set-up and through manual scoring, assessments were made during the
23 same day to facilitate direct comparisons.
24
25
26
27
28
29

30
31
32
33 **Fig. 1.** Description of experimental procedures. (A) Timeline of experimental procedures. The
34 two-stage 6-OHDA bilateral lesion procedure allowed for repetitive assessment of motor
35 symptoms at gradually more severe stages of Parkinsonism over extended time periods. (B)
36 Spontaneous locomotion was evaluated in two testing chambers designed to capture different
37 types of locomotive behavior, including both horizontal and vertical locomotion in both set-ups.
38 Left: Tower – two examples of typical movement bouts between bars located at different heights
39 are illustrated, blue lines denote the tracked movement traces and the red ellipses the position of
40 the thorax as estimated by the image system. 3D image shows the total amount of locomotion
41 displayed during a typical 5 min recording period in a healthy individual (Time represented in
42 color code ranging from dark blue (t=0) to red (t=5 min), Right: Box - side and top view,
43 respectively (color codes as in Tower).
44
45
46
47
48
49
50
51
52
53

54
55
56
57 **Fig. 2.** Manual scoring of motor impairments. (A) Total motor disability score of individual
58 animals during ten weeks following the first unilateral lesion (n=4, yellow), 32 weeks following
59
60
61
62
63
64
65

1
2
3
4 the second lesion (n=6), and under additional treatment with the dopamine synthesis inhibitor
5 AMPT (6h after 2x250mg/kg AMPT; n=6). On the y-axis, 48 points represents the highest
6 possible total score when the partial scores of the 16 categories are added and zero corresponds
7 to pre-lesion behavior for each individual. The average score during each testing period and
8 condition is represented by the thick horizontal lines. (B) Normalized scores of motor
9 impairment after the second lesion divided by symptom category and testing period, week: 1-8,
10 9-16, and 17-24 after lesion, and under the additional effect of AMPT (week 1-32 after lesion;
11 mean values shown, error bars represents S.E.M.). Significant differences in average scores were
12 found week [1-8] vs. [9-16] (α , $p<0.01$), [1-8] vs. [17-24] (β , $p<0.01$), [1-8] vs. [AMPT] (*,
13 $p<0.05$), [9-16] vs. [AMPT] (#, $p<0.01$), and [17-24] vs. [AMPT] (d , $p<0.05$; ANOVA for
14 repeated measures ($p<0.05$) with post hoc Bonferroni-corrected paired tests).
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 **Fig. 3.** General activity in home cage before and after 6-OHDA lesions, measured in two animals
31 during 72h-recordings using accelerometers. (A) Green panels show data collected during
32 baseline conditions prior to lesioning surgery, yellow and red panels represent the activity
33 displayed after unilateral and bilateral lesions, respectively. (B) Average activity recorded during
34 the active periods of the day (4am-6pm) during the three different conditions (baseline, unilateral
35 and bilateral lesion). On the x-axis, day zero corresponds to the day of the second lesion, days -
36 50 to 0 to unilateral lesion, and earlier than day -50 represents baseline recordings. (C) The
37 average activity displayed during circadian cycle for baseline and bilateral lesion conditions.
38
39
40
41
42
43
44
45
46

47 **Fig. 4.** Behavioral testing in Tower. Quantification of spontaneous horizontal and vertical
48 locomotion in the tower testing set-up reveals clearly different behavior in the four different
49 stages of Parkinsonism. (A) Example of vertical (green trace) and horizontal (brown trace)
50 displacement of the animal during a 5 min recording session (height over ground for the different
51 bars are denoted on the axis to the left, tracking data quantized to the levels are shown in the
52 thick lines and the original tracking data are shown as thinner lines). (B) Summary of the average
53 horizontal and vertical distance travelled in all recordings (median, 25% and 75% percentiles
54 shown in boxes, whiskers denote range). Note the successive decrease in distance travelled in the
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 more severe PD models [green=intact (I), yellow=hemilesion (H), red=bilateral lesion (B),
5 black= bilateral lesion + AMPT (A)]. (C) A change in the preference for bars located relatively
6 higher up to bars at lower levels with increasing severity of Parkinsonism. The relative amount
7 of time spent on the respective level is indicated by colored bars. (D) Transition matrices
8 describing the probability that the animal will move from a certain level (row) to another level
9 (column). Levels are denoted from G to 7, where G is ground and 7 is the highest bar, each
10 treatment group is normalized to the total number of transitions observed in that condition. It can
11 be noted that animals move less frequently more than one level at a time and between the higher
12 bars with more severe parkinsonian symptoms.
13
14
15
16
17
18
19
20
21
22



23
24 **Fig. 5.** Behavioral testing in Box. Quantification of spontaneous locomotive behavior in the box
25 set-up reveals marked differences between the different degrees of Parkinsonism. (A) The
26 average horizontal and vertical and distance travelled per minute for animals grouped according
27 to severity of Parkinsonism. (B) Differences in bout duration, distance, speed and frequency,
28 shown for the different groups in histograms representing the relative frequency of observed
29 parameter values in four equally sized intervals of the full range for the respective parameters.
30 (C) Bout frequency and total distance travelled shown for all recorded sessions divided
31 according to lesion group (mean and SD indicated by horizontal line and box, respectively). Note
32 the robust reductions following both the first and second lesion which persist throughout each >8
33 week long testing period. [Color code: green=intact (I), yellow=hemilesion (H), red=bilateral
34 lesion (B), black= bilateral lesion + AMPT (A)].
35
36
37
38
39
40
41
42
43
44
45
46
47

48
49 **Fig. 6.** Histological confirmation of dopaminergic lesions. (A) Examples of tyrosine
50 hydroxylase (TH) immunolabeling in the caudate (Cd), putamen (Put). Immunohistochemistry of
51 TH showed intense labeling of Cd-Put in both hemispheres in control animals (top panel).
52 Lesions performed in the left hemisphere induced a pronounced loss of labeling in Cd/Put on this
53 side (middle panel). In bilaterally lesioned animals, both sides of Cd/Put were strongly affected,
54 showing much weaker TH-staining (bottom panels) compared to controls. (B) Examples of TH
55 immunolabeling in the substantia nigra (SN). In SN, TH-labelling of cell bodies of midbrain
56
57
58
59
60
61
62
63
64
65



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

dopaminergic neurons in the intact brain is evident but is strongly reduced in lesioned hemispheres. (C) Quantitative summary of TH-immunolabeling of terminals in the caudate-putamen (top) and of cell-bodies in substantia nigra (bottom) confirming extensive dopaminergic lesions.




Figure(s)




Figure(s)




Figure(s)






Figure(s)



Figure(s)



Figure(s)



Table(s)

Animal	Lesion	Manual PD assessment	Tower test	Box test	Activity in home cage
1 - Betó	bilateral	X	X		
2 - Dedé	bilateral	X		X	
3 - Max	bilateral	X			
4 - Tom	bilateral	X	X		
5 - Kaká	bilateral		X	X	
6 - Pele	bilateral	X	X	X	X
7 - Zeca	bilateral	X	X	X	X
8 - Deco	unilateral	X	X		
9 - Kadu	unilateral	X	X		